

High resolution imaging of CO outflows in OMC-2 and OMC-3

Jonathan P. Williams

Institute for Astronomy, 2680 Woodlawn Drive, Honolulu, HI 96822

jpw@ifa.hawaii.edu

R. L. Plambeck

Astronomy Department, University of California, Berkeley, CA 94720

rplambeck@astro.berkeley.edu

Mark H. Heyer

Department of Astronomy, LGRT 619, University of Massachusetts, 710 North Pleasant Street, Amherst, MA 01003

hey@astro.umass.edu

ABSTRACT

A large scale, high resolution map of CO (1{0) emission toward the OMC-2 and OMC-3 star forming regions is presented. The map is a mosaic of 46 fields using the Berkeley-Illinois-Maryland Array (BIMA) and covers $10^\circ \times 15^\circ$ at $10''$ resolution. These data are combined with single dish FCRAO observations and analyzed to identify and determine the properties of nine protostellar outflows. The BIMA data alone almost completely resolve out the cloud emission at central velocities and only recover $1/20$ of the flux in the high velocity gas showing that outflows are generally broadly dispersed over 1° angular scales. All nine identified outflows emanate from known Class 0 or borderline Class 0/I sources, are associated with knots of shocked H_2 emission, and have short dynamical times. It is suggested that only the youngest, most spatially compact, and energetic outflows have been found and that more distributed high velocity gas undetected by BIMA is due to older outflows continuing through the Class I phase of protostellar evolution. The mechanical energy injection rate into the cloud is estimated to be $1.5 L_\odot$ which is comparable to the turbulent energy dissipation rate. Outflows appear capable, therefore, of sustaining cloud turbulence but a high star formation rate is required in implying a short cloud lifetime 5 Myr .

Subject headings: ISM : individual(OMC-2, OMC-3) | ISM : kinematics and dynamics | stars: formation

1. Introduction

The accretion of gas onto a protostar is accompanied by collimated jets observed in the optical and infrared (Mundt & Fried 1983; Schwartz et al. 1988) and swept up molecular gas outflows observed at millimeter wavelengths (Snell, Loren, & Plambeck 1980). The resulting transfer of momentum and kinetic energy into a protostar's immediate surroundings may limit further accretion and thereby determine the final stellar mass (Shu, Adams, & Lizano 1987). The combined effect of multiple outflows averaged over space and time may also regulate future star formation on larger scales by maintaining cloud turbulence (McKee 1989). This paper addresses the latter issue by analyzing the energy injection rate into a cloud by a collection of very young outflows and the spatial distribution of high velocity gas.

Molecular outflows are most powerful in the earliest, Class 0, stage of protostellar evolution (Bontemps et al. 1996). A chain of such young protostars was found by Chin et al. (1997; hereafter C97) through mapping dust continuum emission at 1300 μ m along a ridge of dense molecular gas north of the Orion nebula. Subsequent maps at 350 μ m and 850 μ m were made by Lis et al. (1998) and Johnstone & Bally (1999) respectively. The ridge is split into two main groups, OM C-2 in the southern half and OM C-3 in the north. The ratio of sub-millimeter to far-infrared luminosity is higher in OM C-3 than in OM C-2 suggesting a trend toward younger protostars with increasing distance from the Orion nebula (C97; Lis et al. 1998). However, all the sources found by C97 are deeply embedded and can be classified as either Class 0 or borderline Class 0/I. Other papers that are pertinent to this work are Zuckerman et al. (1997) imaging of the $\lambda = 1.0$ S(1) line of shock excited H_2 by Yu, Bally, & Devine (1997; hereafter Y97) and, more recently, Stanke, McCaughrean, & Zinnecker (2002; S02). Y97 found 80 knots of H_2 emission which they connected into twelve jets from young protostars. The positions of these knots are used as a guide to deciphering the molecular outflows in the maps presented here. S02 mapped a much larger region in Orion A and confirmed the positions of the Y97 knots in OM C2/3. Yu et al. (2000; hereafter Y00) followed up their H_2 imaging study with molecular line observations of several jets. C97 also presented molecular line observations. In both cases, the relatively large beam sizes ($> 24''$) of these singledish studies together with the high density of protostars and multiple, overlapping outflows prevented a firm association of molecular outflow gas with H_2 jets. As et al. (2000; hereafter A00) mapped CO (1{0) and denser gas tracers with the largest singledish telescope operating at 3 mm, the Nobeyama 45 m, to achieve maps of the outflows at $17''$. They found eight outflows and their map is the most comparable to this work.

Because of the high protostellar density in the region, it would be desirable to map the outflows at comparable resolution to the dust continuum maps that identified the young protostars. Only interferometers can achieve such a resolution, $\sim 10''$, in the low frequency

lines that the outflows generally excite. The ten 6 m antennae that make up BIMA¹ provide excellent uv coverage and a large primary beam which result in an efficient instrument for creating high resolution, large scale images of the millimeter sky. This paper presents a multi-field BIMA mosaic of the CO (1{0) emission in the OMC2/3 region and shows the network of young molecular outflows from the clustered protostars in unprecedented detail. The following sections present the observational technique used to produce the map, general results from the data and a description of each of the nine individual outflows that were identified. The paper concludes with a discussion of the implications of this work for outflow lifetimes, energetics, and cloud turbulence.

2. Observations

A 46-field mosaic was observed with BIMA four times from October 1998 to March 1999, twice each in the compact C and D configurations. Baselines ranged from 6 m to 81 m. The total on-source integration time was 14 hours, or slightly less than 20 minutes per field. Amplitude and phase were calibrated using 4 minute observations of 0530+135 interleaved with each 23 minute integration on source (30 seconds per field). Observations of the bright quasar 3C 454.3 at the start of each track showed the passband to be flat and no correction was applied. The flux density scale was set by observing Mars during the 1998 observations resulting in a flux for 0530+135 of 2.0 Jy. The CO line was placed in the upper sideband centered in a 256 channel window with 25 MHz bandwidth with a corresponding velocity resolution of 0.25 km s⁻¹. The total continuum bandwidth was 600 MHz but the integration time per field was too short to obtain more than a noisy detection of the brightest sources in the C97 map. The data were reduced in the MIRIAD software package (Wright & Sault 1993) and the final CLEANed map had a spatial resolution of 13''.4 × 8''.8 at a position angle of 1:9 and a (boxcar smoothed) velocity resolution of 0.5 km s⁻¹. The final map covers 10° × 15° at a spatial dynamic range, $A_{\text{map}}/A_{\text{beam}} = 5800$.

The 46 pointing centers were arranged in a hexagonal grid spaced by $\theta = 50''$; $\phi = 86''$. This resulted in a separation between pointing centers of 100'', about the same as the primary beam of the 6 m antennae. Such a beam sampled mosaic is sensitive to compact structures and is suited to mapping large areas with minimal observing overhead. A comparison between a beam sampled and Nyquist sampled mosaic of CO outflows in NGC 1333 showed negligible differences (Plambeck & Engargiola 2000). The noise level, 0.6 K (1 σ) per 0.5 km s⁻¹ velocity channel, was nearly uniform across the final map with a sharp falloff in

¹ Operated with support from the National Science Foundation under grants AST-9981308 to UC Berkeley, AST-9981363 to U. Illinois, and AST-9981289 to U. Maryland.

sensitivity from 80% to 30% over 30° at the map borders.

Singledish data were taken at the Five College Radio Astronomy Observatory 14 m telescope (FCRAO²) in March 1999 with the 16 element, focalplane array SEQUOIA and a system of autocorrelation spectrometers at a spectral resolution of 78 kHz. The observing was carried out using standard position-switching procedures resulting in a Nyquist sampled map with an rms noise of 0.3 K per 0.5 km s^{-1} channel. Antenna temperatures were converted to fluxes using a gain of 44.5 Jy K^{-1} and added to the BIMA data to fill in the short spacing information missing from the interferometer map through joint maximum entropy deconvolution.

The addition of the singledish data partially compensates for the spatial filtering properties of the interferometer and makes possible flux recovery of extended objects. Interferometers null out features that are uniform over angular sizes $f = b_{\text{min}} / \theta < 90^\circ$ where $b_{\text{min}} = 6 \text{ m} = 2.3 \text{ k}$ is the shortest projected baseline separation and f is a factor that theoretical considerations suggest may be $< 1=2$ (Wilner & Welch 1994). However, simulations by Helfer et al. (2002) show that the non-linear deconvolution process can effectively extrapolate structural information to larger values of f . Experience suggests that D-array observations are sensitive to angular sizes $> 60^\circ$, i.e. $f = 2=3$. Since the beam size of the FCRAO 14 m at 115 GHz is 47° , the combined BIMA+FCRAO dataset provides information on small scale and large scale structures but poorly represents features on angular scales in the range $60^\circ - 95^\circ$.

3. Results

3.1. Overview

Average spectra over the mapped region for each of the two datasets are shown in Figure 1. In contrast to the singledish map, large areas of the interferometer map have very little emission so the average intensity is much lower. Spectral profiles are also very different; the FCRAO spectrum is centrally peaked with weak extended line-wing emission, but the BIMA spectrum has a central dip and peaks in the line-wings. This is due to the spatial filtering properties of interferometers as described above. In this case, it can be concluded that the bulk of the cloud emission at $v = 9$ to 12.5 km s^{-1} is relatively smooth over angular scales $> 1^\circ$ but that the line-wing emission arises, at least in part, from more compact spatial

²FCRAO is supported in part by the National Science Foundation under grant AST-0100793 and is operated with permission of the Metropolitan District Commission, Commonwealth of Massachusetts

features (i.e. collimated outflows). Similar singledish/interferometer profiles were observed toward NGC 1333 and HH 7-11 by Plambeck & Engargiola (2000).

The BIMA data accounts for only 0.5% of the flux of the total cloud emission, as measured from the FCRAO data, demonstrating how smooth the CO emission is. Even in the line-wings, however, the BIMA data only accounts for 4.7% of the emission so a large fraction of the outflow emission must also be spatially extended over 1° angular scales. For this reason outflow properties are calculated from the combined BIMA+FCRAO dataset that contains the flux of the singledish map. The broad spatial extent of the high velocity gas has important implications on its origin: either outflows can rapidly distribute their momentum and energy over large regions or most of the line-wing emission arises from a cascade of larger scale processes such as Galactic shear (Fleck 1981).

Although the BIMA data accounts for only $1/20$ of the flux of the line-wing emission, the spatial filtering properties of the interferometer are advantageous in isolating the small scale features of collimated outflows. Low and high velocity BIMA emission is plotted in Figure 2. Several linear features can be seen and some clear outflows, MMS5 and MMS10 being the best examples, with separated blue-red lobes are apparent. Nine outflows were found in these data and they are shown schematically in the map. All appear to originate from one of the C97 protostars. The Y97 H₂ knots are also plotted in the Figure and were an essential guide in the assignment of the line-wing emission to different outflows. Most H₂ knots were found to be associated with CO line-wing emission and those that were not appear to be due to outflows in the plane of the sky. Similarly most of the CO features in Figure 2 can be assigned to an outflow from a C97 protostar but there are also many clumps of high velocity gas whose origin remains unclear.

3.2. Individual outflows

The identification of each of the nine outflows was made from analysis of channel maps and position-velocity slices in conjunction with the positions of the Y97 and S02 H₂ knots. Physical properties are listed in Table 1³ ordered by the C97 driving source where MMS stands for "millimeter source" and FIR for "far-infrared". For each outflow the mass, momentum and energy were determined by first outlining the extent of the blue and red lobes in position and velocity space using interactive cursor-based IDL routines and then employing equations 6 and 9 of Cabrit & Bertout (1990). The combined BIMA+FCRAO dataset was used for the calculations and, following Y00, CO line-wing emission was taken to be opti-

³A distance to the cloud of 450 pc is assumed (Brown, de Geus & de Zeeuw 1994).

cally thin and at an excitation temperature of 30 K. Eyeball estimates of sizes and velocity gradients were made from the channel maps and position-velocity slices shown in Figures 3{7.

MM S2/3: This east-west outflow runs close to the plane of the sky (Figure 3). There is a very weak red lobe coincident with a string of H_2 knots and a compact blue lobe host to a cluster of H_2 knots. Because of its orientation, this outflow is one of the longest identified in these maps and only a very slight velocity gradient could be measured across it. MM S2 and MM S3 are too close to unambiguously identify one over the other as the driving source of this outflow. A large clump of high velocity gas lies to the west that may be an expanding shell or loosely collimated outflow from any of MM S2/3, 4, 5 or 6 but it is not associated with H_2 emission and could not be conclusively connected to any source.

MM S5: This is a small east-west outflow with blue and red peaks straddling MM S5 along the same direction as a line of 3 H_2 knots (Figure 3). The position velocity map (Figure 7) along this line shows the second highest velocity gradient measured out of the nine flows and suggests a near pole-on orientation. Two additional H_2 knots point to a north-south jet from MM S6 (see also S02) but the CO emission is too confused in this area to definitively identify the corresponding outflow.

MM S7: Also known as IRAS 05329-0505 this source powers a bright reflection nebula and drives a well collimated, one-sided (west) optical jet HH 294 (Reipurth, Bally, & Devine 1997). Y 97 found a lone H_2 knot $\sim 2^\circ$ to the east that they suggested was the end of a long outflow from FIR 1c. The data here suggest that this knot is in fact associated with MM S7 through a CO filament that appears to be an outflow in the plane of the sky (Figure 4). The line-wing map in Figure 2 shows no high velocity gas associated with MM S7 but careful inspection of channel maps revealed a linear feature connecting the H_2 emission in HH 294 with the eastern knot. This putative outflow is just distinguishable above a broad plateau of cloud emission in a small range of velocities near the systemic motion of the cloud. Its direction is slightly different from the HH 294 optical jet but agrees with the position angle of a small 3.6 cm radio jet observed with the VLA (Reipurth et al. 2003). Because of its orientation in the plane of the sky and immersion in the general cloud emission, no velocity gradient could be measured along the outflow and only conservative estimates to its size and mass could be made.

MM S8: The identification of this outflow is the most uncertain of the nine. Channel maps and position-velocity cuts hint at an outflow around MM S8 that connects several Y 97 H_2 knots (Figure 5) but there is confusion from high velocity CO emission around MM S9 and the dense chain of H_2 knots from the well collimated jet that it powers. As originally noted by Y 00, the powerful CO outflow around MM S8 discussed in C 97 was misidentified due to

erroneous plotting of source positions and is a combination of flows originating from M M S9 and M M S10.

M M S9: A prominent chain of H_2 knots reveal a long east-west jet originating from M M S9 (Y 97, S02). The B M A data show associated, collimated, high velocity $v > 13 \text{ km s}^{-1}$ CO (Figure 5). On the eastern side of the source, the association of both H_2 knots and gas is less clear but a long, high velocity collimated flow is discernable and best seen in Figure 2. The flow occurs over a narrow range of velocities and suggests an orientation in the plane of the sky, consistent with its projected length, almost 1 pc, by far the longest of the nine flows in Table 1.

Two compact cores of low velocity gas lie to the west of M M S9 and were connected to the high velocity core north of M M S10 by both Y 00 and A 00. However, the greater resolution of these data clearly demonstrate that the H_2 jet is associated with the linear high velocity core $30''$ further south. A more diffuse red lobe is also found toward the blue cores which suggests that this feature may be a pole on flow from an unidentified source or an expanding shell from an older outflow.

M M S10: This is a strong, clearly defined outflow running northeast-southwest (Figure 5). Y 00 suggest that the red lobe here is due to the outflow from M M S9. While there is likely to be some contribution from that flow, the higher resolution of these data show a clear relationship in space and velocity (Figure 7) between M M S10 and the strong blue and red lobes on either side of it. The H_2 knots appear to be associated with the edges of the CO lobes rather than directly lining up with a single jet from the source itself. This is one of the more massive flows identified in the cloud.

F IR 1b/c: This outflow occurs in a highly confused area of the maps where the source density is high and there are multiple, overlapping flows. Because of the large numbers of high velocity CO clumps and scattered H_2 knots here the identification of this outflow is uncertain. The evidence for an outflow is based on the blue and neighboring red lobes on either side of F IR 1c and F IR 1b and H_2 knots in close, but not exact, proximity (Figures 5, 6). A position velocity cut along a line through the two sources is unable to distinguish one over the other as the driving source (Figure 7).

F IR 2: This is a very weak but high velocity flow running north-south. The two lobes almost overlap spatially (Figure 6) but extend over a wide range in velocity (Figure 7) resulting in an extremely high velocity gradient and suggesting a pole-on orientation. H_2 knots are found in the vicinity of F IR 2 but their association with this flow is uncertain and they may well be an extension of the powerful M M S10 outflow.

F IR 3: A compact chain of H_2 knots and intense, collimated, high velocity CO emission

reveal strong outflow activity. This source is a binary with $4''$ separation (Pendleton et al. 1986) and appears to drive two criss-crossed flows. Channel maps show two blue lobes and a single, long, red lobe (Figure 6). The position-velocity map in Figure 7 shows the two blue lobes more clearly. Their red lobe counterparts can be seen in the two lowest contour levels of the merged high velocity emission in the same Figure. The sum of these two flows inject the most mass, momentum, and energy into the cloud of the nine flows discussed in this work.

In addition, a potential tenth flow is seen in the BIMA map (dashed line in Figure 2). This connects a single H_2 knot in the southwest with a chain of high velocity clumps that appear to connect back to FIR 3. It was, however, not possible to distinguish this flow clearly in the combined BIMA + FCRAO dataset due to its length, lack of strong velocity gradient, and the confusion of background cloud emission.

4. Discussion

The OMC-2 and OMC-3 regions are highly active star forming centers with multiple, overlapping outflows. High resolution, large scale CO mapping allows more secure identification of the flows and their driving sources and makes possible the determination of fundamental physical properties such as mass, momentum, and energy. In practice, low and high velocity clumps are found to be widely distributed across the map and, particularly for the flows in the plane of the sky, it was essential to use the Y97 and S02 H_2 maps to assign these features to outflows driven by a given source.

The nine identified CO outflows line up with 50 of the 60 Y97 H_2 knots in the mapped region. There were no clearly identifiable bipolar CO outflows that were not associated with detectable molecular hydrogen shocks. This is partly, but not completely, due to the way in which the high velocity gas is assigned to outflows. The additional information provided by the CO data, however, shows that the extremes of some of the jets found by Y97 appear to be misidentified. For example, in Y97 the H_2 knots 74/77, 76 and 79 define the ends of jets I, E and J respectively but the CO data show that these knots are more likely to be associated with flows from closer submillimeter sources, MMS10, MMS7 and either MMS9 or 10 respectively. These jets are likely, therefore, to be significantly smaller than previously thought. On the other hand, the MMS2/3 and MMS9 flows that lie in the plane of the sky appear to have a measurable effect on the CO emission over a sizeable fraction of a parsec in length as has been well established for optical jets (Reipurth & Bally 2001).

The large scale effect of outflows may also be apparent in the "missing flux" in the

BIMA map due to structures that are larger than $1''$. Over 95% of the line-wing emission is resolved out by the interferometer demonstrating that the high velocity gas is broadly distributed. Similarly, a map of the CO line-width at $10''$ resolution made from the combined dataset is generally featureless across the cloud away from a few peaks toward individual sources such as FIR 3 and some H_2 knots along the MMS9 flow.

The fact that the line-wing emission is widely distributed complicates the definition of the boundaries of the outflows and the measurement of their physical properties. Additional uncertainties are the optical depth of the CO line-wing emission and the inclination of the flow to our line of sight. Since there are no ^{13}CO observations at the same resolution, optically thin emission was assumed, providing a lower limit to the mass and kinematic estimates. No inclination correction was applied to the individual flow momenta and energies listed in Table 1 since it is difficult to determine accurately. Assuming, however, a mean inclination of 45° for the outflow sample, total values of the mass, momenta and energy are $1.4 M_\odot$, $5.4 M_\odot \text{ km s}^{-1}$ and 2.9×10^{44} erg respectively. The variation in masses of individual outflows is about a factor of five but the velocities do not greatly differ and the range of momenta and energies is only a factor of six. There is no clear correlation between outflow properties and the C97 driving source mass or luminosity.

Velocity gradients, $dV = dR / (V_{\text{red}} - V_{\text{blue}}) = L$, measured by eyeballing the position-velocity diagrams in Figure 7, are given in Table 1. Velocity gradients for the two long outflows from MMS2/3 and MMS9 that lie nearly in the plane of the sky could not be accurately measured. The inverse of the gradient in $\text{km s}^{-1} \text{ pc}^{-1}$ is approximately the dynamical time in Myr. The median dynamical time in this sample is 2×10^4 yr which is similar to the statistical lifetime of Class 0 protostars in the Ophi cluster (Andre & Montmerle 1994). However, the identification of the outflows here suffers from a bias toward either high velocity gradients or sharp, collimated features that stand out against the general cloud emission. The former corresponds to short dynamical timescales and the latter is a probable indication of youth (Mason & Chemin 1993). It is not necessarily surprising, then, that all nine flows described in this paper appear to be driven by a young Class 0, or borderline Class 0/I, protostar in the C97 catalog.

The dynamical time is only a lower limit to the age of an outflow (Parker, Padman, & Scott 1991) but is probably a good estimator for the young outflows observed here (Mason & Chemin 1993). If only the very youngest outflows, $t_{\text{out}} \approx 2 \times 10^4$ yr, have been identified, but (less well collimated) outflow activity continues throughout the Class I phase, $t_{\text{I}} \approx 10^5$ yr (Wilking et al. 1989), then the broadly distributed line-wing emission that was resolved out in the BIMA map can be accounted for by outflows alone. In principle, they should be apparent in a combined single-dish + interferometer map such as the one presented in this paper, but higher signal-to-noise data are required to pick them out from the general

cloud background. This paper has focused on the high velocity gas that can be identified with shocked H_2 emission but there are several other high velocity clumps in Figure 2 that were not associated with outflows. These unassigned clumps may be parts of older outflow shells which have broadened and slowed sufficiently such that the shocks with the ambient medium are too weak to excite the H_2 2:12 μ m line. Earlier single dish maps of CO in O₃ Orion indeed found several large scale, slowly expanding shells that were interpreted as resulting from young stellar outflows (Heyer et al. 1992).

It has long been suggested that outflows can maintain cloud turbulence (e.g., Norman & Silk 1980). The agreement between the ratio of Class 0 to Class I lifetimes, and the flux ratio between collimated and broad outflows, suggests an approximately uniform energy input into the cloud that can be estimated from the observed CO outflows here,

$$L_{\text{ow}} = \frac{E_{\text{ow}}}{t_{\text{ow}}} = \frac{2.9 \cdot 10^{44} f_{\text{CO}} \text{ erg}}{2 \cdot 10^4 \text{ yr}}, \quad 5 \cdot 10^{32} f_{\text{CO}} \text{ erg s}^{-1};$$

where f_{CO} is a factor that corrects for the assumption of optically thin CO line-wing emission in the calculation of the outflow masses and energies. Cabrit & Bertout (1990) show that f_{CO} lies in the range 5–10. The cloud loses energy through turbulent decay. Simulations of magnetohydrodynamic cloud turbulence show rapid dissipation with a timescale,

$$t_{\text{diss}} = \frac{3.9}{M_{\text{ms}}} t; \quad ;$$

where $\lambda_d = \lambda_J$ is the driving wavelength of the turbulence in units of the Jean's length, M_{ms} is the rms Mach number of the turbulence, and $t = (3 \cdot 32 G)^{1/2}$ is the free-fall timescale (MacLow 1999). The mass of the OMC-2/3 complex is $1100 M_\odot$ (Lis et al. 1998). The average temperature and 1-d velocity dispersion can be measured from the FCRAO spectrum in Figure 1 and are equal to 20 K and 1.5 km s^{-1} respectively. The equivalent radius, measured from the projected area, is 0.9 pc, and implies an average number density, $n_{H_2} = 5 \cdot 10^3 \text{ cm}^{-3}$. These numbers imply a total kinetic (turbulent) energy $E_{\text{turb}} = 7.4 \cdot 10^{46} \text{ erg}$, Mach number $M_{\text{rms}} = 5$, and free-fall time $t = 5.2 \cdot 10^5 \text{ yr}$. Parameterizing by λ_d , the dissipation time is $t_{\text{diss}} = 4.1 \cdot 10^6 \text{ yr}$, and the turbulent energy dissipation rate is,

$$L_{\text{turb}} = \frac{E_{\text{turb}}}{t_{\text{diss}}} = 6 \cdot 10^{33} \text{ erg s}^{-1};$$

The average cloud density and temperature imply a Jean's length, $\lambda_J = 0.2 \text{ pc}$. If outflows sustain the turbulence, the driving wavelength can be no greater than their size $\lambda_d \leq \lambda_J$. The average outflow length in Table 1, corrected for an average inclination angle of 45° , is $\lambda_J = 0.5 \text{ pc}$. Thus $\lambda_d = \lambda_J \approx 2.5$. Shorter wavelength "harmonics" may also be present, however, and MacLow (1999) argues that the driving wavelength must be less

than the Jean's length, i.e. $\beta < 1$, for turbulence to support the cloud. Taking $\beta \approx 1$ and the maximum CO optical depth correction $f_{\text{CO}} = 10$, the data are consistent with an approximate balance between outflow energy injection rate and turbulent energy dissipation rate,

$$L_{\text{out}} \approx L_{\text{turb}} \approx 6 \times 10^{33} \text{ erg s}^{-1} = 1.5 L_{\odot}.$$

To maintain the cloud turbulence over longer timescales than an individual protostar's outflow lifetime requires a star formation rate $\approx 9 \text{ stars} = 2 \times 10^4 \text{ yr}^{-1} = 2.3 \times 10^{-4} M_{\odot} \text{ yr}^{-1}$ for an average protostellar mass of $0.5 M_{\odot}$ based on the Scalo (1986) IMF. At this rate the OMC-2/3 complex has enough mass to continue forming low mass stars for $5 \times 10^6 \text{ yr}$ where $\epsilon = M_{\text{out}}/M_{\text{cloud}} < 1$ is the overall star formation efficiency. Even if the cloud ultimately converts most of its mass to stars, therefore, the high rate of turbulent energy dissipation restricts its lifetime to a few Myr. This is shorter than predicted for more massive GMCs (Blitz & Shu 1980; Williams & McKee 1997) but consistent with the rapid cloud formation and evolution scenario proposed by Hartmann, Ballesteros-Paredes & Bergin (2001) and the observed small age spread of protostars in nearby star forming regions (Hartmann 2001).

JPW is supported by NSF grant AST-0134739 and thanks Bo Reipurth and Joan Najita for helpful discussions.

5. References

- Andre, P., & Montmerle, T. 1994, *ApJ*, 420, 837
 Aso, Y., Tatematsu, K., Sekimoto, Y., Nakano, T., Umemoto, T., Koyama, K., & Yamamoto, S. 2000, *ApJS*, 131, 465 (A00)
 Blitz, L., & Shu, F.H. 1980, *ApJ*, 238, 148
 Bontemps, S., Andre, P., Terebey, S., & Cabrit, S. 1996, *A & A*, 311, 858
 Brown, A.G.A., de Geus, E.J., & de Zeeuw, P.T. 1994, *A & A*, 289, 101
 Cabrit, S., & Bertout, C. 1990, *ApJ*, 348, 530
 Chini, R., Reipurth, B., Ward-Thompson, D., Bally, J., Nymann, L.-A., Sievers, A., & Billawala, Y. 1997, *ApJ*, 474, L135 (C97)
 Fleck, R.C. 1981, *ApJ*, 246, L151
 Hartmann, L. 2001, *AJ*, 121, 1030
 Hartmann, L., Ballesteros-Paredes, J., & Bergin, E.A. 2001, *ApJ*, 562, 852
 Helfer, T.T., Vogel, S.N., Lugten, J.B., & Teuben, P.J. 2002, *PASP*, 114, 350
 Heyer, M., Morgan, J., Schloerb, F.P., Snell, R.L., Goldsmith, P.F. 1992, *ApJ*, 395, 99
 Johnstone D., & Bally, J. 1999, *ApJ*, 510, L49

- Lis, D. C., Serabyn, E., Keene, J., Dowell, C. D., Benford, D. J., Phillips, T. G., Hunter, T. R., & Wang, N. 1998, *ApJ*, 509, 299
- MacLow, M. M. 1999, *ApJ*, 524, 169
- Masson, C. R., & Chemin, L. M. 1993, *ApJ*, 414, 230
- McKee, C. F. 1989, *ApJ*, 345, 782
- Mundt, R., & Fried, J. W. 1983, *ApJ*, 274, L83
- Norman, C., & Silk, J. 1980, *ApJ*, 238, 158
- Parker, N. D., Padman, R., & Scott, P. F. 1991, *MNRAS*, 252, 442
- Pendleton, Y., Wemer, M. W., Capps, R., & Lester, D. 1986, *ApJ*, 311, 360
- Plambeck, R. L., & Engargiola, G. 2000, in *Imaging at Radio through Submillimeter Wavelengths*, eds. J. G. Mangum & S. J. E. Radford, *ASP Conf. Series*, 217, 354
- Reipurth, B., & Bally, J. 2001, *Ann. Rev. Astron. Astr.*, 39, 403
- Reipurth, B., Bally, J., & Devine, D. 1997, *AJ*, 114, 2708
- Reipurth, B., Rodriguez, L. F., Anglada, G., & Bally, J. 2003, in prep.
- Scalo, J. 1986, *Fund. Cosmic Phys.*, 11, 1
- Schwartz, R. D., Jennings, D. G., Williams, P. M., & Cohen, M. 1988, *ApJ*, 334, L99
- Shu, F. H., Adams, F. C., & Lizano, S. 1987, *A&A*, 25, 23
- Snell, R. L., Loren, R. B., & Plambeck, R. L. 1980, *ApJ*, 239, L17
- Stanke, T., McCaughrean, M. J., & Zinnecker, H. 2002, *A&A*, 392, 239 (S02)
- Wilking, B. A., Lada, C. J., & Young, E. T. 1989, *ApJ*, 340, 823
- Williams, J. P., & McKee, C. F. 1997, *ApJ*, 476, 166
- Wilner, D. J., & Welch, W. J. 1994, *ApJ*, 427, 898
- Wright, M. C. H., & Sault, R. J. 1993, *ApJ*, 402, 546
- Yu, K., Bally, J., & Devine, D. 1997, *ApJ*, 485, L45 (Y97)
- Yu, K., Billawala, Y., Smith, M. D., Bally, J., & Butner, H. M. 2000, *AJ*, 120, 1974 (Y00)




Figure 1: Average spectra over the entire mapped region for the FCRAO and BIMA data. The BIMA spectrum is scaled by a factor of 10 and plotted in the heavier linetype. The lower panel zooms in on the lower intensity emission and demonstrates that the interferometer data preferentially selects the high and low velocity gas from the small scale features in outflows.




Figure 2: Line-wing emission from the BIMA mosaic map (interferometer data only). Blue and red contours show the emission integrated from $v = 5.5$ to 9.0 km s^{-1} and 12.5 to 16.0 km s^{-1} respectively. Contour starting levels and increments are the same ($8, 4 \text{ K km s}^{-1}$) for both velocity ranges. The labeled star symbols show the location of the C97 protostars and dots show the position of the Y972:12 H_2 knots. Heavy dark lines show the outflows that are discussed in this paper. The solid contour outlining the map is the 30% sensitivity level of the mosaic. The $8.3'' \times 13.4''$ beam is plotted in the lower right corner.




Figure 3: Line-wing emission from the combined BIMA+FCRAO mosaicked map toward the northern OMC-3 region containing sources MMS1-6. Coordinates are offset from $\alpha = 5^{\text{h}}35^{\text{m}}26^{\text{s}}$ and $\delta = 50^{\circ}47'7''$ (J2000). The dashed contours (colored blue in electronic edition) show the low velocity gas integrated from $v = 5.0$ to 7.5 km s^{-1} and solid contours (colored red in electronic edition) the high velocity gas integrated from $v = 13.5$ to 16.5 km s^{-1} . Contour starting levels and increments are the same ($8, 4 \text{ K km s}^{-1}$) for both velocity ranges. As in Figure 2, the star symbols show the location of the C 97 protostars and dots show the position of the Y 97 2:12 m H₂ knots. The labeled heavy solid lines show the outflows that are discussed in this paper and along which the position-velocity maps are defined in Figure 7.




Figure 4: Contours of BIMA+FCRAO integrated emission from $v = 11.5$ to 13.5 km s^{-1} in the region around MMS7. Because the velocity of this flow is similar to that of the bulk of the cloud emission the total emission is very strong and contouring begins at 33 K km s^{-1} . The contour step is 3 K km s^{-1} . The solid dark line schematically marks the position of this flow but no velocity gradient across it could be reliably measured.




Figure 5: As Figure 3 for the southern part of OMC-3 containing sources MMS8-10. Dashed contours (colored blue in electronic edition) are for $v = 4.0$ to 8.0 km s^{-1} and solid contours (colored red in electronic edition) are for $v = 13.0$ to 16.0 km s^{-1} . Contours begin at 12 K km s^{-1} and increment at 4 K km s^{-1} .




Figure 6: As Figure 3 for the northern part of OMC-2 containing sources FIR 1-6. Dashed contours (colored blue in electronic edition) are for $v = 4.0$ to 7.5 km s^{-1} and solid contours (colored red in electronic edition) are for $v = 13.5$ to 17.0 km s^{-1} . Contours begin at 12 K km s^{-1} and increment at 4 K km s^{-1} .




Figure 7: Position-velocity maps along the cuts defined in Figures 3, 5, 6. The position increases from the leftmost (east) portion of the cut in each case. The long outflows from MMS2/3 and MMS9 that lie predominantly along the plane of the sky have low velocity gradients and are not shown. The contour starting level and increment are 2 K for all maps. The location of the Chini et al. protostars along the cut (at an assumed $v = 10.5 \text{ km s}^{-1}$) are shown by the star symbols and labeled in the two cases where more than one source lies along the cut.

TABLE 1
Outflow properties

	M ($M_{\odot}$)	P ($M_{\odot} \text{ km s}^{-1}$)	E (10^{43} erg)	L (pc)	$dV=dR$ ($\text{km s}^{-1} \text{ pc}^{-1}$)
MM S2/3					
blue lobe	0.13	0.38	1.26		
red lobe	0.08	0.24	0.78		
total	0.21	0.62	2.04	0.54	21
MM S5					
blue lobe	0.03	0.10	0.40		
red lobe	0.02	0.09	0.34		
total	0.05	0.19	0.74	0.13	100
MM S7					
plane of sky	0.04			0.38	
MM S8					
blue lobe	0.02	0.07	0.23		
red lobe	0.04	0.13	0.44		
total	0.06	0.20	0.67	0.20	55
MM S9					
plane of sky	0.22			0.87	
MM S10					
blue lobe	0.16	0.55	2.08		
red lobe	0.14	0.49	2.04		
total	0.30	1.04	4.12	0.39	34
FIR 1b/c					
blue lobe	0.11	0.28	0.79		
red lobe	0.05	0.16	0.52		
total	0.16	0.44	1.31	0.16	66
FIR 2					
blue lobe	0.02	0.08	0.29		
red lobe	0.03	0.10	0.39		
total	0.05	0.18	0.68	0.08	190
FIR 3					
blue lobe	0.12	0.42	1.67		
red lobe	0.21	0.74	3.32		
total	0.33	1.16	4.99	0.27	66

Appears to be the sum of two overlapping flows.