

Hybrid stars with color superconductivity within a nonlocal chiral quark model

H. Grigorian

Fachbereich Physik, Universität Rostock, D-18051 Rostock, Germany
Department of Physics, Yerevan State University, 375025 Yerevan, ArmeniaD. Blaschke^yFachbereich Physik, Universität Rostock, D-18051 Rostock, Germany
Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, RussiaD. N. Aguilera^zFachbereich Physik, Universität Rostock, D-18051 Rostock, Germany
Instituto de Física Rosario, Bv. 27 de febrero 210 bis, 2000 Rosario, Argentina

The equation of state for quark matter is derived for a nonlocal, chiral quark model within the mean field approximation. Special emphasis is on the occurrence of a diquark condensate which signals a phase transition to color superconductivity and its effects on the equation of state. We present a formula for the Bag pressure, which is density dependent in case when the quark matter is color superconducting. We calculate the quark star configurations by solving the Tolman-Oppenheimer-Volkoff equations and demonstrate the effects of diquark condensation on the stability of hybrid stars for different form factors of the quark interaction.

PACS numbers: 04.40.Dg, 12.38.Mh, 26.60.+c, 97.60.Jd

Keywords: Relativistic stars: structure, stability, and oscillations; Quark-gluon plasma; Nuclear matter aspects of neutron stars; Neutron stars

I. INTRODUCTION

The investigation of color superconductivity in quark matter [1, 2] which was revived by applying nonperturbative QCD motivated interactions [3, 4] finds most of its justification in the possible importance for the physics of compact star interiors [5] and related observable phenomena like neutron star cooling [6, 7, 8], gamma-ray bursts [9, 10, 11, 12], gravitational wave signals for compact star mergers [13] and others [14]. Since calculations of quark pairing predict values of the energy gap

100 MeV and corresponding critical temperatures for the phase transition to the superconducting state are expected to follow the BCS relation $T_c = 0.57$ for spherical wave pairing, quark matter in compact stars should always be in one of the superconducting phases.

The question arises whether conditions in compact stars allow the occurrence of quark matter and the formation of stable configurations of hybrid stars. In order to give an answer to this question one has to rely on models for the equation of state which necessarily introduce free parameters and therefore some arbitrariness in the results [15]. In particular, the question whether color superconductivity shall be realized in the 2-Flavor Superconductivity (2SC) or Color Flavor Locking (CFL) phase in the compact star interior has been discussed controversially [16, 17]. We will restrict our further discussion to dynamical models of the NJL type with their parameters adjusted by fitting hadron properties in the vacuum before extrapolating to finite temperatures and densities within the Matsubara formalism.

Within those models it has been shown that strange quark matter occurs only at densities well above the deconfinement transition, for chemical potentials which are barely reached in the very center of a compact star [18, 19, 20, 21, 22]. The interesting and much investigated CFL phase could thus play only a marginal role for the physics of compact stars. The stability of the so obtained hybrid star configurations, however, appears to depend sensitively on details of the model, including the hadronic phase.

In the present paper we are investigating this dependence in a systematic way by employing a nonlocal chiral quark model which allows to vary the form factor of the interaction kernel while describing the same set of hadronic vacuum properties. We provide a polynomial formula of our quark matter equation of state (EoS) which proves useful for applications to compact star phenomenology, as e.g. the cooling [23] and rotational evolution [24] or the merging [13] of neutron stars. In order to compare the results of the present work for hybrid star configurations with observational constraints, we pick the example of the compact object RX J185635-3754, for which limits for both the mass and the radius have been reported [25, 26]. A further restriction in the mass-radius plane of possible stable configurations comes from the constraint given by the surface redshift measurement of EXO 0748-676 [27].

II. EQUATION OF STATE OF HYBRID STAR MATTER IN EQUILIBRIUM

A. Quark matter with color superconductivity

We consider the grand canonical thermodynamic potential for 2SC quark matter within a nonlocal chiral

^Electronic address: hovik@darss.pg.uni-rostock.de^yElectronic address: david.blaschke@physik.uni-rostock.de^zElectronic address: deborah@darss.pg.uni-rostock.de

quark model [12] where in the mean field approximation the mass gap $\bar{m}_f$ and the diquark gap $\bar{m}_{\bar{f}\bar{g}}$ appear as order parameters and a decomposition into color (c = r, b, g) and flavor (f = u, d, s) degrees of freedom can be made.

$$q(f; \bar{f}; g; \bar{g}; T) = \sum_{c, \bar{c}} X_{c, \bar{c}}^{f, \bar{f}} (f; \bar{f}; g; \bar{g}; T); \quad (1)$$

where T is the temperature and μ_{fc} the chemical potential for the quark with flavor f and color c .

The contribution of quarks with given color and flavor f to the thermodynamic potential is

$$\begin{aligned} & q_{f, \bar{f}}(f; \bar{f}; g; \bar{g}; T) + \frac{c}{v_{\text{vac}}} = \frac{2}{24 G_1} + \frac{2}{24 G_2} \\ & \frac{1}{2} \sum_{\mathbf{p}} \int_0^1 dq q^2 f! [c(E_f(q) + \mu_{fc}); T] + \\ & ! [c(E_{\bar{f}}(q) - \mu_{\bar{f}c}); T] g; \end{aligned} \quad (2)$$

where G_1 and G_2 are coupling constants in the scalar meson and diquark channels, respectively. The dispersion relation for unpaired quarks with dynamical mass function $m_f(q) = m_f + g(q)$ is given by

$$E_f(q) = \sqrt{q^2 + m_f^2(q)}; \quad (3)$$

In Eq. (2) we have introduced the notation

$$! [c; T] = T \ln [1 + \exp \frac{c}{T}] + \frac{c}{2}; \quad (4)$$

where the first argument is given by

$$c(\mathbf{p}) = \frac{P}{1 + \frac{c^2}{p^2}}; \quad (5)$$

When we choose the green and blue colors to be paired and the red ones to remain unpaired, we have

$$c = g(q) (c_b + c_g); \quad (6)$$

For a homogeneous system in equilibrium, the minimum of the thermodynamic potential q with respect to the order parameters $f, \bar{f}, g$ and $\bar{g}$ corresponds to a negative pressure; therefore the constant $\frac{c}{v_{\text{vac}}} = \frac{c}{v_{\text{vac}}}$ is chosen such that the pressure of the physical vacuum vanishes.

The nonlocality of the interaction between the quarks in both channels qq and $q\bar{q}$ is implemented via the same form factor functions $g(q)$ in the momentum space. In our calculations we use the Gaussian (G), Lorentzian (L) and cutoff (NJL) type form factors defined as

$$g_G(q) = \exp(-q^2 \frac{2}{G}); \quad (7)$$

$$g_L(q) = [1 + (q^2 \frac{2}{L})^2]^{-1}; \quad (8)$$

$$g_{\text{NJL}}(q) = (1 - q^2 \frac{2}{\text{NJL}}); \quad (9)$$

The parameter sets (quark mass m , coupling constant G_1 , interaction range λ) for the above form factor models (see Tab. I) are fixed by the pion mass $m_\pi = 140$ MeV, pion decay constant $f_\pi = 93$ MeV and the constituent quark mass $m_0 = 330$ MeV at $T = 0$ [32]. The

diquark coupling constant G_2 is a free parameter of the approach which we vary as $G_2 = G_1$.

Following Ref. [28] we introduce the quark chemical potential for the color c , μ_{qc} and the chemical potential of the isospin asymmetry, μ_I , defined as

$$\mu_{qc} = (\mu_{uc} + \mu_{dc})/2 \quad (10)$$

$$\mu_I = (\mu_{uc} - \mu_{dc})/2; \quad (11)$$

where the latter is color independent.

The diquark condensation in the 2SC phase induces a color asymmetry which is proportional to the chemical potential μ_I . Therefore we can write

$$\mu_{qc} = \mu_q + \frac{8}{3} (c_b + c_g - 2c_r); \quad (12)$$

where μ_q and μ_I are conjugate to the quark number density and the color charge density, respectively.

As has been shown in [29] for the 2SC phase the relation $\mu_u = \mu_d = \mu$ holds so that the quark thermodynamic potential is [30]

$$\begin{aligned} & q(\mathbf{p}; \bar{\mathbf{p}}; q; \bar{q}; T) + \frac{c}{v_{\text{vac}}} = \frac{2}{4G_1} + \frac{2}{4G_2} \\ & \frac{1}{2} \sum_{\mathbf{p}} \int_0^1 dq q^2 f! [r(q + \frac{2}{3}\mu_I); T] + \\ & ! [r(q - \frac{2}{3}\mu_I); T] + ! [r(q + \frac{2}{3}\mu_I + \mu_I); T] + \\ & ! [r(q - \frac{2}{3}\mu_I + \mu_I); T] g \\ & \frac{2}{2} \sum_{\mathbf{p}} \int_0^1 dq q^2 f! [b(E(q) - q, \frac{1}{3}\mu_I); T] + \\ & ! [b(E(q) + q + \frac{1}{3}\mu_I); T] + \\ & ! [b(E(q) - q - \frac{1}{3}\mu_I) + \mu_I; T] + \\ & ! [b(E(q) + q + \frac{1}{3}\mu_I) + \mu_I; T] g; \end{aligned} \quad (13)$$

where the factor 2 in the last integral comes from the degeneracy of the blue and green colors ($b = g$).

The total thermodynamic potential contains besides the quark contribution q also that of the leptons l

$$\begin{aligned} & l(\mathbf{p}; \bar{\mathbf{p}}; q; \bar{q}; T) = q_X(\mathbf{p}; \bar{\mathbf{p}}; q; \bar{q}; T) \\ & + \sum_{\text{lepton}} l(\mathbf{p}; T); \end{aligned} \quad (14)$$

where latter are assumed to be massless, ideal Fermi gas

$$l(\mathbf{p}; T) = \frac{1}{12\pi^2} \mu^4 - \frac{1}{6} \mu^2 T^2 - \frac{7}{180} \mu^4 T^4; \quad (15)$$

At the present stage, we do include only contributions of the first family of leptons in the thermodynamic potential.

In the same way, the number densities for the different particle species j and the energy density are given by

$$n_j = n_{j>0} + (1 - \delta_{j>0}) n_{j=0} \quad (27)$$

$$\epsilon = \epsilon_{>0} + (1 - \delta_{>0}) \epsilon_{=0} \quad (28)$$




FIG. 2: Volume fraction of the phase with nonvanishing diquark condensate obtained by a Gellendinning construction of a charge-neutral mixed phase. Results are shown for three different form factors introduced in the text.

The formulas (25)–(28) define a complete set of thermodynamic relations and can be evaluated numerically. In the next section we present the results in a form analogous to the Bag model which has been widely used in the phenomenology of quark matter.

III. QUARK STAR EOS AND FIT FORMULAS

We calculate the quark matter EoS within this non-local chiral model [12] and display the results for the pressure in a form reminiscent of a bag model

$$P^{(s)} = P_{id}(\mu_B) - B^{(s)}(\mu_B); \quad (29)$$

where $P_{id}(\mu_B)$ is the ideal gas pressure of quarks and $B^{(s)}(\mu_B)$ a density dependent bag pressure, see Fig. 3. The occurrence of diquark condensation depends on the value of $\mu = G_2/G_1$ and the superscript $s = 2$ fS; N g indicates whether we consider the matter in the superconducting mixed phase ($\mu = 1$) or in the normal phase ($\mu = 0$), respectively.

According to heuristic expectations, the effect of this diquark condensation (formation of quark Cooper pairs) on the EoS is similar to the occurrence of bound states and corresponds to a negative pressure contribution (Fig. 3).

For phenomenological applications of the quark matter EoS (29) we provide a polynomial fit of the bag pressure

$$B^{(s)}(\mu_B) = \begin{cases} \sum_{k=0}^{10} a_k^{(s)} (\mu_B - \mu_c^{(s)})^k; & \mu_B > \mu_c^{(s)} \\ P_{id}(\mu_B); & \mu_B < \mu_c^{(s)} \end{cases} \quad (30)$$




FIG. 3: Bag pressure for different form factors of the quark interaction in dependence on the baryon chemical potential for $\mu = 0$ and for $\mu = 1$. For the latter the superconducting phase is realized.

The coefficients $a_k^{(s)}$ as well as the critical chemical potential $\mu_c^{(s)}$ of the chiral phase transition depend on the choice of the form factor (see Tabs. II, III).

The dependence of the diquark gap on the chemical potential can be represented in a similar way by the polynomial

$$(\mu_B) = \begin{cases} \sum_{k=0}^6 b_k (\mu_B - \mu_c^{(s)})^k; & \mu_B > \mu_c^{(s)} \\ 0; & \mu_B < \mu_c^{(s)} \end{cases} \quad (31)$$

where the coefficients b_k for the different form factors are given in Tab. IV.

The volume fraction also can be approximated by polynomials in the following form

$$(\mu_B) = \begin{cases} \sum_{k=0}^8 c_k (\mu_B - \mu_c^{(s)})^k; & \mu_B > \mu_c^{(s)} \\ \sum_{k=0}^1 c_k (\mu_B - \mu_c^{(s)})^k; & \mu_c^{(s)} > \mu_B > \mu_c^{(s)} \\ 0; & \mu_B < \mu_c^{(s)} \end{cases} \quad (32)$$

The coefficients c_k and the chemical potentials $\mu_c^{(s)}$ are given in the Table V for different form factors.

IV. HADRONIC EQUATION OF STATE AND PHASE TRANSITION

At low densities, quarks will be confined in hadrons and an appropriate EoS for dense hadron matter has to be chosen. For our discussion of quark-hadron hybrid star configurations in the next Section, we use the relativistic mean field (RMF) model of asymmetric nuclear matter including a non-linear scalar field potential and the meson (nonlinear Walecka model), see [34]. The quark-hadron phase transition is obtained using the Maxwell construction, see Refs. [35, 36] for a discussion. The resulting EoS is shown in Fig. 4 for the case $\mu = 1$ (left panel) when the quark matter phase is superconducting and for $\mu = 0$ (right panel) when it is normal.




FIG. 4: EoS for strongly interacting matter at zero temperature under compact stars constraints for the coupling parameter $\alpha = 1$ (left panel) and $\alpha = 0$ (right panel). Dashed line: relativistic mean-field model for hadronic matter; solid, dash-dotted and dotted lines correspond to quark matter with Gaussian, Lorentzian and NJL form factor functions, respectively.




FIG. 5: Baryon number density in units of the nuclear saturation density as a function of baryon chemical potential μ_B . Left panel: $\alpha = 1$. Right panel: $\alpha = 0$. Line styles correspond to the previous figure.

When comparing the three quark model form factors under consideration, the hardest quark matter EoS is obtained for the Gaussian, and therefore the critical pressure and corresponding critical energy densities of the deconfinement transition are the smallest, see Table II. The same statement holds for the case $\alpha = 0$, when the quark matter phases are normal, see right panel of Fig. 4.

According to the Maxwell construction of the deconfinement phase transition, there is a jump in the energy density, as is shown in Fig. 4.

The corresponding jumps in the baryon densities at the critical chemical potentials $\mu_B^{(H)}$ are given in Table VI, see also Fig. 5 for the behavior of $n_B(\mu_B)$ for all three

form factors and both cases of the diquark coupling, $\alpha = 1$ (left panel) and $\alpha = 0$ (right panel).

The EoS of hybrid stellar matter for temperature $T = 0$ is relevant also for calculations of compact star cooling, since the star structure is insensitive to the temperature evolution for $T < 1$ MeV.

V. CONFIGURATIONS OF HYBRID STARS

In this Section we consider the problem of stability of cold ($T = 0$) hybrid stars with color superconducting quark matter core. The star configurations are defined from the well known Tolman-Oppenheimer-Volkoff equations [37], written for the hydrodynamic equilibrium of a spherically distributed matter fluid in General Relativity, see also [34],

$$\frac{dP(r)}{dr} = -\frac{[\rho(r) + P(r)][m(r) + 4\pi r^3 P(r)]}{r[r - 2m(r)]}, \quad (33)$$

where the mass enclosed in a sphere with distance r from the center of configurations is defined by

$$m(r) = 4\pi \int_0^r \rho(r') r'^2 dr'. \quad (34)$$

These equations are solved for a set of central energy densities, see Figs. 6 – 9. An approximate criterion for the stability of star configurations is that masses should be rising functions of the central energy density $\rho(0)$.




FIG. 6: Mass-radius and mass-central energy density relation for compact stars configurations according to the EoS shown in the left panel of Fig. 4. Hybrid stars with Gaussian or Lorentzian quark matter models give stable branches.

Our calculations show that for the Gaussian and Lorentzian form factors one can have stable configurations with a quark core, either with (Fig. 6) or without (Fig. 7) color superconductivity whereas for our parameterization of the NJL model (cuto form factor) the configurations with quark cores are not stable. For the Gaussian form factor case the occurrence of color superconductivity in quark matter shifts the critical mass of the hybrid star




FIG. 7: Same as Fig. 6 for the EoS of the right panel of Fig. 4.

from $1.04 M_{\odot}$ to $0.62 M_{\odot}$ and the maximal value of the hybrid star mass from $1.83 M_{\odot}$ to $1.79 M_{\odot}$. For the Lorentzian form factor the branch of stable hybrid stars with 2SC superconducting quark cores lies in the mass range between $1.55 M_{\odot}$ and their maximum mass $1.72 M_{\odot}$.




FIG. 8: Radius-mass relation for Gaussian form factor including the mass dependence of the quark core radius in both cases, one with and the other without 2SC phase.

In Figs. 8 and 9 we demonstrate that these models fulfill the observational constraints from the isolated neutron star RX J185635-3754 [25, 26] and from the observation of the surface redshift for EXO 0748-676 [27].

The Lorentzian model with normal quark matter has marginally stable quark cores with radii less than 2 km, in the mass range $1.91 - 1.92 M_{\odot}$, see Fig. 9.

VI. CONCLUSION

We have investigated the influence of the diquark condensation on EoS of quark matter and obtained the crit-




FIG. 9: Same as Fig. 8 for Lorentzian form factor.

ical densities of phase transition to hadronic matter for different form factors of quark interaction.

We find that the charge neutrality condition requires that the quark matter phase consists of a mixture of 2SC condensate and normal phase. The volume fraction of the condensate phase amounts to 65% – 85% depending on the form factor function of the interaction. In the present work we did not consider muons in the quark matter phase. Their occurrence would increase the volume fraction of the superconducting phase by about a 5%, helping to stabilize the 2SC phase.

We have shown that for our set of form factors the NJL model gives no stable quark core hybrid stars. The occurrence of the superconducting 2SC phase in quark matter supports the stability of the quark matter phase.

Comparison of the quark core neutron star mass-radius relation with the mass and radius of the recently observed 'small' compact object RX J185635-3754 and with the constraints from the observation of the surface redshift for the object EXO 0748-676 shows that our model perfectly obeys those constraints.

These studies can be viewed as a preparatory step before more fundamental nonperturbative interactions can be provided, e.g. by QCD Schwinger-Dyson Equation studies [38, 39, 40].

Acknowledgements

We thank our colleagues for discussions and interest in our work, in particular during the NATO workshop in Yerevan, Armenia. Special thanks go to M. Buballa and D. Rischke for important remarks on a previous version of this work.

The research of D.N. Aguilera has been supported by DFG Graduiertenkolleg 567 "Stark korrelierte Vielteilchensysteme", by CONICET PIP 03072 (Argentina), by DAAD grant No. A/01/17862 and by the Hans Stifting of the University of Rostock. H.G. acknowledges support by DFG under grant No. 436 ARM 17/5/01 and by the Virtual Institute of the Helmholtz Association

-
- [1] B. C. Barrois, Nucl. Phys. B 129 (1977) 390.
- [2] D. Bailin and A. Love, Phys. Rept. 107 (1984) 325.
- [3] R. Rapp, T. Schafer, E. V. Shuryak and M. Velkovsky, Phys. Rev. Lett. 81 (1998) 53 [arXiv:hep-ph/9711396].
- [4] M. A. Alford, K. Rajagopal and F. W. Ilczek, Phys. Lett. B 450 (1999) 325.
- [5] D. Blaschke, N. K. Glendenning and A. Sedrakian, \Physics Of Neutron Star Interiors." Springer Lecture Notes in Physics 578 (2001).
- [6] D. Blaschke, T. Klahn and D. N. Voskresensky, Astrophys. J. 533 (2000) 406 [arXiv:astro-ph/9908334].
- [7] D. Page, M. Prakash, J. M. Lattimer and A. Steiner, Phys. Rev. Lett. 85 (2000) 2048 [arXiv:hep-ph/0005094].
- [8] D. Blaschke, H. Grigorian and D. N. Voskresensky, Astron. Astrophys. 368 (2001) 561 [arXiv:astro-ph/0009120].
- [9] D. K. Hong, S. D. H. Hsu and F. Sannino, Phys. Lett. B 516 (2001) 362 [arXiv:hep-ph/0107017].
- [10] R. Ouyed, eConf C 010815 (2002) 209.
- [11] D. N. Aguilera, D. Blaschke and H. Grigorian, arXiv:astro-ph/0212237.
- [12] D. Blaschke, S. Fredriksson, H. Grigorian and A. M. Oztas, arXiv:nucl-th/0301002.
- [13] R. Oechslin, K. Uryu, G. Poghosyan and F. K. Thielmann, arXiv:astro-ph/0401083.
- [14] K. Rajagopal, \Colour Superconductivity", Beatenberg, High-energy physics (2001) 345-384.
- [15] M. A. Alford and S. Reddy, Phys. Rev. D 67 (2003) 074024 [arXiv:nucl-th/0211046].
- [16] M. A. Alford and K. Rajagopal, JHEP 0206 (2002) 031 [arXiv:hep-ph/0204001].
- [17] A. W. Steiner, S. Reddy and M. Prakash, Phys. Rev. D 66 (2002) 094007 [arXiv:hep-ph/0205201].
- [18] C. Gocke, D. Blaschke, A. Khatryan and H. Grigorian, arXiv:hep-ph/0104183.
- [19] M. Buballa and M. Oertel, Nucl. Phys. A 703 (2002) 770 [arXiv:hep-ph/0109095].
- [20] F. Neumann, M. Buballa and M. Oertel, Nucl. Phys. A 714 (2003) 481 [arXiv:hep-ph/0210078].
- [21] I. Shovkovy, M. Hanauske and M. Huang, Phys. Rev. D 67 (2003) 103004 [arXiv:hep-ph/0303027].
- [22] M. Baldo, M. Buballa, F. Burgio, F. Neumann, M. Oertel and H. J. Schulze, Phys. Lett. B 562 (2003) 153 [arXiv:nucl-th/0212096].
- [23] H. Grigorian, D. Blaschke and D. N. Voskresensky, \Cooling evolution of hybrid stars with two-flavor color superconductivity," Preprint MPG-VT-UR 231/02.
- [24] G. S. Poghosyan, H. Grigorian and D. Blaschke, Astrophys. J. 551 (2001) L73 [arXiv:astro-ph/0101002].
- [25] M. Prakash, J. M. Lattimer, A. W. Steiner and D. Page, Nucl. Phys. A 715 (2003) 835 [arXiv:astro-ph/0209122].
- [26] S. Zane, R. Turolla and J. J. Drake, arXiv:astro-ph/0302197.
- [27] J. Cottam, F. Paerels and M. Mendez, Nature 420 (2002) 51 [arXiv:astro-ph/0211126].
- [28] M. Huang, P. f. Zhuang and W. q. Chao, Phys. Rev. D 67 (2003) 065015 [arXiv:hep-ph/0207008].
- [29] M. Frank, M. Buballa and M. Oertel, Phys. Lett. B 562 (2003) 221 [arXiv:hep-ph/0303109].
- [30] O. Kiriya, S. Yasui and H. Toki, Int. J. Mod. Phys. E 10 (2001) 501 [arXiv:hep-ph/0105170].
- [31] N. K. Glendenning, Phys. Rev. D 46 (1992) 1274.
- [32] S. M. Schmidt, D. Blaschke and Y. L. Kalinovsky, Phys. Rev. C 50 (1994) 435.
- [33] J. Berges and K. Rajagopal, Nucl. Phys. B 538 (1999) 215 [arXiv:hep-ph/9804233].
- [34] N. K. Glendenning, \Compact Stars: Nuclear Physics, Particle Physics, And General Relativity," (Springer, New York & London, 2000).
- [35] D. N. Voskresensky, M. Yasuhira and T. Tatsumi, Phys. Lett. B 541 (2002) 93 [arXiv:nucl-th/0109009].
- [36] D. N. Voskresensky, M. Yasuhira and T. Tatsumi, Nucl. Phys. A 723 (2003) 291 [arXiv:nucl-th/0208067].
- [37] J. R. Oppenheimer and G. M. Volko, Phys. Rev. 55 (1939) 374.
- [38] A. Bender, D. Blaschke, Y. Kalinovsky and C. D. Roberts, Phys. Rev. Lett. 77 (1996) 3724 [arXiv:nucl-th/9606006].
- [39] C. D. Roberts and S. M. Schmidt, Prog. Part. Nucl. Phys. 45 (2001) S1 [arXiv:nucl-th/0005064].
- [40] A. Maas, B. Guter, R. Alkofer and J. Wambach, arXiv:hep-ph/0210178.

Form		G_1^{-2}	m	$T_c (\mu = 0)$	$T_c^{(S)} (T = 0)$	$T_c^{(N)} (T = 0)$
Factor	[GeV]		[MeV]	[MeV]	[MeV]	[MeV]
Gauss.	1.025	3.7805	2.41	174	965	991
Lor.	0.8937	2.436	2.34	188	999	1045
NJL	0.9	1.944	5.1	212	1030	1100

TABLE I: Parameter sets (μ, G_1^{-2}, m) of the nonlocal chiral quark model for different form factors discussed in the text. The last three columns show the critical temperatures at vanishing chemical potential and the critical chemical potentials with and without diquark condensate at vanishing temperature, respectively.

k	$a_k^{(N)} [\text{GeV}^{-1-k} \text{fm}^{-3}]$		
	Gaussian	Lorentzian	NJL
0	$7.2942 \cdot 10^{-2}$	$9.0218 \cdot 10^{-2}$	$1.1071 \cdot 10^{-1}$
1	$2.5122 \cdot 10^{-2}$	$7.6973 \cdot 10^{-2}$	$3.0219 \cdot 10^{-1}$
2	$9.1152 \cdot 10^{-2}$	$6.8728 \cdot 10^{-1}$	$1.2820 \cdot 10^{+0}$
3	$1.6402 \cdot 10^{-1}$	$3.7260 \cdot 10^{+0}$	$4.0634 \cdot 10^{+1}$
4	$5.9621 \cdot 10^{-3}$	$1.1862 \cdot 10^{+1}$	$3.0828 \cdot 10^{+2}$
5	$5.1899 \cdot 10^{-1}$	$2.2342 \cdot 10^{+1}$	$1.2301 \cdot 10^{+3}$
6	$9.0892 \cdot 10^{-1}$	$2.4464 \cdot 10^{+1}$	$2.9441 \cdot 10^{+3}$
7	$6.5617 \cdot 10^{-1}$	$1.4374 \cdot 10^{+1}$	$4.3677 \cdot 10^{+3}$
8	$1.7810 \cdot 10^{-1}$	$3.5006 \cdot 10^{+0}$	$3.9376 \cdot 10^{+3}$
9	0	0	$1.9774 \cdot 10^{+3}$
10	0	0	$4.2442 \cdot 10^{+2}$

TABLE II: Coefficients for Bag function fit formula for the normal phase case, for different form factors Eq. (30).

k	$a_k^{(S)} [\text{GeV}^{-1-k} \text{fm}^{-3}]$		
	Gaussian	Lorentzian	NJL
0	$6.5168 \cdot 10^{-2}$	$7.5350 \cdot 10^{-2}$	$8.4897 \cdot 10^{-2}$
1	$1.4638 \cdot 10^{-1}$	$2.8766 \cdot 10^{-1}$	$2.8604 \cdot 10^{-1}$
2	$1.8020 \cdot 10^{+0}$	$3.2215 \cdot 10^{+0}$	$1.0708 \cdot 10^{+0}$
3	$9.8125 \cdot 10^{+0}$	$1.6528 \cdot 10^{+1}$	$2.8157 \cdot 10^{+1}$
4	$3.0515 \cdot 10^{+1}$	$4.9352 \cdot 10^{+1}$	$1.6904 \cdot 10^{+2}$
5	$5.4888 \cdot 10^{+1}$	$8.7023 \cdot 10^{+1}$	$5.4828 \cdot 10^{+2}$
6	$5.6610 \cdot 10^{+1}$	$8.9237 \cdot 10^{+1}$	$1.0896 \cdot 10^{+3}$
7	$3.1096 \cdot 10^{+1}$	$4.9210 \cdot 10^{+1}$	$1.3643 \cdot 10^{+3}$
8	$7.0479 \cdot 10^{+0}$	$1.1275 \cdot 10^{+1}$	$1.0518 \cdot 10^{+3}$
9	0	0	$4.5653 \cdot 10^{+2}$
10	0	0	$8.5443 \cdot 10^{+1}$

TABLE III: Coefficients for Bag function fit formula for the superconducting mixed phase, for different form factors, Eq. (30).

k	$b_k [\text{GeV}^{1-k}]$		
	Gaussian	Lorentzian	N JL
0	$9.33 \cdot 10^{-2}$	$1.04 \cdot 10^{-1}$	$3.30 \cdot 10^{-2}$
1	$2.13 \cdot 10^{-1}$	$1.63 \cdot 10^{-1}$	$1.05 \cdot 10^0$
2	$4.27 \cdot 10^{-2}$	$9.19 \cdot 10^{-2}$	$3.42 \cdot 10^0$
3	$1.14 \cdot 10^{-2}$	$1.92 \cdot 10^{-1}$	$6.13 \cdot 10^0$
4	$5.27 \cdot 10^{-3}$	$8.88 \cdot 10^{-2}$	$5.20 \cdot 10^0$
5	0	0	$1.55 \cdot 10^0$
6	0	0	$1.03 \cdot 10^{-1}$

TABLE IV: Coefficients for the diquark condensate χ formula, for different form factors, Eq. (31).

k	$a_k [\text{GeV}^{-k}]$		
	Gaussian	Lorentzian	N JL
$c^{(s)} [\text{MeV}]$	995	1054	1095
	for $c^{(s)} < B < c^{(v)}$		
0	$1.29 \cdot 10^{-1}$	$8.56 \cdot 10^{-2}$	$7.00 \cdot 10^{-3}$
1	$1.66 \cdot 10^{-1}$	$1.15 \cdot 10^{-1}$	$8.51 \cdot 10^0$
	for $c^{(v)} < B$		
0	$6.28 \cdot 10^{-1}$	$7.17 \cdot 10^{-1}$	$5.60 \cdot 10^{-1}$
1	$3.36 \cdot 10^{-1}$	$2.83 \cdot 10^{-1}$	$2.47 \cdot 10^0$
2	$3.94 \cdot 10^{-1}$	$3.43 \cdot 10^{-1}$	$7.67 \cdot 10^0$
3	$6.57 \cdot 10^{-2}$	$1.19 \cdot 10^{-2}$	$1.06 \cdot 10^{-1}$
4	$8.78 \cdot 10^{-3}$	$2.50 \cdot 10^{-2}$	$5.34 \cdot 10^0$
5	0	0	$6.25 \cdot 10^{-1}$
6	0	0	$7.45 \cdot 10^{-1}$

TABLE V: Coefficients for the volume fraction Eq. (32) and their valid ranges, for different form factors.

	$n_B^{(Q)} [n_0]$		$n_B^{(H)} [n_0]$		$\mu_B^{(H)} [\text{MeV}]$	
	$= 1$	$= 0$	$= 1$	$= 0$	$= 1$	$= 0$
Gaussian	1.84	2.14	1.20	1.68	1005	1059
Lorentzian	3.16	3.66	2.25	2.79	1149	1252
N JL	4.79	5.76	3.02	3.71	1303	1455

TABLE VI: Limiting densities of the coexistence region between quark (Q) and hadron (H) matter phases for different form factors ($n_0 = 0.16 \text{ fm}^{-3}$ is the nuclear saturation density) in the first two columns. The third column shows the critical baryon chemical potential at the phase transition $\mu_B^{(H)} = \mu_B^{(Q)}$, see Fig. 5. The subcolumns indicate the cases of superconducting ($= 1$) and normal ($= 0$) quark matter.