# Cluster Lensing of the CM B

Chris Vale<sup>a;1</sup>, A lexandre Am blard<sup>b;2</sup>, M artin W hite<sup>a;b;3</sup>

<sup>a</sup>D epartm ent of P hysics, U niversity of C alifornia, B erkeley, CA, 94720 <sup>b</sup>D epartm ent of A stronom y, U niversity of C alifornia, B erkeley, CA, 94720

#### A bstract

We investigate what the lensing inform ation contained in high resolution, low noise CMB temperature maps can teach us about cluster mass proles.We create lensing elds and Sunyaev-Zel'dovich elect maps from N-body simulations and apply them to primary CMB anisotropies modeled as a Gaussian random eld.We exam ine the success of several techniques of cluster mass reconstruction using CMB lensing inform ation, and make an estimate of the observational requirements necessary to achieve a satisfactory result.

Key words: Cosm ology:Cosm ic M icrow ave Background,Cosm ology:Theory, Cosm ology:Gravitational Lensing,Cosm ology:Large-Scale Structure of Universe PACS:98.80 k

### 1 Introduction

The study of anisotropies in the Cosm ic M icrowave Background (CMB) has proven to be a gold-m ine for cosm ology. The primary anisotropies on scales larger than  $10^0$  have now been probed with high delity by W MAP (Bennett et al. 2003) over the whole sky, leading to strong constraints on our cosm ological m odel. W ithin the next few years this activity will be complemented by high angular resolution, high sensitivity observations of secondary anisotropies by the SZA <sup>4</sup>, APEX-SZ experiment<sup>5</sup>, the South Pole Telescope (SPT <sup>6</sup>) and the

<sup>&</sup>lt;sup>1</sup> E-m ail: cvale@ astro berkeley.edu

<sup>&</sup>lt;sup>2</sup> E-m ail: am blard@ astro.berkeley.edu

<sup>&</sup>lt;sup>3</sup> E-m ail: m w hite@ astro berkeley.edu

<sup>&</sup>lt;sup>4</sup> http://astro.uchicago.edu/sza/

<sup>&</sup>lt;sup>5</sup> http://bolo.berkeley.edu/apexsz/

<sup>&</sup>lt;sup>6</sup> http://astro.uchicago.edu/spt/

A tacam a Cosm ology Telescope (ACT  $^7$ ) which are aim ing to make arom inute resolution m aps with 10 K sensitivity (or better) at m illim eter wavelengths.

On the angular scales the dom inant secondary anisotropy is expected to be the C om pton scattering of cold CM B photons from hot gas along the line of sight, known as the therm al Sunyaev-Zel'dovich (SZ) e ect ((Sunyaev & Zel'dovich 1972, Sunyzev & Zel'dovich 1980); for recent reviews see Rephaeli (1995) and B irkinshaw (1999)). The therm alSZ e ect can be spectrally distinguished from prim ary CM B anisotropies given enough sensitivity and frequency coverage, and we shall not consider it in this work. At slightly lower am plitudes are the kinetic SZ e ect and gravitational lensing, both of which leave the CM B spectrum unaltered but modify the spatial correlations and statistics of the signal. These e ects can in principle provide useful constraints on re-ionization models (the kSZ e ect, e.g. (Zhang, Pen & Trac 2003)) and allow us to map the dark matter back to the surface of last scattering (lensing, e.g. (Seljak & Zaldarriaga 1999, Zaldarriaga & Seljak 1999, Hu 2001, Hirata & Seljak 2003, O kam oto & Hu 2003)).

In this paper we want to consider gravitational lensing of the CM B by galaxy clusters. This was rst studied by Seljak & Zaldarriaga (2000), which is the starting point for our work. Relatively little other work has been done on this phenom enon, notable exceptions being the work of Cooray (2003) who described a m ethod to m easure the equation of state of the dark energy, Bartelm ann (2003) who gave CM B lensing as an example of num erical techniques and Holder & Kosowsky (2004) whose aim was quite sim ilar to the work presented here.

O ur goal is to study how well, and in what manner, we can reconstruct the cluster prole, or an integrated quantity such as the mass, from the lensing induced distortion in the CMB temperature eld<sup>8</sup>, or conversely to understand the impact of large collapsed structures on the statistics of the CMB. The principle advantage of CMB lensing over traditional lensing of galaxies is that the source redshift is almost perfectly known. The main disadvantage is that it is presents a single, xed source plane. Lensing also represents auxiliary science that can be done with already planned or funded instrum ents, at little or no additional cost. As such it is worth investigating in some detail.

It has become well known that lensing su ers from severe projection e ects (Reblinsky & Bartelmann 1999, Metzler, W hite & Loken 2001) so we shall here consider how well the projected mass pro les can be constructed from

<sup>&</sup>lt;sup>7</sup> http://www.hep.upenn.edu/ angelica/act/act.htm l

<sup>&</sup>lt;sup>8</sup> W e shall only consider lensing of the tem perature anisotropies in this work, neglecting polarization. This is partly motivated by the fact that some of the upcoming experiments will not be polarization sensitive, and partly to keep the calculation under control.

CM B lensing, leaving aside the question of how well such pro les can be deprojected to get 3D quantities. In order of decreasing desirability we would like to reconstruct the convergence map (projected density) of every cluster in the eld; compute the total convergence (mass) of every cluster in the eld; compute the pro le of Stacked' clusters or compute some integral of the Stacked' pro le.W e shall investigate each of these in turn.

The plan of the paper is as follows. In x2 we introduce the lensing form alism, largely following Seljak & Zaldarriaga (2000), and introduce our notation.We im plement the S& Z program me in x3, where we show how the procedure works in a simple toy model of an ideal cluster lensing a pure, known CMB gradient. Then we begin to add com plications in x4, looking in particular at the fact that the CMB is not a pure gradient, the contam ination from kSZ (which is highly correlated with the lensing structures) and the non-G aussianity of the lensing eld and nally at the elects of noise. We summarize with our conclusions in x5. Some details of the simulations we use to make mock observations are given in an Appendix.

## 2 The Theory of Cluster Lensing of the CMB

In this section we review the e ect of cluster lensing on the CMB.Our goal is to explore to what extent the mass and the mass prole of a cluster may be constrained using information from high resolution temperature maps of the CMB if the cluster's redshift and position on the sky are known.We introduce the formalism of CMB lensing but provide only a brief sum mary of equations directly relevant here; see Bartelmann & Schneider (2001) for a comprehensive review of weak lensing. Throughout this paper we work in the weak lensing limit, assume a at CDM universe, adopt units where where the speed of light c = 1, and work in com oving coordinates.

#### 2.1 Weak Lensing of the CMB

We begin by examining the gravitational lensing of light rays that originate at the surface of last scattering by inhom ogeneities in the intervening matter distribution. We dene the primordial CMB temperature eld at the surface of last scattering as T<sup>(0)</sup> at an angular position on the sky <sup>0</sup>. Lensing by large scale structures such as clusters will cause CMB light rays that originate at a position <sup>0</sup> to be de ected by an angle to an observed position on the sky , so that the observed temperature T () is

 $T() = T(^{0}) = T()$  (1)

It is simple to derive a mathematical expression for the dejection angle in the weak lensing limit. The total dejection angle of a source at position  $_{\rm s}$  as seen by an observer at = 0 is

$$= \frac{2}{s_{0}} \int_{0}^{2s} d(s_{s}) r_{?}$$
(2)

where r<sub>?</sub> denotes the spatial gradient perpendicular to the path of the light ray, is the three-dimensional peculiar gravitational potential, and is the radial com oving coordinate.

As we discuss below, we will be interested in the dependence of the lensing angle on the matter distribution. To see this dependence, we de ne the convergence  $\frac{1}{2}r$  where r is the angular gradient operator. Then

$$Z_{s}$$
  
d g(; s)r<sup>2</sup> (3)

where we have invoked the Born and Limber approximations (see Jain, Seljak & W hite (2000) and Vale & W hite (2003) for a discussion) and made use of the lensing kernel

$$g(; s) = \frac{(s)}{s}$$
(4)

Them atter density is related to the three-dimensional gravitational potential through

$$r^{2} = 4 G_{0} - a$$
 (5)

where all quantities are dened with respect to comoving coordinates, G is the gravitational constant,  $_0$  is the mean density of the present day universe, and

 $=_0$  1 is the relative m ass overdensity. It is worth noting that if only local lensing information is available, equation (5) will be uncertain up to an overall constant. This is the source of the so called m ass sheet degeneracy. We are now in a position to relate to the matter density by combining equations (3) and (5)

() 
$$4 G_0 d g(; s) \frac{(;)}{a}$$
 (6)



Fig. 1. The convergence m aps for isolated and non-isolated clusters as found in our simulations. The color bar shows the value of the (dimensionless) convergence  $\ .$ 

If the lensing e ect is primarily due to a single structure whose size is much less than its comoving distance , one can make use of the thin lens approximation. Then the convergence is simply related to the projected two dimensional mass density R d of the lens, so that equation (6) becomes

() 
$$4 G_0 g(; s) (; )$$
 (7)

W e m ake use of equation (7) to create the convergence eld from the N-body simulations (e.g.Figure 1), as detailed in the Appendix. From the convergence eld we compute the de ection angle via

$$= r_{2} r^{2}$$
 (8)

and hence the lensed tem perature eld.

#### 2.2 Lensing by an Ideal Cluster

It is instructive to exam ine the lensing e ect of an isolated cluster in the absence of other secondary anisotropies, foregrounds, instrum ent e ects, or lensing by other structures, all of which we include later. If the de ections are sm all, we may expand the right hand side of equation (1) to linear order, so that

 $T() \quad \tilde{T}() \qquad \tilde{T}() \tag{9}$ 

W e will assume that equation (9) holds both here and in x3 for the purpose of illustrating some basic ideas. However, we note that this approximation



Fig. 2. (left) A 1-d cut of the lensing signal T () for a circularly symmetric ideal cluster lensing the CMB. The red line shows the unlensed CMB, the blue line shows the lensed CMB with the characteristic kink' near the origin and the green line shows the di erence. (right) The signal in 2D for a circularly symmetric ideal cluster lensing a constant gradient. Note the dipolar nature of the lensing signal.

is not actually necessary, and we shall dispense with it altogether in x4. It's useful to consider the de ection angle due to a spherically symmetric cluster at com oving distance

$$= 4G \frac{(s)^{M}}{s} (10)$$

where = (x; y) is now de ned as the angular displacement from the center of the cluster, is the absolute value of , and M () is the mass of the cluster within a radius . To rst order, the cluster's lensing e ect is to rem ap the CMB radially away from its center, creating a step like wiggle in the CMB gradient centered on the cluster (Seljak & Zaldarriaga 2000). We give an example of this behavior for a typical large cluster from our simulations in Figure (2). As expected from equation (9), the magnitude of the e ect is proportional to both the local gradient of the CMB and the de ection angle along that gradient.

The lensing angle cannot be solved for using equation (9) without more information; you can't in general measure a scalar eld and expect to solve for a scalar eld and a vector eld! If progress is to be made, som e assumptions must be made about , rT'(), or both. W e begin by noting that, in the absence of secondary anisotropies, the CM B is expected to have little power on small angular scales, so that <math>rT'() may be slow ly varying in relevant regions near the cluster's center. W e exam ine this statem ent more carefully below, but for now we follow Seljak & Zaldarriaga (2000) and make use of this

idea to model the primordial CMB gradient in small regions near the cluster as a constant whose direction of steepest ascent can without loss of generality be taken as the y-axis, so that

$$T'() T'_{y0 y}$$
 (11)

where  $T_{y0}$  is the slope of the primordial CMB along the y-axis and we have ignored an overall constant.

We de ne the lensing signal due to a cluster T() as the di erence between the lensed and unlensed CMB temperature

T() T() T() T()(12)

C om bining this with equations (9) and (11) and de ning the de ection due to lensing by the cluster along the y-axis as  $_{\rm v}$  gives

$$\Gamma() \quad T() \quad \tilde{T}_{y0 y} \quad y() \quad \tilde{T}_{y0} \tag{13}$$

In Figure (2) we show T () for a circularly symmetric isolated cluster lensing a constant gradient. The signal crudely resembles a dipole in appearance as you would expect from equation (9), though it falls o as M () = away from the cluster as predicted by equation (10).

The lensing angle  $_{y}()$  is not generally considered measurable because the original position of the background in age isn't known. However, far away from the cluster, the e ect of lensing must be small, and the lensed and unlensed CMB must be roughly equal. If the approximation of equation (11) is valid at this distance, then it will be possible to measure T () away from the cluster, determ ine  $T_{y0}$ , and solve for  $_{y}()$ . A swe show in Figure (4), the convergence prole can be well reconstructed from this information. We note that the reconstruction is degenerate for density uctuations that change  $_{x}$  but don't alter  $_{y}$ . This degeneracy is similar to that of the mass sheet degeneracy, but it applies to any line of constant density in () that happens to lie in the direction of the y-axis. We not the error due to this degeneracy to be quite small, and it is of course identically zero for a circularly symmetric cluster prole, where  $2 = \frac{2}{x} + \frac{2}{y}$ 

It is clear from F igure (4) that reconstructing the convergence pro le of a cluster from CMB temperature m aps is certainly possible under the following highly articial conditions:

No foregrounds

No instrument e ects

N o CM B secondary an isotropies other than lensing The clusters are isolated The CM B is a pure gradient of constant slope

In x3, we begin to include these e ects in the context of a toy m odel.

3 Results From a Toy M odel

In this section, we make use of a toy model of cluster lensing of the CMB in order to investigate to what extent the issues raised in the bulleted points listed at the end of x2 will impact on our ability to reconstruct the convergence proles of clusters using high resolution CMB temperature maps. We then present results from this toy model for two general cases: individual clusters and \stacked" average clusters.

3.1 Signal and Noise in the Toy Model

In the toy model, we assume that the primordial CMB in a small region near a cluster is a known quantity, which we then model as a gradient. This allows us to bypass the step of estimating the unlensed CMB from the actual maps (we address this issue in x4), which will be both lensed and noisy, and instead to directly measure the signal as de ned in equation (12) plus a noise term N (), which includes all other elects, so that

$$T_{t}() = T() + N()$$
 (14)

where we de ne the measured signal in the toy model as T<sub>t</sub>. A coording to equation (10), the de ection angle M ()=, and from equation (13) the signal in any circular annulus is proportional to times the slope of the CMB gradient  $T_{y0}$  times the area of the annulus. Then the total signal to noise inside a circle of radius scales as

$$\frac{S}{N} = \frac{T_{y0}}{\frac{1}{2}}^{Z} d M ()$$
(15)

for G aussian noise. Since the model relies on the approximation that the CMB gradient is constant (equation 11), which can only be valid on small scales, we have chosen to impose a cut-o radius of  $4^{\circ}$  from the center of the cluster. That is, we assume a measurement of T<sub>t</sub> for <  $4^{\circ}$ , and use no information at larger radii. We note that we do address the use of lensing information for reconstruction on larger scales elsewhere (Amblard, Vale & White 2004).

The toy signal T<sub>t</sub> is derived using an input convergence m ap m ade by raytracing through our N-body simulation. This convergence m ap is then used to make the de ection eld  $_{\rm y}$  using Fourier m ethods, and T is made by remapping T( <sup>0</sup>) into T () according to equation (1), where T( <sup>0</sup>) is a gradient of constant slope. We provide a description of the N-body simulation and the creation of the convergence m ap in an appendix, and for now note only that the simulated clusters are in general neither isolated nor ideal, as can be seen in the convergence pro les of Figure (1).

A fler creating a map of the toy signal, we next introduce two noise components to the model. The rst, the kinetic SZ, is highly correlated with the position of the cluster, and because it has the same spectral dependence as the CMB itself, it cannot be removed using multi-frequency measurements. We then add a G aussian \white noise" component, which is uncorrelated with the location of the cluster, and can be thought of as instrument noise, and unless stated otherwise, we smooth with a 0.75 beam (FW HM), such as is expected for APEX-SZ.We do not include noise sources which we expect to be small and uncorrelated with the cluster, such as other CMB secondaries, nor do we include point sources, dust, or the thermal SZ, which can in principle be removed or at least reduced by making use of their spectral dependence. In any real experiment, treatment of these elects will not be perfect, so excluding them entirely is somewhat optim istic.

# 3.2 Reconstruction of Cluster Pro les

In this section we address the issue of the reconstruction of cluster convergence pro les within the toy m odelboth for individual clusters and \stacked" average clusters. W e shall see that both instrum ent noise, and m ore permiciously the kinetic SZ, signi cantly degrade the reconstruction for individual clusters.

Let us begin by looking at the reconstruction including kinetic SZ for a typical large m ass, high-z cluster in F igure (3). Note that when the kinetic SZ is added the reconstruction is signi cantly altered, because som e of the kinetic SZ signal ism is interpreted as lensing. This e ect is particularly troublesom e because it is spatially correlated with the lensing signal, spectrally indistinguishable from it, non-G aussian, and only loosely correlated with the m ass or therm al SZ signal from the cluster.

In the last panel of F igure (3), we show a reconstruction with no kinetic SZ but now instead including G aussian random noise at a level of 3 K -arcm in, roughly one third of that expected for the highest resolution m aps from APEX - SZ and consistent with m ore ambitious future projects such as ACT and SPT. Even at this noise level, m ost of the features of the convergence m ap are lost,



Fig. 3. The reconstructed convergence maps for a typical cluster of  $M_{200} = 2.35 \quad 10^{14} \, h^{-1} \, M$  and  $z = 1. W \, e$  show the input map, the reconstruction with kinetic SZ, then with 3 K-arcm in of instrument noise and no kinetic SZ.



Fig. 4. The convergence reconstruction for a large (M  $_{200} = 7 \quad 10^{14} \, h^{-1} \, M$  ), high redshift (z = 0:95) cluster. The left panel is the true convergence sm oothed by 0: $^{0}$ 75, the center panel is the the reconstructed convergence using the toy m odel and no noise, and the right panel is the reconstructed convergence including the kinetic SZ. In this unusual case, the kinetic SZ dom inates the reconstruction com pletely, causing the center of the cluster to appear as large and negative.

and it is evident that high quality reconstruction of cluster convergence m aps for typical clusters is not achievable within the context of the experimental parameters we are considering. A strategy designed to integrate deeply on cluster locations would have to be adopted.

W hile the situation depicted in Figure (3) is typical, in some cases the kinetic SZ can be completely dom inant, as we show in Figure (4). In this particular case the cluster happens to be rotating, so that the kinetic SZ signal has a dipole-like structure sim ilar to that produced by lensing. D epending on the relative orientation of the kinetic SZ lobes and the direction of the CM B gradient, these signals can enhance or overwhelm the lensing signal. It is even possible to reconstruct a large negative convergence right at the cluster's location!

W hile this \disaster" cluster is clearly beyond hope, the reconstruction of a typical cluster m ight be improved if we mask out pixels which are likely to



Fig. 5. The kinetic SZ (left) for a typical (non-isolated) cluster and the therm al SZ (center) that you could try to use to mask it. The right panel is an example of a typical masked map, where the masked region is shown in black and the unmasked region is now (for clarity) the absolute value of the kinetic SZ. The therm al SZ is not a perfect tracer of the kinetic SZ, resulting in map with many perfectly good pixels removed, and residual kinetic SZ still included. The color bar in each case shows T in K.

contain a large kinetic SZ component, which can be done by using the therm al SZ to roughly estimate the likelihood of a large magnitude kinetic SZ. This is essentially the idea proposed in Seljak & Zaldarriaga (2000). Figure (5) illustrates why this is not in general helpful; most clusters are not isolated, and many pixels must be excised. To make matters worse, the therm al SZ is not a perfect tracer of the kinetic SZ, so many perfectly good pixels are thrown away while others with large kinetic SZ signals are included.

O byiously, even if most clusters are not suitable candidates, we may be able to select some that are and focus on those. We have so far been thwarted in our reconstruction e orts by the kinetic SZ and instrument noise. If we ignore the latter for the time being, we realize right away that, although rare, clusters do sometimes form in relative isolation, as depicted in Figure (1). Perhaps these will be suitable?

Indeed, the level of contam ination from the kinetic SZ is dram atically lower, and mostly associated with the cluster itself. We may now mask out the center of the cluster and have a reasonable expectation that the kinetic SZ contam ination will be under control, and indeed that does prove to be the case here. However, we have speci cally gone out of our way to select an isolated cluster, using the therm al SZ, and this has introduced a strong selection bias. The absence of other structures is marvelous for controlling the impact of the kinetic SZ, but the cluster is now located in a large void which dram atically reduces the projected mass below what one would expect from an average location. In essence we have introduced a mass sheet degeneracy. This is depicted in F igure (8). Here, we present the actual detection angle along the y-axis. The dot-dash curve is the lensing angle that would result if the cluster were the only object in the universe, the solid line is the result including the cluster plus projection



Fig. 6. The cumulative convergence for our stack of 61 clusters with  $2 < M = 10^{14} h^{-1} M < 3$  and 1 < z < 1.5 as a function of radius. Shown here are the input (red line), the reconstruction including kinetic SZ (green line), and reconstruction including kinetic SZ and noise (blue line).

e ects, and the dashed line is the actual de ection angle, where structures far away from the cluster of interest contribute to the de ection angle even though they don't alter the convergence (this last is a degeneracy that arises because we have access only to the y-axis de ections). Thus, even though we \cheated" and simply used the y-axis de ection angle directly, rather than m easuring it, we are still unable to reconstruct the clusters m ass from this inform ation due to a strong m ass sheet degeneracy that in fact is consistent with a cluster m ass estim ate of zero at the virial radius, with absolutely no noise introduced, simply because of the confusion introduced from lensing by other structures.

O ur investigation of the likely success of reconstruction of individual cluster m ass pro les has shown that it will prove a more di cult problem than one m ight have hoped. It is worth noting that some of the di culties we have encountered have come because we have used full eld lensing and SZ sim ulations, rather than clusters that have been 'cut out' of simulations and then used to lens voids. D espite this technical di erence though, our results so far are qualitatively in agreem ent with those of H older & K osow sky (2004). They show in their F ig. 4 that they reconstruct the cluster m ass to be essentially zero at the virial radius. W e shall return to this point, and som e of the reasons



Fig. 7. The convergence map (left) of our stack of 61 clusters with  $2 < M = 10^{14} h^{-1} M$  < 3 and 1 < z < 1.5, it's reconstruction (center) including the kinetic SZ, and then again (right) with both Gaussian noise and kinetic SZ included.

behind it, in the next section.

A nother approach, called stacking, is to consider a group of clusters binned according to relevant observables (e.g. tem perature and redshift), cut out regions in the signal maps within a speci ed area centered on each cluster, stack them one on top of another, and then com pute the average. The principal advantage of this technique is that essentially every unbiased source of confusion is reduced by this kind of averaging as N, where N is number of clusters used. This includes the important cases of the kinetic SZ, since the baryons are a priori equally likely to be moving toward or away from the observer, and of projection e ects that proved so troublesome in the case of \isolated" clusters. If our cluster signal grows as M , our signal-to-noise is enhanced by stacking more relatively low mass clusters as long as the mass function is steeper than M<sup>2</sup> per mass interval considered. This is the case for the high mass clusters rather than fewer high mass clusters.

In Figure (7) we show the reconstruction of a stack m ade from 61 interm ediate m ass, high redshift clusters. First, consider the case where the only source of noise is the kinetic SZ. The convergence prole of the stack is circularly symmetric to a good approximation, which greatly reduces the mass line artifact that can arise in the reconstruction of nite elds, and it becomes feasible to reconstruct more of the central region. As in the case of single clusters, adding G aussian instrument noise of 1.25 K -arcm in to the stack (consistent with the anticipated 10 K -arcm in for APEX-SZ averaged over 61 clusters) introduces substantial uncertainty to the convergence prole, but the total convergence as a function of radius is reasonably constrained, as can be seen in Figure (6). Thus, there is some promise that one might be able to provide reasonable m assetim ates for cluster stacks, which might then be used to constrain the mass-tem perature relation.



Fig. 8. The de ection angles for an isolated cluster that occurred naturally in our simulations. The dot-dash line shows the de ection angle for the cluster if there were no other matter in the universe. The solid line shows the lensing from the projected convergence prole at the cluster's location on the sky. Since lensing is dependent on the relative mass overdensity , the void acts as a negative mass sheet. Finally, the dashed line is the actual de ection angle. Structures away from the center of the cluster do contribute to de ections, som etim es quite substantially.

The reconstruction of cluster convergence pro les using weak lensing of the CMB is a technique that clearly shows prom ise for cluster stacks if the approximations of the toy model can be trusted. Some potentially important issues are not addressed by this simple model, the most important of which is the fact that the unlensed CMB is not a known quantity, so the signal as described in equation (13) is not directly measurable. We exam ine this issue in the next section.

## 4 Beyond the Toy M odel

In the previous section we showed that using high resolution m aps of the CM B tem perature to reconstruct stacked cluster convergence proles is a technique that shows some promise when examined within the context of a simple toy m odel. By employing a toy m odel we have brushed aside a number of potentially important complications, perhaps the most glaring of which is having so



Fig. 9. The lensing signal estim ated using the ansatz of Seljak & Zaldarriaga (2000) (Leff.) The lensed CMB (blue) is used as an estim ate of the unlensed CMB on either side of the cluster, and the unlensed CMB (dashed orange) is estim ated by connecting the dots. The di erence between the two is then the estim ate of the signal (red). (R ight) A generic feature of this method of reconstruction is its system atic underestim ate of the magnitude of the signal. The actual signal (dot-dash orange) is included for comparison.

far ignored the uctuations intrinsic to the prim ordial CMB itself. A lthough ourm ain focus in this section will be on stacked clusters, we nonetheless begin with some discussion in the context of individual clusters as a natural lead in to understanding this issue.

First recall the toy model ansatz of Seljak & Zaldarriaga (2000) where the CMB temperature gradient can be approximated as constant over relevant length scales, and the CMB can be measured far from the cluster where the kinetic SZ and lensing by the cluster are both small. The unlensed CMB gradient can then be simply determ ined, as we illustrate in Figure (9), by bonnecting the dots' between the CM B on either side of the cluster. However, the CMB cannot be well approximated as a gradient on scales greater than a few arom inutes even for carefully selected portions of the CMB, so to use this technique to get a good t for the unlensed CMB at the center of the cluster you are forced to set the lensing signal (and therefor the de ection angle) to zero at roughly a few arom inutes from the center of the cluster. O byjously the lensing e ect of large clusters extends well beyond this range, so the 'connect the dots' m ethod system atically underestim ates the cluster's mass (Equation 10) as you go farther from the cluster's center, culm inating in a total cluster m ass estim ate equal to zero on scales of a few arom inutes; that is, roughly at the virial radius. This is a completely generic feature of the method, and is easily seen in the second panel of Figure (9) or in Figure (4) ofHolder & Kosowsky (2004).

A betterm ethod of estimating the unlensed CMB is certainly of interest. One

obvious in provement is to extend on the original proposal of Seljak & Zaldarriaga (2000) by making use of som ething other than a simple gradient. For example, one could t the CMB far from the cluster with a second order polynom ial (but note that nearby structure may cause problems). A lternatively, you could use a W iener lter, as was recently suggested by Holder & K osow sky (2004). Unfortunately, although the t is som ewhat im proved, these m ethods appear to su er from the same draw back as the original, and lead to estimates of the virial mass that are system atically low and often consistent with zero.

A more sophisticated approach would be to iterate the torto include a model for both lensing and the unlensed CMB. This is the approach we take. Specifically we assume that the cluster is spherically symmetric and has an NFW prole, and we model the unlensed CMB locally as a 2nd degree polynom ial in 2D (the above mentioned nearby structure contamination will average out when stacking). We consider a range of masses for the assumed NFW cluster and compute the lensing de ection angle for each. This is used to \delens" the CMB map. The resulting map which best ts, in the <sup>2</sup> sense, a second degree polynom ial (in 2D) is chosen. A second degree polynom ial has the advantage of being even about the center of the cluster, while the lensing e ect is odd, and on scales of a few arcm inutes tting an unlensed CMB fairly well. As a result, bad' ts will often be glaring.

We initially tested this method on clusters but out 'from simulations and with no instrum ent noise, kinetic SZ, etc., and were able to routinely estim ate the mass to within 10%, and to within a few percent for our ducial 61 cluster stack. However, the method incurs a bias toward underestimating the mass that becomes more signicant as you include regions further from the center of the cluster. The bias occurs due to the failure of the 2D polynom ial to accurately represent the CMB in the region around a cluster. One look at Figure (9) will convince the reader of the origin of the bias. The departure of the CMB from a constant gradient is also odd about the center of the cluster, so that while lensing by a cluster will degrade the ability of the polynom ial to t CMB near the cluster, it will actually help it t better farther away, and the two e ects compete. Given the need for as much usable area as possible to overcome other di culties, it is likely that a 2D polynom ial is ultimately not the best choice for the t. One way to correct for this is to determ ine the expectation value of the odd component in the unlensed CMB and include this in the tting procedure. A Itematively with enough clusters to beat down the kSZ contam ination, a higher resolution, higher sensitivity observation would be able to work at smaller radius where the bias is much reduced.

# 5 Conclusion

In this paper we have investigated in som e detail the prom ise of cluster lensing of the CMB. This idea was rst introduced by Seljak & Zaldarriaga (2000) and will soon become observationally feasible with the imm inent comm issioning of the APEX-SZ telescope. We verify that the method of Seljak & Zaldarriaga works well given their assumptions, but note that these assumptions are not well satis ed in practice.

In particular we highlight the role of the kinetic SZ signal as an important contam inant which is spatially correlated with the cluster and spectrally indistinguishable from the lensing signal itself. The kSZ uctuations, being non-G aussian and signal-correlated, are also an issue for reconstruction of largescale structure as we discuss further in Am bland, Vale & W hite (2004). If the unlensed CM B can be adequately estim ated, we show that the stacked proles and totalm asses are reasonably well constructed by CM B lensing, in contrast to proles orm asses of any individual cluster. We elucidate some of the issues in x4.

F inally we make some comments about the role of polarization in lensing. It has been emphasized before that the inclusion of polarization information can dram atically enhance the prospects for large-scale structure reconstruction from lensing of the CMB. This is because lensing induces a B-m ode polarization signal which is otherwise absent for purely scalar, primary uctuations. The large intrinsic signal, which is a source of hoise' for lensing reconstruction, is thus absent.

It is possible that the addition of polarization inform ation could enhance the prospects for cluster lensing also. To see whether the elects we have identied are mitigated by polarization inform ation requires a detailed calculation. The signal levels for polarization are much smaller, and the spatial structure complicated as for temperature. The kSZ elect, which is one of our major contaminants, is also polarized. The dominant polarization signal comes in at order Q where is the cluster optical depth and Q the local CMB intensity quadrupole at the location of the cluster. The resulting eld is a mix of E – and B -m ode signals with the polarization pointing in the direction of the quadrupole cold lobe. Depending on the variation of the quadrupole with distance and position in the eld, and on the cluster though it might be possible to signil cantly reduce the kSZ contamination through modeling. We leave a detailed investigation of this question to future work.

#### A cknow ledgem ents:

C.V. would like to thank the organizers of the workshop \Cosmology with Sunyaev-Zel'dovich cluster surveys" held in Chicago in September 2003 for allowing him the chance to present some of this work. A dditionally we would like to thank T. Chang, J. Cohn, D. Holz, B. Jain, A. Lee, G. Smoot, M. Takada and M. Zaldarriaga for helpful discussions about these results. The simulations used here were performed on the IBM -SP2 at the National Energy Research Scientic C om puting Center. This research was supported by the NSF and NASA.

### References

- Amblard, A., Vale, C., White, M., 2004, in preparation
- Bennett C L., et al, 2003, ApJS, 148, 1 [astro-ph/0302208]
- Bartelm ann, M., & Schneider, P., 2001, Phys. Rep., 340, 291-472
- Bartelm ann M., 2003, preprint [astro-ph/0304162]
- Birkinshaw M ., 1999, Phys. Rep., 310, 98
- Cooray, A., 2003, ApJ, 596, L127
- Hirata, C.M., Seljak, U., 2003, PhysRevD, 67, 43001
- Holder, G., & Kosowsky, A., 2004, preprint [astro-ph/0401519]
- Hu, W, 2001, ApJL, 557, L79
- Jain B., Seljak U., W hite S.D. M., 2000, ApJ, 530, 547 [astro-ph/9901191]
- Metzler C., White M., Loken C., 2001, ApJ, 547, 560 [astro-ph/0005442]
- O kam oto, T., Hu, W., 2003, PhysRevD, 67, 83002
- Reblinsky K., Bartelm ann M., 1999, A&A, 345, 1
- Rephaeli, Y., 1995, ARA & A, 33, 541
- Schulz A., W hite M., 2003, ApJ, 586, 723 [astro-ph/0210667]
- Seljak U., Zaldarriaga M., 1996, ApJ, 469, 437 [astro-ph/9603033]
- Seljak, U., Zaldarriaga, M., 1999, Physical Review Letters, 82, 2636
- Seljak, U., & Zaldarriaga, M., 2000, ApJ, 538, 57-64 (2000ApJ...538...57S)
- Sunyaev R A., Zel'dovich Ya.B., 1972, Comm. Astrophys. Space Phys., 4, 173
- Sunyaev R A , Zel'dovich Ya.B., 1980, ARA & A, 18, 537

Vale C., W hite M., 2003, ApJ, 592, 699 [astro-ph/0303555]
W hite M., 2002, ApJS, 143, 241 [astro-ph/0207185]
Yan R., W hite M., Coil A., 2004, ApJ, in press [astro-ph/0311230]
Zaklarriaga, M., Seljak, U., 1999, PhysRev D, 59, 123507
Zhang, P., Pen, U., Trac, H., 2003, [astro-ph/0304534]

# A The simulated map

We construct maps of lensing convergence and the thermal and kinetic SZ e ect making use of a large, high-resolution N-body simulation of the CDM cosm ology (speci cally M odel 1 of Yan, W hite & Coil (2004)). In this appendix we give some details of how this was done.

# A.1 The N-body simulation

To construct the maps we need some information on the spatial distribution and evolution of the mass in our model. We obtain this from an N-body simulation. The simulation modeled a large volume of the universe, a periodic cube 300 h<sup>-1</sup> M pc on a side, to ensure a good sampling of the clusters of interest to us. We considered only the dark matter component which was modeled using  $512^3$  particles of mass 1:7  $10^{0}$ h<sup>-1</sup> M . For computation of the therm al and kinetic SZ e ects we assume that any baryonic component would trace the dark matter, a reasonable approximation on the scales of interest to us.

The simulation was started at z = 60 and evolved to the present using the TreePM code described in W hite (2002), with the full phase space distribution dum ped every 100h <sup>1</sup> M pc between redshifts 2 > z > 0. It is this range of redshifts which dom inates the lensing and SZE signal on the angular scales of interest to us. The gravitational softening used is of a spline form, with a \P lum m er-equivalent" (com oving) softening length of 20h <sup>1</sup> kpc. All of the relevant cluster-scale halos contain several thousand particles to begin to resolve sub-structure. The simulation was performed on 128 processors of the IBM -SP2 at NERSC, took nearly 4000 time steps and approximately 100 wall clock hours to complete.

The maps were made in essentially the same manner as in Schulz & W hite (2003) to which we refer the reader for discussion of the various approxim ations. The past lightcone was constructed by stacking the interm ediate stages of the simulation between redshifts 2 > z > 0. In order to avoid multiply sam – pling the same large scale structures, each 300h <sup>1</sup> M pc box has been random by re-oriented in one of the six possible orientations, and has furtherm ore been shifted by a random amount, perpendicular to the line-of-sight, making use of the periodic boundary conditions. There are three time dum ps perbox length. Each 300h <sup>1</sup> M pc volume in the stack is made up of three segments, each segment evolved to a later epoch than the previous one by the time it takes light to travel 100h <sup>1</sup> M pc. We chose 100h <sup>1</sup> M pc as the sam pling interval because it is large enough that edge e ects are minimal, yet ne enough that the line of sight integrals are well approximated by sum s of the thirds as the oldest, cyclically permuting the other two. This approach preserves the continuity of large-scale structure over distances of 300h <sup>1</sup> M pc w ithout com promising the resolution in time evolution.

W e produced m aps of the lensing and SZ e ects. Each m ap was 7:5 on a side, with  $2048^2$  square pixels each  $0.22^0$  on a side. The same particle distribution and random seeds were used to construct each of the m aps, so that they represent the same patch of sky in each quantity. W e now discuss each m ap in turn.

## A.2 Convergence () maps

The e ect of lensing was computed from maps of the convergence, , assuming the weak lensing approximation (see x2). This is valid except near the very center of the cluster, and so adequate for our purposes. The convergence was computed from the density contrast along the past lightcone as

$$= \frac{3}{2} \prod_{m at} H_0^2 d_{g()} - \frac{1}{a}$$
 (A.1)

where  $g() = (_s ) = _s$  is the lensing kernel, is the com oving distance and  $_s' 9h^1$ Gpc is the com oving distance to the last scattering surface.We have assumed the universe is spatially at. The (projected) density contrast within each 100 h<sup>1</sup> Mpc slice was computed from the dark matter distribution using a spline kernel interpolation with a sm oothing length equal to the force softening in the simulation.

# A.3 Compton-y and bm aps

Because the simulation contains no gas we use a sem i-analytic model to include the gas physics. First we assume that the gas closely traces the dark m atter. This is likely a good approximation in all regions except the innerm ost O (100)kpc of the cluster, which for clusters at cosm ological distances will be unresolved by the experiments of interest. (e.g. 100kpc subtends only  $0.26^{\circ}$  at z = 0.5.) W e ignore the presence of cold gas and stars in the ICM, assuming that the mass in hot gas is  $_{\rm b} = _{\rm m}$  of the total. Second, each cluster is assumed isothermal. We assign to each particle in a group a temperature proportional to its mass to the 2=3 power.

W e generate C om pton-y m aps by integrating for each pixel

$$y = \int_{T}^{Z} n_e \frac{k_B T_e}{m_e c^2} dl \qquad (A 2)$$

Here  $_{T}$  is the Thompson scattering cross section, and  $n_{e}$ ,  $m_{e}$  and  $T_{e}$  are the electron number density, mass and temperature respectively. We assume that within the clusters the gas is fully ionized. The contribution from each particle is distributed over the pixels with a spline weighting and a size equal to the sm oothing length of the simulation as for the maps above. The temperature uctuation at frequency is then obtained from the y-maps by

$$\frac{T}{T} = y x \frac{e^{x} + 1}{e^{x} - 1} 4$$
 (A.3)

where  $x = h = k_B T_{CM B}$  ' =56.84G H z is the dimensionless frequency and the second expression is valid in the Rayleigh-Jeans lim it. In what follows we shall assume the low-frequency lim it unless otherwise stated.

The C om pton-b m aps are produced in an alm ost identical m anner, replacing  $kT = m_e c^2$  with v=c, where v is the line-of-sight velocity. The spectrum is an undistorted black body, so the tem perature perturbation is simply T=T = b.

## A.4 Primary CMB anisotropies

W hen needed we generate prim ary CM B anisotropies as a random realization of a Gaussian eld with power spectrum computed from CM B fast (Seljak & Zaldarriaga 1996). We generate random phases in momentum space, and assign am plitudes to each of the k-m odes using a distribution whose average value is the am plitude in the CM B power spectrum. We have used the at sky approximation, in which the k-m ode in momentum space corresponds to `value in the CM B power spectrum.