# CONSTRAINING THE MAGNETIC EFFECTS ON HIROTATION CURVES AND THE NEED FOR DARK HALOS

F.J.Sanchez-Salcedo

Instituto de Astronom a, UNAM, C iudad Universitaria, Aptdo. 70 264, C P. 04510, M exico C ity, M exico jsanchez@ astroscu.unam m x

and

M . Reyes-Ruiz Instituto de Astronom a, UNAM, Aptdo. 877, Ensenada, B.C. 22800, Mexico D raft version February 19, 2022

## ABSTRACT

The density proles of dark halos are usually inferred from the rotation curves of disk galaxies based on the assumption that the gas is a good tracer of the gravitational potential of the galaxies. Some authors have suggested that magnetic pinching forces could alter signi cantly the rotation curves of spiral galaxies. In contrast to other studies which have concentrated in the vertical structure of the disk, here we focus on the problem of magnetic con nem ent in the radial direction to bound the magnetic e ects on the H i rotation curves. It is shown that azim uthalm agnetic elds can hardly speed up the H i disk of galaxies as a whole. In fact, based on virial constraints we show that the contribution of galactic magnetic elds to the rotation curves cannot be larger than 10 km s<sup>1</sup> at the outerm ost point of H i detection, if the galaxies did not contain dark matter at all, and up to 20 km s<sup>1</sup> in the conventional dark halo scenario. The procedure to estimate the maximum e ect of magnetic elds is general and applicable to any particular galaxy disk. The inclusion of the surface term s, nam ely the intergalactic (therm al, magnetic or ram) pressure, does not change our conclusions. O ther problem s related with the magnetic alternative to dark halos are highlighted. The relevance of magnetic elds in the cuspy problem of dark halos is also discussed.

Subject headings: galaxies: halos | galaxies: kinem atics and dynam ics | galaxies: m agnetic elds | galaxies: spiral

## 1. introduction

The rotation curves of spiral galaxies, especially dwarf galaxies and low surface brightness galaxies, are used to derive the proles of the dark halos which are very useful as tests for the standard cold dark matter scenario (e.g., de B lok & Bosm a 2002). The usual assumption is that the dynam ics of the neutral atom ic gas is a good tracer of the radial gravitational force. It can be shown that although spiral waves and non-circular motions may contribute to produce substructure in the rotation curves (e.g., de B lok, Bosm a & M cG augh 2003), they do not alter the rotation curves signi cantly for the galaxies selected in these investigations. G alactic magnetic elds could also a ect the gas dynam ics in spiral galaxies (P iddington 1964; P eratt 1986; B attaner et al. 1992; Sanchez-Salcedo 1997; B enjam in 2000; Beck 2002, 2003).

The random sm all-scale component of the galactic m agnetic elds acts as a pressure, giving support to the disk and, therefore, leading to a rotation (slightly) slower than the gravitational circular speed. By contrast, some congurations of the large-scale magnetic eld can give rise to a faster circular velocity. N elson (1988) and Battaner et al. (1992) proposed the so-called \m agnetic alternative" to dark matter in which the contribution of the m agnetic pinch could be large enough to explain the observed rotation curves of spiral galaxies without the necessity of dark matter. This is an interesting possibility because some universal properties such as the disk-halo conspiracy or the H i-dark matter constant relation, rst noticed by Bosm a (1978), would have a natural explanation in this scenario (Sanchez-Salcedo 1996a), as well as the truncation of stellar disks (B attaner, F lorido, Jim enez-V icente 2002). However, after the publication of the paper by B attaner et al. (1992), a plethora of possible di culties faced by the model was pointed out by di erent authors (Cuddeford & B inney 1993; Jokipii & Levy 1993; Persic & Salucci 1993; Vallee 1994; K atz 1994; P fenniger, C om bes, & M artinet 1994; Sanchez-Salcedo 1996b, 1997). In an extensive review, B attaner & F lorido (2000) argue that all the shortcom ings can be overcom e and, therefore, m agnetism can explain the way in which galaxies rotate, w ithout the help of dark m atter.

A certain MHD con quration has the ability of producing a faster rotation of the whole plasm a in the system, i.e. magnetic con nement, when the net contribution of the magnetic elds to the virial theorem is negative (in the conventional notation). In an isolated system, it is well known that the magnetic eld contribution to the virial theorem, applied to a large enough volume, is positive or zero, but never negative, re ecting the net expansive tendency of magnetic elds (e.g., Shafranov 1966). Jokipii & Levy (1993) used this argum ent to conclude that in the presence of magnetic elds, galaxies would need even more dark matter than in the unmagnetized case. One could argue that, since the virial theorem uses global (volum eintegrated) variables, it is possible that magnetism has a \centrifugal" action in the vertical direction but centripetal in the radial direction (radial con nem ent) and,

therefore, the expansive tendency of magnetic elds manifests only in the vertical direction. We advance that this claim is very wrong. An alternative way out is to assume that galaxies are not at isolation but embedded in the ubiquitous (magnetized) intergalactic medium which would be responsible for the required radial con nem ent.

M otivated by these ideas and given the importance of a correct interpretation of the rotation curves for C osm obgy, we will try to put bounds on the e ects of magnetic elds on the H i rotation curves. How ever, we do not restrict ourselves to discuss the viability of explaining rotation curves without dark halos; the orbits of G alactic globular clusters, the satellite galaxies and possibly the dynam ics of rem ote stars in the outer G alactic halo, which are not a ected by magnetic elds, strongly argue in favor of a dark halo around the M ilky W ay (e.g. Freem an 1997). The main issue is to see whether galactic magnetic elds can alter appreciably the H i rotation curves, especially in external galaxies for which the dark matter content is alm ost entirely derived from the rotation curve of the gas. Only if the answer is negative, we can be con dent that the pro-

les of dark halos, the dark matter content in galaxies as well as other interesting empirical correlations found between, for instance, halo parameters and lum inosity are reliable.

O ur investigation will be focus on the radial equilibrium con guration of magnetized rotating disks. For reasons that should become clear later, the analysis of the radial equation of motion puts stringent bounds on the maximum magnetic e ects to the rotation curves; this being free of other complicating assumptions that appear in studies of the vertical equilibrium con guration such as the external pressure, the vertical dependence of the magnetic eld within the height of the H i thin disk, or the attening of the dark halo, which are all very uncertain.

The paper is organized as follows. In  $x^2$  we describe the ingredients of a simplified model in which magnetic fields could give rise to a faster rotation in the outer parts of H i disks. In x3 the maximum elect on the rotation curves is studied. Other different to the magnetic alternative are discussed in x4. The case of magnetic fields producing a slower rotation and their role as a possible remedy to the cuspy problem of dark halos is left to x5 and the nal remarks are given in x6.

# 2. ingredients of the magnetic model and basic theory

In this Section, we describe some basic concepts useful for understanding the remainder of the paper. The issue of the problem of radial con nem ent will be addressed in x3.

## 2.1. The One-dimensionalCase

In presenting the role of the magnetic tension in the H i rotation curves, the equations are simplied by assum - ing an equilibrium, steady-state, axisymmetric disk of a slightly ionized gas with a purely azim uthal large-scale magnetic eld B (e.g., Battaner et al. 1992; Sanchez-Saloedo 1997). The radial component of the equation of motion of the gas at z = 0 in cylindrical coordinates reads

$$v^{2}(R) = R \frac{d}{dR} + \frac{R}{dR} \frac{dP}{dR} + \frac{1}{8} \frac{d(R^{2}B^{2})}{dR};$$
 (1)

where (R) is the gas density at z = 0, v (R) the circular speed of the gas, (R) the gravitational potential created by all masses in the system, and P (R) is the gas pressure consisting of the thermal, turbulent, convective, and cosm ic ray components plus also the magnetic pressure arising from the random magnetic eld component, but does not include the magnetic pressure arising from the large-scale magnetic eld. All these variables should be taken at them idplane of the disk. In equation (1) it was assumed v  $v_{\rm R}$ ;  $v_z$ , and thus we kept only the dom inant term s associated with v . In terms of the total pressure  $P_{\rm tot}$  (R) = P (R) + B $^2$  (R)=8 , it can be rewritten as:

$$v^{2}(R) = R \frac{d}{dR} + \frac{R}{2} \frac{dP_{tot}}{dR} + \frac{B^{2}}{4}$$
: (2)

The second term of the RHS is the radial pressure force that acts inwards or outwards depending on whether  $P_{tot}$  increases outwards or inwards. The third term is the magnetic tensile force which alw ays in poses a net inward force.

We should rem ind that ordered galactic-scale m agnetic elds are not expected to be a general feature in all disk galaxies, especially in dwarf galaxies. A lthough observations of m agnetic elds in these galaxies are scarce, they do not indicate m agnetic elds with a global sym m etry, but a m uch m ore chaotic galactic-scale eld than in norm al spirals (e.g., Chyzy et al. 2000). Therefore, m odels based on a regular azim uthal m agnetic eld m ay not be an adequate representation for allofthem. In fact, indirect observations suggest that regular m agnetic elds m ay be ignorable, at least in som e dwarf galaxies. The dom inance of spherical holes in the H i of the nearby dwarf galaxy IC 2574 has lead W alter & Brinks (1999) to conclude that large-scalem agnetic elds are dynam ically unim portant in the outer parts of this galaxy.

As it is shown in Appendix A, P (R) is always a decreasing function of R in approximately-isothermal, exponential galactic disks, even if the disk were embedded in an ambient medium with high pressure. Therefore, the asymmetric drift always produces a force pointing outwards and hence it can never contribute to enhance the circular speed of the gas. Moreover, the asymmetric drift is almost negligible in rotating galactic disks except in very low-mass galaxies such as DDO 50 (Carignan, Beaulieu & Freeman 1990; de Blok & Bosma 2002; Bureau & Carignan 2002). Thus we will also ignore it, hence dP<sub>tot</sub>=dR (1=8)dB<sup>2</sup>=dR, and the force balance equation is simplied to

$$v^{2}(R) = v_{c}^{2}(R) + \frac{1}{8 R} \frac{d(R^{2}B^{2})}{dR};$$
 (3)

where  $v_c^2$  Rd =dR is the gravitational circular velocity. Note that the magnetic pressure arising from the largescale eld is kept in our analysis because it cannot be neglected in general. The role of the magnetic pressure term associated with the random (small-scale) magnetic eld component is discussed in detail in x5. It should be also clear that the vertical gradient of P is not negligible when considering the vertical con guration of the disk, since it is the main agent responsible for the vertical support against gravity.

From Eq. (3) we see that for a purely azim uthal magnetic eld decaying radially not faster than R  $^{1}$ , the magnetic tension force produces radial con nem ent. Such conguration is usually called a linear or Z-pinch. The resulting inward magnetic force, the pinch, accounts for the required centripetal force responsible for an anom alous faster rotation. Our aim is to constrain the maximum e ect of the Z-pinch in galactic disks from very general grounds. It is worth mentioning that the Z-pinch con guration is not used in tokam aks as interchange and other instabilities quickly destroy the magnetic con guration (e.g., Roberts 1967). Restricted to purely azim uthal elds, the force-free magnetic con guration occurs for a eld pro le  $B = B_0 (R_0 = R)^2$ , where  $B_0$  is the magnetic strength at som e radius  $R_0$ . For that pro le, the outward magnetic pressure force is in balance with the inward tensile force at any radius. This pro le is, how ever, rather unphysical because it is singular at R = 0.

O bservations of galactic magnetic elds suggest that the radial component of the eld,  $B_R$ , is comparable to B at least in the optical disk and hence neglecting  $B_R$  may not be a good approximation. One could argue, however, that the magnetic pitch angle becomes very small in the outermost parts of galactic disks, where measurements of the eld orientation are scarce. Since  $B_R$  is not well-suited to produce radial con nement, the most optimistic and simplest situation, in order to maximize the magnetic effects on the rotation curves, occurs when the eld is taken as purely azimuthal. As we are interested in putting upper limits on the role of magnetic elds we will assume throughout this paper that  $B_R = 0$  in the outer disk to give the model asmuch leew ay as possible, as it is assumed in the magnetic alternative.

An important contribution to P is the turbulent pressure. For a certain component of the interstellar medium, the averaged turbulent pressure is given by  $v_t^2 = 2$ , where  $v_t$ is the one-dimensional turbulent velocity dispersion. For 6{10 km s<sup>1</sup> (e.g., Boulares & the H i components,  $v_t$ Cox 1990) and is nearly constant with galactocentric radius and also from galaxy to galaxy (Sellwood & Balbus 1999, and references therein). The magnetic eld is said to be in equipartition with the turbulent motions when the gas-to-magnetic eld pressure ratio is 1, or equivalently, when the turbulent velocity dispersion coincides with the Alfven speed. In the literature, one often nds the statem ent that dynam o action in a thin disk saturates at equipartition eld strengths (e.g., Ruzmaikin, Sokolo & Shukurov 1988; Shu 1992; Poezd, Shukurov & Sokolo 1993; Beck et al. 1996; Kulsrud 1999). Therefore, it is interesting to estimate those values of v ve for which the large-scale magnetic eld would be in superequipartition; this is done in the next. Denoting by  $v_A$  the A liven velocity associated with the regular eld, we can write  $v^2 \quad v_{L}^2$  (8 R )  $^1d(R^2B^2)=dR = v_{A}^2 + (R=8) dB^2=dR$ . In the likely case that the strength of the large-scale m agnetic eld does not increase with galactocentric radius it im plies v<sup>2</sup>  $v_{\!\scriptscriptstyle C}^2$  .  $v_{\!\scriptscriptstyle A}^2$  . Hence, for changes in the rotation velocity of  $v^2 - v_c^2 > v_t^2$ , the regular magnetic eld should be above equipartition with the turbulent motions. The above condition is equivalent to  $v > v_t^2 + v_c^2$  or, in

termsofv v:

v 
$$W > (v_t^2 + v_c^2)^{1=2}$$
  $W = U + \frac{1}{2} \frac{v_t^2}{v_c^2}$   $W = \frac{1}{2} \frac{v_t^2}{v_c}$  (4)

For  $v_c$  100 km s<sup>1</sup>, variations of v  $v_c$  & 0.5 km s<sup>1</sup> already require magnetic elds over equipartition. Such a situation is not very appealing to most dynam o theorists. How ever, in order to present our results as generic as possible and since there are some observations that could suggest magnetic energy densities above the turbulent ones in the disk of galaxies (e.g., Beck 2002, 2003) and in the Galactic center (e.g., Moss & Shukurov 2001), we will not employ equipartition when discussing the magnetic e ects on the rotation curves.

So far, the usual way to proceed was to integrate Eq. (1) out to a certain radius in order to derive the required radial pro le for B (R) to explain the observed rotation curve v, given the volume density at z = 0 and the gravitational circular velocity (including a contribution of the dark halo as in Sanchez-Salcedo (1997) or only the lum inous com ponents as in Battaner et al. (1992)). The question that arises is whether these local solutions can be matched with boundary conditions at in nity or not. This issue is very delicate and it will be proven very useful to put bounds on the in uence of galactic magnetic elds in the gas dynam ics at large scales. Before that, how ever, it is illustrative to describe how galaxies can reach a con guration with a faster rotation in order to have a complete picture of the underlying physics. This is brie y described in the next subsection.

# 2.2. The Phase of Magnetic Contraction

The assumption that the magnetic eld is mainly azin uthal in m ediately suggests that it com m enced to grow after the galactic disk has formed. In the absence of magnetic tension, the initial galactic disk or protodisk should have a rotation speed given by the mass distribution  $v^2 = v_c^2 = Rd = dR$ . If the rotation curves are signi cantly a ected by the magnetic eld, one should expect that, as the magnetic eld is amplied no matter by what process, the disk initially in gravitational force balance, will experience a radial contraction caused by the growing magnetic tension, leading to a disk with a faster rotation due to angular m om entum conservation. W e will refer to this process as the magnetic contraction phase (MCP). If the mass loss by out ows is unimportant, the MCP is the only way capable of spinning up the disk because the total angular momentum stored in the elds (/ r [E B]dV) is negligible in both the initial protodisk and in the nalcon guration. O ther processes, such as turbulent viscosity, are very ine cient to produce the desired e ect.

The conservation of angularm on entum in plies that, for a galaxy sim ilar to the M ilky W ay with a dark halo and a rotation speed of 220 km s<sup>1</sup> at 30 kpc, an increase of 10 km s<sup>1</sup> in the velocity speed requires a radial contraction of 1.3 kpc at that radius. The contraction m ust be huge in the magnetic alternative; see x4.1.

3. galactic disks with faster rotation. the problem of the radial confinement

## 3.1. Isolated Galaxies

In the previous section it was pointed out that there is a pinching e ect only if B does not decrease faster than 1=R. However, this decay is untenable at large R in real astrophysical system s as it will be shown in the next and, therefore, it is inevitable that beyond a certain radius, the m agnetic tension produces a force pointing outwards (see Fig.1 for a visualization). In fact, the slowest radialm agnetic decay occurs in the ideal case of having strict cylindrical geom etry with currents,  $J_z$ , owing from z =1 to z = 1 (otherwise, if currents are conned to a certain region, B must decrease with radius at least as fast as a dipole eld for larger: B  $K = r^3$ , where K is a constant). Under strict cylindrical geom etry  $^1$ , the condition of closed current loops is equivalent to imposing  ${}^{\rm R}_{0} \, J_z R^{\,0} dR^{\,0}$ ! 0 at large R (hereafter, this will be our de nition of isolated system in a cylindrical system). From Ampere's law, this requirement implies that B < C = R at large R, being C a constant. So there must exist a certain radius denoted by R kep at which the magnetic eld starts to decay as 1=R or faster.  $R_{kep}$  is therefore the radius at which the rotation speed of the gas coincides with the local K eplerian velocity. If  $R_{HI}$  is the radius of the last point of H i detection, we require R  $_{\rm kep}$  > R  $_{\rm H~I}$  in order to have an anom alous faster rotation along the observed disk.

The system will be in a con guration of equilibrium at  $R > R_{kep}$  only if the inward gravitational force is higher than the outward magnetic force

$$R\frac{d}{dR} > \frac{1}{8 R} \frac{d(R^2B^2)}{dR}$$
(5)

If this condition is not ful led the disk will expand radially due to the magnetic forces. This condition can be written as

$$B (R) > \frac{1}{R} R_{kep}^{2} B^{2}; kep \qquad 8 \qquad R v_{c}^{2} dR ; (6)$$

where B  $_{;kep}$  is the magnetic eld strength at R  $_{kep}$ . Since a dependence B C = R at large R is unphysical (see above), an upper lim it to B  $_{;kep}$  can be established from Eq. (6):

$$B^{2}_{;kep} = \frac{8}{R^{2}_{kep}} \left[ \frac{R}{R_{kep}} \right]^{2}_{R_{kep}} (R;0) R v_{c}^{2}(R) dR;$$
(7)

where  $R_B$  is the radius where B 0. In practice we will take  $R_B$  ! 1 to estimate the maximum contribution. The maximum value for B ;kep corresponds to a situation in which there is balance at  $R > R_{kep}$  between gravitational and magnetic forces and thus the material outside the  $R_{kep}$  circle has no rotation at all.

The upper limit derived above has an easy interpretation. In principle, one could achieve very large di erences  $v = v_{\rm i}$  if one would make the density arbitrarily small than the A liven speed would be arbitrarily large. However, according to Eq. (5), this is not possible globally because the m agnetic eld has to be held in by the radial weight of the plasm a in the radial gravitational eld since the net e ect of magnetic elds is never con ning.

It is custom any and natural to discuss this kind of constraints using the virial theorem which says that the net nature of magnetic elds is expansive or null (but never con ning) in both the radial and vertical directions (Appendix B). More speci cally, for toroidal magnetic elds and strict cylindrical geom etry, the total rotational energy of an isolated system should be the same with or without magnetic elds, keeping the mass distribution xed (so that (R); see Appendix B for further details). By imposing that constraint, a similar upper limit for B<sub>kep</sub> is found (Appendix C), except by a factor ( $_{\rm B}$ ), between 1{2, that takes into account possible three-dimensional ects.

It is convenient to consider rst the extrem e and hypothetical case that spiral galaxies do not host dark m atter halos. For a galaxy with a mass in gas and stars M, (R) GM = R in the outer parts and

$$B_{;kep}^{2} B_{cr}^{2} 8_{B} \frac{GM}{R_{kep}^{2}} R_{R}(R;0) dR:$$
 (8)

The assumption that (R) GM = R is satisfactory for many high surface density galaxies (e.g., van Albada et al. 1985) but it is not necessarily fullled in some low surface brightness galaxies (LSB) and dwarfs for which the gas surface density is not exponential and its contribution to the potential may be comparable or even higher than that from the stellar component at radii  $R_{\rm H\,I}$  for certain values of the adopted stellarm ass-to-lum inosity ratio. How ever, we are con dent that our conclusions would not change for these galaxies even adopting the most unfavorable assumptions.

In the optim istic and rather unlikely situation that B were constant with R out to  $R_{kep}$ , i.e., B (R) =  $B_{cr}$  at R <  $R_{kep}$ , the characteristic variation  $v^2$   $\psi^2_{c}$  would be  $v^2 \qquad \psi^2_{c} \qquad B^2_{cr}=(4)$ . By using Eq. (8), it is possible to estimate the maximum relative variation of the circular speed of the gas at a radius R <  $R_{kep}$ :

$$\frac{v^2}{v_c^2} \frac{v_c^2}{R} = 2 \quad B \quad \frac{R}{R_{kep}} \quad \frac{R_g}{R_{kep}} \quad \exp \quad \frac{R}{R_g} \frac{R_{kep}}{R_g} ;$$
(9)

where we have adopted a radial exponential decay for (R;0) with scale  $R_g$  (e.g., B land-H awthorn, Freem an & Q uinn 1997). Note that  $(v^2 \qquad v_c^2) = v_c^2$  is not sensitive to  $P_{ext}$  because even though the density at the midplane increases with  $P_{ext}$ , equation (9) does not depend on the absolute value of the density but only on  $R_g$ . The main source of uncertainty stems from the value of  $R_{kep}$ , but the most optimistic situation occurs when  $R_{kep}$   $R_{H I}$ , so we will assume for the discussion that both radii are identical,

30 kpc for common spiralgalaxies. In order to estimate  $v^2 = v_c^2$  from Eq. (9), we will assume that the mass of gas in the disk beyond R<sub>HI</sub> and within  $\dot{z}_{\rm J} < h_1$ , being h<sub>1</sub> the characteristic scale height of the gas disk (see Appendix C for further details), is 0.5 percent the totalmass of the disk. This seems to be the case of dwarf galaxies (van den Bosch et al. 2001), but it is probably a too generous value for normal spiral galaxies. The latter assumption corresponds to a radial scale length for the gas, R<sub>g</sub>, of

5.5 kpc. Putting these values into Eq. (9), we obtain  $v^2 = v_c^2 = 0.36$  B at R = 30 kpc. If we take for

<sup>&</sup>lt;sup>1</sup> In such a situation, the m agnetic eld is always purely azim uthal, consistent with the assumptions given in x2 and, hence, the essential physics of the m echanism giving rise to greater azim uthal velocities of the H i disk, is entirely contained within the one-dimensional radial m odels.

reference values  $v_c = 100$  km s<sup>1</sup> at 30 kpc, the maximum change in the rotation velocity, v v v, ranges between 15 to 30 km s<sup>1</sup> for  $_B = 1$  to 2, respectively.

This value for v should be regarded as a generous estim ate for two reasons. First, the azim uthal eld is not expected to be approximately constant while the volume density of gas at z = 0 varies by a factor of 50{100 (Diplas & Savage 1991). Secondly, the adopted equilibrium con guration, namely v (R) = 0 at any R >  $R_{kep}$ , cannot be reached in practice, starting from a rotating disk in equilibrium with the gravitational force, due to the conservation of angularm om entum. In fact, the m agnetic eld begins to grow in the disk as discussed in x2.2, creating a magnetic force directed outward beyond  $R_{kep}$ . This region will undergo a continuous outward drift, while the inner disk ( $R < R_{kep}$ ) must be subjected to the required contraction phase, generating a gap in the surface density of the disk. In order to quantify this e ect, we assume for simplicity that the outer disk ( $R > R_{kep}$ ) su ers an approximately self-similar expansion with factor > 1, i.e.,  $(\mathbf{R}) = (_0 = ^2) \exp(\mathbf{R} = \mathbf{R}_g) \text{ at } \mathbf{R} > \mathbf{R}_{kep}, \text{ in the nal}$ equilibrium con guration. From the conservation of angularm om entum, it can be shown that the right-hand side of Eq.(9) is now a factor  $^{2}$ =( 1) sm aller (Appendix C). Hence, the optim ized situation occurs for = 2 and then  $v_c^2$ )= $v_c^2$  < 0.1 <sub>B</sub> at R = 30 kpc. This implies that  $(v^2)$ v < 10 km s<sup>1</sup> at the periphery of H i disks in norm al spiral galaxies under the assumption that they contained only stars and the observed gas. Therefore, it is not possible to banish dark halos by magnetic elds because one would require much larger values of v .

For disk galaxies with a dark halo, as postulated in the conventional scenario, it is possible to follow the same procedure to estimate the maximum e ect of magnetic elds when  $v_c = cte$ , instead of  $v_c^2 = GM = R$  (see Appendix C). Similar arguments suggest that the expected local, and probably transient, variations in the rotation speed larger than 20 km s<sup>1</sup> are very unlikely even in models including dark halos (see also x2.2 and Sanchez-Saloedo 1997).

In the next subsections, it will be shown that these upper limits for v also apply even when galaxies are not isolated system sbut are embedded in an ambient medium, thereby making these estimates very robust.

### 3.2. Non-Isolated Galaxies

It could be argued that the model described by Eq. (1) is too sim plistic because other term s not considered in that equation, as a radial ow or other components of the magnetic eld, might be important, regarding the radial connem ent, beyond  $R_{\rm H\,I}^2$ . In fact, unlike the radial component of the eld, the z-component m ay contribute to the radial con nem ent of the disk and will be included in our analysis. In addition, the hydrodynam ical pressure by a radial in ow due to intergalactic accretion of matter may also play a role. Since system s embedded in an ambient m edium usually present sharp boundaries, it is convenient to use an integral form of the equations as that given by the virial theorem .

The scalar virial theorem for elds with  $B_R = 0$  and strict cylindrical symmetry (i.e., 0=0 z = 0=0 = 0) applied to a nite cylindrical volume V, bounded by a cylin-

der S of height L, can be written as:  $\frac{7}{7}$ 

2T+2 P dV 2P<sub>S</sub>V+W+M <sub>cyl</sub> (v r) (v dS)
$$\frac{1}{2}$$
;  
v (10)

where T is the kinetic energy of the gas, W its gravitational energy in the gravitational eld created by all masses in the system, P the gas pressure as de ned in x2.1, P<sub>S</sub> the gas pressure at the boundary surface, r is a coordinate vector, I d<sup>2</sup>I=dt<sup>2</sup> with I the trace of the m om ent of inertia tensor and M <sub>cyl</sub> is given by

$$M_{cyl} = \frac{1}{4} \int_{V}^{L} B_{z}^{2} dV = B_{z;s}^{2} + B_{;s}^{2} V ; \quad (11)$$

(e.g., Fiege & Pudritz 2000 and Appendix B). Throughout this paper the subscript S refers to quantities valuated at the radial surface of our cylindrical volum e with radius  $R_s$ , and the subscript byl' stands for the cylindrical case. This form for the virial theorem is valid for any arbitrary helical eld ( $B_R = 0$ ) provided cylindrical geometry; no other additional assumption is required. The surface integral terms including the z = L=2 surfaces are taken into account in Eq. (11). The integration of the magnetic terms can be found in Appendix B.For any further details in the derivation, we refer the reader to the paper by Fiege & Pudritz (2000).

We can identify two new vias of radial con nem ent: by ram pressure [last term of LHS of Eq. (10)] or by an external magnetic eld. The ambient pressure ( $P_S$ ), on the other hand, only produces vertical con nem ent as discussed in Appendix A. If these new terms are able to produce radial con nem ent then the constraint given by Eq. (7) could be somehow relaxed. The possible role of these terms will be discussed in two separate sections.

#### 3.2.1. Ram Pressure

So far, we have neglected the terms involving  $v_R$ , because H iobservations show that  $v_R$  is very sm all compared to v for the galaxies under consideration. Beyond the edge of the observed H i disk, the value of  $v_R$  is uncertain. W e consider now the surface integral of the momentum stress in Eq. (10), which m ay have a negative sign for the case of strong m ass accretion into the disk. In fact, it can be shown that even though I is expected to be negative, the combination I=2  $_{S}$  (v r) (v dS) could be still negative, showing the con ning e ect of the ram pressure. (For example, in the one-dimensional case of a supersonic ow that slows down to rest by crashing perpendicularly to a rigid plane wall and forms a strong adiabatic shock, I=2  $_{R}$  (v r) (v dS) = (2=3) (v r) (v dS).)

I=2  $_{\rm S}$  (v r) (v dS) = (2=3) (v r) (v dS).) Since spherical accretion cannot con ne a disk, the inow should be extremely collim ated parallel to the galactic plane, i.e.  $v_{\rm R}$   $v_{\rm z}$ ; v, for which there is no evidence. Moreover, this collim ation is very unlikely to occur if the disk forms from a sphere of gas. G as cools and parcels of gas acquire circular orbits, conserving approximately angularm omentum, when they collapse to settle into a disk, leading to  $v_{\rm R}$  ! 0 (see R ognvaldsson 1999 for results based on numerical simulations). Furtherm ore, in the hypothetical case that such in ow were possible by som e unknown m echanism, the accretion speed should be so high that an

5

 $^2$  W ithin the R<sub>HI</sub> circle, observations suggest that other term s are unim portant for those galaxies under consideration.

accretion shock would be unavoidable, transform ing ram pressure into therm alpressure and probably generating a hydraulic jum p in the edge of the disk and other observable features such as H em ission. The hydraulic jum p will divert the ow around the disk decreasing dram atically its con nem ent ability.

# 322. Con nement by the Intergalactic Magnetic Field

A coherent external (intergalactic) m agnetic eld could in principle contribute to the con nem ent under certain circunstances. It is important to distinguish here between the intracluster and the wider intergalactic eld. As we are dealing with galaxies in the eld, we are interested in the amplitude of the latter one. At present, setting observational constraints to the strength and topology of the intergalactic magnetic elds is a challeging task (see Ryu, Kang & Bierm ann 1988; Kronberg et al. 2001; and Widrow 2002 for a review) and, therefore, they are very uncertain. C oherent m agnetic elds on scales m uch greater than the typical sizes of galaxies could be as large as 0:1 G in the optim istic situation that extra ampli cation through batteries, dynam os or other routes of m agnetization w ere operative.

W e assume the ideal situation that initially the pregalactic magnetic eld is uniform and lls the protogalaxy making a nite angle with the rotation axis of the galaxy. A s the disk is forming by accretion of gas, the magnetic eld lines are advected with the ow. It is well-known in studies of the collapse of interstellar clouds that, under ux-freezing conditions, this compression produces that the total magnetic contribution to the virial theorem is positive (see, e.g., Chapt. 24 of Shu 1992 and Chapt. 11 of Mestel 1999) and, therefore, the con nem ent by the external eld is impeded. In order to have con nem ent, the disk must be able to get rid of the magnetic ux.

In the strict cylindrical sym m etry, the external m agnetic eld must be vertical and then Eq. (11) with B  $_{\rm S}$  = 0 tells us that magnetic con nement is only possible if the volum e-averaged vertical com ponent of the magnetic eld within V is strictly smaller than the magnetic eld at the boundary. Since  $v_R$  is directed inward and  $@v_R = @R < 0$ in the MCP,  $\beta_z$  j increases at every point of the disk due to the inward advection (see also Reyes-Ruiz & Stepinski1996; Moss & Shukurov 2001); under those conditions con nement is not viable. This problem could be alleviated if magnetic eld lines were expelled from the disk by am bipolar and turbulent di usion. However, a thorough num erical study by Howard & Kulsrud (1997) shows that  $\beta_z$  j is also increased by ambipolar shearing terms to a value considerably larger than its initial value. In the lack of a mechanism able to reduce  $\beta_z$  jw ithin disks, pregalactic magnetic elds do not help to con ne galactic disks. This example clearly illustrates why self-con nem ent is so rare in nature.

M odels based on the con nem ent by anchored elds to the galactic nucleus will present a similar problem, as  $\beta_z j$  is expected to decline outwards.

# 4. the magnetic alternative

Battaner & Florido (1995, 2000) and Battaner, Florido & Jim enez-Vicente (2002) pointed out that: (1) the magnetic alternative to dark halos does not su er from the problem of excessive aring of the H i disk and (2) the existence of truncated stellar disks is explained satisfactorily in the magnetic alternative. Putting aside the problem of radial con nement discussed in the previous sections, it is stimulating to see whether alternative scenarios to conventional ideas have the potential of explaining other phenomena in a natural way. If this occurs, alternative m odels could attrack much more attention. For instance, som e empirical correlations of galactic warps have led to C astro-Rodr guez et al. (2002) to suggest tentatively that outer rotation curves could not be caused by the presence of a massive halo. In the next sections we will show that the aring problem remains unsolved and that the existence of truncated stellar disks cannot be considered a prediction of the magnetic alternative.

# 4.1. The MCP and Implications for the Truncation of Stellar D isks

If galaxies contained only stars and the observed gas, the initial protodisk of mass M<sub>i</sub> should follow approximately a Keplerian rotation,  $v^2 = GM_i = R$ , at large R, and evolves to the present-day exponential disk of m ass M f and radial scale length  $R_d$ , due to magnetic contraction. The peak velocity of the naldisk is  $v_{max} = 0.62^{\circ} \text{ GM}_{f} = R_{d}$  (e.g., Binney & Trem aine 1987). For sim plicity, we assume that beyond 2:27R<sub>d</sub> the rotation curve of a present galaxy is at with circular velocity  $v_{max}$ . During the MCP, a ring of gas bcated initially at a radius  $R_{ip}$  <u>must squeeze</u> up to a nal radius  $R_{f}$ , such as  $v_{max}R_{f} = GM_{i}R_{i}$  from angular momentum conservation. Therefore, the initial radius is given by the simple expression  $R_i = 0.38 (M_f = M_i) (R_f = R_d) R_f$ . For a norm al spiral galaxy like the Milky Way, with a nearly at rotation curve up to the last measured point, typically at 30 kpc, and with  $R_d$ 4 kpc, the MCP should be able to squeeze the disk at least from 85 kpc 30 kpc in radius, provided the mass loss in out ows to is negligible, corresponding to a typical radial velocity of 3 km s<sup>1</sup> for one Hubble time. A coording to x3, this contraction is an unfeasible process due to the global expansive nature of magnetic elds.

The level of contraction m ay be signi cantly reduced if a substantial fraction of the m ass is lost in winds following early epoch of star bursts, i.e. M<sub>f</sub> < M<sub>i</sub>. Recently, Zhao (2002) has estimated that less than 4% of the total mass of the G alaxy has been lost during its lifetime. Even assuming an amount of mass loss 3 times larger than this value, the magnitude of the contraction of the disk is still huge, from 72 kpc to 30 kpc, again for  $R_d = 4$  kpc.

The MCP represents a serious threat to the suggestion by Battaner, Florido, Jin enez-Vicente (2002) that, since stars do not feel the galactic magnetic eld, the escape of stars form ed beyond a certain galactocentric radius could explain the observed truncation of stellar disks. The calculations by Battaner et al. (2002) are rooted in a scenario that does not consider the MCP, with the gas disks being in anom alous rotation from the very beginning. If one tries to include the MCP in the model, the stellar distribution function depends on di erent assumptions that concern the details of the MCP, such as the radial velocity of the contraction, or the relationship between star form ation rate, surface density and magnetic eld strength. C onsequently, the escape radius depends on time. If the disk was contracted from 80 to 30 kpc in radius, it could be even more natural to construct a model in which the stellar surface density decays with R slowly enough, with no stellar truncation radius.

Even assuming that stars in the outer disk were not formed until the gas disk stopped its contraction, which is in contrast with observations that show old stars in the outer disk (e.g., Ferguson & Johnson 2001), the planar velocity dispersions of the stars should increase dram atically with R and this is not observed (e.g., Pfenniger et al. 1994; Vallee 1994).

### 4.2. The Flaring Problem

In the past years, much of the challenge involved in m aking the magnetic alternative viable concerned the excessive aring of the disk, which is caused by the vertical magnetic pressure (Cuddeford & Binney 1993). In a twodimensional analysis, Battaner & Florido (1995) predict a scale height of 3{35 kpc at a radius of 15 kpc and of 7 kpc at 30 kpc for the H i disk of M 31. These values are still too large by a factor of 3{5 if com paring with the observed H i scale height observed in the Galaxy by D iplas & Savage (1991) and Burton (1992).

This discrepancy in the scale height is the least of our worries. An inspection of the radial prole of the magnetic strength (g. 3 of Battaner & Florido 1995) reveals that there is something deeply wrong in their calculations. As shown in x2.1, the magnetic eld cannot decrease faster than 1=R in order to have a disk with a faster rotation. Therefore, a magnetic eld of 5 G at 15 kpc implies a eld strength larger than 2.5 G at 30 kpc. The magnetic eld estim ated by Battaner & Florido (1995, 2000) decays faster than 1=R.

If the dark matter contribution to the rotation curve is null, the required magnetic elds in the outer parts of typical spiral galaxies are so intense ( 5{10 G) that the problem of excessive aring is unresolved. The aring problem is exacerbated when the magnetic pressure of the small-scale magnetic eld is included because it also contributes to the vertical expansion of the disk. This pressure is observed to be comparable to the magnetic pressure of the regular eld at least in the optical disk of the Galaxy. On the other hand, the excessive aring is also aggravated if the magnetic pitch angle is not zero. We conclude that the aring problem by itself is su cient to reject the magnetic alternative.

# 5. galactic disks with slower rotation: the cuspy halo problem

In this Section, we will discuss the role of magnetic elds as a remedy for the cuspy problem of dark halos. It is interesting to see whether the magnetic pressure of the random component, as well as certain con gurations of a global-scale magnetic eld (not necessarily azim uthal) could help to explain the discrepancy between the proles of dark halos obtained in cold dark matter (CDM) simulations, which becomes very cuspy towards the center (e.g., Navarro, Frenk & White 1996, 1997), and observations of the halos of dwarf galaxies and LSB, which are better tted with a core-dom inated halo (see de Blok, Bosma & M cG augh 2003, and references therein). O ne possibility is that galaxies are indeed embedded in cuspy halos follow – ing the prole found by Navarro, Frenk and W hite (1996; hereafter NFW) but the combination of asymmetric drifts and m agnetic elects wipe out completely the inner cusp, leading to rotation curves that m in ic a halo with a core.

The magnetic pressure of the random component gives some support to the disk, reducing the rotation speed. Since it is expected to be in equipartition with the gas turbulent pressure, corrections due to the magnetic pressure should be of the order of the asymmetric driff, 2{3 km s<sup>1</sup> (e.g., de B lok & Bosm a 2002), which are smaller or comparable to observational uncertainties, typically 4{6 km s<sup>1</sup>. This contribution, how ever, could be larger in a certain radial portion of the disk if the magnetic pressure decays faster than  $v_t$ . W ritting the A liven speed associated to the random eld as  $v_A^2 = g(R)v_t^2$ , with the turbulent velocity dispersion  $v_t$  constant with R, and dg=dR < 0, the rotation curve changes according to the relation:

$$v^{2} = v_{c}^{2} + g(R) \frac{v_{t}^{2}}{2} \frac{d\ln}{d\ln R} + \frac{d\ln g}{d\ln R} :$$
 (12)

Following de Blok & Bosm a (2002), we de ne R in as the radius of the innerm ost sam pled point of the rotation curve, and assume that at  $R > R_{in}$ ,  $g(R) = (R_{in}=R)^n$  for sim plicity. Doing so, we are imposing that at  $R_{in}$ ,  $v_A$ ¥. The second term of the RHS of Eq. (12) is of the order of the asymmetric drift (comparable to observational uncertainties), whereas the last term is  $n\sqrt{2}R_{in}^{n}=2R^{n}$ . As the observed rotation curve diers from the NFW prole along a signi cant range of galactocentric radius, say at R R <sub>in</sub> 500 pc, then 0 < n 1, im plying that  $nv_t^2 R_{in}^n = 2R^n < v_t^2 = 2$  50 km<sup>2</sup> s<sup>2</sup> (for  $v_t = 10$  km s<sup>1</sup>). Consequently, this e ect would be only signi cant very close to the galactic center where v<sub>c</sub> 10 km s<sup>1</sup>, a region with large observational uncertainties. Sum ming up, the contribution from the total pressure to the circular velocity is less than  $5\{6 \text{ km s}^1$ . This result agrees with the fact that CO, H and H i rotation curves reveal the same kinem atics for the central regions of some galaxies (e.g., Bolatto et al. 2002), which suggests that m agnetic elds do not contribute signi cantly to the support of the gas disks. In particular, the CO rotation curves are ex-

gas disks. In particular, the CO rotation curves are expected to trace the motion of the highest density regions, including molecular clouds, which are thought to be hardly a ected by the galactic magnetic elds.

In conclusion, magnetic e ects are not su cient to solve the cuspy halo problem satisfactorily.

# 6. final remarks

It has been recognized for decades that the galacticm agnetic eld provides signi cant vertical support to the interstellarm edium. The role of the magnetic tension in the vertical hydrostatic con guration was considered by C ox (1988) and Boulares & C ox (1990), whereas the role of the tension in the plane of the disk was studied by Nelson (1988), Battaner et al. (1992) and Benjam in (2000) [see also P iddington 1964]. The idea that large-scale m agnetic elds can explain the rotation curves of spirals without invoking dark matter has been revived by Battaner & F lorido (2000) and Battaner et al. (2002).

All the previous work aim ed at giving bounds to m agnetice ects on gas dynam ics are based on the aring of the disk by the vertical magnetic gradients. Here we propose that the radial analysis provides stringent constraints, being free of other com plicated assumptions such as the unknown external pressure. We have shown that magnetic

elds have a net expansive (or null, but never con ning) effect not only in the vertical direction but also in the radial direction. W ithout dark m atter halos, the contribution of the magnetic eld to the rotation curves at the periphery of spiral galaxies is not expected to be larger than 10 km s<sup>1</sup>, which is insu cient to banish dark halos by magnetic elds. This result is valid even though an external pressure, intergalactic magnetic elds or ram pressure are taken into account. In addition, the huge contraction of the disk required in the magnetic alternative would spoila simple magnetic explanation for the observed truncation of stellar disks. Moreover, the problem of excessive aring in the outer disk by the magnetic pressure is unresolved and may be aggravated if the random component of the magnetic eld is included, or if the eld is not purely toroidal. By contrast, there is observational evidence that the dwarf, dark-m atter dom inated galaxy IC 2574 does not possess a dynam ically-in portant galactic-scale m agnetic eld. O ur rst conclusion is that the magnetic alternative is not viable.

Even though there is no doubt about the existence of dark matter halos around galaxies, it is desirable to constrain the magnetic e ects on the H i rotation curves for a correct interpretation of the observations. It is suggested that magnetic elds can contribute to produce wiggles in the H i rotation curves, which could be likely associated with non-axisymmetric waves in the disk, as may be the case of NGC 1560. However, the magnetic tension is unlikely to produce substructure in the rotation curve ofm assive spiral galaxies above 20 km s<sup>1</sup>. In Sanchez-Salcedo (1997), the detailed shape of the rotation curve of this dwarfgalaxy was tted reasonably well combining a magnetic eld of strength 1 G plus an isotherm al dark halo. Finally, we have considered magnetic elds as a possible solution to the cuspy problem of dark halos. We conclude that they are insu cient to explain the discrepancy between the observations and the predictions of CDM sim ulations.

We thank J. Canto and A. Hidalgo for discussions, and the anonym ous referees and the Editor for critical comments. FJSS.acknowledges nancial support from CONACYT project 2002-C 40366.

### APPENDIX

# a. why an external pressure is unimportant to study the radial configuration

Consider a three-dimensional axisymmetric gas disk with surface density (R) supported mainly by rotation and embedded in an ambient medium of pressure  $P_{ext}$ . The transition between the gas disk and the external medium occurs at a vertical height  $z_s$  (R). From the vertical hydrostatic equilibrium, the radial gradient of the total (turbulent plus magnetic) gas pressure  $P_{tot}$  at the midplane of the disk is given by

$$\frac{dP_{tot}(R;0)}{dR} = \frac{d}{dR} P_{ext} + \begin{pmatrix} Z & Z_{S}(R) \\ 0 & \frac{\theta}{\theta z} dz \end{pmatrix} = \frac{d}{dR} \begin{pmatrix} Z & Z_{S}(R) \\ 0 & \frac{\theta}{\theta z} dz \end{pmatrix}$$
(A1)

where (R;z) is the gravitational potential. The pressure exerted by the ambient medium squeezes the disk along z, increasing  $P_{tot}(R;0)$  but barely  $dP_{tot}(R;0)=dR$ , and thereby making the radial con nem ent extrem ely di cult. This can be visualized in the following manner. Ignoring for a moment magnetic elds, an external pressure will produce some radial con nem ent if it is possible to reach a con guration with QP = QR and  $P_{rad} > 0$  just by increasing  $P_{ext}$ :

$$\frac{d}{dR} \int_{0}^{Z_{z_{s}}(R)} \frac{\theta}{\theta z} dz > 0:$$
 (A2)

For a gas density decaying along z with a scale height h, for instance  $/ \exp(\frac{2}{2}=2h^2)$  (for  $j_2j < z_s$ ), Eq. (A2) is satis ed if the scale height increases with galactocentric radius at least as fast as h  $z_s = R^1 - 1$ , where l is an exponent that varies from 2{3 for a gravitational potential with a at and a K eplerian circular speed, respectively. However, this requirem ent cannot be fulled in approximately isotherm algalactic disks with the surface density decreasing outwards, because the maximum variation of h occurs in the limit of large external pressure and goes as h  $R^1$ .

There are other physical di culties in the boundaries of the disks if they were con ned by P<sub>ext</sub>. From dimensional grounds, it is expected that the gas pressure P (R;0), comprising m ainly therm aland turbulent contributions as de ned in x2.1, should be greater than the external pressure, as a consequence of turbulent m otions and m ixing by R ayleigh-T aylor type instabilities in the disk. The total pressure at z = 0 is

$$P_{tot}(R;0) = P(R;0) + \frac{B^2(R;0)}{8} > P_{ext} + \frac{B^2(R;0)}{8}:$$
 (A3)

In the periphery of the disk, pressure con nem ent requires  $P_{tot}(R;0) = P_{ext}$ , thus Eq. (A 3) in plies B<sup>2</sup> (R;0)=8  $P_{ext} < P$  (R;0), i.e. well below equipartition and, therefore, they are not su cient to a ect appreciably the rotation curves, according to our discussion in x2.1.

## b. derivation of the magnetic terms in the virial theorem

Here we discuss the magnetic terms that appears in the virial theorem. The same derivation can be found in Fiege & Pudritz (2000). Consider a distribution of a electrically conducting plasma with velocity v in a gravitational eld (R)

9

created by all masses in the system. The scalar virial theorem applied to a nite volume V, bounded by a surface S, can be written as:

$$2T + U + M_{cyl} + W$$
 (v r) (v dS)  $\frac{1}{2} \frac{d^2 I}{dt^2}$ ; (B1)

where T is the kinetic energy of the plasm a within V, W the gravitational term given by  $W = \frac{1}{v}$  r r dV and

т

$$U = 3 P dV Pr dS;$$
(B2)

$$M = \frac{1}{8} \int_{V}^{2} B^{2} dV + \frac{1}{4} \int_{S}^{2} (r B) (B dS) \frac{1}{8} \int_{S}^{2} B^{2} r dS;$$
(B3)

$$= r^2 dV; (B 4)$$

where r is a coordinate vector (e.g., Shu 1992).

In the case of having cylindrical symmetry (i.e., @=@z = @=@ = 0) and for elds with no radial component,  $B_R = 0$ , a more simple expression for  $M_{cyl}$ , which denotes  $M_{under}$  cylindrical symmetry, can be derived when the virial theorem is applied to a nite cylindrical volume; this is done in the following. Since  $B_R = 0$ , the second term in the RHS of Eq. (B3) contributes only at z = L=2 surfaces, where L is the vertical length of the volume V. The component of the eld perpendicular to these surfaces is  $B_z$ , so that

$$\frac{1}{4} \int_{S}^{Z} (r B) (B dS) \frac{L}{4} \int_{0}^{Z} 2 R B_{z}^{2} dR = \frac{1}{4} \int_{V}^{Z} B_{z}^{2} dV:$$
(B5)

The last term of Eq. (B3) comprises an integral along the vertical surface of the cylinder  $I_1$  and a contribution of the cap surfaces,  $I_2$ , given by

$$I_{1} = \frac{1}{4} (B_{;s}^{2} + B_{z;s}^{2})V; \qquad (B6)$$

and

$$I_{2} = \frac{L}{8} \int_{0}^{Z_{R_{s}}} 2 R B^{2} dR = \frac{1}{8} \int_{V}^{Z} B^{2} dV :$$
 (B7)

C on bining the three contributions the nalexpression for M  $_{\rm cyl}$  reads as

7

$$M_{cyl} = \frac{1}{4} \int_{R}^{2} dV = B_{z;s}^{2} + B_{;s}^{2} V :$$
(B8)

Similar manipulations allow us to rewrite U as  $U = 2_v P dV = 2P_S V$ .

For a cold galactic disk, the therm all energy is negligible compared to the kinetic energy. If, in addition, there is no radial accretion ows and the disk can be assumed to be in a steady state, the virial theorem is reduced to

$$2T + M_{cvl} + W = 0$$
: (B9)

A ssum ing that galaxies are isolated system s,  $B_{z,S}V$  and  $B_{,S}V$  go to zero at large R in plying that always  $M_{cyl} = 0$ , re ecting the in possibility of magnetic connement in the radial direction. In fact, for a given distribution of mass, W is xed and if  $M_{cyl}$  is positive, the total kinetic energy should be smaller than in the unmagnetized case in order to satisfy the virial constraint. This does not exclude the possibility that some parts of the disk could present a faster rotation (larger kinetic energy) as long as rotation is slower than that given by gravity in other parts of the disk. It might be that we are observing just the part of faster rotation and the material with a slower rotation lies outside the last point of H i detection, if it is fully ionized by the metagalactic UV background (see Fig. 1 for a visualization). For the case of purely azim uthal elds  $M_{cyl} = 0$ , so 2T + W = 0. Therefore, given a distribution of mass, all the congurations in equilibrium must have the same T regardless of magnetic elds. Note that  $M_{cyl} = 0$  does not imply necessarily that the magnetic con guration is force-free. In the particular case of a force-free magnetic con guration, it can be shown that  $M_{cyl} = 0$  at any arbitrary volum e and hence magnetic elds are neither expansive nor con ning.

## c. constraining the azimuthal magnetic field strength in isolated galaxies

Consider an isolated disk galaxy comprising of gas and stars with a given total surface density (R). The gas is moving in pure circular orbits (i.e.  $v_z = v_R = 0$ ) and threaded by a purely azim uthal magnetic edd. The distribution of mass (stars+ gas+ dark matter) will create a gravitational potential at z = 0,  $_0$  (R). The gravitational circular velocity (not the observed rotation velocity) is dened as  $v_c^2$  Rd  $_0$ =dR. We will assume for simplicity that along the thickness of the H i disk the rotation curve is constant, i.e.  $v_z = v_z = 0$ . The disk is supposed to be rotating with an anom alous rotation speed v (R) >  $v_c$ (R) out to the radius  $R_{kep}$ .

We split up the rotational energy, T, in two parts  $T = T_0 + T$ , where  $T_0$  is the kinetic energy as if the disk, as it is, were rotating in force balance with the gravitational potential  $_0$  without magnetic elds (keeping all the rest of the

variables of the disk (R) and so on unchanged). More speci cally, T between two arbitrary radii, say R<sub>1</sub> and R<sub>2</sub>, and within  $jzj < h_1$ , where  $h_1$  is the characteristic scale height of the disk, is given by:

$$T = T_0 + 2 \qquad R_1 \qquad R_$$

where (R;z). According to our discussion in Appendix B,  $M_{cyl} = 0$  for pure azimuthal elds, and thus the total rotational energy of the disk must be the same with or without magnetic elds. Therefore, T = 0 between R = 0 and 1, implying

$$Z_{R_{kep}} Z_{h_1} \qquad Z_{1} Z_{h_1} = R (v^2 v_c^2) dz dR = R (v^2 v_c^2) dz dR : (C2)$$

$$0 \qquad 0 \qquad R_{kep} 0$$

Since we have assumed that  $v(R) > v_c(R)$  at  $R < R_{kep}$ , this equation implies that  $v(R) < v_c(R)$  outside the  $R_{kep}$  circle. Using Eq. (1), the LHS of the equation above reads  $Z_{R_{kep}} Z_{h_1}$   $1 Z_{R_{kep}} Z_{h_1} d(R^2 R^2)$ 

$$\int_{0}^{P^{2}} \int_{0}^{h_{1}} R(v^{2} - v_{c}^{2}) dz dR = \frac{1}{8} \int_{0}^{2} \int_{0}^{R_{kep}^{2}} \int_{0}^{h_{1}} \frac{d(R^{2}B^{2})}{dR} dz dR = \frac{B}{8} h_{1}R_{kep}^{2}B_{ikep}^{2};$$
(C3)

where B ;kep is the eld at R kep and B is a dimensionless coe cient less than 1 de ned by the relation  $\binom{R_{h_1}}{0} B^2 (R_{kep};z) dz = B h_1 B^2 (R_{kep};0)$ . A nalogously, the RHS of Eq. (C2) can be written as:

where is a dimensionless parameter less than 1. Thus, the value of B ;kep turns out to be

$$B_{jkep}^{2} = \frac{8}{R_{kep}^{2}} \int_{R_{kep}}^{B} (R;0)R(v_{c}^{2}(R) - v_{c}^{2}(R)) dR:$$
(C5)

The factor  $_{B}$  =  $_{B}$  takes into account possible three-dimensional elects. For a disk with a constant scale height with radius, one would expect =  $_{B}$  1, whereas for a disk that area, it could be a little bit larger, depending on the value adopted for  $h_{1}$ . As  $h_{1}$  ! 0, =  $_{B}$  goes to 1, whereas for a value of order of the scale height, ' 400 pc, =  $_{B}$  2 could be considered as an optim istically high value.

The maximum magnetic e ect on the rotation velocity within the  $R_{kep}$  circle occurs when  $B_{;kep}$  takes its maximum value; this happens when v (R) = 0 for  $R > R_{kep}$  7

$$B_{jkep}^{2} B_{cr}^{2} = \frac{8}{R_{kep}^{2}} \sum_{R_{kep}}^{2} (R;0)R v_{c}^{2}(R) dR:$$
(C 6)

 $B_{\rm cr}$  is an upper lim it to the magnetic  $\,$  eld strength at R  $_{\rm kep}$  .

In the standard dark matter scenario and for galaxies with a at gravitational circular speed,  $v_c(R) = constant$ , the maximum relative contribution of the magnetic eld, assuming generously that B (R) = B<sub>cr</sub> at R < R<sub>kep</sub> (see x3.1), is:

$$\frac{v^2}{v_c^2}\frac{v_c^2}{v_c^2} = 2 \quad B \quad \frac{R_g}{R_{kep}} \quad 1 + \frac{R_g}{R_{kep}} \quad \exp \quad \frac{R}{R_g} \quad ; \qquad (C7)$$

where an exponential law with scalelength R  $_{\rm g}$  was adopted for  $\$  (R ;0) beyond R  $_{\rm kep}$  .

In the other extrem e case that no contribution to the gravitational potential com es from a dark component,  $v_c^2$  (R) GM = R at large R, where M is the mass of the lum incuss matter in the disk. The corresponding maximum relative variation is given by

$$\frac{v^2}{v_c^2} \frac{v_c^2}{v_c} = 2 \quad B \quad \frac{R}{R_{kep}} \quad \frac{R_g}{R_{kep}} \quad \exp \quad \frac{R}{R_g} = \frac{R_{kep}}{R_g} \quad (C8)$$

If the disk is initially, in the lack of magnetic elds, rotating with velocity  $v_c$ , the growing magnetic tension during the MCP will produce a contraction of the inner disk and a radial drift of the outer disk. Let us assume that the disk beyond  $R_{kep}$  expands in a self-similar fashion and achieves a naldensity prole  $f(R;0) = (0 = 2) \exp((R = R_g))$ , with

> 1 the scale factor. T the conservation of angular momentum in plies that  $v(R) = {}^{1}v_{c}(R = )$ , where  $v_{c}(R = )$  is the circular velocity at R =. In the conventional dark halo scenario,  $v_{c} = constant$  so that  $v(R) = v_{c} =$ . Substituting these relations into Eq. (C 5), we get

$$B_{;kep}^{2} = \frac{8 (2 1)_{B} v_{c}^{2}}{{}^{2}R_{kep}^{2}} \int_{R_{kep}}^{2} f(R;0) R dR:$$
(C9)

Therefore, the relative e ect on the rotation curves starting with a disk in gravitational force balance is

$$\frac{v^2}{v_c^2} \frac{v_c^2}{v_c^2} = 2 B \frac{2}{2} \frac{1}{R_{kep}} \frac{R_g}{R_{kep}} = 1 + \frac{R_g}{R_{kep}} \exp \frac{R_g}{R_g} :$$
(C10)

Sim ilar manipulations allow us to calculate the sam e relation but assuming that galaxies do not contain massive dark halos. The corresponding calculation will give

$$\frac{\mathbf{v}^2}{\mathbf{v}_c^2} \frac{\mathbf{v}_c^2}{\mathbf{R}} = 2 \quad \mathbf{B} \quad \frac{1}{2} \quad \frac{\mathbf{R}}{\mathbf{R}_{kep}} \quad \frac{\mathbf{R}_g}{\mathbf{R}_{kep}} \quad \exp \quad \frac{\mathbf{R} \quad \mathbf{R}_{kep}}{\mathbf{R}_g} \quad : \tag{C11}$$

- Battaner, E., Garrido, J. L., Membrado, M., & Florido, E. 1992, Nature, 360, 652
- Battaner, E., & Florido, E. 1995, M N R A S, 277, 1129 Battaner, E., & Florido, E. 2000, Fund. Cosm ic Phys., 21, 1
- Battaner, E., Florido, E., & Jim enez-Vicente, J. 2002, A&A, 388, 213
- Beck, R., Brandenburg, A., Moss, D., Shukurov, A., & Sokolo, D. 1996, ARA&A, 34, 155
- Beck, R. 2002, in From Observations to Self-Consistent M odelling of the ISM in Galaxies, ed. M. de Avillez, & D. Breitschwerdt, Ap& SS, (astro-ph/0212288), in press
- Beck, R. 2003, in How does the Galaxy work?, ed. E. J. Alfaro, E.Perez, & J.Franco (Kluwer, Dordrecht), (astro-ph/0310287), in press
- Benjamin, R.A. 2000, RM xAC, 9, 256
- Binney, J., & Tremaine, S. 1987, Galactic Dynamics (Princeton Univ. Press: Princeton)
- Bland-Hawthorn, J., Freeman, K.C., & Quinn, P.J. 1997, ApJ, 490, 143
- Bolatto, A.D., Simon, J.D., Leroy, A., & Blitz, L. 2002, ApJ, 565, 238
- Bosma, A. 1978, PhD Thesis, University of Groningen
- Boulares, A., & Cox, D.P. 1990, ApJ, 365, 544
- Bureau, M., & Carignan, C. 2002, AJ, 123, 1316
- Burton, W.B. 1992, in The Galactic Interstellar Medium, Saas-Fee Advanced Course 21, ed. D. Pfenniger & P. Bartholdi (Berlin: Springer), 1
- Carignan, C., Beaulieu, S., & Freeman, K.C. 1990, AJ, 99, 178
- Castro-Rodr guez, N., Lopez-Corredoira, M., Sanchez-Saavedra, M. L., & Battaner, E.2002, A & A, 391, 519
- Chyzy, K.T., Beck, R., Kohle, S., Klein, U., & Urbanik, M. 2000, A&A, 355, 128
- Cox, D. P. 1988, in Supernova Remnants and the Interstellar Medium, ed. R.S. Roger, T.L. Landecker (Cambridge: Cambridge University Press), 73
- Cuddeford, P., & Binney, J. 1993, Nature, 365, 20
- de Blok, W.J.G., & Bosma, A. 2002, A&A, 385, 816
- de Blok, W.J.G., Bosma, A., & McGaugh, S. 2003, MNRAS, 340, 657
- Diplas, A., & Savage, B.D. 1991, ApJ, 377, 126
- Ferguson, A.M. N., & Johnson, R.A. 2001, ApJ, 559, L13
- Fiege, J.D., & Pudritz, R.E. 2000, MNRAS, 311, 85

- Freem an, K.C. 1997, in Dark M atter and V isible M atter in G alaxies, ASP Conference Series, ed.M. Persic, P. Salucci, 242
- Howard, A.M., & Kulsrud, R.M. 1997, ApJ, 483, 648
- Jokipii, J.R., & Levy, E.H. 1993, Nature, 365, 19
- Katz, J. I. 1994, Ap& SS, 213, 155
- Kronberg, P.P., Dufton, Q.W., Li, H., & Colgate, S.A. 2001, ApJ, 560,178
- Kulsrud, R.M. 1999, ARA&A, 37, 37
- M estel, L. 1999, Stellar M agnetism (C larendon P ress: O x ford)
- M oss, D ., & Shukurov, A . 2001, A & A , 372, 1048
- Navarro, J.F., Frenk, C.S., & White, S.D.M. 1996, ApJ, 462, 563 (NFW)
- Navarro, J.F., Frenk, C.S., & White, S.D.M. 1997, ApJ, 490, 493 Nelson, A.H. 1988, MNRAS, 233, 115
- Peratt, A.L. 1986, EEE Trans. Plasm a Sci., vol. PS-14, no. 6, 763

Persic, M ., & Salucci, P. 1993, M N R A S, 261, L21

Pfenniger, D., Combes, F., & Martinet, L. 1994, A&A, 285, 79

- Piddington, J.H. 1964, MNRAS, 128, 345
- Poezd, A., Shukurov, A., & Sokolo, D. 1993, MNRAS, 264, 285
- Reyes-Ruiz, M., & Stepinski, T.F. 1996, ApJ, 459, 653
- Roberts, P. H. 1967, An Introduction to Magnetohydrodynamics (Longm ans, London)
- Rognvaldsson, O.E. 1999, PhD Thesis, University of Copenhaguen Ruzmaikin, A.A., Shukurov, A.M., & Sokolo, D.D. 1988, Magnetic Fields of G alaxies (K luwer, D ordrecht)
- Ryu, D., Kang, H., & Bierm ann, P.L. 1998, A & A, 335, 19

Sanchez-Salcedo, F.J. 1996a, PhD Thesis, Universidad de Granada Sanchez-Salcedo, F.J. 1996b, ApJ, 467, L21

- Sanchez-Salcedo, F.J. 1997, M NRAS, 289, 863
- Sellwood, J.A., & Balbus, S.A. 1999, ApJ, 511, 660
- Shafranov, V.D. 1966, in Review s of Plasm a Physics, vol. I, ed. M.A. Leontovich (Consultants Bureau, New York)
- Shu, F.H. 1992, The Physics of A strophysics: G as D ynam ics, vol. II (Sausalito: University Science Books)
- Vallee, J.P. 1994, ApJ, 437, 179
- van Albada, T.S., Bahcall, J.N., Begeman, K., & Sancisi, R. 1985, ApJ, 295, 305
- van den Bosch, F.C., Burkert, A., & Swaters, R.A. 2001, MNRAS, 326, 1205
- Walter, F., & Brinks, E. 1999, AJ, 118, 273
- W idrow, L.M. 2002, Reviews of M odern Physics, 74, 775
- Zhao, H.S. 2002, MNRAS, 336, 159

11



Fig. C1. Schem atic diagram showing the e ect of the magnetic forces in a vertical cross-section of the gas disk in the magnetic alternative (panel a). R<sub>size</sub> is the radius of the gas disk. Beyond R<sub>H I</sub>, the outerm ost radius with H i detection, the gas could be fully ionized. The big arrows indicate the net e ect of the magnetic elds onto the gas disk. Beyond R<sub>kep</sub>, the magnetic forces point outwards in all directions. The expected rotation velocity versus radius is shown in panel b (solid line), together with the gravitational circular rotation speed given by baryonic matter (dashed line). In both panels R has the same scale.