# Occultations of HIP and UCAC2 stars downto 15<sup>m</sup> by large TNO in 2004-2014

#### © 2004 D.V. Denissenko

Space Research Institute (IKI), Moscow, Russia

Received 27 February 2004

Occultations of stars brighter than  $15^{m}$  by largest TNOs are predicted. Search was performed using the following catalogues: Hipparcos; Tycho2 with coordinates of 2838666 stars taken from UCAC2 (Herald, 2003); UCAC2 (Zacharias et al., 2003) with 16356096 stars between 12.00 and 14.99 mag to the north from -45 declination. Predictions were made for 17 largest numbered transneptunian asteroids, recently discovered 2004 DW and 4 known binary Kuiper Belt objects. 64 events occuring at solar elongation of 30° and more are selected, including exceptionally rare occultation of  $6.5^{m}$  star by double (66652) 1999 RZ<sub>253</sub> on 2007 October 4th. Observations of these events by all available means are extremely important since they can provide unique information about the size of TNOs and improve their orbits dramatically.

#### INTRODUCTION

Over 800 Transneptunian Objects (TNO) are discovered since 1992 with 67 of them being numbered and 7 having proper names as of February 2004. No reliable measurements of their sizes have been obtained so far. Angular sizes even of the largest objects are about  $0.04"\pm0.01"$  (Quaoar) which is at the limit of Hubble Space Telescope resolution. Most size estimates are based on the indirect methods strongly dependant on assumptions. Albedoes 0.04-0.08 are generally supposed for TNOs, but they are definitely varying among different objects.

The only direct method to measure sizes of single TNOs at present is observing stellar occultations by them. Over 500 occultations of stars by Main Belt asteroids have been observed so far, with 17% of them having 5 or more chords measured. For some objects size and shape were determined with an equivalent angular resolution of 0.002".

#### OCCULTATIONS BY TNO

TNOs are typically at 30-50 AU distances from the Earth corresponding to 0.2"-0.3"parallax. This means that occultation will only happen somewhere on the Earth if the geocentric path of the object passes within 200-300 mas from the star. Combined with a very slow angular motion of TNO (0.1-1 mas per second of time) it makes rate of stellar occultations by them about three orders of magnitude smaller than by main belt asteroids and about 100 times less than that of Jovian Trojans, approximately inversely proportional to  $(a-1)^2$ .

Estimates show that any given TNO will occult a star brighter than  $15^{\rm m}$  on average approximately every 4 or 5 years. For  $\sim 10^5$  HIPPARCOS stars brighter than  $10-11^{\rm m}$  we should be expecting one occultation in 10 years by 20 largest transneptunian objects, and only one event per century for  $\sim 10^4$  stars brighter than  $6.5^{\rm m}$ .

### STAR CATALOGUES ACCURACY

Error in coordinates of occulted star will significantly increase the uncertainty of both event time and position. At 40 AU distance, 1 milliarcsecond is corresponding to 30 km. 0.25" error in star coordinates will shift the occultation path by ~1 Earth radius. Since the positions of Transneptunian objects are known to 0.25" uncertainty, 0.05" accuracy of the star coordinates will be acceptable. Before 2003, coordinates were known to milliarcecond accuracy for ~100000 stars in HIPPARCOS catalogue. Tycho2 catalogue including ~2 Mln stars usually brighter than  $12^{m}$  is essentially a by-product of the same HIPPARCOS project. It has systematic and statistical errors typically within 0.1"-0.2" but reaching 0.5"-1.0" in some fields. Thus Tycho2 is not satisfactory for predicting occultations by TNO.

Situation in astrometry was revolutionized in 2003 with the release of  $2^{nd}$  USNO CCD Astrograph Catalogue UCAC2 (Zacharias et al.) with coordinates and proper motions for 48 Mln stars between  $8^{m}$  and  $16^{m}$ . Position errors of  $12^{m}$  stars in UCAC2 are 15-20 mas and 40-50 mas for  $15^{m}$ . This catalogue does not cover areas to the north from +45..+52 declination, but that is not a problem since no objects involved in current search will reach ±45 declination during next 10 years.

PREDICTIONS FOR 2004-2014

Occultations from 2004 Jan 27 to 2014 Dec 31 were computed for the following objects:

a) Recently discovered largest known TNO 2004 DW with absolute mag  $H_0=2.2-2.4$ ;

b) 17 numbered objects with  $H_0 < 5.5$ 

2.6 (50000) Quaoar 3.2 (28978) Ixion 3.3 (55565) 2002 AW<sub>197</sub> 3.3 (55636) 2002 TX<sub>300</sub> 3.6 (55637) 2002 UX<sub>25</sub> 3.7 (20000) Varuna 4.2 (42301) 2001 UR<sub>163</sub> 4.5 (19308) 1996 TO<sub>66</sub> 4.7 (26375) 1999 DE<sub>9</sub> 4.7 (38628) Huya 4.8 (24835) 1995 SM<sub>55</sub> 4.9 (19521) Chaos 4.9 (47171) 1999 TC<sub>36</sub> (binary) 5.2 (26181) 1996 GQ<sub>21</sub> 5.3 (55638) 2002 VE<sub>95</sub> 5.4 (15874) 1996 TL<sub>66</sub> 5.4 (48639) 1995 TL<sub>8</sub>

c) 3 numbered binary TNO

5.8 (26308) 1998 SM<sub>165</sub> 5.9 (66652) 1999 RZ<sub>253</sub> 6.6 (58534) 1997 CQ<sub>29</sub>

Single body diameters were calculated from  $H_0$  by the formula log D[km]=3.52-0.2\*H\_0. This corresponds to assumed albedo of 0.16 and is a conservative estimate. For (50000) Quaoar this formula gives 1000 km which is in a good agreement with the lower limit from Hubble direct imaging (1260±300 km). If albedo is 0.04, actual size of TNO will be exactly twice the value listed in table. For binary asteroids with two bodies of the same size and albedo diameter of each component will be 0.71\*D.

Orbits of selected TNO from 2004 Jan. 27th version of astorb.dat (Bowell, 2004) were integrated through the end of 2014 and all geocentric conjunctions within 0.5" to HIP and UCAC2 stars were found. For the year 2004 occultations by 36 largest unnumbered objects with  $H_0$ <5.5 were also computed and 9 events have been found. Mean value of current ephemeris uncertainty (CEU) was 0.26" for the numbered objects and 1.0" for unnumbered ones.

Results of the search are presented in table. Intervals of time (rounded to 5 minutes) are given when occultation may happen on the night part of Earth with 0.25" uncertainty at both sides already taken into account. Due to the different speed of TNOs relative to Earth (from 25 km/s at opposition to 5 km/s and even less near the stationary point) these interval may vary from ~10 minutes to 1.5 hour in case of Varuna occultation in 2012. For double asteroids extra 0.25" were added since orbital distances in these pairs are also estimated as 6-8 thousand km.

Exceptional occultation of  $6.5^{m}$  K0 star HIP 111398 = HD 213780 = SAO 146161 = ZC 3313 in Aquarius by binary object (66652) 1999 RZ<sub>253</sub> in 2007 should be noted. Additional observations of this TNO on the largest telescopes are extremely important to predict the separation and positional angle of components at the epoch of the event. International campaign involving observers in Australia, New Zealand, China, Japan, eastern Russia and Hawaii should be organized to measure the sizes and shape of this binary system.

Finder charts and preliminary path plots for occultations listed here are available at http://hea.iki.rssi.ru/~denis/TNOocc.html or by E-mail request to denis@hea.iki.rssi.ru.

# **OBSERVING METHODS**

Observations can be performed by several techniques and different kinds of equipment. In principle, highest temporal resolution is not necessary for occultations by TNO. With event durations about several dozen seconds and even 2-3 minutes in some cases, 1-3 sec integration time will already give the unprecedented chord measurement accuracy. This is why one can use CCD imaging, CCD in drift-scan mode, photometers, high-sensitivity videocameras and even visual

observations with stopwatch. CCD readout time should be reduced to minimum by binning pixels and exposing small window instead of full frame. Optimal exposure time should be selected depending on the magnitude of occulted star and detector sensitivity. Precise absolute timing of star disappearance and reappearance is not critical since with 5-10 minutes uncertainty of event time even 1-second accuracy will improve the orbital position by two orders of magnitude. For bright occultations fast photometers can provide the measurements of occulted star diameter at unprecedented angular resolution. Since TNOs are moving at 0.1-1 mas/sec rate (500-5000 times slower than the Moon), new close binaries can be discovered which are impossible to resolve by any other methods.

# DISCUSSION

Search was intentionally limited to the largest objects with ephemeris uncertainty of 0.3" or better. Although smaller objects might (and definitely will!) occult brighter stars, detection probability of those events would be much smaller compared to those listed here. It is preferrable to concentrate on 2-5 "special" events each year. Success can only be achieved by the joint efforts planned well in advance. Worldwide observations by as many telescopes as possible are necessary. Prediction of events till 2014 will allow to make changes in tight observing schedules at the large scopes. Additional observations of selected TNOs to improve ephemeris and prediction accuracy are also encouraged. New large objects will definitely be found in the nearest future. Predictions of occultations by them will be posted at author's site together with updates on those listed here.

# APPLICATIONS

Positive observation of stellar occultation by TNO will make significant impact in several areas of astronomy at once. It will allow to: - measure the size and shape of these distant objects directly which is impossible by other means;

- give indications on their composition, and together with estimates of masses in binary systems directly determine their density;

- improve orbital position by 10-50 times assisting possible space missions to TNO;

- provide unique information valuable for understanding the theory of Solar system evolution;

- discover probable close binary objects with the separations between components of the order of 1000 km.

#### ACKNOWLEDGEMENTS

I would like to thank Alexander Yascovich for lending the copies of UCAC2 catalogue and Andrei Plekhanov for writing LinOccult program which was used to plot the maps of occultation paths posted on the website.

#### REFERENCES

- Bowell E., http://www.lowell.edu/users/elgb (2004)
- Brown M. E., Trujillo C. A., IAU Circular 7807 (2002)
- Dunham, D. W.; Goffin, E.; Manek, J.; Federspiel, M.; Stone, R.; Owen, W., Mem. Soc. Astr. Ital., Vol. 73, No. 3, p. 662 (2002)
- Herald, D., Cat.1291....0H (2003)
- Marsden B. G., Minor Planet Electronic Circular 2004-D15 (2004)
- Noll K., Stephens D., IAU Circular 7824 (2002)
- Noll K., Stephens D., IAU Circular 8143 (2003)
- Stone, R. C.; McDonald, S. W.; Elliot, J. L.; Bowell, E., The Astron. Journ., Vol. 119, No. 4, pp. 2008-2017 (2000)
- Trujillo C. A., Brown M. E., IAU Circular 7787 (2002)
- Zacharias, N.; Urban, S. E.; Zacharias, M. I.; Wycoff, G. L.; Hall, D. M.; Germain, M. E.; Holdenried, E. R.; Winter, L., Cat.1289....0Z (2003)

# OCCULTATIONS OF STARS BY TNO 2004-2014

Date	of ev	vent	. Time	Asteroid	size	Dur	Star	R.A. (2000.0) Dec
Year	Mon D	DD	(UT)	name	km	sec	Name	hh mm ss.sss dd mm ss.ss Ma
2004	Jan 2	29	23:35-23:55	2003 AZ84	568	23.4	2UCAC 36806754	07 12 19.883 +14 04 20.02 14.
2004	Feb 1	14	22:10-22:35	2001 XR254	255	14.2	TYC 1343-00785-1	06 54 14.687 +21 37 48.10 11.
2004	Mar C	7	03:15-03:30	2000 CO105	268	14.7	2UCAC 39141168	08 23 16.013 +20 33 01.42 12.
2004	Mar C	8	02:10-02:25	1999 CC158	241	11.6	2UCAC 40185292	09 15 58.264 +23 38 56.76 13.
2004	Mar 2	20	13:35-13:45	1996 TL66	275	10.5	2UCAC 36232084	03 00 28.306 +12 40 40.89 13.
2004	Mar 2	27	00:20-00:40	2002 KW14	302	21.2	2UCAC 24855750	15 34 29.721 -18 18 28.41 13.
2004	Apr C	)7	10:30-11:00	2001 KD77	252	24.7	2UCAC 23658681	16 58 51.823 -20 36 51.67 11.
2004	Jun 2	26	19:55-20:05	Chaos	347	11.6	2UCAC 40137180	04 08 53.436 +23 56 34.41 13.
2004	Jun 2	29	18:25-19:10	1997 CQ29	158	9.5	2UCAC 34467578	11 04 07.504 +07 39 54.56 14.
2004	Aug (	8	23:15-23:30	2001 QD298	257	10.5	2UCAC 24989527	21 52 30.098 -18 08 13.32 14.
2004	Nov 1	15	09:00-09:25	2000 CN105	340	32.5	2UCAC 35937214	10 34 08.639 +11 59 33.61 14.
2005	Mar 1	13	21:25-21:35	2002 UX25	631	21.6	2UCAC 34953654	01 37 42.194 +09 22 06.65 13.
2005	May C	7	01:05-01:40	1999 RZ253	219	14.9	2UCAC 28298467	22 29 21.656 -10 02 41.04 14.
2005	Sep 2	29	07:50-08:10	1995 SM55	363	16.6	2UCAC 39255131	02 07 19.531 +21 06 25.09 14.
2005	Oct 2	24	08:05-08:30	1998 SM165	229	9.4	2UCAC 33162437	01 31 09.061 +04 23 37.99 14.
2005	Dec 2	27	09:15-09:35	Huya	380	17.9	2UCAC 30864992	14 20 27.168 -02 54 42.58 11.
2006	Jun 1	18	20:25-20:50	1999 KR16	229	17.4	2UCAC 28922119	14 18 38.914 -08 29 05.17 13.
2006	Jul 2	26	07:55-08:25	Ixion	759	55.9	2UCAC 22945926	16 34 14.971 -22 04 37.89 13.
2006	Oct (	7	19:30-19:45	Ixion	759	36.0	2UCAC 22946030	16 35 05.045 -22 11 08.86 13.
2007	Aug (	)4	04:20-04:35	2002 VE95	288	15.8	2UCAC 35331886	04 14 47.942 +10 08 18.19 13.
2007	Oct (	)4	11:10-11:50	1999 RZ253	219	11.2	HIP 111398	22 34 06.604 -09 36 30.64 6.
2008	Jan 2	22	06:55-07:30	Chaos	347	26.0	2UCAC 40810575	04 21 06.919 +25 33 26.70 10.
2008	Feb 1	11	04:20-04:40	Varuna	603	28.8	TYC 1913-00670-1	07 18 50.100 +25 43 19.31 11.
2008	Sep 1	10	12:25-13:15	Quaoar	1000	149.5	2UCAC 26249129	17 05 49.913 -15 20 44.26 13.
2008	Sep 3	30	07:20-07:40	Ixion	759	44.6	2UCAC 22496420	16 44 05.610 -23 18 07.11 13.
2008	Oct (	7	19:40-19:55	Ixion	759	37.3	2UCAC 22496543	16 44 33.933 -23 19 19.45 12.
2008	Dec C	)7	01:55-02:20	Varuna	603	29.0	2UCAC 40846256	07 29 48.171 +25 40 07.71 14.
2009	Mar 1	10	08:25-08:40	2002 VE95	288	19.7	2UCAC 34960757	04 17 00.297 +09 18 45.63 13.
2009	May C	01	13:50-14:15	Quaoar	1000	54.1	2UCAC 26252549	17 18 12.542 -15 24 49.90 14.
2009	Jul 1	18	04:40-04:50	2002 VE95	288	10.8	HIP 21308	04 34 30.274 +10 00 58.46 7.
2009	Aug 2	22	10:40-11:15	2002 UX25	631	68.5	2UCAC 35326247	02 09 00.114 +10 07 56.40 14.
2009	Sep 2	25	11:05-11:30	1995 TL8	275	15.1	TYC 1223-00579-1	02 40 23.728 +15 46 01.74 11.

Date	of e	event	: Time	Asteroid	size	Dur	Star	R.A. (2000.0) Dec
Year	Mon	DD	(UT)	name	km	sec	Name	hh mm ss.sss dd mm ss.ss Mag
2009	Oct	09	10:15-10:35	2002 TX300	724	28.6	2UCAC 41650964	00 37 13.610 +28 22 22.98 13.1
2009	Oct	15	17:55-18:10	Quaoar	1000	45.8	2UCAC 26023237	17 12 33.132 -15 30 54.97 15.0
2010	Feb	19	22:55-23:20	Varuna	603	32.3	TYC 1914-00301-1	07 29 22.470 +26 07 23.23 11.0
2010	Jul	19	01:35-02:05	Ixion	759	42.4	2UCAC 22032118	16 54 25.862 -24 24 22.86 11.5
2011	Feb	22	02:25-02:50	1995 TL8	275	18.3	2UCAC 37301105	02 39 37.120 +15 41 48.75 13.6
2011	Jun	01	11:05-11:25	Ixion	759	30.3	2UCAC 21584537	17 03 39.446 -25 01 27.82 14.7
2011	Aug	03	22:35-23:10	1999 RZ253	219	10.8	2UCAC 29384710	22 59 38.963 -07 02 53.38 14.6
2011	Nov	04	00:35-00:50	Quaoar	1000	35.9	2UCAC 26027484	17 24 23.771 -15 38 43.34 14.6
2011	Dec	18	11:40-12:50	1999 TC36	347	40.7	2UCAC 32137136	01 35 51.911 +01 13 15.69 14.1
2012	Apr	16	04:20-04:55	Ixion	759	55.9	TYC 6816-00401-1	17 12 26.791 -25 34 10.85 10.8
2012	May	01	03:35-03:55	1999 RZ253	219	10.2	2UCAC 29759182	23 04 54.428 -06 29 06.50 14.9
2012	Jun	01	17:55-18:15	Quaoar	1000	41.3	2UCAC 26259184	17 32 11.415 -15 24 30.06 13.5
2012	Jun	04	01:10-01:30	1996 GQ21	302	14.0	2UCAC 28386513	15 19 25.744 -09 50 28.94 14.8
2012	Oct	07	04:30-04:45	2002 AW197	724	30.2	2UCAC 32701289	09 34 01.570 +02 31 43.27 14.7
2012	Oct	28	19:50-21:25	Varuna	603	144.8	2UCAC 41016878	07 53 29.509 +26 05 51.86 13.4
2013	Jan	29	06:10-06:30	Varuna	603	24.2	2UCAC 41187592	07 47 50.499 +26 31 18.27 14.8
2013	Feb	18	11:40-12:50	Chaos	347	93.9	2UCAC 41496855	04 49 59.491 +27 50 52.79 14.4
2013	May	28	18:35-18:55	1999 KR16	229	10.1	2UCAC 29288802	15 13 08.695 -07 07 13.91 11.7
2013	Jun	02	22:20-22:35	Ixion	759	30.4	2UCAC 21139770	17 14 19.280 -26 08 23.27 14.4
2013	Oct	23	13:20-13:35	Ixion	759	31.3	2UCAC 21137306	17 10 57.419 -26 07 23.91 14.2
2013	Dec	06	09:10-09:20	Huya	380	11.0	TYC 5025-00448-1	15 39 28.493 -05 37 11.16 10.1
2013	Dec	20	14:05-14:40	2002 TX300	724	55.4	TYC 2276-01081-1	00 52 46.890 +31 03 39.18 11.4
2013	Dec	27	17:20-17:50	1995 SM55	363	23.4	2UCAC 42006839	02 40 01.674 +29 27 23.90 13.3
2014	Jan	24	01:35-01:50	Quaoar	1000	34.9	2UCAC 26041319	17 43 03.235 -15 45 10.63 14.6
2014	Mar	01	16:30-16:45	2004 DW	1200	45.9	TYC 5476-00882-1	09 58 22.549 -08 16 55.27 12.1
2014	Mar	10	15:25-15:55	1997 CQ29	158	6.3	2UCAC 32542503	11 58 36.918 +02 28 27.68 11.9
2014	Apr	15	01:50-02:00	1996 TL66	275	9.9	2UCAC 34961430	04 26 07.463 +09 23 13.78 14.7
2014	Jul	24	14:20-15:10	Huya	380	51.0	TYC 5024-00589-1	15 36 41.149 -04 36 29.35 11.7
2014	Aug	14	21:40-22:40	1996 GQ21	302	48.1	HIP 75882	15 29 58.355 -10 03 19.73 9.1
2014	Sep	13	08:35-09:20	2002 VE95	288	27.1	2UCAC 34784210	05 31 16.439 +08 55 17.49 12.3
2014	Oct	20	14:45-15:25	1999 RZ253	219	12.4	2UCAC 29941401	23 09 49.521 -05 59 47.15 14.5
2014	Nov	8 0	22:00-22:15	Quaoar	1000	35.4	2UCAC 26039148	17 40 27.392 -15 41 59.91 13.9