

V. A. Sirota, A. S. Ilyin, K. P. Zybin, A. V. Gurevich.

The growth of supermassive black holes in the centers of galaxies: absorption of stars and activity of galactic nuclei.

Abstract

The evolution of a supermassive black hole in the center of galaxy is considered. We analyze the kinetic equation describing relaxation processes associated with stellar encounters. The initial distribution function of stars is assumed to be isothermal, and it changes in accordance with consumption of stars by the black hole. We calculate the consumption rate and obtain the law of growth of supermassive black hole, i.e. the dependence of the black hole mass on time and on the velocity dispersion of stars. The obtained values of mean black hole mass and mean rate of stars consumption by a black hole are in reasonable agreement with observational data.

1. Introduction.

The massive compact concentrations of mass in the centers of galaxies are believed to be supermassive black holes with masses $10^6 - 10^8$ solar mass ($M_\odot$). The recent observations [1] provide significant evidences for this hypothesis.

There are two global approaches to the origin of these black holes: they could either form together with the galaxy or even earlier [2], or grow up during galaxy evolution, capturing matter from the surrounding. We throughout the paper consider the second approach.

Up to the present time there has been no consistent theory how the growth of black holes in the galactic centers goes on.

Black hole can capture luminous (stars and gas) and cold dark matter from the central parts of galactic core (bulge). Consumption rate of stars has been investigated in [3, 4, 5, 6, 7]. (The first statement of the problem of distribution and absorption of particles in the Coulomb potential has been considered in [8]). In [5, 6, 7] the diffusion in two-dimensional energy-momentum space has first been discussed as the main process. However, all these authors considered the dynamics of stars in the Coulomb potential of black hole, whereas, as it has already been pointed out in [6], in the case of giant black holes the consumption rate is determined by distances where it is negligible and the stars move in their own self-consistent potential field. On the other hand, in all these papers stationary solution has been considered; as we will show below, the solution of the real problem is non-stationary.

The kinetic theory describing dynamics of non-baryon cold dark matter (CDM) alone (without baryon fraction and central black hole) has been developed in [9]. It has been shown that the dark matter should have formed quasi-stationary spherical structures with singular density distribution, bound by gravitational forces. These structures are now observed as galactic halos. The baryon matter should have fallen after recombination in the potential wells formed by dark matter, beginning formation of galaxies and forming seed black holes in the centers of the wells [10]. This picture explains well why the black

holes are situated exactly in the dynamical centers of galaxies [11]: the potential is very smooth in the central part, and the black hole could not reach the center unless it would be formed just in the center, in the potential well already existing.

The contemporary infall of CDM particles into the black hole caused by gravitational interaction with stars and the corresponding growth of black hole has been investigated in [12]. It was shown that this process leads to formation of black holes whose masses are consistent with the most part of supermassive black hole masses measured nowadays.

In the present paper we analyze the mutual dynamics of cold dark matter and stars in central parts of a galaxy. In accordance with previous papers, we find that the most important process leading to the black hole growth is (for both components) far encounters causing the relaxation in the energy-momentum space, or, to be more precise, in the space of adiabatic invariants. The influence of black hole appears both in gravitational potential (i.e. the view of adiabatic invariants) and { even stronger { in boundary condition.

We derive the equations describing time-dependence of stars- and dark matter- distribution functions, and calculate the coefficients in one particular assumption of isothermal distribution of stars and the potential dominated by stars, either (Section 2).

In Section 3 we discuss the boundary condition caused by presence of black hole, and the influence of the black hole potential.

Further, we find an approximate solution of the equation for stars, ignoring presence of dark matter and gas (Section 4). The result shows that black hole mass should grow in time as $t^{1/2}$, and is proportional to v^3 , where v is the rms velocity in the inner regions of bulge.

In discussion, we review briefly the results of the paper. We show that the mean energy released due to stellar matter absorption is in reasonable agreement with the observed AGN activity.

2. Two-component dynamics.

We denote by $f(\mathbf{r};\mathbf{v})$ and $f_d(\mathbf{r};\mathbf{v})$ the distribution functions of stars and dark matter, respectively. This means that fraction of mass of sort contained inside the interval $d^3\mathbf{r}$, $d^3\mathbf{v}$ is $f d^3\mathbf{r}d^3\mathbf{v}$.

The mutual dynamics of dark matter particles and stars including diffusion in the momentum space is described by kinetic equation with Landau scattering integral in the right part [13]:

$$\frac{df}{dt} = St[f];$$

or

$$\frac{\partial f}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{r}} f + \mathbf{r} \cdot \nabla_{\mathbf{v}} f = 2 G^2 \frac{\partial}{\partial v_k} \int d^3\mathbf{v}^0 w_{kp} M f \frac{\partial f}{\partial v_p} - M \frac{\partial f}{\partial v_p^0} f \quad (1)$$

Here greek indexes denote the sort of particles, M denotes the mass of particles, $f(\mathbf{r};\mathbf{v})$ corresponds to scattered particles, $f(\mathbf{r};\mathbf{v}^0)$ corresponds to scatterers,

$$w_{kp} = u^2_{kp} \quad u_k u_p = u^3; \quad u = v^0 - v;$$

u is the relative velocity, γ is gravitational Coulomb logarithm. The gravitational

potential $\Phi(r)$ is determined by Poisson law,

$$\Delta \Phi = 4\pi G \rho(r) = 4\pi G \int_0^\infty f(r;v) d^3v;$$

where $\rho(r)$ is mass density of fraction.

Comparing the terms in (1) one can see that scattering of both dark matter and stars by stars is $M = M_{dm}$ times more effective than scattering by dark matter particles, and in the case of dark matter the second term is $M_{dm} = M$ times smaller than the first. So we rewrite the equation (1) in the form

$$\frac{df}{dt} = 2\pi G^2 M \frac{\partial}{\partial v_k} W_{kp} \frac{\partial f}{\partial v_p} - \frac{\partial}{\partial v_k} W_k f \quad (2)$$

$$\frac{df_d}{dt} = 2\pi G^2 M \frac{\partial}{\partial v_k} W_{kp} \frac{\partial f_d}{\partial v_p} \quad (3)$$

where

$$W_k = \int_0^\infty \frac{\partial f}{\partial v_p^0} w_{kp} d^3v^0; W_{kp} = \int_0^\infty f w_{kp} d^3v^0; \quad (4)$$

Rewriting equation in terms of adiabatic invariants

If the potential is spherically symmetric, one can introduce Hamiltonian variables $(I; \theta)$, where $I = (I; m; m_z)$ are adiabatic invariants (m is absolute value of angular momentum, m_z is its projection on z axis, I is radial action; all per unit of mass):

$$I = \frac{1}{2} \int_{r_-(E,m)}^{r_+(E,m)} v_r dr = \frac{1}{2} \int_{r_-(E,m)}^{r_+(E,m)} \frac{dr}{\sqrt{2(E - \Phi(r) - \frac{m^2}{2r^2})}} \quad (5)$$

$$E = \frac{v^2}{2} + \Phi(r); m = r v_\theta; E - \Phi(r) - \frac{m^2}{2r^2} = 0;$$

and θ are corresponding angles. One of the advantages of these variables is that the adiabatic invariant I of a particle (unlike the energy E) does not change during slow changes of the potential.

In these variables, the left side of the kinetic equation does not contain derivatives over action variables I_k [12, 14]:

$$\frac{\partial f(I; \theta)}{\partial t} + \sum_k \dot{I}_k(I) \frac{\partial f}{\partial I_k} = St_I; [f]$$

We now make use of the fact that the frequency of collisions is much smaller than the frequency of orbital motion. After averaging over the angle variables θ_k the second term in the equation vanishes, and the equations (2),(3) then take the form (see [14, 12] for detailed derivation)

$$\frac{\partial f(I)}{\partial t} = \frac{\partial}{\partial I_k} R_{kp} \frac{\partial f}{\partial I_p} - \frac{M}{M} \frac{\partial}{\partial I_k} (R_k f); \quad (6)$$

where

$$R_k = 2 G^2 M \frac{1}{(2)^3} \int d^3 \frac{\partial I_k}{\partial v_p} W_p ; \quad (7)$$

$$R_{kp} = 2 G^2 M \frac{1}{(2)^3} \int d^3 \frac{\partial I_k}{\partial v_m} \frac{\partial I_p}{\partial v_n} W_{mn} ;$$

We assume that in the vicinity of the center the distribution function is spherically symmetric, i.e. does not depend on the direction of angular momentum. Then, there is a cylindrical symmetry in the space of adiabatic invariants. Making use of

$$R_3 = \frac{m_z}{m} R_2 ; \quad R_{31} = \frac{m_z}{m} R_{21} ; \quad R_{32} = \frac{m_z}{m} R_{22} ;$$

we can exclude m_z from the equation (6). Eventually, we obtain the equations for distribution functions of stars and of dark matter:

$$\frac{\partial f}{\partial t} = \frac{\partial}{\partial I} R_{11} \frac{\partial f}{\partial I} + R_{12} \frac{\partial f}{\partial m} R_1 f + \frac{1}{m} \frac{\partial}{\partial m} m R_{21} \frac{\partial f}{\partial I} + R_{22} \frac{\partial f}{\partial m} R_2 f \quad (8)$$

$$\frac{\partial f_d}{\partial t} = \frac{\partial}{\partial I} R_{11} \frac{\partial f_d}{\partial I} + R_{12} \frac{\partial f_d}{\partial m} + \frac{1}{m} \frac{\partial}{\partial m} m R_{21} \frac{\partial f_d}{\partial I} + R_{22} \frac{\partial f_d}{\partial m} \quad (9)$$

We see that evolution of distribution functions is described by axially symmetric diffusion in three-dimensional space of adiabatic invariants.

We note that equations (6), (8) have a divergent form, i.e.

$$\frac{df}{dt} = \frac{\partial}{\partial I_k} S_k$$

The flux of matter through any surface in the momenta space is given by the integral of $\vec{S}$ over the surface. For example, the flux through the plane $I = 0$ is $\int dm \int dm_z S_I = 2m \int S_I dm$. The boundary $I = 0$ is not physical, hence flux of matter over any part of it should be zero. Thus, we obtain the boundary condition

$$S_I \Big|_{I=0} = R_{11} \frac{\partial f}{\partial I} + R_{12} \frac{\partial f}{\partial m} R_1 f \Big|_{I=0} = 0 \quad (10)$$

In absence of a black hole, the other boundary condition would be

$$S_m \Big|_{m=0} = 0 :$$

The second boundary condition imposed by presence of black hole will be considered in Section 3.

Isothermal distribution function

One important example is the Maxwellian stars distribution function with arbitrary time-independent potential:

$$f_M = f_0 e^{-E/\sigma^2}$$

This function corresponds to the equilibrium state, since it satisfies equation (2). This means that both parts of equation (8) are equal to zero. Moreover, flux of matter over any surface in the momenta space must be zero, hence all components of the flux S must vanish. This puts two relations on coefficients $R_k; R_{kp}$:

$$R_{11} \frac{\partial f_M}{\partial I} + R_{12} \frac{\partial f_M}{\partial m} - R_1 f_M = 0; \quad R_{21} \frac{\partial f_M}{\partial I} + R_{22} \frac{\partial f_M}{\partial m} - R_2 f_M = 0 \quad (11)$$

In the particular case of potential dominated by stars, there is the only parameter characterizing the distribution function:

$$f_{is} = f_0 e^{-E/v^2}; \quad v_{is}^2 = 2^{-2} \ln r; \quad f_0 = \left(\frac{h}{2} \right)^{5=2G} i^{-1} : \quad (12)$$

This function is called isothermal, the parameter v^2 corresponds to velocity dispersion.

We now calculate the coefficients $R_k; R_{kp}$ for isothermal distribution function, $f = f_{is}$. All manipulations needed for the procedure are described in Appendix. The result is

$$\begin{aligned} R_{22} &= 0.5GM - R; \\ R_{12} &= 0.6R(1 - v^2); \\ R_{11} &= R(0.4 - v^2 + 0.8v^2 + 0.4); \\ R_1 &= \frac{R}{I + 0.6m} (0.7 - v^2 - 0.8v^2); \\ R_2 &= 1.3 \frac{R}{I + 0.6m} v^2; \\ &= 0.3 \frac{m}{I + 0.6m} \end{aligned} \quad (13)$$

where approximate expressions for functions are:

$$\begin{aligned} v^2 &= 1 + 0.35 r^{0.6}; \\ v^2 &= 0.5 - 0.8 \ln \left(\frac{r}{r_0} \right) - 1; \quad v^2 = 0.5 + 0.33 r^{0.9}; \end{aligned}$$

The isothermal distribution function can also be expressed in terms of Hamiltonian variables (see Appendix):

$$f_{is}(I; m) = f_0 e^{-E/v^2}, \quad \frac{0.3f_0 v^2}{(I + \frac{1}{2}m)^2} \quad (14)$$

3. The black hole, and setting the problem.

The scattering of dark matter particles by stars and their consumption by a black hole has been discussed in [12]. It has been assumed in that paper that growth of black holes is fully determined by dark matter. Here we discuss an opposite assumption of full absence of dark matter. From now on, we discuss only dynamics of stars. Accordingly, we hereafter omit the index in f .

Black hole can effect the dynamical evolution of surrounding matter in two ways: consuming matter from the loss-cone and changing the gravitational potential. We now consider the first effect as principal one, and later on will briefly discuss the second.

The black hole with mass M_{bh} disrupts (and after that consumes) all stars that come nearer than the tidal radius $r_t = \left(\frac{6 M_{bh}}{\rho} \right)^{1/3}$, where ρ is the density of the star. We assume it to be approximately equal to solar density. In terms of variables $(I; m)$ this means that during one oscillation period all stars inside the loss-cone

$$m_t = \frac{q}{2GM_{bh}r_t} \approx 6 \cdot 10^{-8} \frac{M_{bh}}{M} \approx 10^{-6} \text{ cm}^2 = s \quad (15)$$

are disrupted. This boundary condition on m has first been pointed out in [6]. On the other hand, if angular momentum is less than m_g :

$$m < m_g = 4GM_{bh}c; \quad (16)$$

the star will anyway fall into the black hole [15], independently of m_t . (If the black hole mass M_{bh} is greater than $3 \cdot 10^6 M_\odot$, the tidal radius becomes smaller than the gravitational radius r_g , and stars become consumed without disruption. For dark matter particles the loss cone is always defined by m_g .) For brevity, we shall hereafter denote the boundary value by m_t , assuming maximal of the two values m_g and m_t : $m_t = \max(m_g, m_t)$.

One can easily estimate the growth of the black hole caused by direct capture of matter from the loss-cone: taking, e.g., the isothermal function (14), we have $\dot{M}_{bh} \approx 2 m_t G$. For black holes with masses more than $10^3 M_\odot$ relative increase of mass $\dot{M}_{bh}/M_{bh}$ is less than 10%. Hence the main growth is connected with diffusion of matter into the loss-cone.

The condition that the particles vanish as they get into the area $m = m_t$ may be expressed in terms of equation (8) by boundary condition

$$f|_{m=m_t} = 0 \quad (17)$$

It is the change of gradient of distribution function $\partial f / \partial m$ near the boundary that causes shift of particles in the momenta space into the loss-cone. This mechanism may lead to consuming much more mass than has initially been inside the loss-cone.

Applying Gauss theorem to the equations (6), (8) we see that the rate of black hole growth dM_{bh}/dt , i.e. loss of whole mass of stars in the surrounding space, is equal to flux through the boundary m_t :

$$\dot{M}_{bh} = S = \int_0^{m_t} d^3I \int_{m_t}^{m_t} dm_z S_m = (2\pi)^3 2m_t \int_0^{m_t} dI R_{22} \frac{\partial f}{\partial m} + R_{12} \frac{\partial f}{\partial I} R_2 f|_{m=m_t} \quad (18)$$

The last two terms are zero because of the boundary condition (17).

So, to find the contemporary mass of black hole and distribution of stars in its vicinity one should solve equation (8) with boundary conditions (10) and (17). The isothermal function (14) is the most natural choice of initial distribution because it is stationary

distribution with self-consistent potential and with no flows. If there were no black hole, the isothermal distribution remained constant. The boundary condition (17) disturbs the equilibrium and makes the distribution change.

Furthermore, we take the same isothermal function to calculate the coefficients $R_k; R_{kp}$ at any moment of time. We suppose that one may neglect the changes in distribution of scatterers, because (7) includes only integrated function f , and changes in f do not effect the coefficients significantly. We will check the convenience of this approximation in Section 4.

Let us now estimate the influence of gravitational potential of black hole. Taking values R_{ik} of the isotherm (13) for an estimate, one can see that to the lifetime of the Universe $t \approx 3 \cdot 10^{10}$ s different members of Φ may produce significant changes of f at scales $m \approx \frac{1}{R} \frac{p}{t} \approx 8 \cdot 10^{-25} \text{ m}^2/\text{s}$, this corresponds to

$$r \approx 1 \text{ pc} \cdot \frac{1}{200 \text{ km/s}} \quad (19)$$

Dynamics of matter at these distances determines contemporary infall into the black hole. This is much smaller than size of bulge ($\approx 1 \text{ kpc}$), so we should not take boundaries of the distribution into account. On the other hand, the black hole potential becomes significant at distances

$$r \approx r_g = \frac{GM_{bh}}{2} \approx 1 \text{ pc} \cdot \frac{M_{bh}}{10^8 M} \cdot \frac{1}{200 \text{ km/s}}$$

We see that for not very massive black holes,

$$M_{bh} \approx 10^8 M \cdot \frac{1}{200 \text{ km/s}}; \quad (20)$$

the region of influence of the black hole potential does not reach up to characteristic distances (19). In this case only a small fraction of particles enters the region, and, because of ellipticity of most orbits, spends there rather small part of time. This is in full conformity with conclusions by [6].

The same arguments are applicable to the region $r < r_{coll} \approx R \cdot \frac{M_{bh}}{M}$ where orbital velocities become comparable to the escape velocity from a typical star, and inelastic scatterings are important [5, 6]. One can easily check that $r_{coll} \approx r_g$ for values under discussion.

4. A symptotic solution.

The equation (8) is difficult to solve. In this section we obtain at least some integrated characteristics that make us able to estimate the total consumption rate S of stars and its evolution in time.

Let us first discuss the vicinity of the black hole boundary, $m_t \approx m \cdot \frac{p}{R_{22}t}$ and $I > m$. Comparing different terms in equation Φ we find out that, e.g., the term containing R_{22} is of the order of $R_{22}f \approx m^2 \cdot \frac{\partial f}{\partial t}$, which means that the picture near the boundary is quasi-stationary at contemporary time. All the other terms except the two (with $R_{11}; R_1$) that are singular at $m \rightarrow 0$ (see (13)) are maximum of the order of

$R_{22}f = \text{Im}$, i.e. much smaller. To get rid of the two remaining terms, we integrate (8) over I and introduce $F(m) = \int_R f(I; m) dI$. Thus, we obtain¹

$$\frac{1}{m} \frac{\partial}{\partial m} m R_{22} \frac{\partial F}{\partial m} = 0 \quad (21)$$

Taking into account the boundary condition at $m = m_t$, we find

$$F(m) = C(t) \ln \frac{m}{m_t}; \quad m \ll \frac{p}{Rt} \quad (22)$$

The unperturbed isothermal function (14) gives a solution far from the black hole boundary,

$$F(m) = 0.5 f_0^{-2} \frac{1}{m}; \quad m \gg \frac{p}{Rt}$$

Substituting (22) into (18), we see that

$$S = (2)^3 2 R_{22} C(t) \quad (23)$$

The constraint on $C(t)$ is that the flux S is equal to the whole loss of mass:

$$S = \frac{d}{dt} \int_{m_t}^{\infty} d^3 Z \int_{m_t}^{\infty} dI \int_{m_t}^{\infty} dm \int_{m_t}^{\infty} dm_z f = (2)^3 \frac{d}{dt} \int_{m_t}^{\infty} 2m F(m) dm$$

Comparing this with (23) and calculating the integral, assuming the two asymptotes are equal at $m = m_1$, we obtain two equations:

$$C(t) m_1 \ln \frac{m_1}{m_t} = 0.5 f_0^{-2};$$

$$\frac{G}{C} m_1^2 \ln \frac{m_1}{m_t} - \frac{1}{2} = 2R$$

If the boundary does not move, $m_t = \text{const}$, then the approximate solution to the problem is

$$m_1(t) = \frac{p}{2 Rt} \ln^{-1/2} \frac{2 \frac{p}{Rt}}{m_t} \quad (24)$$

$$C(t) = \frac{f_0^{-2}}{4 \frac{p}{Rt}} \ln^{-1/2} \frac{2 \frac{p}{Rt}}{m_t} \quad (25)$$

Now, knowing the dependence of flux $S = \dot{M}_{bh}$ on time, we calculate the expression for mass of black hole.

$$M_{bh} = (2)^3 f_0^{-2} \frac{p}{Rt} \int_{m_t}^{\infty} \frac{1}{\ln \frac{2 \frac{p}{Rt}}{m_t}} dm$$

¹We neglect here small values of I , for which the terms with R_{12}, R_2 may also be significant (although, no more than of the same order as the term with R_{22}). However, it is easy to check that in the vicinity of the boundary (and hence, everywhere) the adiabatic invariant of each star could only grow with time (i.e., $S_I < 0$), and thus, contribution of small values I in the flux S and in the integrated function F is not large.

Substituting value of f_0 , we find

$$M_{bh} \approx \frac{2.6 \frac{p}{Rt}}{G \ln \frac{2 \frac{p}{Rt}}{m_t}} \quad (26)$$

The remarkable fact is that the result depends only weakly on the boundary m_t . This is why the solution (24)–(26) remains the same (up to the same accuracy) if we account the change of m_t with time ((15) or (16)) according to change of the black hole mass (26).

Taking $t = 15, 3 \cdot 10^8$ s (the Universe lifetime, and mass of star M equal to solar mass, we obtain the mass of black hole at arbitrary time:

$$M_{bh} \approx 1.2 \cdot 10^6 \frac{1}{200 \text{ km/s}} \left(\frac{t}{3 \cdot 10^8 \text{ s}} \right)^{3/2} \quad (27)$$

The contemporary consumption rate of stars by black hole is (23)

$$S = \frac{M_{bh}}{2t} \approx 6 \cdot 10^6 \text{ year}^{-1} \frac{1}{200 \text{ km/s}} : \quad (28)$$

It decreases in time as $t^{-1/2}$, and corresponds to $3.6 \cdot 10^6$ erg/s at contemporary time if $v = 200 \text{ km/s}$. Occasionally, at $v = 200 \text{ km/s}$ tidal and gravitational limits of momentum m_t and m_g corresponding to the mass (27) are close to each other,

$$m_g \approx 2.7 \cdot 10^3 \frac{p}{Rt} \frac{1}{200 \text{ km/s}} ; m_t \approx 4 \cdot 10^3 \frac{p}{Rt} \frac{1}{200 \text{ km/s}} :$$

The corresponding m_1 is

$$m_1 \approx 0.8 \frac{p}{Rt} :$$

We note that mass of black hole is no more than half the mass that was initially (when there was isothermal distribution) in the area $m = m_1$. The same is correct for any time (excluding the very beginning). This proves our initial assumption that change of distribution function does not affect significantly the coefficients R_p .

Possible constraint.

Let us now discuss briefly possible constraints on diffusion model. All equations beginning with (6) are averaged over orbital period. The boundary condition (17) also assumes that all particles with average angular momentum less than m_t become captured into the black hole. This means that during one period the change of trajectory is negligible. But for very elongated and large trajectories change of angular momentum of particle during one orbital period is comparable with the boundary angular momentum m_t . Hence, whether the star will or not be captured by black hole during one period depends not on the mean trajectory, but on its small fluctuations caused by individual collisions.² In terms of adiabatic invariants this may be formulated as follows.

²The effect was first pointed out in [4, 6]. It was then formulated in terms of critical energy, or critical radius in real space.

The diffusion approximation and equation (8) are applicable at any point of the momenta space. However, the boundary condition (17) may not be applied at very large values I (i.e., for very long orbits). Namely, mean change of angular momentum Δm during one period $T(I; m_t)$ is of the order of $m \frac{q}{RT(I; m_t)}$, hence the constraint on (17) is

$$q \frac{1}{RT(I; m_t)} - q \frac{1}{RT(I_{crit}; m_t)} = m_t$$

On the other hand, we have seen in Section 3 that main changes in the distribution function, and hence main contribution to the flux occur at scales $I \sim \sqrt{Rt}$ where t is the age of the black hole. This means that the consumption rate of stars is not disturbed by influence of I_{crit} if

$$I \gg \sqrt{Rt} \gg I_{crit}$$

. At sufficiently large values $I \gg m_t$ the orbital period is approximately equal to $T(I; m_t) \approx I^2$ (see Appendix). Substituting m_t from (15) we obtain that one anyway should not take the existence of I_{crit} into account if the black hole mass is greater than

$$M_{bh} \gg \frac{1}{50} \frac{t}{3} \frac{200 \text{ km/s}}{10^3 \text{ s}}^{3/8} \approx 10^6 M_\odot \quad (29)$$

Nowadays, this corresponds to minimal $M_{bh} \approx 2 \cdot 10^6 M_\odot$. Comparing the time dependence in (29) with that in (27) we find that (if $\dot{M}$ has not changed dramatically) the masses of the black holes that have grown as a result of the described relaxation mechanism have always been greater than the critical limit as far as galaxies exist. On the other hand, for black holes with sufficiently small masses additional investigations of influence produced by I_{crit} are needed.

5. Discussion

In the paper we derived the equations describing kinetic relaxation of two-component (stars + dark matter) system in the vicinity of supermassive black hole, in self-consistent gravitational potential. The equations have been expressed in terms of Hamiltonian variables.

We have analyzed the growth of black hole caused by consumption of stars, without taking non-baryon dark matter and luminous gas into account. We have seen that for black holes smaller than $10^6 M_\odot$ (29) individual collisions near pericentre might play crucial role. For black holes larger than $10^8 M_\odot$ (20) one should take the potential of the black hole into account. However, the most part of discovered black holes [11] have masses in the range between $10^6 M_\odot$ and $10^8 M_\odot$ where the main mechanism leading to consumption of stars is diffusion in the momenta space, or in the space of adiabatic invariants.

Analogous mechanism has been discussed previously in a set of papers [3, 4, 5, 6, 7]. But all of them have examined stationary distribution in the Coulomb potential of the black hole. We have considered the evolution in self-consistent potential and found that there is no stationary state in presence of black hole. Also in previous papers only diffusion along one axis (either m or E) was taken into account. The advantage of Hamiltonian variables $m; I$ is that in this variables the diffusion along m axis contains both shifts

along m - and E -axes in the $(E; m)$ plane and m makes the main contribution into the flux on the black hole. The diffusion along I axis plays less important role in the absorption process, being merely responsible for change of form of stars trajectories. Flux along I -axis increases smoothly as m decreases, reaching its maximum at the black hole boundary $m = m_t$, where $S_I = S_m = 0.6$ (i.e., the particles in the $I-m$ plane intersect the black hole boundary $m = m_t$ at the same angle independently on I).

The solution (27) obtained in Section 4 gives reasonable estimate for contemporary characteristic black hole mass which lies in the middle of observed range of masses. The estimate for flux of matter onto a black hole (28) also corresponds to average observed activity of galactic centers.

Equation (27) also shows that M_{bh} should change very slowly in time, $M_{bh} \propto \frac{P}{t} (1+z)^{3-4}$. Hence, the corresponding difference between black hole masses in far and nearby galaxies is too small and hidden by other effects. On the other hand, the obtained M relation differs from that derived from observations (from 3.7 in [16] to 5.3 in [17]) However, this discrepancy is not critical, since black hole masses depend not on one parameter only, but also on the bulge mass [11], mass of halo [18, 12], amount of interstellar gas etc. For example, the contribution of dark matter could increase the resulting black hole mass in those galaxies where amount of dark matter in the central parts is significant.

In the previous paper [12] we have discussed the growth of black hole caused by consumption of dark matter scattered by stars. The result was that the characteristic black hole mass is

$$M_{bh(dark)} = 8 \cdot 10^6 \frac{M_H^{1/2}}{10^{12} M} \frac{R_H^{9/14}}{100 \text{ kps}} \frac{t^{4/7}}{200 \text{ km/s}} \frac{1}{3 \cdot 10^8 \text{ s}}; \quad (30)$$

where M_H and R_H are the mass and the radius of the dark matter halo.

This result and solution (27) correspond to two limit cases describing the black hole growth caused either by dark or by luminous matter consumption. We see that contributions of stars and dark matter into the black hole growth are comparable for galaxies where densities of dark and luminous matter in the bulge are of the same order. But dark matter plays in this case a dominant role. This, in particular, means that in galaxies where observations show very small amount of dark matter the black hole mass should be in average smaller than in others. With account of real variations of all parameters (27) and (30) cover the range

$$5 \cdot 10^5 \leq M_{bh} \leq 2 \cdot 10^6 \quad (31)$$

which includes the most part of observed black holes.

The common point of view is that the consumption of stars by a black hole in the galactic centre is the main source of energy providing activity of the galactic nuclei [19, 20]. To verify this hypothesis we compared the observed X-ray luminosity of the galactic centres with the central black hole masses for 46 galaxies (Table 1). From the Table it follows that:

1. The most part of the black holes have masses in the range (31) restricted by (27) and (30).

2. The observed intensity of X-ray emission is in a reasonable agreement with the mean energy released as a result of stars consumption (28).
3. The analysis of the data shows very strong variations in observed luminosity of galactic nuclei. But no correlation with the black hole masses was found. This indicates that the luminosity of one nucleus should vary strongly in time. In other words, the duration of energy emission following a star absorption is much less than the mean time interval between two absorption events $T_{\text{M}} = S \approx 2000$ years (28). From observed variations (3-4 orders of magnitude), assuming an exponential fall of emission in time, we find an estimate $\tau = 10 - 200$ years. It follows that the most part of galaxies containing supermassive black holes are not active at present (as, e.g., Milky Way). On the other hand, maximal intensity of emission caused by star consumption should be much more than the average value S obtained in (28).
4. In Fig.1, the histogram of average luminosity versus the black hole mass is presented. The histogram is based on Table 1. It shows clearly that the luminosity breaks sharply at $M_{\text{bh}} > 3 \cdot 10^6 M_{\odot}$. This corresponds to the fact mentioned in Section 3 that for such black holes the gravitational radius is greater than the tidal radius, and stars sink into the black hole without disruption and energy emission. The observed cutoff proves that it is the infall of stars (not gas etc.) that provides the main contribution in the emission activity. On the contrary, in quasars with large redshifts the main mechanism causing energy emission may be consumption of gas [22].

Thus, in this paper and in [12] we present the theory of growth of supermassive black holes in the galactic centers caused by Coulomb scattering on stars and further consumption of stars and dark matter. The theory is in reasonable agreement with observations.

We thank Prof. A.M. Cherepashchuk and Prof. A.V. Zasov for valuable discussions. The research was supported by RAS Program "Mathematical methods in nonlinear dynamics" and by the President of RF grant "Support of Scientific schools" N Sh-1603.2003.2.

Table 1.

The X-ray luminosities and the central black hole masses of galactic nuclei.³

Name	L , erg/s	M_{bh} , $M_{\odot}$
3C 120 (Mrk 1506)	9.772 10^{43} 1.343 10^{44}	2.3 10^6
Ark 120 (Mrk 1095)	7.805 10^{43}	1.84 10^6
Circinus	2.47 10^{43}	1.3 10^6
Fairall 9	6.688 10^{43}	8.0 10^6
IC 1459	8.398 10^{43} 1.051 10^{44}	3.7 10^6
IC 4329A	3.748 10^{43} 2.541 10^{44}	5 10^6
UGC 3973 (Mrk 79)	2.594 10^{43}	5.2 10^6
Mrk 110	1.625 10^{43}	5.6 10^6
Mrk 335	2.04 10^{43} 4.916 10^{43}	6.3 10^6
Mrk 509	1.188 10^{43}	5.78 10^6
Mrk 590 (NGC 863)	3.853 10^{43} 8.884 10^{43}	1.78 10^6
NGC 205 (M 110)	< 1.172 10^{38}	< 9.3 10^6
NGC 598 (M 33)	1.46 10^{38} 1.902 10^{38}	< 1.5 10^6
NGC 1068 (M 77)	1.315 10^{43} 5.124 10^{43} 8.81 10^{43}	1.6 10^6
NGC 3115	1.773 10^{38}	9.1 10^6
NGC 3227	1.371 10^{43} 7.127 10^{43}	3.9 10^6
NGC 3516	5.578 10^{43} 1.084 10^{44}	2.3 10^6
NGC 3608	1.174 10^{43}	1.1 10^6
NGC 3783	8.515 10^{43} 7.384 10^{43}	9.4 10^6
NGC 3998	1.425 10^{43}	5.6 10^6
NGC 4051	7.609 10^{43} 7.624 10^{43}	1.3 10^6
NGC 4151	9.025 10^{43} 5.578 10^{43} 4.784 10^{43}	1.53 10^6

³The black hole masses are adapted from [11], the X-ray luminosities { from [21]. Only galaxies with measured X-ray luminosities are taken from the sample of [11].

NGC 4203	7:471	10	< 1.2	10
	6:424	10		
NGC 4258 (M 106)	2:914	10	4.1	10
NGC 4261 (3C 270)	1:575	10	5.2	10
	1:434	10		
NGC 4291	6:128	10	1.5	10
	6:769	10		
	4:28	10		
NGC 4342	2:077	10	3.4	10
NGC 4374 (M 84)	6:317	10	1.6	10
	6:657	10		
	6:138	10		
NGC 4459	4:169	10	6.5	10
NGC 4473	1:109	10		
	5:089	10	1.0	10
NGC 4486 (M 87)	3:257	10	3.4	10
NGC 4593	6:535	10	8.1	10
NGC 4594 (M 104)	3:361	10	1.1	10
NGC 4649	1:002	10	2.0	10
	1:587	10		
NGC 4697	9:141	10	1.2	10
NGC 4945	5:447	10	1.1	10
NGC 5548	2:182	10	1.23	10
	2:154	10		
	2:778	10		
NGC 6251	1:71	10	5.4	10
NGC 7469	1:699	10	6.5	10
	3:34	10		
	2:071	10		
PG 0026+ 129	2:798	10	4.5	10
PG 0052+ 251	4:766	10	2.2	10
PG 1211+ 143	1:098	10	4.05	10
PG 1411+ 442	5:049	10	8.0	10
PG 1426+ 015 (M rk 1383)	8:507	10	4.7	10
PG 1613+ 658 (M rk 876)	1:652	10	2.41	10
PG 1617+ 175 (M rk 877)	6:188	10	2.73	10

Appendix.

Calculating the coefficients R_k, R_{kp} in (7) and the adiabatic invariant in the case of isothermal stellar distribution function.

A number of simplifications may be made for any isotropic distribution function. We first discuss the coefficients $W_{kp}; W_k$ introduced in (4). If the distribution function f is isotropic at any point of coordinate space, i.e. does not depend on momentum, but only on energy, then tensor W_{kp} depends on only one vector v , and hence takes the form

$$W_{kp} = A(E; r) \delta_{kp} + B(E; r) \frac{v_k v_p}{v^2}; \quad (32)$$

where, calculating convolution W_{kk} and product $W_{kp} v_p$, we find

$$A = \frac{8}{3} \int_0^\infty dE^0 f(E^0) \begin{cases} \frac{3}{2} \frac{v^0}{v} - 1 & \frac{v^0}{3v^2} & ; E^0 < E \\ 1 & ; E^0 > E \end{cases}$$

$$A - B = \frac{8}{3} \int_0^\infty dE^0 f(E^0) \begin{cases} \frac{v^0}{v^3} & ; E^0 < E \\ 1 & ; E^0 > E \end{cases}$$

$$v = \frac{1}{2} \frac{1}{(E - (r))}; \quad v^0 = \frac{1}{2} \frac{1}{(E^0 - (r))}$$

and similarly

$$W_k = \frac{v_k}{v} D(E; r); \quad (33)$$

$$D(E; r) = \frac{8}{3} \int_0^\infty dE^0 \frac{\partial f}{\partial E^0} v \begin{cases} (v^0 - v)^3 & ; E^0 < E \\ 1 & ; E^0 > E \end{cases}$$

To calculate the coefficients R_k, R_{kp} in (7) it is useful to introduce new variables $\epsilon = f_1; g = fE; m g$. Then, since $I_i = I_i(E; m)$, we have from (7)

$$R_{ij} = \frac{\partial I_i}{\partial \epsilon_a} \frac{\partial I_j}{\partial \epsilon_b} R_{ab}^0; \quad R_i = \frac{\partial I_i}{\partial \epsilon_a} R_a^0;$$

For brevity, we denote

$$\epsilon_0 = 2 G^2 M; \quad \langle \epsilon \rangle = \frac{1}{(2\pi)^3} \int d^3$$

Then, making use of (32) and (33), we obtain the expressions for $R_a^0; R_{ab}^0$:

$$R_1^0 = \epsilon_0 \langle v D \rangle; \quad R_2^0 = \epsilon_0 m \langle D = v \rangle; \quad (34)$$

$$R_{11}^0 = \epsilon_0 \langle v^2 (A - B) \rangle; \quad R_{12}^0 = \epsilon_0 m \langle A - B \rangle; \quad R_{22}^0 = \epsilon_0 \langle A r^2 - B m^2 = v^2 \rangle$$

Note that because of spherical symmetry of all coefficients, only one integral remains in the averaging:

$$\langle \epsilon \rangle = \frac{2}{T(E; m)} \int_0^\infty \frac{dr}{v_r}; \quad T(E; m) = \int \frac{dr}{v_r}; \quad v_r = \sqrt{\frac{2}{E - (r)} - \frac{m^2}{2r^2}}$$

All the previous results are applicable for any isotropic function. Now we substitute the isothermal distribution function (12). For calculations it is useful to introduce dimensionless variables:

$$= \frac{m}{2} e^{\frac{E-2r^2}{m^2}}; \quad x = re^{\frac{E-2r^2}{m^2}}$$

The condition $v_r^2 = 2(E - m^2/2r^2) > 0$ leads to

$$0 < x < 1; \quad 0 < \frac{p}{1-2e} < \dots$$

Then, for example, period of oscillations is

$$T(E; m) = \frac{p}{2} \int_0^1 \frac{dr}{\sqrt{E - \frac{m^2}{2r^2}}} = \frac{1}{m} e^{\frac{E-2r^2}{m^2}} T(x); \quad T(x) = \int_0^1 \frac{dx}{\sqrt{\ln x - \frac{1}{2}x^2}}$$

The remarkable fact is that $T(x)$ is almost constant! It grows very slowly from 1.77 at $x=0$ to 1.9 at $x=1$ - maximum possible.

The averaging procedure now can be rewritten as

$$\langle \dots \rangle = \frac{2}{T(E; m)_r} \int_0^1 \frac{dr}{v_r} (\dots) = \frac{1}{T(x)_0} \int_0^1 \frac{dx}{\sqrt{\ln x - \frac{1}{2}x^2}} (\dots)$$

We also can calculate A, B and D for the isothermal distribution function:

$$A = \frac{8}{3} \frac{f_0}{r^2} (2H + x^2) - H + x^2; \quad A - B = \frac{8}{3} \frac{f_0}{r^2} (2H + x^2);$$

$$D = v = \frac{8}{3} \frac{f_0}{r^2} (2H + x^2); \quad v^2 = 4x^2 \ln x; \quad H(x) = \int_x^1 y dy \frac{\ln y}{\ln x}$$

Note that

$$A - B = -2D = v$$

which leads to (11).

From (34) we now find

$$\begin{aligned} R_{11}^0 &= -2R_1^0; \quad R_{12}^0 = -2R_2^0 \\ R_1^0(E; m) &= -4 \int_0^1 e^{\frac{E-2r^2}{m^2}} \frac{8}{3} f_{0-1}(r) dr \\ R_2^0(E; m) &= -m \int_0^1 e^{\frac{E-2r^2}{m^2}} \frac{8}{3} f_{0-2}(r) dr \\ R_{22}^0(E; m) &= -2 \int_0^1 \frac{8}{3} f_{0-22}(r) dr \end{aligned}$$

where

$$f_{0-2}(x) = \langle 2H(x) - x^2 \rangle + 1; \quad f_{0-1}(x) = \langle \ln x - 2H(x) \rangle + \langle \ln x \rangle$$

$$\sigma_{22}(\epsilon) = \langle \frac{1}{2} \ln x (2H + x^2) - H + x^2 \rangle = \frac{1}{2} \langle 4H - x^2 + 1 + 3H - (x^2 \ln x) \rangle$$

Up to at least 10^{-4} one can use approximate expressions:

$$\sigma_{22}(\epsilon) \approx 1 + 0.35 \epsilon^{0.6};$$

$$\sigma_{11}(\epsilon) \approx 0.5 - 0.8 \ln \left(\frac{\sigma_{22}(\epsilon) - 1}{0.5} \right) \approx 0.5 - 0.3 \ln \epsilon^{0.6};$$

$$\sigma_{22}(\epsilon) \approx 0.87$$

Finally, we calculate the adiabatic invariant I (5):

$$I(E; m) = \frac{2}{\pi} e^{E=2\epsilon^2} \int_0^{z_1} \frac{dx}{\ln x - \epsilon^2 x^2} \quad (z_1 = \ln x - \epsilon^2 x^2)$$

Calculating the integral we obtain

$$I(E; m) \approx \frac{2}{\pi} e^{E=2\epsilon^2} (0.9 - \epsilon^2) = \frac{2}{\pi} (0.9 - \epsilon^2) m$$

Hence,

$$I_m^0 \approx \frac{2}{\pi}; \quad I_E^0 \approx \frac{I + \frac{2}{\pi} m}{2\epsilon^2};$$

The distribution function can now be expressed in terms of adiabatic invariants:

$$f(I; m) = f_0 e^{-E=\epsilon^2}, \quad f_0 = \frac{2}{\pi} \frac{0.9^2}{(I + \frac{2}{\pi} m)^2}$$

Another useful formula is the expression for the period $T(E; m)$ formulated in terms of $I; m$:

$$T(I; m) = \frac{T(\epsilon)}{\pi} e^{E=2\epsilon^2}, \quad \frac{2}{\pi} \frac{I + \frac{2}{\pi} m}{0.9}.$$

References

- [1] Chakrabarti S.K. (Ed.) *Observational Evidence for Black Holes in the Universe* (Astrophys. and Space Sci. Library, Vol. 234) (Dordrecht: Kluwer Acad. Publ., 1999); Kaper L., van den Heuvel E.P.J., Woudt P.A. (Eds) *Black Holes in Binaries and Galactic Nuclei: Diagnostics, Demography, and Formation: Proc. of the ESO Workshop, Garching, Germany, 6-8 Sept. 1999, in Honour of Riccardo Giacconi* (ESO Astrophys. Symposia) (Berlin: Springer, 2001)
- [2] Dokuchaev V.I., Eroshenko Yu.N. 2002, astro-ph/0209324v1
- [3] Peebles P.J.E. 1972, *ApJ*, 178, 371
- [4] Bahcall J.N., Wolf R.A. 1976, *ApJ*, 209, 214
- [5] Frank J., Rees M.J. 1976, *MNRAS* 176, 633

- [6] Lightman A.P., Shapiro S.L. 1977, *ApJ*, 211, 244
- [7] Dokuchaev V.I., Ozernoy L.M. 1977, *Sov. Phys. JETP* . 73, 5, 1587
- [8] Gurevich A.V. 1964, *Geomagnetism and Aeronomy*, 4, 247
- [9] Gurevich A.V., Zybin K.P. 1995, *Sov. Physics-Uspokhi* 38 (7) 687-722 ;
Gurevich A.V., Zybin K.P. 1988, *Sov. Phys. JETP* (67), 1957
- [10] Gurevich A.V., Zybin K.P. 1990, *Sov. Phys. JETP* (70), 10
- [11] Cherepashchuk A.M. 2003, *Sov. Phys. Uspokhi*, 173 (4), 345
- [12] Ilyin A.S., Zybin K.P., Gurevich A.V. 2003, *Sov. Phys. JETP* 98 1, 1-16
(astro-ph/0306490)
- [13] Lifshitz E.M., Pitaevskii L.P., *Physical Kinetics*. Oxford: Pergamon press. 1981
- [14] Budker G.I., Belyaev S.T., *Plasma Physics and Controllable fusion reaction*. 1958 (in russian)
- [15] Landau L.D., Lifshitz E.M., *The classical Theory of Fields*. Oxford: Pergamon press. 1981
- [16] Gebhardt K., Bender R., Bower G., Dressler A., Faber S.M. et al. 2000 *A & A*, 539, L13
- [17] Ferrarese L., Merritt D. 2000, *ApJ*. 539 L9
- [18] Baes M., Buyle P., Hau G.K.T., Dejonghe H. 2003, *MNRAS* 341, L44
- [19] Shapiro S.L. 2003, *Carnegie Observatories Astrophysical Series*, Vol. 1;
astro-ph/0304202 v1
- [20] Merritt D., Ferrarese L. 2001, astro-ph/0107134v2
- [21] NASA/IPAC Extragalactic Database (NED), <http://nedwww.ipac.caltech.edu>
- [22] Vestergaard M., 2004, astro-ph/0401430

This figure "fig1.png" is available in "png" format from:

<http://arxiv.org/ps/astro-ph/0403023v1>