

Large-scale wind structures in O B supergiants: a search for rotationally modulated H α variability?

T. Morel,^{1,2} Y. S. V. M archenko,³ A. K. Pati,⁴ K. K uppuswamy,⁵ M. T. Carini,³ E. W ood³ and R. Zimm erman³

¹ Istituto Nazionale di Astrofisica, Osservatorio Astronomico di Palermo G. S. Vain a, Piazza del Parlamento 1, I-90134 Palermo, Italy

² Inter-University Centre for Astronomy and Astrophysics (IUCAA), Post Bag 4, Ganeshkhind, Pune, 411 007, India

³ Department of Physics and Astronomy, Western Kentucky University, 1 Big Red Way, Bowling Green, KY 42101(3576, USA

⁴ Indian Institute of Astrophysics (IIA P), III block, Koramangala, Bangalore, India

⁵ Vainu Bappu Observatory, Indian Institute of Astrophysics (IIA P), Kavalur, Angayam 635 701 India

Accepted ????. Received ???; in original form ???

ABSTRACT

We present the results of a long-term monitoring campaign of the H α line in a sample of bright O B-supergiants (O 7.5{B 9) that aims at detecting rotationally modulated changes potentially related to the existence of large-scale wind structures. A total of 22 objects were monitored during 36 nights spread over 6 months in 2001{2002. Co-ordinated broad-band photometric observations were also obtained for some targets. Conspicuous evidence for variability in H α is found for the stars displaying a feature contaminated by wind emission. Most changes take place on a daily time-scale, although hourly variations are also occasionally detected. Convincing evidence for a cyclical pattern of variability in H α has been found in 2 stars: HD 14134 and HD 42087 (periodic signals are also detected in other stars, but independent confirmation is required). Rotational modulation is suggested from the similarity between the observed recurrence time-scales (in the range 13{25 days) and estimated periods of stellar rotation. We call attention to the atypical case of HD 14134 which exhibits a clear 12.8-d periodicity both in the photometric and in the spectroscopic data sets. This places this object among a handful of early-type stars where one may observe a clear link between extended wind structures and photospheric disturbances. Further modelling may test the hypothesis that azimuthally-extended wind streams are responsible for the patterns of spectral variability in our target stars.

Key words: stars: early-type { stars: supergiants { stars: emission-line { stars: winds, outflows { stars: rotation.

1 INTRODUCTION

One of the major legacies of the International Ultraviolet Explorer (IUE) was to establish the ubiquity of UV line-profile variability in O B stars. The pattern of variability predominantly consists of extra absorption features (the so-called ‘discrete absorption components’; hereafter DACs) moving blueward across the (unsaturated) absorption troughs of P Cygni lines (e.g., Kaper et al. 1996). The repetitive nature of DACs and the fact that the recurrence time-scale is commensurate with the (otherwise ill-defined) stellar rotational

period suggest that rotation plays a key role in inducing the patterns of variability (e.g., Kaper et al. 1999). UV time-series with a long temporal baseline (i.e., sampling several rotational periods) have revealed that O B stars exhibit a great diversity of behaviours that cannot always be strictly related to the stellar rotation period (e.g., Howarth, Prinja & Massa 1995; Prinja, Massa & Fullerton 1995; Prinja, Massa & Fullerton 2002). Nevertheless, this global phenomenon has been successfully interpreted in several cases in the framework of a model in which the changes are caused by the rotational modulation of large-scale, azimuthally-extended wind streams (e.g., Fullerton et al. 1997).

The hydrodynamical simulations of Cranmer & Owocki (1996) suggest that large-scale photospheric perturbations (e.g., bright ‘spots’ causing a local enhancement of the radiative driving force) may lead to the development of these

² Based on observations collected at the Vainu Bappu Observatory (Kavalur, India).

Y email: morel@astropa.unipa.it

structures. Although the underlying physical mechanism remains elusive, pulsational instability can be invoked as a trigger of the wind variability (e.g., Fullerton, Gies & Bolton 1996). Large-scale or ‘surface’ magnetic field structures could be a viable alternative in view of the growing evidence for (weak) magnetic fields in OB stars (e.g., Henrichs, Neiner & Geers 2003). Photospheric perturbations arising from these two phenomena are likely to have a significant impact on the global wind morphology (e.g., ud-Doula & Owocki 2002).

With the demise of IUE, further progress on this issue is likely to stem in the foreseeable future from time-resolved studies of line variability in the optical. Firstly, intensive UV observations have been carried out over several rotation periods for few stars (Mossa et al. 1995; Prinja et al. 1998; de Jong et al. 2001). For practical reasons, an optical survey is much more amenable at collecting high-quality observations covering several cycles for a large, representative sample of OB stars. Secondly, in contrast to the UV resonance lines, optical recombination lines are (at least partly) formed in OB supergiants in the dense, strongly accelerating part of the outflow close to the stellar core. In virtue of their density-squared dependence, they are good probes of the large density gradients expected to prevail in the co-rotating interaction regions (Cranmer & Owocki 1996). Revealing a cyclical pattern of variability in these lines would give further credence to the idea that the origin of anisotropic outflows in OB stars is deep-seated and causally linked to processes taking place at the stellar photosphere. Here we discuss the temporal behaviour of H β , which is typically formed at $r \approx 1.5 R_*$ in O-type stars (e.g., Prinja, Fullerton & Crowther 1996). Although the line-formation region is likely to be more extended in B-type supergiants, this transition still probes the few inner radii of the stellar wind.

Previous surveys have been very useful in documenting H β line-profile variability in OB stars (e.g., Ebbets 1982), but frequently suffered from a poor temporal sampling hampering the study of the line-profile variations on a rotational time-scale (up to several weeks in the case of B supergiants). Despite the persistent and numerous searches for optical line-profile variability in OB stars, evidence for rotational modulation has been found in only few cases (e.g., Mot & Michaud 1981; Kaper et al. 1997, 1998). In the UV and optical domains the time-scales associated with the changes are often identical, within the uncertainties (e.g., Howarth et al. 1995; Kaper et al. 1999; de Jong et al. 2001; Kaufer, Prinja & Stahl 2002). This suggests that H β variability and DACs diagnose the same underlying physical phenomenon.

The long-term monitoring of the H β line presented in this paper aims at better establishing the incidence of rotationally modulated variability in OB supergiants. To potentially link the photospheric and wind activity, coordinated broad-band photometric observations were also carried out for a subset of this sample. Our survey primarily targets early B-type stars and can thus be regarded as complementary to others that concentrate on O-type (Kaper 1998; Kaper et al. 1998) or late B- to early A-type supergiants (Kaufer et al. 1996). The results presented in the following supersede previous preliminary reports (Morel et al. 2004a, b).

2 TARGET SELECTION AND TEMPORAL SAMPLING OF THE OBSERVATIONS

Our targets are drawn from a magnitude-limited ($V < 7.5$) sample of early-type supergiants (O7.5(B9) with evidence for an emission-like H β profile or, in the case of stars with an absorption feature, for previous claims of a wind-related variability (e.g., filling-in of the profile). Known close binaries were discarded from this list in order to avoid contamination from the wind collision effects (e.g., Thaller et al. 2001). The wide binary orbit of HD 37742 rules out the formation of a bow shock created from the interaction with an O-type companion (Hummel et al. 2000). In our data only HD 13854 and HD 47240 present significant velocity variations of the photospheric H ϵ 6678.15 line which might be attributed to binary motion (see below). Dwarfs and giant stars were not included in this list, as H β is of photospheric origin in the vast majority of these objects. The only exception was the O7.5 giant HD 24912 (Per) because of previous claims of a cyclical, wind-related pattern of variability in H β (Kaper et al. 1997; de Jong et al. 2001). For the other stars in our list, no clear evidence for cyclical line-profile variations in optical wind lines was found prior to our survey. Some physical parameters of our program stars are given in Table 1.

The basic requirement of this survey was to achieve a time sampling allowing a proper assessment of the variability pattern on a rotational time-scale (typically 1–2 weeks). Although the rotational periods of our program stars are unknown, the critical and projected rotational velocities can be used, along with the estimates of the stellar radii, to set a lower and an upper limit on this quantity. As can be seen in Table 2, the estimated rotational periods span a wide range of values (1.2–144 d). Our 22 program stars were monitored during a total of 36 nights spread over 6 months, with 1–2 spectra obtained every night. This long-term monitoring was necessary to cover at least one rotational cycle for the slowly rotating objects. We also obtained intranight observations, expecting significant hourly changes in some cases (e.g., HD 24912). In general, this temporal sampling appears to be adequate for our purpose (see Table 3). For 5 stars, however, the observations were too sparse to reliably investigate a potential cyclical behaviour (HD 14818, HD 21291, HD 119608, HD 151804, and HD 152236). No period search was performed in this case (see Section 5.2), but a qualitative description of the changes is presented in Section 4.

3 OBSERVATIONS AND REDUCTION PROCEDURE

3.1 Spectroscopy

Our observations were obtained during 4 observing runs conducted between 2001 November and 2002 May at the 40-inch telescope of the Vainu Bappu Observatory (Kavalur, India). The spectrograph was equipped with a TK1k CCD chip. We used the 651 l/m grating blazed in the 1st order at 7500 Å, which yields at H β a reciprocal dispersion of 1.45 Å pixel⁻¹ (we used a 120 μ m-wide slit), thus covering the spectral region 5810–7205 Å. The slit of the spectrograph was rotated for HD 37742 in order to avoid contamination from the visual companion. The speckle interferometric ob-

Table 1. Physical parameters of the program stars: spectral types and wind terminal velocities, v_1 , from Howarth et al. (1997) and references therein; effective temperatures from the calibration of Humphreys & McElroy (1984); stellar luminosities calculated from the calibrated M_V values and bolometric corrections of Humphreys & McElroy (1984); stellar masses from Lamers (1981) (the numbers in brackets are estimates based on his values for similar spectral types and luminosity classes). This source was used because it constitutes the most homogeneous database, but we warn the reader that these values may be significantly revised (see, e.g., Herrero, Puls & Najarro 2002); stellar radii determined from the effective temperatures and luminosities; mass-loss rates, $\dot{M}$, calculated from the empirical relation for galactic OBA supergiants of Lamers & Cassinelli (1996) (the value for HD 24912 was derived from their relation for giants); projected rotational velocities, $v \sin i$, from Howarth et al. (1997); critical rotational velocities, v_{crit} , calculated following Howarth & Prinja (1989) (the values in brackets are poorly determined because of an uncertain stellar mass).

Name	V (mag)	Spectral type	T_e (K)	$\log(L/L_\odot)$	$\dot{M}$ ($M_\odot$)	R ($R_\odot$)	$\dot{M}$ ($10^{-6} M_\odot \text{ yr}^{-1}$)	v_1 (km s^{-1})	$v \sin i$ (km s^{-1})	v_{crit} (km s^{-1})
HD 13854	6.49	B1 Iab	20,260	5.18	(22)	32	0.76	920	97	(328)
HD 14134 (V 520 Per)	6.55	B3 Ia	16,300	5.24	24	52	1.45	465	66	265
HD 14818 (10 Per)	6.26	B2 Ia	18,000	5.40	(30)	52	2.13	565	82	(293)
HD 21291	4.25	B9 Ia ^a	10,250	4.86	19	85			30 ^b	195
HD 24398 (Per)	2.88	B1 Ib	20,260	4.90	21	23	0.23	1295	67	396
HD 24912 (Per)	4.04	O7.5 III (n) (f)	35,400	5.40	28	13	0.22	2330	213	550
HD 30614 (Cam)	4.30	O9.5 Ia	29,900	5.86	40	32	5.03	1590	129	350
HD 31327	6.09	B2 Ib	18,000	4.76	(20)	25			60	(377)
HD 37128 (Ori)	1.70	B0 Ia	28,600	5.78	38	32	3.15	1910	91	362
HD 37742 (OriA)	1.7	O9.7 Ib	29,250	5.40	25	20	1.05	1860	124	421
HD 38771 (Ori)	2.04	B0.5 Ia	23,100	5.50	35	35	1.37	1525	83	379
HD 41117 (2 Ori)	4.64	B2 Ia	18,000	5.40	32	52	2.36	510	72	305
HD 42087 (3 Gem)	5.76	B2.5 Ib	17,150	4.72	(19)	26	0.20	735	71	(359)
HD 43384 (9 Gem)	6.29	B3 Iab	16,300	4.88	(22)	35	0.30	760	59	(331)
HD 47240	6.18	B1 Ib	20,260	4.90	(21)	23	0.31	960	103	(396)
HD 52382	6.51	B1 Ib	20,260	4.90	(21)	23	0.33	900	74	(396)
HD 53138 (o ² CMa)	3.00	B3 Ia	16,300	5.24	23	52	0.78	865	58	258
HD 58350 (CMa)	2.40	B5 Ia	13,700	5.02	20	58	0.91	320	50	238
HD 91316 (Leo)	3.84	B1 Iab	20,260	5.18	22	32	0.63	1110	75	328
HD 119608	7.50	B1 Ib	20,260	4.90	(21)	23	0.34	880	74	(396)
HD 151804	5.24	O8 Iaf	33,500	6.06	(60)	32	11.3	1445	104	(416)
HD 152236 (1 Sco)	4.77	B0.5 Ia+	19,700 ^c	6.05 ^c	(35)	91 ^c	5.36	390	74	(99)

^a From the SIMBAD database.

^b From Abt, Levato & Grosso (2002).

^c From Rivinius et al. (1997).

servations of Mason et al. (1998) exclude the existence in HD 24912, HD 30614 and HD 151804 of bright ($m < 3$ mag), close companions ($0.035'' < \theta < 1.5''$). Apart from HD 37742, a search in the Hipparcos database reveals only one star in our sample with a close visual component (HD 42087: $H_p = 2.58$ mag and $\theta = 0.6''$).

The IRAF² tasks were used to carry out standard reduction procedures (i.e., bias subtraction, flat-fielding, extraction of the spectra, and wavelength calibration). Spectra of FeNe lamps were taken immediately before and/or after the stellar exposure. The spectra were subsequently put in the heliocentric velocity frame. The H γ line profiles were found to be significantly affected by telluric lines, as the observations were often carried out under unfavourable atmospheric conditions (e.g., high humidity). Therefore, we have obtained spectra of the rapid rotator HD 65810 (A1V) to create a library of telluric lines that would be used on the target stars. Since this procedure proved unsatisfactory, we

used instead the high-resolution telluric spectrum of Hinkle et al. (2000) degraded to our resolution. The water vapour lines in the pseudo-continuum regions around H γ were interactively shifted and scaled with the IRAF task telluric until the residuals between the observed spectra and the template telluric spectrum were minimized. The spectra were finally continuum-rectified by fitting a low-order spline3 polynomial to the line-free spectral regions on both sides of H γ . We generally obtained three consecutive exposures to allow a more robust rectification of the spectra and filtering of cosmic ray events. Any individual exposure with evidence for an imperfect continuum normalization was excluded during the co-adding operation. The typical signal-to-noise ratio (S/N) at H γ in the combined spectrum is quoted for each star in Table 3, and lies in the range 125–335 per pixel in the continuum. Large velocity variations of the photospheric HeI 6678 line, which might arise from binary motion, were found in HD 13854 and HD 47240 (15 and 30 km s^{-1} , respectively). The spectra were hence realigned and put in the same reference frame. The remaining stars may have companions (e.g., HD 37742), but the radial velocity variations are small, i.e., comparable to the typical uncertainties in the wavelength calibration ($\sim 5 \text{ km s}^{-1}$). Hence, all the discussed spectral variations should be intrinsic to the star.

² IRAF is distributed by the National Optical Astronomy Observatories, operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

Table 2. Summary of period search (all periods in days). $P_{\min}$ and $P_{\max}$: lower and upper limits on the stellar rotational period (determined from the critical and projected rotational velocities, along with an estimate of the stellar radius); P (phot): periodicities in the Hipparcos data (var: stars displaying photometric variability but without evidence for periodicity; non-var: stars without significant variability). The light curves are shown in Fig. 2 (and in Fig. 5 for HD 14134); P (literature): Literature values for the periodicities in H (Kaper et al. 1997, 1998; de Jong et al. 2001; Markova 2002). The numbers in italics refer to the periodicities in the UV wind line profiles (Kaper et al. 1999; de Jong et al. 2001). Multiple values refer to periods derived at different epochs. Numbers in brackets indicate that the periodicity found is longer than the time span of the IUE observations. For the H periodicities derived from our observations, P (H), '2D' and 'EW' indicate that the period is present in the H pixel-to-pixel or EW time-series, respectively.

Name	$P_{\min}$	$P_{\max}$	P (phot)	P (literature) ^a	P (H) ^b
HD 13854	(4.9)	16.5	5.644		1.047 (EW)
HD 14134 (V 520 Per)	10	40.1	12.823		22.2 (2D), 12.5 (2D and EW)
HD 14818 (10 Per)	(8.9)	31.8	2.754		No period search performed
HD 21291	22	144	26.76	[1,2,3]	No period search performed
HD 24398 (Per)	2.9	17.3	non-var		No periods found
HD 24912 (Per)	1.2	3.17	non-var	2.1 0.1, 1.96 0.11, 2.086 [4,5,6] 1.9 0.3, 1.0 0.1, 2.0 0.2	2.197 ? (2D and EW)
HD 30614 (Cam)	4.6	12.5	var	5.6, 7, 10 [5,6,7,8,9]	No periods found
HD 31327	(3.3)	20.8	non-var		No periods found
HD 37128 (Ori)	4.4	17.6	var	[8,10]	18.2, 0.781 (both 2D)
HD 37742 (OriA)	2.3	7.97	1.407	6 [5,6,8] (6.3 2.1), (6.1 2.7)	> 80 (2D), 50 (2D), 13.3 (EW) 1.136 (2D), 0.877 (2D) 4.76, 1.047 (both 2D)
HD 38771 (Ori)	4.7	21.4	var	[7,8,11]	
HD 41117 (2 Ori)	8.6	36.3	2.869	[1,8,12,13]	> 80 (2D), 40 (2D and EW), 0.957 (2D), 0.922 (2D)
HD 42087 (3 Gem)	(3.7)	18.5	6.807		25 (2D and EW)
HD 43384 (9 Gem)	(5.3)	29.7	13.7		No periods found
HD 47240	(2.9)	11.3	2.7424		No periods found
HD 52382	(2.9)	15.7	var		No periods found
HD 53138 (α^2 CMa)	10	45.7	var	[8,14]	No periods found
HD 58350 (CMa)	12	58.2	var	[8]	No periods found
HD 91316 (Leo)	4.9	21.3	var ^c	[12,13,15]	No periods found
HD 119608	(2.9)	15.7	non-var		No period search performed
HD 151804	(3.9)	15.5	non-var ^d	7.3 [6,16]	No period search performed
HD 152236 (1 Sco)	(46)	62.3	var ^e	[17,18]	No period search performed

^a The numbers in square brackets refer to previous investigations of H variability in our program stars: [1] Rosendhal (1973); [2] Denizman & Hack (1988); [3] Zeinalov & Rzaev (1990); [4] de Jong et al. (2001); [5] Kaper et al. (1997); [6] Kaper et al. (1998); [7] Ebbets (1980); [8] Ebbets (1982); [9] Markova (2002); [10] Cherrington (1937); [11] Rusconi et al. (1980); [12] Underhill (1960); [13] Underhill (1961); [14] van Helden (1972); [15] Smith & Ebbets (1981); [16] Prinja et al. (1996); [17] Sterken & Wolf (1978); [18] Rivinius et al. (1997).

^b The typical uncertainties for the quoted periods have been derived from the FWHM of the peaks in the PS and amount to: 20% (30 days ($P > 50$ d)), 14% (10% ($10 < P < 25$)), 0.01% (0.2 days ($1 < P < 5$)), and 0.01 days ($P < 1$ d)).

^c Multi-periodic variable according to Koen (2001).

^d Photometric variable according to Balona (1992), and references therein.

^e Irregular, multi-periodic photometric variable (Sterken, de Groot & van Genderen 1997).

3.2 Photometry

Coordinated B- and I-band (Johnson) photometric observations were carried out with the 24-inch telescope of the Bell Observatory operated by the Western Kentucky University (USA). Six stars were observed during 7 nights in January, February and April 2002: HD 13854, HD 14134, HD 14818, HD 42087, HD 47240, and HD 52382. A typical observation (repeated up to 3 times per night) comprises up to 10 short-exposure (1/10 sec) CCD images of the target and the surrounding field (0.53'' pixel⁻¹ scale, providing an 4.4'' × 6.6'' field).

The standard processing routines were performed with IRAF tasks and included bias and flat field correction, followed by measurements of all available comparison stars in the fields surrounding the targets. Pre-selection of appropriate comparison stars was based on their brightness and non-variability. All the differential measurements performed

on the stars from individual frames were combined into normal observation, thus producing 1/3 data points per night for each star. This resulted in overall accuracy 0.01 mag per data point for each comparison star. This relatively low accuracy was mainly dictated by the lack of appropriately bright comparison stars, the problem being especially acute for HD 14818, HD 42087, and HD 47240. For HD 42087, in particular, there was only one very faint comparison star in the field, resulting in a much lower accuracy (0.02 mag). These data are not discussed in the following.

4 OVERVIEW OF THE SURVEY AND QUANTITATIVE ANALYSIS

The H time-series are shown in Fig. 1, along with the temporal variance spectra (TVS; Fullerton et al. 1996). As can be seen, all stars display significant variability across most

Table 3. Temporal sampling of the spectroscopic observations. N : total number of spectra obtained; T : total time span of the observations; M in (t), M ean (t), and M ax (t) : m in im u m , m ean, and m ax im u m tim e intervals between consecutive spectra, respectively.

Na m e	Run ^a	Nb. nights	N	< S=N >	T (d)	M in (t) (d)	M ean (t) (d)	M ax (t) (d)
HD 13854	A ,B ,C	16	19	210	71.1	0.07	3.95	24.0
HD 14134 (V 520 Per)	A ,B ,C	17	22	205	67.1	0.07	3.19	24.0
HD 14818 (10 Per)	C	7	7	170	12.0	0.93	2.01	4.06
HD 21291	C	7	7	190	8.01	0.93	1.33	2.97
HD 24398 (Per)	A ,B	9	13	305	38.8	0.07	3.24	24.9
HD 24912 (Per)	A ,B ,C	17	23	250	75.8	0.07	3.45	26.0
HD 30614 (Cam)	A ,B ,C	16	19	240	70.9	0.07	3.94	26.0
HD 31327	A ,B	10	12	265	38.9	0.10	3.54	23.0
HD 37128 (Ori)	A ,B ,C ,D	17	21	335	129	0.05	6.44	48.9
HD 37742 (Ori A)	A ,B ,C ,D	23	27	275	133	0.10	5.10	48.9
HD 38771 (Ori)	A ,B ,C ,D	19	22	280	130	0.15	6.17	49.8
HD 41117 (² Ori)	A ,B ,C ,D	22	25	220	131	0.11	5.44	49.9
HD 42087 (3 Gem)	A ,B ,C ,D	20	20	210	135	0.86	7.09	49.8
HD 43384 (9 Gem)	A ,C ,D	17	17	220	133	0.87	8.29	51.9
HD 47240	A ,C ,D	16	16	195	133	0.91	8.85	51.9
HD 52382	A ,C ,D	17	17	180	164	0.85	10.2	56.0
HD 53138 (α^2 CM a)	A ,B ,C ,D	20	22	230	164	0.10	7.79	50.9
HD 58350 (CM a)	C ,D	14	17	210	104	0.08	6.48	48.8
HD 91316 (Leo)	A ,B ,C ,D	19	20	200	162	0.07	8.50	48.8
HD 119608	C ,D	7	7	135	104	1.09	17.3	55.9
HD 151804	D	7	7	125	42.0	0.99	6.99	30.9
HD 152236 (¹ Sco)	D	6	6	165	42.0	1.02	8.39	31.0

^a A : 2001 Nov. 18 { Dec. 04; B : 2001 Dec. 27 { 28; C : 2002 Jan. 21 { Feb. 03; D : 2002 Mar. 24 { May 07.

of the H β profile (the extent of the variability in velocity space is well correlated with the wind terminal velocity: $v \approx 0.3 v_1$). One can note that morphological evidence for a H β feature partly formed in the outflow (i.e., a line profile clearly departing from a pure photospheric one) is systematically accompanied by much more prominent changes. The detected variability is qualitatively similar to the cases reported by Ebbets (1982). Noteworthy is the emission-like episode experienced by HD 14134 and HD 43384. The dramatic variations previously reported in HD 21291 (Denizman & Hack 1988; Zeinalov & Rzaev 1990) and HD 91316 (Smith & Ebbets 1981) are not seen in our data.

Most changes take place on a daily time-scale, although significant hourly variations are sometimes observed. These subtle, short-term changes should be considered with some caution. In some cases, we cannot exclude the possibility that they largely arise from an imperfect continuum normalization and/or removal of telluric features (Section 3.1). To better assess the reality of these short-term variations, we use a ‘quality flag’ defined as the number of co-added consecutive exposures (indicated for each spectrum in Fig. 1). Hourly changes between spectra rated as ‘1’ should be regarded with some suspicion (e.g., HD 14134), whereas they are very likely to be real between spectra rated as ‘3’ (e.g., HD 37128). Our data support the hourly changes found in HD 38771 by Rusconi et al. (1980), but not the variations on a time-scale of 30 minutes reported for HD 53138 by van Helden (1972).

One should bear in mind that the observed variations in H β may not be straightforwardly associated to wind activity. It is likely that time-dependent changes, possibly arising from pulsations, as commonly observed in early-type super-

giants (e.g., de Jong et al. 1999; Kaufer et al. 1997), also affect the underlying photospheric profile. Although our modest spectral resolution precludes a detailed study, we choose to illustrate the temporal changes in photospheric lines by means of He I 6678.^x With the exception of HD 152236 which occasionally exhibits a P Cygni profile, this feature appears to be primarily of photospheric origin for all the stars in our sample. The changes affecting this line are generally significant (see Fig. 1), and are likely paralleled in the H β photospheric component. Because of substantial pressure broadening in B-type supergiants, such variations will extend in velocity space well beyond the projected rotational velocity (e.g., Ryans et al. 2002). In many cases the observed He I 6678 variability barely exceeds the detection thresholds, which prevents us from linking the patterns observed in H β and He I 6678. The limited spectral resolution also precludes a straightforward interpretation of the variability as arising either from a variable, incipient emission or from temporal fluctuations in the shape of the absorption profile (e.g., HD 31327). A direct comparison between the patterns of variability in He I 6678 and H β will be presented for two illustrative cases in Section 6.1.

To quantify the level of spectral and photometric variability, we define the two activity indices a_{phot} and $a_{\text{H}\beta}$ (Table 4). The former was calculated both from Hipparcos data and from our photometric observations, and is defined as: $a_{\text{phot}}^2 = \frac{2}{\text{obs}} \left\{ \frac{2}{\text{instr}} \right\}$, where obs and instr are the scatter of the

^x We do not discuss the variations that may affect the potentially interesting He I 5876 line because of difficulties in defining the neighbouring continuum, as well as edge effects.




Figure 1. H α time-series for HD 13854, HD 14134, HD 14818, HD 21291, HD 24398, HD 24912, HD 30614, and HD 31327. The spectra are displayed in the stellar rest frame. Consecutive spectra are shifted by 0.15, 0.10, 0.10, 0.07, 0.10, 0.10, 0.10, and 0.10 continuum units, respectively. The mean profile is overplotted as a dashed line. The numbers to the left-hand and right-hand sides of the upper panels give the number of consecutive exposures used to form the corresponding spectrum and the mean Julian date of the observations (H JD (2,452,000), respectively). The two bottom portions of each panel show the TVS of H α and He I 6678 (Fullerton et al. 1996), along with the threshold for a significant variability at the 99.0 per cent confidence level (dashed line).




Figure 1. H I time-series for HD 37128, HD 37742, HD 38771, HD 41117, HD 42087, HD 43384, HD 47240, and HD 52382. Consecutive spectra are shifted by 0.12, 0.13, 0.08, 0.20, 0.10, 0.12, 0.15, and 0.10 continuum units, respectively.




Figure 1. $H\alpha$ time-series for HD 53138, HD 58350, HD 91316, and HD 119608. Consecutive spectra are shifted by 0.10 continuum units in all cases.

observations and the instrumental noise, respectively (see Marchenko et al. 1998). The spectral variability index, a_{pv} , is the fractional amplitude of the line-profile variations, i.e., the amplitude of the changes normalized by the strength of the feature (see equation [15] of Fullerton et al. 1996). This index was calculated both for $H\alpha$ and for He I 6678.

5 PERIOD SEARCH

We performed a period search analysis by calculating the power spectra (hereafter PS) using the technique of Scargle (1982) on: (a) the photometric Hipparcos data (our observations are not amenable to a period search), (b) the pixel-to-pixel $H\alpha$ time-series, (c) the equivalent widths (EWs) measured within fixed wavelength intervals encompassing the whole of $H\alpha$ (even in the case of P Cygni profiles). Despite the fact that the EW and pixel-to-pixel line-profile variations are not, strictly speaking, two independent quantities, a period search in the two data sets helps assessing the reality of the detected signals. Subsequent correction of the frequency spectrum by the CLEAN algorithm was performed in order to remove aliases and spurious features induced by the uneven spacing of the data in the time domain (Roberts, Lehar & Deher 1987).

5.1 Photometric data set

Periodicities in the range 1–27 days have been found for 9 stars in our sample. Several stars display significant variability at the 0.03–0.07 mag level without, however, evidence for cyclical changes. A complex, multi-periodic behaviour cannot be ruled out in this case. As can be seen in Fig. 2, our photometric observations of HD 13854, HD 14134, HD 14818, and HD 47240 support the periodic nature of the variations suggested by the Hipparcos data. This demonstrates the remarkable long-term (years) coherency in the patterns of variability: folded with the appropriate periods, practically all the new data points stay well within the scatter limits (mainly intrinsic, considering the typical accuracy of the Hipparcos photometry), without any adjustment of the initially chosen zero phases.

5.2 Spectroscopic data set

The significance of the peaks in the PS can be, in principle, rigorously estimated after normalization by the variance of the data (e.g., Scargle 1982). In the case of unevenly spaced time-series, however, the false alarm probability (hereafter FAP) is only indicative and cannot be used as a robust indicator of the presence of periodic signals in the data (see, e.g., Antokhin et al. 1995). While assessing the reality of the periods found in our spectroscopic data set, we therefore took into account additional criteria. Namely, we inspected the period-folded EW/line-profile data, and compared the




Figure 1. H I time-series for HD 151804 and HD 152236. Consecutive spectra are shifted by 0.20 and 0.40 continuum units, respectively.

EW and 2D pixel-to-pixel PS (even though the line-profile variations are obviously not necessarily accompanied by significant EW changes). The existence of a cyclical pattern of variability is suggested for 9 stars out of the 17 included in the analysis. These cases are individually discussed below. No indication of a cyclical behaviour was found in HD 24398, HD 30614, HD 31327, HD 43384, HD 52382, HD 53138, HD 58350, and HD 91316. Kaper et al. (1998) and Markova (2002) report variations on a time-scale of about one week in HD 30614. A search for periodic modulations of the UV lines in HD 53138 proved inconclusive, in line with our non-detection (Périn et al. 2002).

HD 13854: Significant power at 0.955 d^{-1} is found in the EW data (Fig. 3), but not in the H I time-series.

HD 14134: This star does not exhibit any evidence for binarity (Abt & Levy 1973). The photometric variability of HD 14134 is relatively well documented (Rufener & Bartholdi 1982; Waelkens et al. 1990; Krzesinski, Pigulski & Kolaczowski 1999; Adelman, Yuce & Engin 2000), but the Hipparcos data provide the first clear evidence for a periodic pattern with $P = 12.823 \text{ d}$ (Fig. 2). Two periodic signals at $\nu_1 = 0.045$ and $\nu_2 = 0.080 \text{ d}^{-1}$ ($P = 22.2$ and 12.5 d , respectively) are unambiguously detected in the 2D PS, with $\text{FAP} < 0.1$ per cent (see Fig. 4). The latter frequency is also found in the EW PS (once again with $\text{FAP} < 0.1$ per cent) and is, within the errors, identical to the value found in the photometric data. The non-detection of ν_1 both

Table 4. Photometric and spectroscopic activity indices.

Name	a_{phot}		a_{pv}	
	This study (mmag)	Hipparcos (mmag)	H (per cent)	He I 6678 (per cent)
HD 13854	26	13	47.3	10.6
HD 14134 (V 520 Per)	22	24	259	11.2
HD 14818 (10 Per)	20	17	34.8	6.01
HD 21291		17	12.5	0 ^b
HD 24398 (Per)		0	5.91	5.57
HD 24912 (Per)		0	14.6	6.53
HD 30614 (Cam)		9	17.8	10.7
HD 31327		0	6.45	7.92
HD 37128 (Ori)		11	81.9	7.66
HD 37742 (OriA)		4	82.7	11.0
HD 38771 (Ori)		9	32.6	7.35
HD 41117 (2 Ori)		15	27.9	12.8
HD 42087 (3 Gem)	20:	13	91.2	8.98
HD 43384 (9 Gem)		20	135	7.69
HD 47240	17	5	40.2	5.89
HD 52382	20	16	32.1	11.0
HD 53138 (α^2 CMa)		21	66.4	9.64
HD 58350 (CMa)		28	44.1	6.90
HD 91316 (Leo)		9	12.8	10.1
HD 119608		0	10.2	0 ^b
HD 151804		0 ^a	5.16	0 ^b
HD 152236 (1 Sco)		22	5.66	19.1

^a Photometric variable according to Balna (1992), and references therein.

^b These stars do not exhibit significant variability in He I 6678 (see Fig. 1).

in the EW and in the photometric data sets leads us to take ν_2 as the fundamental frequency. The two signals are likely to be simple harmonics ($\nu_2 = 2\nu_1$). As can be seen in Fig. 5, maximum H I emission nearly coincides with maximum light. We use the photometric ephemeris when folding the EWs ($P = 12.823$ days and $T_0 = 2,447,867.8$), as this period is undistinguishable, but more accurate, than the value derived from the spectroscopic data.

The clearly phase-locked nature of the variations is reminiscent of some early-type stars displaying evidence for a decentered, dipole magnetic field (e.g., Stahl et al. 1996). A single circular polarization spectrum of HD 14134 was therefore obtained on 29 July 2003 using the MUSEOS spectropolarimeter at the 2-m Bernard Lyot telescope (Pic du Midi Observatory). The peak S/N was 250 (weather conditions were relatively poor). The mean profiles obtained using the Least-Squares Deconvolution procedure (LSD; Donati et al. 1997) show no significant circular polarization within spectral lines. The measured longitudinal magnetic field is $45 \pm 145 \text{ G}$ (G.A. Wade, private communication).

HD 24912: de Jong et al. (1999) discussed periodic changes in photospheric lines that may arise from pulsations ($P = 3.45$ hours). This star presents a strong case for a persistent, cyclical variability in the UV wind line-profiles with $P = 2$ days (Kaper et al. 1997). From an analysis of an extensive data set of UV and optical observations, de Jong et al. (2001) reported a 2.086-d modulation of the line profiles, notably in H I. In our data we find indication for a period tantalizingly close to, but apparently significantly different




Figure 2. Hipparcos light curves for stars displaying periodic variations. Zero phase is arbitrarily assigned to the start of the observations. The size of the symbols approximately corresponds to 2- σ error bars (if not shown). Our B- and I-band observations are overplotted as filled triangles and circles, respectively (the points are vertically shifted by a constant value to match the Hipparcos data). In this case the size of the symbols corresponds to 4- σ error bars.

from this value: $P = 2.20$ d (with a typical uncertainty of 0.03 d). There is also some evidence for long-term changes in the pattern of variability. While cyclical variations are suggested here in the red wing of H α (Fig. 4), such changes predominantly occurred in the blue wing on October 1994 (de Jong et al. 2001). Although the periodic signal found in our data is present both in the H α line-profile and EW time-series, we warn that its formal significance is low in the 2D PS (Fig. 4), while the phase-locked nature of the EW variations is only




Figure 3. EW of the H α line as a function of phase (data for HD 14134 in Fig. 5). Zero phase is arbitrarily set at maximum emission (or, alternatively, at minimum absorption). The uncertainties have been calculated following Chalabaev & Maillard (1983).

marginal (Fig. 3).

HD 37128: Prinja et al. (2002) discussed the UV line-profile variations in this star, but the limited time span of the IUE observations (~ 17 hours) precluded the detection of a cyclical behaviour on a rotational time-scale. Three signals are detected at the 99.0 per cent confidence level in the pixel-to-pixel H α PS at: $\tau_1 = 0.055$, $\tau_2 = 1.070$, and $\tau_3 = 1.280$ d $^{-1}$ (Fig. 4). τ_2 can be identified as the 1-d alias of τ_1 . Interestingly, the highest peaks in the EW PS correspond to the two remaining signals, although with low significance level, as confirmed by the poorly coherent patterns of variability. Our data therefore suggest the existence of two periods at $P = 18.2$ and 0.78 d, but at a low confidence level.

HD 37742: Line-profile variations on a time-scale of about 6 days as suggested both in H α and in UV wind lines (Kaper et al. 1998, 1999). Several significant peaks (all with FAP < 1 per cent) are found in the pixel-to-pixel PS at: $\tau_1 = 5 \times 10^{-3}$, $\tau_2 = 0.020$, $\tau_3 = 0.880$, and $\tau_4 = 1.148$ d $^{-1}$ (Fig. 4). Only τ_4 is found in the EW data set, but this signal is likely spurious, as judged from the lack of coherent variations. Alternatively, the EW variations appear reasonably phase-locked when folded with $\tau_5 = 0.075$ d $^{-1}$ (Fig. 3). This signal is also found in the pixel-to-pixel PS, but the confidence level is low.

HD 38771: Two signals are found in the 2D PS (see Fig.4): at $\tau_1 = 0.210$ (FAP 5 per cent) and $\tau_2 = 0.955 \text{ d}^{-1}$ (FAP 1 per cent), but there is no evidence for periodicity in the EW time-series.

HD 41117: Several peaks are found in the 2D PS (all with FAP < 1 per cent) at: $\tau_1 = 5 \times 10^{-3}$, $\tau_2 = 0.025$ (and its first harmonic at 0.050 d^{-1}), $\tau_3 = 1.045$, and $\tau_4 = 1.085 \text{ d}^{-1}$ (Fig.4). The highest peaks in the EW PS are found at $\tau_5 = 0.025$, $\tau_6 = 0.065$, and $\tau_7 = 0.205 \text{ d}^{-1}$. Among these, τ_6 and τ_7 can be ruled out from the lack of phase-locked EW variations. Despite the detection of τ_5 both in the 2D and in the EW PS, the poorly coherent pattern of variability casts serious doubt on the existence of this 40-d signal.

HD 42087: This star does not exhibit significant radial velocity variations (Mayer et al. 1999). The highest peak in the 2D PS is found at $\tau_1 = 0.040 \text{ d}^{-1}$, and appears both in the emission component and in the absorption trough of the H β Cygni profile (Fig.4). Although its formal significance is relatively low (FAP 5 per cent), the existence of a 25-d periodicity is supported by the EW data (Fig.3).

HD 47240: There is a tentative indication that the changes in SiIV 1400 are repetitive over a 3- to 5-d time-scale (Prinja et al. 2002). An ill-defined, long-term periodic signal is found in the EW data set at $\tau_1 = 0.0075 \text{ d}^{-1}$, along with smaller peaks at $\tau_2 = 0.105$ and $\tau_3 = 0.247 \text{ d}^{-1}$. The two latter frequencies are likely spurious, as judged from the poorly coherent nature of the phased EW variations. Interestingly, τ_1 is also found in the pixel-to-pixel H β PS with a confidence level close to 95 per cent (Fig.4). However, both the close correspondence with the total time span of the observations (see Table 3) and the sparsely sampled EW data (Fig.3) lead us to discard it from further consideration. Furthermore, the H morphology is typical of a fast-rotating star seen nearly equator-on (Conti & Leep 1974; Petrenz & Puls 1996). It is therefore unlikely that spectral changes operating on such a long time-scale would be associated to rotation. Variations intrinsic to a circumstellar disc-like enhancement cannot be ruled out a priori, but they would manifest themselves as V/R variations of the line emission peaks, which are not observed (Fig.4).

6 DISCUSSION

6.1 Incidence of cyclical line-profile variations

The results of our period search are summarized in Table 2. Several of the listed periods must be regarded as tentative, especially those detected either in the 2D or in the EW PS. Independent confirmation is clearly needed in this case. We also recall that HD 13854 and HD 47240 might be part of a close, massive binary system (Section 3.1). Since our survey was not designed to detect short-term periodicities, the found periods close to one day must also be regarded with some caution. They might arise from rotational modulation only for the fastest rotators. Conversely, they might be associated with the pseudo-periodic ejection of 'pu' s' of

material. Such an outward propagation of density perturbations has been proposed to account for the recurrent changes (with a time-scale of 20 days) in the H β line of the B0.5 hypergiant HD 152236 (Rivinius et al. 1997). Changes on a time-scale of about one day might be compatible with the characteristic flow time through the H β line-formation region in our program stars, provided that these putative, spatially-localized density enhancements follow a τ -type velocity law with an exponent close to unity.[†] Although such an interpretation cannot be completely ruled out, it should be noted that HD 152236 is considered as a candidate luminous blue variable and is likely not representative of the stars in our sample (Sterken et al. 1997). Pulsations can also operate on a 1-d time-scale, but we find no trivial relation between the spectroscopic and photometric periods (the latter presumably related to pulsations, as argued below).

Summarising, we consider that the two stars with a relatively long-term periodic modulation seen both in the EW and in the line-profile data sets present the strongest cases for a cyclical behaviour: HD 14134 and HD 42087 (with P 12.8 and 25.0 d, respectively). The phase-related patterns of variability are remarkably similar (see Fig.6). In addition, the line-profile changes in H β and HeI 6678 also seem to be correlated in both stars, with excess emission in H β being accompanied by a weakening of HeI 6678. As can be seen in Fig.5, this may also apply to weak metal lines such as C II 6578.05 and C II 6582.88 (note that variations in the continuum level may be invoked in the case of HD 14134 in view of the large, phase-locked photometric variations). Since a significant amount of incipient wind emission is not expected for stars with such relatively low mass-loss rates (Table 1), this phenomenon may point to a link between the wind variability and photospheric disturbances. Higher resolution data are, however, needed to reveal an organized pattern of variability in photospheric lines, as observed in other B-type supergiants (Kaufer et al. 1997, 2002).

The lack of cyclical variations for several stars in our sample can have a physical origin or may be ascribed to the limitations of the current data set (e.g., insufficient time sampling). Concerning the latter point, it is suggestive that the stars with a detected period in H β are among those displaying the highest level of variability. The large amplitude of the variations exhibited by most stars suggests that an imperfect removal of telluric features or S/N issues are probably irrelevant. One may be more concerned by the fact that our observations were relatively few and spread out over several months. If we accept the idea that large-scale wind streams give rise to the variability, substantial variations in the global wind morphology might be expected on a time-scale of months/years (as in the case of the B0.5 Ib star HD 64760; Fullerton et al. 1997). A lack of coherency in the pattern of variability on a monthly time-scale would thus not be surprising, especially for the fastest rotators in our sample. It is unlikely in this case that a strong periodic signal would appear in the data. Another issue which may lead to a loss of periodic signal is the complex pattern of variability

[†] Here we do not consider the outward propagation of small-scale clumps of material resulting from the unstable nature of radiatively-driven winds, as this would lead to a stochastic pattern of variability (e.g., Eversberg, Lepine & Moe 1998).




Figure 4. Pixel-to-pixel power spectra of the H α line series. The number of iterations and gain of the CLEAN algorithm were typically set to 2000 and 0.6, respectively. All the PS are variance-normalized to provide a uniform assessment of significance. The solid contours are drawn for intensities of 12.9, 10.6, and 8.0 (this corresponds to significance levels of 99.9, 99.0, and 95.0 per cent, respectively). The dotted contours are drawn for intensities of 7.0, 5.0, 3.0, and 1.0. In each diagram the frequency range is chosen to show more clearly the detected signals; no significant power peaks are found outside this frequency domain. For HD 24912, the dashed line indicates the 2.086-d period found by de Jong et al. (2001). The bottom section of each panel shows the mean spectrum (in the stellar rest frame), while the side panel presents the PS summed over the shown velocity range.




Figure 5. Left-hand panels: light curve of HD 14134 binned to 0.1-phase resolution (symbols as in Fig. 2. Here note that zero phase is arbitrarily set at maximum light) and EWs of the H line, as a function of phase. Right-hand panels: phase-averaged H and He I 6678 profiles. In all cases we use the photometric ephemeris: $P = 12.823$ days and $T_0 = 2,447,867.8$.

presented by H in B supergiants (see, e.g., Kaper et al. 1996; de Jong et al. 2001; Prinja et al. 2001), emphasizing the need for high spectral/time resolution.

6.2 Evidence for rotational modulation?

Detailed studies of the line-profile variations in the UV domain have inferred a rather simple geometrical structure for the wind of some B supergiants, with generally two (or four) symmetrical, spatially-extended wind streams (e.g., Fullerton et al. 1997). Consequently, one generally expects the recurrence time-scale of the line-profile variations to be an integer fraction of the stellar rotational period, rather than the period itself. In HD 24912, for instance, the period found in H can be identified, within the uncertainties, with one half of the rotation period (de Jong et al. 2001). The fact that the periodicity found in the spectroscopic data set for HD 42087 is longer than the estimated maximum rotational period is therefore puzzling in this respect (Table 2). Apart from rotation, however, there is a clear indication for the existence of an additional (perhaps dominating) line-broadening mechanism in B-type supergiants (plausibly tangential macro-turbulence; Howarth 2004). The projected rotational velocities are thus likely to be grossly overestimated for the stars in our sample, with the result that the true rotational periods may be significantly longer than anticipated (Ryans et al. 2002). The modulation time-scale of H in HD 14134 and HD 42087 is commensurate with the probable rotation periods, but rotational modulation cannot be firmly established in view of these uncertainties. For the former, the 12.8-d period found is close to the estimate of the minimum rotational period (~ 10 days), which would imply that this star is a fast rotator viewed almost equator-on. However, because in HD 14134 the $v \sin i$ is typical of stars of the same spectral type (Howarth et al. 1997), this 12.8-d time-scale is more likely to be an integer fraction of the rotation period.

The prominent changes affecting the P Cygni absorption component of the H line in HD 14134 and HD 42087

(Fig. 6) may diagnose variations in the amount of wind material seen projected against the stellar disk because of an azimuthal dependence of the mass-loss rate. Note in this respect that in HD 14134 maximum H emission and maximum light do not seem to exactly coincide (see Fig. 5). The existence of such a phase offset (which needs to be confirmed thanks to a more accurate ephemeris) would be consistent with a picture in which photospheric perturbations lead to the development of large-scale, spiral-shaped wind structures (Cranmer & Owocki 1996). Let us suppose a bright photospheric region causing a local enhancement of the mass-loss rate. One observes a classical P Cygni profile when this 'overloaded' portion of the wind is projected against the stellar disk. However, the motion of this wind feature towards the stellar limb because of rotation will cause the growth of H emission. A small phase lag (i.e., < 0.25) would be expected if the spot has significant azimuthal extent. A substantial modulation of the local mass-loss rate may also be invoked to explain the large, phase-locked EW variations seen in other stars (e.g., HD 13854; Fig. 3).

6.3 Relation to photospheric variability

The hypothesis that the development of azimuthally structured outflows in early-type stars is intimately linked to processes taking place at the photosphere (Cranmer & Owocki 1996) has received considerable observational support (e.g., Henrichs, Kaper & Nichols 1994; Reid & Howarth 1996). Much more controversial, however, is the nature of the 'seed' perturbations, with pulsational and magnetic activity as the most straightforward candidates. In the light of our results, we discuss below the relevance of these two scenarios in B-type supergiants.




Figure 6. Grey-scale plots of the residuals of H and He I 6678, as a function of phase, for HD 14134 ($P = 12.823$ d) and HD 42087 ($P = 25.0$ d). The residual profiles are the individual spectra binned to 0.05-phase resolution minus the mean spectrum. Excess emission (or, alternatively, weakening of absorption) appears brighter in these plots. We use the photometric ephemeris in the case of HD 14134, while zero-phase is arbitrarily fixed at maximum emission for HD 42087. The middle portion of each panel shows the superposition of the binned profiles. The TVS, along with the horizontal dashed line indicating the 99.0 per cent confidence level for significant variability, are displayed in the lower portion of each panel. The grey-scale plots are displayed in the stellar rest frame.

6.3.1 Pulsations

The Hipparcos data have revealed the existence of a large population of B-type supergiants with daily, quasi-periodic light variations. These stars might constitute the extension to higher luminosities of the slowly pulsating B stars, suggesting the mechanism excitation of multiple, non-radial gravity modes arising from a metal opacity bump at $T \approx 2 \times 10^5$ K in stellar interiors (e.g., Waelkens et al. 1998). Both the amplitude and the time-scale of the photometric variations are compatible to what is observed in our sample. To investigate this issue in more detail, we plot in Fig. 7 our stars on a HR diagram, along with the theoretical instability strips for non-radial g modes calculated without mass loss and rotation (Pamyatnykh 1999). The stars with a clear cyclical behaviour (a sample evidently biased towards the stars with large photometric amplitudes) do not appear to fall in the predicted domains of pulsational instability. The NLTE spectral synthesis of Kudritzki et al. (1999) suggests systematically higher effective temperatures and luminosities for some stars included in our study. However, such a temperature offset may not be sufficient to bring these stars in the instability zone. Furthermore, only the late B-type

stars in our sample seem to follow the period-luminosity relation for g modes (see Fig. 3 of Waelkens et al. 1998). Massive stars are also believed to experience a variety of strange-mode and resonance instabilities (Kiriakidis, Fricke & Glatzel 1993), but there is no indication that the photometric variations are related to this phenomenon (Fig. 7). The ubiquity of pulsational instability in this part of the HR diagram (e.g., Fullerton et al. 1996; Kaufer et al. 1997) leads us to consider pulsations as most likely responsible for the photometric variations, although it is clear from the above discussion that the pulsation modes have yet to be identified in this case. The apparent dichotomy between the photometric properties of O and B supergiants seen in Fig. 7 (the former exhibiting significantly lower variability amplitudes) suggests that the pulsations are of different nature in these 2 class of objects (e.g., higher modes are excited in O-type stars).

The complete absence of the photometric periods in the H time-series (HD 14134 excluded) seems at first glance to argue against pulsations as driver of the wind variability (once again under the assumption that the light variations can be ascribed to this mechanism). As can be seen in Fig. 7, the stars displaying the highest level of H variability




Figure 7. Upper panel: the circles show the positions of the program stars in the HR diagram (some stars have been slightly shifted along the ordinate axis for the sake of clarity). The size of the symbols is directly proportional to the amplitude of the photometric variations in the Hipparcos data, a_{phot} (see Table 4). The squares and triangles show the position of galactic OBA-type stars (van Genderen et al. 1989) and blue supergiants in NGC 300 (Bresolin et al. 2004), respectively. The size of the symbols is proportional in this case to the amplitude of the variations in V-band. Filled and open symbols correspond to stars with and without evidence for photometric periodicities, respectively. Lower panel: positions of the program stars in the HR diagram with the size of the symbols being proportional to the fractional amplitude of the line-profile variations in H γ , a_{lpv} . The evolutionary tracks (dotted lines) have been calculated for solar metallicity and for the various initial masses indicated to the right-hand side of the panels. Domains of the strange mode oscillations are overplotted as solid lines, while thick lines correspond to very unstable phases (from Kiriakidis et al. 1993). The pulsational instability strip of non-radial g-type modes is shown as (nearly vertical) solid lines. The models have been calculated using the OPAL opacities, a hydrogen abundance, $X = 0.70$, and a heavy element abundance, $Z = 0.02$ (from Pamyatnykh 1999).

are not preferentially found close to the zones of pulsational instability. This hypothesis can be investigated further by looking for a correlation between the amplitude of the photometric and H γ changes, as parametrized by the activity indices a_{lpv} and a_{phot} (the latter calculated from Hipparcos data; Section 4). Although there is a hint of a positive correlation (FAP 2.3 per cent), this trend disappears when the 4 stars displaying line-profile variations that cannot be un-

ambiguously related to wind activity, but can be of purely photospheric origin, are excluded (i.e., HD 21291, HD 24398, HD 31327, and HD 119608). In addition, there is also no correlation between a_{lpv} (H γ) and a_{lpv} (He I 6678).

The lack of a clear statistical relationship between the levels of activity at the photosphere and at the base of the outflow (as diagnosed by the photometric changes/He I 6678 and H γ , respectively) suggests that the same physical

mechanism (i.e., presumably pulsations) is not responsible for the photospheric and wind activity. This conclusion relies, however, on statistical grounds and should be in principle examined on a star-to-star basis: here we note the example of HD 64760, where the same periodic signal seems to appear in photospheric and wind lines (Kaufer et al. 2002). Although detailed models are still lacking, one may also expect a complex interplay between any pulsation signal and wind changes (e.g., beating of some pulsation modes). This is likely to result in a loose (if any) relationship between photometric and H α changes, for instance.

6.3.2 Magnetic fields

A growing amount of evidence suggests the existence of weak magnetic fields in early-type stars, although firm detections are still scarce and might be restricted to some atypical cases. Spectropolarimetric observations of the very young O4.6 star γ Orionis C have revealed a relatively strong magnetic field, 1.1 ± 0.1 kG (Donati et al. 2002). The existence of a weaker field in the B1 IV star ϵ Cephei (with a polar component of 360 ± 30 G) has also been reported recently. This field appears to be sufficient to control the outflow at large distances from the star (out to $9R_*$; Donati et al. 2001). Magnetic activity in OB stars is also hinted at by X-ray observations of the O9.5 II star HD 36486 (α Ori) and HD 37742 (Waldron & Cassinelli 2000; Miller et al. 2002) which challenge the generally accepted view that the X-ray emission in early-type stars solely arises from outwardly propagating shocks developing as the result of the inherently unstable nature of radiation-driven winds (see also Feigelson et al. 2002). Recent theoretical studies indicate that a magnetic field of a moderate strength can be produced via dynamo action at the interface between the convective core and the outer radiative envelope, and brought up to the surface on a time-scale typically smaller than the main-sequence lifetime (Charbonneau & MacGregor 2001; MacGregor & Cassinelli 2003).

Apart from our observations of HD 14134 (Section 5.2), HD 24912 is the only star in our sample for which a magnetic field has been searched for via spectropolarimetric observations. A 3- σ upper limit of 210 G in the longitudinal component was derived (de Jong et al. 2001). It is worth noting that this does not set stringent constraints on the potential role played by magnetic activity, as even relatively weak fields ($B < 100$ G) are suspected to significantly perturb line-driven flows (Owocki 1994). This issue has been recently worked out in more detail by ud-Doula & Owocki (2002) who studied the influence of a large-scale magnetic field with a dipole configuration on the radiatively-driven outflow of an early-type star. They found that the disturbance imposed by the magnetic forces can be parametrized by a confinement parameter, $\eta = B_{\text{eq}}^2 R_*^2 / M v_1$, where B_{eq} is the equatorial field strength at the stellar surface. The magnetic field appears to significantly impact on the wind morphology and dynamics even for values as low as $\eta = 0.1$, while the emergent flow is channeled along closed magnetic loops for $\eta > 1$. For the physical parameters typical of our program stars (e.g., HD 14134; see Table 1), we obtain $\eta = 1$ for $B_{\text{eq}} = 70$ G. In this case the Alfvén radius would be about $2 R_*$, i.e., comparable to the extent of the H α line-formation region (ud-Doula & Owocki 2002). This rough estimate shows that

magnetic activity cannot be ruled out as driver of the wind activity at this stage. A much stronger field is unlikely in HD 14134, considering our spectropolarimetric observations together with the lack of strong X-ray emission ($L_X [0.2 - 4.0 \text{ keV}] < 33.0 \text{ ergs s}^{-1}$; Gribb et al. 1992). Fields of the same magnitude can also be expected in other stars from our sample in view of their relatively low X-ray luminosities (Berghofer, Schmitt & Cassinelli 1996).

7 CONCLUSION

Our long-term monitoring has confirmed the ubiquity of H α line-profile variations in OB supergiants. The changes result from the combined effect of a varying amount of incipient wind emission and an intrinsically variable underlying photospheric profile, with the relative contribution of these two phenomena varying from one star to another. As a consequence of the large amplitude of the H α variations observed, a precise determination of the stellar atmospheric parameters will not be generally possible from profiling of the Balmer lines in snapshot spectra. Taken at face value, for instance, the EW variations shown in Fig. 3 would translate in the most extreme cases into a ± 0.8 dex uncertainty in the derived mass-loss rate (Puls et al. 1996; equation [43]). Although it remains to be seen whether a sophisticated modelling would result in drastically different figures, the large H α changes intrinsic to the early-type supergiants may be one of the most important factors limiting the accuracy with which these objects can be used as extragalactic distance indicators (see discussion in Kudritzki 1999).

This survey has revealed the existence of cyclical H α line-profile variations in HD 14134 and HD 42087. The low detection rate (2 cases out of 17) may have several explanations, including observational limitations or a loss of coherency in the patterns of variability. HD 14134 is of particular interest, as it is one of the very few early-type stars clearly exhibiting the same pattern of variability in the photometric and spectroscopic modes. The existence of azimuthally structured outflows in our program stars might explain the dramatic H α changes, but firm conclusions must await detailed modelling (see Harries 2000 for the first effort in this direction). Sensitive searches for close companions (e.g., via long-baseline optical interferometric observations) are also needed to firmly rule out a variability arising from wind interaction.

We do not find evidence for a direct coupling between the wind activity and base perturbations arising from pulsations, but we caution that our data are not well suited to examine this aspect in detail. It is conceivable that in several instances the globally anisotropic nature of the outflow primarily results from the putative existence of small-scale magnetic loops anchored at the stellar photosphere (or alternatively a presumably fossil, large-scale dipole field). A major observational effort is currently undertaken to detect such weak magnetic fields in early-type stars (e.g., Henrichs et al. 2003). Firm detections can be expected in the foreseeable future. They would place the above conclusions on a quantitative footing and directly address the relevance of this scenario.

8 ACKNOWLEDGMENTS

We are indebted to G. A. Wade and the TBL-MuSiCoS team for obtaining a single circular polarization spectrum of HD 14134 during the mission 'Precision Studies of Stellar Magnetism Across the HR Diagram' (July 2003). This work greatly benefited from the comments of the anonymous referee. The authors would like to acknowledge the support from the Applied Research and Technology Program at W KU and the assistance of the undergraduate telescope operators at W KU who participated in the observing program: A. Atkinson, T. Monroe and W. Ryle. It is a pleasure to thank the staff of the VBO for their excellent support. We are also grateful to the VBO chairman for the generous allotments of observing time. We wish to thank A. F. J. M. Offert for useful comments. This research made use of NASA's Astrophysics Data System Bibliographic Services and the Simbad database operated at CDS, Strasbourg, France.

REFERENCES

- Abt H. A., Levy S. G., 1973, *ApJ*, 184, 167
 Abt H. A., Levato H., Grosso M., 2002, *ApJ*, 573, 359
 Adelman S. J., Yuce K., Engin S., 2000, *IBVS*, # 4946
 Antokhin I. I., Bertrand J.-F., Lamontagne R., Mot A. F. J., Matthews J., 1995, *AJ*, 109, 817
 Balona L. A., 1992, *MNRAS*, 254, 404
 Berghofer T. W., Schmitt J. H. M. M., Cassinelli J. P., 1996, *A & A*, 118, 481
 Bresolin F., Pietrzynski G., Gieren W., Kudritzki R. P., Przybilla N., Fouque P., 2004, *ApJ*, 600, 182
 Chalabaev A., Maillard J. P., 1983, *A & A*, 127, 279
 Charbonneau P., MacGregor K. B., 2001, *ApJ*, 559, 1094
 Cherrington E. F., 1937, *ApJ*, 85, 139
 Conti P. S., Leep E. M., 1974, *ApJ*, 193, 113
 Cranmer S. R., Owocki S. P., 1996, *ApJ*, 462, 469
 de Jong J. A., Henrichs H. F., Schrijvers C., Gies D. R., Telting J. H., Kaper L., Zwarthoed G. A. A., 1999, *A & A*, 345, 172
 de Jong J. A. et al., 2001, *A & A*, 368, 601
 Denizin L., Hack M., 1988, *A & A*, 75, 79
 Donati J.-F., Semel M., Carter B. D., Rees D. E., Cameron A. C., 1997, *MNRAS*, 291, 658
 Donati J.-F., Wade G. A., Babel J., Henrichs H. F., de Jong J. A., Harries T. J., 2001, *MNRAS*, 326, 1265
 Donati J.-F., Babel J., Harries T. J., Howarth I. D., Petit P., Semel M., 2002, *MNRAS*, 333, 55
 Ebbets D., 1980, *ApJ*, 235, 97
 Ebbets D., 1982, *ApJS*, 48, 399
 Eversberg T., Lepine S., Mot A. F. J., 1998, *ApJ*, 494, 799
 Feigelson E. D. et al., 2002, *ApJ*, 574, 258
 Fullerton A. W., Gies D. R., Bolton C. T., 1996, *ApJS*, 103, 475
 Fullerton A. W., Massa D. L., Prinja R. K., Owocki S. P., Cranmer S. R., 1997, *A & A*, 327, 699
 Giallo F., Sciortino S., Micela G., Vaiana G. S., Hamden Jr. F. R., 1992, *ApJS*, 81, 795
 Harries T. J., 2000, *MNRAS*, 315, 722
 Henrichs H. F., Kaper L., Nichols J. S., 1994, *A & A*, 285, 565
 Henrichs H. F., Neiner C., Geers V. C., 2003, in van der Hucht K., Herrero A., Esteban C., eds, *Proc. IAU Symp. 212, A Massive Star Odyssey: From Main Sequence to Supernova*, Astron. Soc. Pac., San Francisco, p. 202
 Herrero A., Puls J., Najaro F., 2002, *A & A*, 396, 949
 Hinkle K., Wallace L., Valenti J., Hammer D., 2000, *Visible and Near Infrared Atlas of the Arcturus Spectrum 3727{9300 Å*, Astron. Soc. Pac., San Francisco
 Howarth I. D., Prinja R. K., 1989, *ApJS*, 69, 527
 Howarth I. D., Prinja R. K., Massa D. L., 1995, *ApJ*, 452, L65
 Howarth I. D., Siebert K. W., Hussain G. A. J., Prinja R. K., 1997, *MNRAS*, 284, 265
 Howarth I. D., 2004, in Maeder A., Eenens P., eds, *Proc. IAU Symp. 215, Stellar Rotation*, Astron. Soc. Pac., San Francisco, in press
 Hummel C. A., White N. M., Elias II N. M., Hajian A. R., Nordgren T. E., 2000, *ApJ*, 540, L91
 Humphreys R. M., McElroy D. B., 1984, *ApJ*, 284, 565
 Kaper L., Henrichs H. F., Nichols J. S., Snoek L. C., Volten H., Zwarthoed G. A. A., 1996, *A & A*, 116, 257
 Kaper L. et al., 1997, *A & A*, 327, 281
 Kaper L., 1998, in Wolf B., Stahl O., Fullerton A. W., eds, *Proc. IAU Colloquium 169, Variable and Non-spherical Stellar Winds in Luminous Hot Stars*, Astron. Soc. Pac., San Francisco, p. 193
 Kaper L., Fullerton A. W., Baade D., de Jong J. A., Henrichs H. F., Zaai P., 1998, in Kaper L., Fullerton A. W., eds, *Proc. ESO workshop, Cyclical variability in stellar winds*, Springer-Verlag, Berlin, p. 103
 Kaper L., Henrichs H. F., Nichols J. S., Telting J. H., 1999, *A & A*, 344, 231
 Kauer A., Stahl O., Wolf B., Gang Th., Gummertsbach C. A., Kovacs J., Mandel H., Szeifert Th., 1996, *A & A*, 305, 887
 Kauer A. et al., 1997, *A & A*, 320, 273
 Kauer A., Prinja R. K., Stahl O., 2002, *A & A*, 382, 1032
 Kiriakidis M., Fricke K. J., Glatzel W., 1993, *MNRAS*, 264, 50
 Koene C., 2001, *MNRAS*, 321, 44
 Kuzinski J., Pigulski A., Kolarowski Z., 1999, *A & A*, 345, 505
 Kudritzki R. P., 1999, in Wolf B., Stahl O., Fullerton A. W., eds, *Proc. IAU Colloquium 169, Variable and Non-spherical Stellar Winds in Luminous Hot Stars*, Astron. Soc. Pac., San Francisco, p. 405
 Kudritzki R. P., Puls J., Lennon D. J., Venn K. A., Reetz J., Najaro F., McCarthy J. K., Herrero A., 1999, *A & A*, 350, 970
 Lamers H. J. G. L. M., 1981, *ApJ*, 245, 593
 Lamers H. J. G. L. M., Cassinelli J. P., 1996, in Leitherer C., Fritze-von Alvensleben U., Huchra J., eds, *ASP Conf. Ser. Vol. 98, From Stars to Galaxies: the Impact of Stellar Physics on Galaxy Evolution*, Astron. Soc. Pac., San Francisco, p. 162
 MacGregor K. B., Cassinelli J. P., 2003, *ApJ*, 586, 480
 Marchenko S. V., et al. 1998, *A & A*, 331, 1022
 Markova N., 2002, *A & A*, 385, 479
 Mason B. D., Gies D. R., Hartkopf W. I., Bagnuolo Jr. W. G., Brummelaar T. T., McAlister H. A., 1998, *AJ*, 115, 821
 Massa D. L. et al., 1995, *ApJ*, 452, L53
 Mayer P., Hanna M. A., Wolf M., Chochold., 1999, *Ap & SS*, 262, 163
 Miller N. A., Cassinelli J. P., Waldron W. L., McFarlane J. J., Cohen D. H., 2002, *ApJ*, 577, 951
 Mot A. F. J., Michaud G., 1981, *ApJ*, 251, 133
 Morel T., Marchenko S. V., Patia K., Kuppaswamy K., Carini M. T., Wood E., Zimmerman R., 2004a, in Maeder A., Eenens P., eds, *Proc. IAU Symp. 215, Stellar Rotation*, Astron. Soc. Pac., San Francisco, in press
 Morel T., Marchenko S. V., Patia K., Kuppaswamy K., Carini M. T., Wood E., Zimmerman R., 2004b, *BASI*, in press
 Owocki S. P., 1994, *Ap & SS*, 221, 3
 Pamyatnykh A. A., 1999, *Acta Astronomica*, 49, 119
 Petrenz P., Puls J., 1996, *A & A*, 312, 195
 Prinja R. K., Massa D. L., Fullerton A. W., 1995, *ApJ*, 452, L61
 Prinja R. K., Fullerton A. W., Crowther P. A., 1996, *A & A*, 311, 264
 Prinja R. K., Massa D. L., Howarth I. D., Fullerton A. W., 1998, *MNRAS*, 301, 926
 Prinja R. K., Stahl O., Kauer A., Colley S. R., Crowther P. A., Wolf B., 2001, *A & A*, 367, 891

- Prinja R. K., Massa D. L., Fullerton A. W., 2002, *A & A*, 388, 587
- Puls J. et al., 1996, *A & A*, 305, 171
- Reid A. H. N., Howarth I. D., 1996, *A & A*, 311, 616
- Rivinius Th. et al., 1997, *A & A*, 318, 819
- Roberts D. H., Lehar J., Deher J. W., 1987, *A J*, 93, 968
- Rosendhal J. D., 1973, *ApJ*, 182, 523
- Rufener F., Bartholdi P., 1982, *A & A S*, 48, 503
- Rusconi L., Sedmak G., Stalio R., Apigny C., 1980, *A & A S*, 42, 347
- Ryans R. S. I., Dufton P. L., Rolleston W. R. J., Lennon D. J., Keenan F. P., Smoker J. V., Lambert D. L. 2002, *MNRAS*, 336, 577
- Scargle J. D., 1982, *ApJ*, 263, 835
- Smith M. A., Ebbets D., 1981, *ApJ*, 247, 158
- Stahlo et al., 1996, *A & A*, 312, 539
- Sterken C., Wolf B., 1978, *A & A*, 70, 641
- Sterken C., de Groot M., van Genderen A. M., 1997, *A & A*, 326, 640
- Thaller M. L., Gies D. R., Fullerton A. W., Kaper L., Wimmer R. 2001, *ApJ*, 554, 1070
- Ud-Doula A., Owocki S. P., 2002, *ApJ*, 576, 413
- Underhill A. B., 1960, *PASP*, 72, 363
- Underhill A. B., 1961, *Pub. Dom. Ap. Obs. Victoria*, 11, 353
- van Genderen A. M. et al., 1989, *A & A S*, 79, 263
- van Helden R., 1972, *A & A*, 21, 209
- Waelkens C. et al., 1990, *A & A S*, 83, 11
- Waelkens C., Aerts C., Kestens E., Grenon M., Eyer L., 1998, *A & A*, 330, 215
- Waldron W. L., Cassinelli J. P., 2000, *ApJ*, 548, L45
- Zeinalov S. K., Rzaev A. K. H., 1990, *Ap&SS*, 172, 217

This paper has been produced using the Royal Astronomical Society/Blackwell Science L^AT_EX style file.