Isabel A lem an¹ and Ruth G ruenwald

Instituto de Astronom ia, Geof sica e Ciências Atmosfericas, Universidade de Sao Paulo, SP, Brazil

ABSTRACT

This paper presents an analysis of the concentration of the hydrogen molecule inside the ionized region of planetary nebulae. The equations corresponding to the ionization and chem ical equilibria of H, H⁺, H, H₂, H₂⁺, and H₃⁺ are coupled with the equations of ionization and therm albalance for a photoionized atom ic gas. Forty di erent reactions related to the form ation or the destruction of these species are included. The presence of dust is taken into account, since grains act as catalysts for the production of H_2 , as well as shield the m olecules against the stellar ionizing radiation. We analyze the elect of the stellar ionizing continuum, as well as of the gas and grain properties on the calculated H_2 m ass. It is shown that a signi cant concentration of H_2 can survive inside the ionized region of planetary nebulae, mostly in the inner region of the recombination zone. The total H_2 to total hydrogen m assuratio inside the ionized region increases with the central star tem perature, and, depending on the PN physical conditions, it can be of 10^{-6} or even higher. The increase of the recombination zone with the stellar the order of tem perature can account for such correlation. This can explain why the H_2 emission is more frequently observed in bipolar planetary nebulae (Gatley's rule), since this kind of object has typically hotter stars. A pplying our results for the planetary nebula NGC 6720, we obtain an H $_2$ to hydrogen m ass ratio similar to the value obtained from the observed H $_2$ line emission.

Subject headings: astrochem istry | ISM :m olecules | planetary nebulae: general

1. IN TRODUCTION

Since the rst detection of molecular hydrogen in NGC 7027 (Tre ers et al. 1976), this molecule has been detected in many planetary nebulae (PNe), in both the infrared (A spin et al. 1993;Kastner et al. 1996;Hora, Latter, & Deutch 1999) and the ultraviolet (Bowers et al. 1995;Herald & Bianchi 2002).

The analysis and interpretation of the infrared em ission due to hydrogen molecules in PNe has been made in the context of: (a) dense neutral clumps inside the ionized region (Beckwith, Persson, & Gatley 1978; Reay, Walton, & Atherton 1988; Tielens 1993; Schild 1995), (b) photodissociation regions (PDRs; Tielens 1993; Natta & Hollenbach 1998; Viciniet al. 1999), as wellas (c) shocks between the expanding shell and the precursor red giant wind (Natta & Hollenbach 1998). Visual inspection of H2 and [N II] line in ages observed by several authors (Schild 1995; Guerrero et al. 2000; A rias et al. 2001; B ohigas 2001) can not exclude, however, that the H_2 em ission is partly (ormostly) produced inside the ionized region. In fact, the presence of grains in these regions, the coexistence of electrons and hydrogen ions, and the corresponding thresholds for the destruction or ionization of the di erent species, suggest that H_2 m olecules can survive inside ionized regions. Interpretation of the far ultraviolet absorption spectra recently obtained by the Far U ltraviolet Spectroscopic Explorer (FUSE) also suggests the existence of a hot component of excited H₂, which could be inside the ionized gas in PNe (Herald & Bianchi 2002).

Form ation on grain surface is the main process for the production of H_2 molecules in the interstellar medium. This formation route was rst proposed to explain the amount of H_2 found in interstellar clouds (Hollenbach & Salpeter

¹Send o print requests to isabel@astro.iag.usp.br.

1971). G rains are condensed in the atm ospheres of asymptotic giant branch and red giant stars (see W hittet 1992 and Evans 1994) and, since these stars correspond to the earlier evolutionary stages of PNe, grains could be present in the ionized region if they are not destroyed by the stellar radiation. Continuum infrared radiation observations and modeling of PNe indicate that dust is present inside the ionized region of PNe (e.g., Natta & Panagia 1981; Harrington, Monk, & Clegg 1988; Lenzuni, Natta & Panagia 1989; Stasinska & Szczerba 1999; Van Hoofet al. 2000). In particular, the distribution of the infrared em ission follows the morphology of infrared lines and continuum radiation emitted by the ionized gas (Barlow 1993; Meixner et al. 1996), also suggesting the presence of grains in the ionized region. A very strong evidence that dust is present in the ionized region com es from the models obtained by Kingdon, Ferland, & Feibelm an (1995) and Kingdon & Ferland (1997), which show depletion of elements that can be present in grains. Som e authors (for example, Stasinska 2002 and references therein) argue, however, that the grains are in clumps and condensations inside the ionized region. If grains are indeed m ixed with the ionized gas, they can act as catalyst for the form ation of H_2 molecules.

The presence of electrons and/or H^+ inside the ionized gas can also produce hydrogen molecule nuclei, through radiative attachment and associative detachment reactions. On the other hand, the threshold for the photodissociation of hydrogen molecules is 14.7 eV, while the photocionization threshold is 15.4 eV. These values are higher than the threshold for the ionization of hydrogen atoms (13.6 eV). Hydrogen molecules could then be shielded from the photons that could dissociate it, surviving inside the ionized region in zones far from the ionizing source.

D etailed analyses of the presence of m olecules have been done for interstellar clouds (for exam – ple, de Jong 1972; B lack & D algamo 1977; V iala, R oue, & Abgrall1988), as well as for photodissociation regions (for exam ple, T ielens & H ollenbach 1985; D raine & B ertoldi 1996). M odels for photodissociation regions have been applied to planetary nebulae, assuming that the incident radiation is litered after crossing the ionized region (e.g., N atta & H ollenbach 1998). These m odels assum e that the ux of the incidente radiation is zero between 13.6 and 54.4 eV. Due to the behavior of the ionizing cross-sections with the photon energy, the correct ionizing spectrum that reaches each point of the ionized region is, how ever, m uch m ore com – plex, and a proper account of the radiation transfer along the nebula m ust be done when analyzing the m olecular chem istry.

Since the rst detection of m olecular hydrogen in PNe, only a few attempts to calculate the H_2 content in ionized regions have been m ade (B lack 1978; G ussie & Pritchet 1988; C eachi-Pestellini & D algarno 1993). In these studies, how ever, one or m ore of the following e ects is not taken into account: the e ect of dust on the radiation transfer, the form ation of H_2 m olecules on grain surface, and the photodissociation in two steps.

An interesting correlation has been obtained between the presence of H_2 line em ission and the morphological type of the planetary nebula. Signi cant H_2 em ission is more frequently observed in PNe having bipolarm orphology (G atley's rule; Zuckerm an & G atley 1988; K astner at al. 1996). This correlation between H_2 em ission and them orphology is not understood yet.

Furtherm ore, som e problem s regarding the line em ission of H₂ and neutral atom s in PNe are not solved yet. The excitation mechanism of the observed H₂ em ission lines, for exam ple, is still uncertain. Som e authors conclude that excitation is due to radiative cascades following the UV pum ping. Shock waves produced by the expanding nebulae are also suggested as a possible excitation mechanism. See, for example, Black, Porter & Dalgamo (1981), Schild (1995), Kastner et al. (1996), Natta & Hollenbach (1998), Hora et al. (1999), Guerrero et al. (2000), and Arias et al. (2001). Another problem is that the emission lines produced in low-ionization regions by neutral atom s like C, N, and O are not well tted by the models (Aller 1984; Clegg et al. 1987; van Hoofet al. 2000; Stasinska & Szczerba 2001). The analysis of these lines is made using photoionization models, where lines of neutral atom s are produced in the outer regions of a photoionized gas, and the presence of molecules is not taken into account. The hydrogen molecule, being the most abundant, can a ect the physical conditions of the gas, since its presence changes the concentration of the di erent species in the gas, as well as the

therm al equilibrium, due to the mechanisms involving H_2 . If molecules are indeed present in the region where lines of neutral C, N, and O are produced, and if their e ect on the gas is not taken into account, the results can be misleading. A detailed study, with a self-consistent modeling of the presence of H_2 molecules, is then in portant to a better know ledge of the chemical composition and physical conditions of PN e.

The aim of this paper is to make a rst estimate of the amount of the H_2 molecule in the ionized regions of planetary nebulae, in order to verify if the presence of this molecule can be signi cant in these regions, as suggested by the observations. If the calculations show that the am ount of H $_2$ is im portant, a next step would be a careful calculation of the level population distribution, in order to properly account for the di erent mechanism sthat depend on the level population. In the present paper, the equations for the chem ical equilibrium of m olecular hydrogen are coupled to the equations of a photoionization code. The concentration of the hydrogen molecule is obtained along the nebula, as well as the total am ount of m olecules inside it. The theoretical models used in our analysis are described in x 2. The results for a standard PN and the e ect of the input param eters on the obtained H_2 mass appear in x 3, while a discussion of the main results is presented in x 4.

2. MODELS

M odels for typical PNe are obtained with the photoionization code AANGABA (Gruenwald & Viegas 1992). The physical conditions of the gas are determined by solving the coupled equations of ionization and therm al balance for a spherically symmetric cloud. Several processes of ionization and recombination, as well as of gas heating and cooling, are taken into account. The energy input to the gas is essentially due to ionization of atom s and atom ic ions by primary (stellar) and secondary (di use) radiation. The processes contributing to the gas cooling are colisional excitation, radiative and dielectronic recombination, thermal ionization and free-free emission. The transfer of the primary and diuse radiation elds is treated in the outward-only approxim ation. Twelve elements (H, He, C, N, O, Mg, Ne, Si, S, Ar, Cl, and Fe) are included. The code was

m odi ed in order to include the chem ical equilibrium of the species H, H^+, H_-, H_2, H_2^+ , and H_3^+ . The corresponding equations were then coupled with the equations for the ionization balance for the atom ic gas. The H_3 m olecule is not included due to its short lifetime (10⁻⁷s; G aillard et al. 1983; B ordas, C osby, & Helm 1990). The therm al balance was solved for the atom s and atom ic ions; the e ect due to H_, H_2, H_2^+ , and H_3^+ on the therm al balance is not taken into account.

Several processes that can form or destroy the included species are taken into account, as for example chemical reactions, photoprocesses, as well as grain surface reaction. The forty reactions included in the hydrogen equilibrium are listed in Table 1. Rate coe cients and cross sections are taken from the literature (see references in Table 1). Special attention was given to the search of the reactions and their corresponding rate coe cients, in order to have a set as complete and accurate as possible for the physical conditions prevailing in PNe. Som e rate coe cients depend on the distribution of the reactants on their di erent energy levels. As said above, we do not calculate in detail the level distribution of the molecules in the present paper; a LTE level distribution for the hydrogen molecules, taking into account the three lowest vibrational levels of the electronic ground state is, how ever, adopted when cross-sections for the photoprocess are provided by the literature for individual levels.

G rains are assumed to be uniform by mixed with the gas, and are heated by the stellar and nebular continuum radiation. The attenuation of the stellar and di use radiation by dust is taken into account; the e ect of scattering by dust on the radiation eld is, however, not included.

2.1. The H₂ Form ation on G rain Surface

The form ation of H_2 on grain surface requires an analysis of the grain survival. We assume that the grain can survive if the grain temperature (T_g) is smaller than the sublimation temperature, which is a characteristic of the grain material.

The grain temperature is calculated assuming radiative equilibrium. The grains are heated by the stellar and the nebular continuum radiation, and emit as blackbodies.

Following Tielens & Hollenbach (1985) and Hol-

lenbach & Salpeter (1971), the rate of H_2 form ation on grain surface, $_g$, is given by the product of the rate of H-grain collisions by probabilistic factors that depend on the grain surface properties and on the gas tem perature, i.e.,

$$_{g} = \frac{Sf}{2} a_{g}^{2} v_{H} n_{g} n (H)$$
 (1)

where a_g is the grain radius, v_H is the therm alvelocity of the H atom s, n_g is the grain density, and n (H) is the neutral atom ic H density. The probabilistic factors S and f are given by Hollenbach & M cK ee (1979). These factors decrease with the grain temperature (T_g) and the gas temperature (T), particularly when T_g is higher than 100K. They also depend on the properties of the grain m aterial.

A new model for the H_2 form ation in grains is suggested by Cazaux & Tielens (2002). In this model, they include the physiobsorption of the H atom (which dominates in temperatures lower than 50-100K), as well as the chem isorption, which would be m one in portant for higher temperatures. H ow ever, as stated by G lover (2003), the subject is still not clear.

From equation (1), the rate of H_2 form ation on grain surface is proportional to the total available surface area of all grains, i.e., $g / a_g^2 n_g$. However, following the most common notation in the literature, we chose to describe the quantity of grains by m eans of the dust-to-gas ratio,

$$\frac{M_{d}}{M_{g}} = \frac{4 a_{g}^{3} gn_{g}}{3m_{H} n_{H} (1 + 4n_{H} e = n_{H})}; \qquad (2)$$

where m_H is the atom ic mass of H, n_H and n_{He} are the total num erical density of hydrogen and helium nuclei, respectively, and $_g$ is the grain speci c density. A ssum ing a M axwellian distribution of velocity for the H atom s, equation (1) can then be rewritten as

$$_{g} / \frac{\text{SfT}^{1=2}M_{d}=M_{g}n_{H}n(H)}{_{g}a_{g}};$$
 (3)

We de ne for convenience (M $_{\rm d}=$ M $_{\rm g})a_{\rm g}^{-1}$ hereafter as .

2.2. M odel Input Param eters

A grid of theoretical models is obtained with input parameters typical of PNe, which includes the incident spectrum, the gas density and the elem ental abundances of the nebula, as well as the dust m aterial, size, and density.

In our models, a blackbody radiation is assumed for the PN central star, characterized by its temperature (T) and lum inosity (L). According to Pottasch (1984), T ranges from 3 10^4 to 5 10^5 K, while L ranges from 10^2 L to 10^4 L.

The nebular gas is characterized by its total hydrogen nuclei density (n_H) , which is assumed constant inside the nebula, and by its elemental abundances. Gas densities typically range from 10^2 to 10^5 cm 3 (Pottasch, 1984). We assume constant elem ental abundances for the atom ic gas along the nebulae, with the same values for all m odels. These are mean values for planetary nebulae taken from Kingsbourgh & Barlow (1994), for He, C, N, O, Ne, S, and Ar, obtained from atom ic line-em ission intensities. For the elements whose abundance is not provided by these authors (Mg, Si, Cl, and Fe), we adopted the values given by Stasinska & Tylenda (1986), in order to make a rough correction for grain depletion. Since the effect of these elements is not very e ective for the gas cooling, their exact proportion in the form of grains will not be im portant.

Typical grain radius ranges from 10^{-3} m to 10m and dust-to-gas mass ratio from 10 5 to 10 1 in PNe (Stasinska & Szczerba 1999). The com position of the ISM grains (Aller 1984; Evans 1994), i.e., silicate and carbon-based compounds, is assum ed here for the PNe grains. In this work, we analyze the presence of am orphous carbon, graphite, silicate, sm oothed ultraviolet silicate, silicon carbide (SiC), as well as neutral and ionized polycyclic arom atic hydrocarbons (PAHs). Optical properties for these astrophysical grains are taken from Rowan-Robinson (1986), Draine & Lee (1984), Laor & Draine (1993), Weingartner & Draine (2001), and Li & Draine (2001). These properties depend on the grain size and com position, as well as on the photon energy. Sublim ation tem peratures and speci c densities are taken from Laor & Draine (1993) and Li & Draine (2000).

We de ne a standard PN with a given set of input parameters, in order to explore the in uence due to each one on the produced amount of H_2 . For this, we vary each parameter along its typical range, while keeping the other parameters xed. The input parameters for the standard PN are the following: T = 150000 K, L = 3000L , $n_{\rm H}$ = 10³ cm ³, $a_{\rm g}$ = 0:1 m, M d=M g = 10 ³, and am orphous carbon grains.

3. RESULTS

3.1. Standard M odel

The calculated concentration of species containing hydrogen nuclei versus the distance from the central star is shown on the upper panel of F igure 1, for the standard PN m odel. The concentrations are given in units of $n_{\rm H}$. The tem perature radial prole for the standard nebula is plotted on the lower panel. The distance is given in units of the outer radius of the ionized region, $R_{\rm max}$, which is de ned as the distance to the central star where the ionization degree, n (H⁺)=n_{\rm H}, reaches 10⁻⁴. For the standard PN, $R_{\rm max} = 0.21$ pc; for a nebula with same input parameters, but T = 50000 K, $R_{\rm max} = 0.17$ pc, and, if the nebula has a density $n_{\rm H} = 10^4$ cm⁻³ and other parameters as for the standard nebula, $R_{\rm max} = 0.04$ pc.

As it can be seen in Figure 1, the hydrogen molecule is mainly ionized in the inner and interm ediate regions. The abundance of the neutral molecule, as well as of the ionized molecule, has its maximum in the inner region of the recombination zone, decreasing slightly in the less ionized regions. This behavior can be explained by the curves of Figure 2, which shows the rates for the most im portant reactions of form ation and destruction of the hydrogen molecule in the standard PN, as a function of the distance from the central star.

In brief, the most important reactions for the form ation of the hydrogen molecule nuclei (H $_2$ or H $_2^+$) are:

$$H + H ! H_2 + e$$
 (4a)

$$2H + grain ! H_2 + grain$$
 (4b)

$$H + H^{+} ! H_{2}^{+} + h$$
 (4c)

$$H^{+} + H ! H_{2}^{+} + e$$
 (4d)

with H produced by the reaction

$$H + e ! H + h$$
 (5a)

Charge exchange reactions are e ective in ionizing or recombining the hydrogen molecule:

$$H_{2}^{+} + H ! H_{2} + H^{+}$$
 (6a)

$$H_2 + H^+ ! H_2^+ + H$$
 (6b)

(6c)

C om paring the distribution of the various species (Fig. 1) with the formation and destruction rates along the standard PN (Fig. 2), the maximum concentration of molecular nuclei can be associated to a semi-ionized region, where H, H^+ and e coexist with signi cant abundance. In fact, as said above, the maximum abundance of molecular hydrogen occurs in the inner recombination zone, and the distribution follows a radial distribution similar to that of H .

G rains are the m ost im portant route to the form ation of H_2 in the outer regions since other reactions include ionized species. Therefore, when grains are not present the concentration of the m olecule decreases in the outer regions.

Regarding the destruction mechanisms, photodissociation is an important process for the destruction of H_2 , as well as of H_2^+ . In particular, the photodissociation in two steps is important for the destruction of H_2 in the whole nebula, since the maximum threshold for this process is lower than that of hydrogen. The direct photodissociation, as well as photoionization of H_2 are ofm inor importance, mainly in the intermediate and outer zones, since the energy threshold is higher than 13.6 eV.

The abundance of H_3^+ is much lower compared with that of other species, and has its maximum in the same region as H_2 and H_2^+ . The formation of thism olecular ion occursm ainly through radiative association of H_2 with H^+ , but the ion-molecule reaction of H_2 and H_2^+ is also important in the recombination zone. D issociative recombination and collisions with H are the main destruction routes of H_3^+ .

3.2. The E ect of Dust

The grain temperature depends on the incident radiation, as well as on the grain optical properties, which, in turn, depend on the grain material and size. The grain temperature decreases steeply in the inner 20% region of the nebula, and decreases slow ly for increasing radius. For example, assuming $a_g = 0.01 \text{ m}$, and all other input param – eters the same as those for the standard model, the tem perature for graphite grains decreases from 500 K, in the inner regions, to 50 K, in the farthest zones, while for $a_g = 0.1 \text{ m}$, the grain tem – perature ranges from 180 K to 26 K. For other materials the tem peratures are slightly lower. G rains can survive inside the entire ionized region, except those m ade of PAH, which sublim ate for distances shorter than $0.01R_{max}$.

D irect stellar radiation is much more in portant for the grain heating than absorption of Ly radiation in the ionized region. The grain tem perature is increased by 10% when Ly radiation is included in the grain heating. Such increase leads to a maximum decrease of only 0.3% in the H $_2$ concentration.

The e ect of the dust-to-gas ratio on the relative abundance of H₂ m olecules is shown in Figure 3. The results presented in this gure correspond to m odels with the same input parameters as in the standard model, except for the dust-togas ratio. In the gure, the behavior of (H_2) with $R = R_{max}$ is shown for dierent values of M d=M d with the lowest curve corresponding to a model without dust. Since the reaction on grain surface is an important reaction of H₂ form ation all along the nebula, increasing M $_d=M_q$ results in higher values of n (H $_2$)=n $_{\rm H}$. It can be also noticed that for $R = R_{max}$ between 0.8 and 0.9 the results do not depend on M $_{d}=M_{q}$. This can be explained by the fact that in this region the dom inant processes of H_2 form ation are the charge exchange and the associative detachm ent reactions. As said above, when few or no grains are present, there is a steep decrease of H_2 m olecules in the outer regions of the nebula, due to the decrease of H , which is mainly produced in regions where the electron density is signi cant.

As discussed in x2.1, the key parameter for the form ation of H_2 in the surface of grains is the total available surface area of grains or, in terms of our input parameters, the dust-to-gas mass ratio divided by the grain radius (). For a given dustto-gas ratio, sm aller grains in ply a higher form ation rate (see eq.[3]) of H_2 molecules. On the other hand, sm aller grains have larger temperatures, leading to lower form ation rates. How ever, the e ect due to the grain temperature is sm all; xing the dust-to-ras ratio, a decrease of the size

xing the dust-to-gas ratio, a decrease of the size of the grain leads to an increase of the abundance of H_2 .

As said in x1, the form ation of H_2 m olecules on the surface of grains has been assumed in the interstellarm edium in order to explain the required amount of H_2 molecules. For the ionized region of PNe, however, the associative detachment reaction, as well as the charge transfer of H_2^+ with H are also important processes for the form ation of H_2 molecules. The dust contributes more signi cantly for the production of H_2 molecules for values of > 100 cm⁻¹.

3.3. Othermodels

The distribution of the various species, as well as that of the reactions rates, along the nebula for models with input parameters other than those of the standard model is similar to that shown in Figures 1 and 2. In all models, the maximum H_2 concentration occurs where the gas temperature is around 7500 K and the ionization degree $n(H^+)=n_H$ is approximately 0.1.

In order to com pare the am ount of H₂ produced in di erent PN e conditions, we de ne the param – eter R_M , which corresponds to the ratio between the total hydrogen molecule nuclei m ass (H₂ + H₂⁺) and the total hydrogen nuclei m ass, i.e.,

$$R_{M} = \frac{\frac{R_{R_{max}}}{R_{0}} 4 \left[n \left(H_{2} \right) + n \left(H_{2}^{+} \right) \right] m_{H_{2}} r^{2} dr}{\frac{R_{R_{max}}}{R_{0}} 4 n_{H} m_{H} r^{2} dr}; (7)$$

where r is the distance from the central star, while R_0 and R_{max} are the inner and the outer radius of the ionized region as de ned above, respectively. The inner radius is chosen as $R_0 = 10^{15}$ cm, but its exact value does not a low the results presented here, provided that it is low or than $0.3R_{max}$.

For the standard model, we obtain a value for R_M of the order of 2 10 ⁶. Increasing the dustto-gas ratio (or decreasing the grain size) by a factor of 10, and keeping the other input parameters xed, the fraction of molecular mass increases by a factor around 2. As discussed above, the main parameter regarding the presence of grains is the available total surface of grains. When dust is not present in the gas, R_M is only 10% lower than in the standard model. In fact, R_M increases with $M_d=M_g$ and/or decreasing a_g , but, as said above, for a value of lower than that for the standard model, the presence of grains does not a ect the obtained results for R_M . D i erent grain m aterials leads to a m aximum di erence in R $_{\rm M}\,$ by less than 25% .

Plots of R_M versus T are shown in Figure 4, for dierent values of L , n_H , and M $_d=M_q$. In this gure, it can be seen that R_M increases with T . This is an interesting result, since hotter stars produce m ore high-energy photons. How ever, since the hydrogen photoionization cross section decreases with the photon energy, high-energy photons can travel farther into the nebula before being absorbed by the gas, producing a more extended recombination zone. The increase in R_M with T can explain why H₂ is more frequently detected in bipolar PNe, since these objects have hotter central stars (Corradi & Schwarz 1995). In fact, the relative thickness of the recombination zone is the crucial factor for $R_{\,M}$, and it implies that $R_{\,M}\,\,$ also increases with the decrease of $n_{\,H}\,\,$ and L , as shown in Figure 4. An increase of the gas density by a factor 10 decreases R_M by a factor of the order of 3, while a change of a factor 100 in the lum inosity leads to a change of R_M by a factor around 24. However, R_M increases more steeply with T than with any other input param eter, as can be seen from Figure 4. An increase of T by a factor 10 leads to an increase of R_M by a factor around 6000.

4. CONCLUDING REMARKS

U sing a photoionization code which includes atom s and ions of twelve elements, as well as H , H_2 , H_2^+ , and H_3^+ , we analyze the presence of the H_2 m olecule over the entire ionized region of PN e, up to the region where the ionization degree decreases to 10⁻⁴. The transfer of the radiation em itted by the star, as well as of that produced by the nebula, is treated in detail in order to obtain a self-consistent distribution of the hydrogen m olecule in the nebula. The presence of grains is taken into account, since they are present in the ionized region and can act as catalyst in the form ation of the H_2 m olecule. The e ect of the various reactions that contribute to the form ation and destruction of H_2 is studied.

O ur calculations show that H_2 m olecules can survive inside the ionized region of PNe, particularly in the inner zone of the recombination region, where the ionization degree is of the order of 0.1. In this region, the form ation rate of H_2 m olecules is enhanced due to the high e ciency of the associative detachment of H and H and the charge exchange between H and H_2^+ , while the destruction is not very e ective due to the shielding provided by H atoms. W hen signi cant super cial area of grains is available (i.e., > 100 cm⁻¹), the grain reaction is the main process of H_2 production in regions of low ionization.

Notice that the grain surface reaction is not the only important production mechanism of H_2 molecules in PNe, as it is in the interstellar medium. The associative detachment of H and H, as well as the charge exchange between H and H_2^+ are also important processes for the form ation of H_2 molecules. In the H recombination zone, where the density of H_2 reaches its higher values, the main processes, even when large amounts of grains are present, are via associative detachment and charge exchange. The H and the H_2^+ ions have sim ilar concentrations, except in the outer region of the PNe, and are abundant in regions where both electrons and H ions are present.

The e ect of the stellar tem perature and lum inosity, as well as that of the gas density, dustto-gas m ass fraction, and grain m aterial, on the obtained abundance of the hydrogen m olecule, is also analyzed. Regarding the presence of grains, the available super cial area is the m ain param eter that controls the m olecule form ation e ciency, while the grain tem perature and com position are of m inor im portance.

W e conclude that $R_{\rm M}$, the H_2 to total hydrogen nucleim ass ratio, increases with the relative thickness of the recombination zone. This explains why $R_{\rm M}$ increases with T and decreases with $n_{\rm H}$ and L , them ain e ect being due to the stellar tem perature. The behavior of $R_{\rm M}$ versus T can explain why the emission of H_2 is more offen detected in PNe with bipolar morphology, since these objects have in general hotter central stars.

The presence of molecular material is generally used as evidence that the region is optically thick. However, if the hydrogen molecule can survive within the hot gas, the detection of H_2 em ission is not an indication that a nebula is optically thick. This point has important consequences for the determ ination of PNe distances, since some distance indicators are obtained assuming that the gas is completely optically thick (i.e., all ionizing photons are absorbed). The same applies to the Zanstram ethod for the determ ination of the stellar tem perature.

As shown in this paper, H_2 m olecules can survive not just in very low, but also in m ild ionization regions inside PNe. Thus, a detailed study of the e ect the presence of the hydrogen m olecule on the gas therm al equilibrium is essential for an accurate picture of the physical processes inside PNe. The presence of H_2 m olecules in m ild ionized gas can solve, for example, som e problem s faced by photoionization m odels in explaining em ission lines from the low-ionization regions. Hydrogen m olecules can a ect the physical conditions of the gas, providing the additional heating required to explain the discrepancies. A more realistic estimate for the gas abundance could also be obtained.

In this work we do not include the self-shielding of H₂ in the calculations, since it would be necessary to obtain the distribution of the molecule in its various energy levels in order to estimate the shielding. The e ect of the self-shielding on the photoprocessess can be very important in the outer regions of PN e where the H₂ column density is high. A pproxim ated m ethods for the treatm ent of the self-shielding e ect on the photodissociation of H_2 are available in the literature as, for example, in Van der W erf (1989) and D raine & Bertoldi (1996). Taking this approxim ated methods into account, the computed H_2 concentration can increase up to two orders of magnitude in the outer parts of the ionized region. Therefore the calculated H₂ concentrations in the present work can be considered as lower lim its.

Shocks are frequently invoked in the literature for explaining the H_2 line emission since pure radiative excitation does not account for the observations. The presence of H_2 m olecules in regions of higher excitation level, excited by UV radiation not fully absorbed by inner regions, could account for the line emission, without invoking shocks.

M olecular and atom ic hydrogen m asses for the R ing N ebula (NGC 6720) were derived from observations by G reenhouse, H ayward, & Thronson (1988), resulting in a m olecular to atom ic gas ratio equal to 7 10 ⁵. A ssum ing the equilibrium between form ation on grain surface and photodissociation, these authors conclude that chem ical equilibrium does not account for the detected quantity of H₂. How ever, from ourm ore detailed m odel and assum ing the parameters for the central star and

the gas given by G reenhouse et al. (1988), we obtain $R_M = 2.4 \quad 10^{-5}$, which is reasonably similar to the value obtained by these authors.

W e are thankfulto S.M. Viegas form any useful comments, to E.Roue for kindly providing the transitions probabilities and dissociation fractions for the hydrogen molecule, and to T.Abel for the helpful discussion about reaction rates. W e are pleased to thank the anonymous referee for his valuable comments and suggestions that helped to im prove the paper. The authors thank FAPESP, CNPq, and Pronex/CNPq for the nancial support of this work.

REFERENCES

- Abel, T.G., Anninos, P., Zhang, Y., & Norman, M.L.1997, New A., 2
- Abgrall, H., Roue, E., Launay, F., Roncin, J.-Y., 1994, Can. J. Phys., 72, 856
- Abgrall, H., Roue, E., & Drira, I. 2000, A&A, 141, 297
- Aller, L. H. 1984, Physics of Thermal Gaseous Nebulae (Dordrecht: Reidel)
- Allison, A.C. & Dalgamo, A.1969, Atom ic Data, 1,298
- A rias, L., Rosado, M., Salas, L., & C ruz-G onzalez, I. 2001, ApJ, 122, 3293
- Arthurs, A., & Hyslop, J. 1957, Proc. Phys. Soc., 70, 849
- Aspin, C., et al. 1993, A&A, 278, 255
- Bates, D.R.& Opik, U.1968, J. Phys. B, 1, 543
- Barbw, M.J. 1993, in IAU Symp. 155, Planetary Nebule, ed.R.W einberger & A.Acker (Leiden: Kluwer), 163
- Beckwith, S., Persson, S.E., & Gatley, I. 1978, ApJ, 219, L33
- Black, J.H. 1978, ApJ, 222, 125
- Black, J.H., & Dalgamo, A. 1977, ApJS, 35, 405
- Black, J. H., Porter, A., & Dalgamo, A. 1981, ApJ, 249, 138

- Bordas, C., Cosby, P.C., & Helm, H. 1990, J. Chem. Phys., 93, 6303
- Bowers, C.W., Blair, W.P., Long, K.S., & Davidsen, A.S. 1995, ApJ, 444, 748
- Bohigas, J. 2001, Rev. Mex. A stron. A strof., 37, 237
- Cazaux, S., & Tielens, A.G.G.M. 2002, ApJ, 575, L29
- Cecchi-Pestellini, C., & Dalgamo, A. 1993, ApJ, 413, 611
- C legg, R.E.S., Harrington, J.P., Barlow, M.J., & W alsh, J.R. 1987, ApJ, 314, 551
- Corradi, R.L.M., & Schwarz, H.E.1995, A&A, 293, 871
- Dalgamo, A.& McCray, R.A. 1973, ApJ, 181, 95
- Datz, S. et al. 1995, Phys. Rev. Lett., 74, 896
- Donahue, M . & Shull, J.M . 1991, ApJ, 383, 511
- Draine, B.T., & Bertoldi, F. 1996, ApJ, 468, 269
- Draine, B.T., & Lee, H.M. 1984, ApJ, 285, 89
- Evans, A. 1994, The Dusty Universe (Chichester: Praxys)
- Gaillard, M.J., de Pinho, A.G., Poizat, J.C., Rem illieux, J., & Saoudi, R.1983, Phys. Rev. A, 28, 1267
- Galli, D., & Palla, F. 1998, A & A, 335, 403
- G lover, S.C.O. 2003, ApJ, 584, 331
- G reenhouse, M . A ., H ayward, T . L ., & Thronson, H . A ., Jr. 1988, ApJ, 325, 604
- G nuenwald, R., & Viegas, S.M. 1992, ApJ, 87, 153
- Guerrero, M. A., Villaver, E., Manchado, A., Garcia-Lario, P., & Prada, F. 2000, ApJS, 127, 125
- Gussie, G., & Pritchet, C. 1988, JRASC, 82, 69
- Harrington, J.P., Monk, D.J., & Clegg, R.E.S. 1988, MNRAS, 231, 577
- Herald, J.E., & Bianchi, L. 2002, ApJ, 580, 434

- Hollenbach, D., & McKee, C.F. 1979, ApJ, 41, 555
- Hollenbach, D., & McKee, C.F. 1989, ApJ, 342, 306
- Hollenbach, D., & Salpeter, E.E.1971, ApJ, 163, 155
- Hora, J.L., Latter, W.B., & Deutch, L.K. 1999, ApJS, 124, 195
- Izotov, Y. I., & Kolesnik, I. G. 1984, Sov. Astr., 28, 15
- de Jong, T. 1972, A & A, 20, 263
- Jura, M., & Dalgamo, A. 1971, A&A, 14, 242
- Kanpas, Z., Anicich, V., & Huntress, W. T., Jr. 1979, J. Chem. Phys., 70, 2877
- K astner, J.H., W eintraub, D.A., G atley, I., M erril, K.M., & Probst, R.G. 1996, ApJ, 462, 777
- Kingdon, J.B., & Ferland, G.J. 1997, ApJ, 477, 732
- K ingdon, J., Ferland, G.J., & Feibelm an, W.A. 1995, ApJ, 439, 793
- Kingsbourgh, R.L., & Barlow, R.L. 1994, MN-RAS, 271, 257
- Laor, A., & Draine, B.T. 1993, ApJ, 402, 441
- Lenzuni, P., Natta, A., & Panagia, N. 1989, apj 345, 306
- Lepp, S., Stancil, P.C., & Dalgamo, A. 2002, J. Phys. B, 35, R57
- Li, A., & Draine, B.T. 2000, ApJ, 554, 778
- M cC all, B. J., et al 2003, Nature, 422, 500
- M eixner, M ., Skinner, C . J., K eto, E ., Zijlstra, A ., Hoare, M . G ., A rens, J. F ., & Jernigan, J. G . 1996, A & A , 313, 234
- Natta, A., & Hollenbach, D. 1998, A & A, 337, 517
- Natta, A., & Panagia, N. 1981, ApJ, 248, 189
- O sterbrock, D.E. 1989, A strophysics of G aseous Nebulae and A ctive G alactic Nuclei (M illValey: University Science Books)

- Pequignot, D., Petitjean, P., & Boisson, C. 1991, A&A, 251, 680
- Pottasch, S. R. 1984, Planetary Nebulae (Dordrecht: Reidel)
- Reay, N. K., W alton, N. A., & Atherton, P. D. 1988, MNRAS, 232, 615
- Rowan-Robinson, M. 1986, MNRAS, 219, 737
- Shapiro, P.R., & Kang, H. 1987, ApJ, 318, 32
- Schild, H. 1995, A & A, 297, 246
- Schneider. I. F., Dulieu, O., Giusti-Suzor, A., & Roue, E. 1995, ApJ, 424, 983
- Schneider. I. F., Dulieu, O., Giusti-Suzor, A., & Roue, E.1997, ApJ, 486, 580

Stancil, P.C. 1994, ApJ, 430, 360

- Stasinska, G . 2002, P roceedings of the X III C anary Islands W inter School of A strophysics, in press (astro-ph/0207500)
- Stasinska, G., & Szczerba, R. 1999, A & A, 352, 297
- Stasinska, G., & Szczerba, R. 2001, A&A, 379, 1024
- Stasinska, G., & Tylenda, R. 1986, A & A, 155, 137
- Theard, L.P., & Huntress, W.T.1974, J.Chem. Phys., 60, 2840
- Tielens, A.G.G.M. 1993, in IAU Symp. 155, Planetary Nebule, ed. R.Weinberger & A. Acker (Leiden: Kluwer), 155
- Tielens, A.G.G.M., & Hollenbach, D.1985, ApJ, 291, 722
- Tre ers,R.R.,Fink,U.,Larson,H.P.,& Gautier, T.N., III 1976,ApJ, 209, 793
- Van der Werf, P. 1989, An Study of HII Regions and Dark Clouds, PhD Thesis (Nederlands: Univ. Groningen)
- Van Hoof, P.A.M., Van de Steene, G.C., Beintem a, D.A., Martin, P.G., Pottasch, S.R., & Ferland, G.J. 2000, ApJ, 532, 384
- Viala, Y.P., Roue, E., & Abgrall, H.1988, A&A, 190, 215

- V icini, B ., N atta, A ., M arconi, A ., Testi, L ., H ollenbach, D ., & D raine, B . T . 1999, A & A , 342, 823
- W eingartner, J.C., & Draine, B.T. 2001, ApJ, 548, 296
- W hittet, D.C.B.1992, Dust in the Galactic Environment (Bristol: IOP Publishing)
- Yan, M., Sadeghpour, H.R., & Dalgamo, A. 1998, ApJ, 496, 1044
- Yan, M., Sadeghpour, H.R., & Dalgamo, A.2001, ApJ, 559, 1194
- Zuckerm an, B., & Gatley, I. 1988, ApJ, 324, 501

T his 2-column preprint was prepared with the AAS ${\rm IAT}_E X$ m acros v5.0.



Fig. 1. Concentrations of the species containing H nuclei (upper panel) and gas tem perature (low er panel), for the standard m odel.



Fig. 2. Rates of the most in portant form ation and destruction reactions of molecular hydrogen, for the standard m odel. H $_2$ form ation (upper left): (solid curve), H_2^+ + H (dashed), and H H + H + H + grain (dot-dashed). H₂ destruction (upper right): H₂ + h (photodissociation in two steps; solid), H₂ + h (direct photodissociation; dashed), $H_2 + H^+$ (charge exchange, dot-dashed), $H_2 + e$ (colisional dissociation, dot-dot-dot-dashed), and $H_2 + h$ (photoionization; dotted). H_2^+ form ation (lowerleft): $H_2 + H^+$ (solid), $H^+ + H^+$ (dashed), and $H + H^+$ (dot-dashed). H_2^+ destruction (lower right): H_2^+ + h (producing H^+ + H; solid), H_2^+ + e (dashed), H_2^+ + H (dot-dashed), and H_2^+ + h (producing $2H^{+} + e$; dotted).



Fig. 3.| E ect of the dust-to-gas mass ratio on the calculated abundance of H_2 . Curves are shown for M_d=M_g equal to 0, 10⁻³, 10⁻², and 10⁻¹; the lower curve corresponds to a model without grains, while upper curves correspond to models with higher values of the dust-to-gas ratio. O ther parameters are the same as in the standard PN model.



Fig. 4. Molecular hydrogen mass fraction versus central stellar temperature. Upper panel: The results are shown for $n_{\rm H}=10^3,\,10^4,\,{\rm and}\,10^5$ cm 3 (lower curves correspond to higher $n_{\rm H}$) for two values of : 10^2 cm 1 (solid curves) and 10^4 cm 1 (dashed). Lower panel: Results for L = 500, 3000, and $10^4 {\rm L}$, with $R_{\rm M}$ decreasing with increasing L . O ther parameters correspond to those of the standard model.

R eaction	Ref.	R eaction	Ref.
P hotoionization		P hotodissociation	
H + h ! H ⁺ + e	1	H ₂ +h ! 2H	12
$H_2 + h ! H_2^+ + e$	2	H ₂ + h ! H ₂ ! 2H	13
-		$H_{2}^{+} + h ! H + H^{+}$	14
Photodetachm ent		$H_{2}^{+} + h$! 2 $H^{+} + e$	15
H + h ! H + e	3		
		R adiative A ttachm ent	
Radiative Recombination		H + e ! H + h	16
H ⁺ + e ! H + h	4		
		C harge Exchange	
D issociative R ecom bination	_	$H_{2}^{+} + H_{1}^{+} + H_{2}^{+}$	17
H ₂ + e ! 2H	5	$H_2 + H^+$! $H + H_2$	7
$H_3 + e ! H_2 + H$	6	$He^{+} + H ! He^{+} H^{+}$	18
$H_{3}^{+} + e ! 3H$	6	$He^{+} + H$! $He^{+} + H^{+}$	19
A coopietize D stacky ont		D adjusting A consistion	
	7		7
$u^+ + u^- + u^+ + o^-$	י ד	$\Pi + \Pi : \Pi_2 + \Pi$ $\Pi_2 + \Pi^+ + \Pi_2$	7
<u> </u>	7	112 + 11 : 11 ₃ + 11	1
Ion-M olecule Reaction		G rain Surface Reaction	
$H_2^+ + H_2 ! H_3^+ + H$	8	2H + grain ! H ₂ + grain	20
N autor lingtion			
	0		21
	9 10		21
п ₂ тп : п ₂ тп u ⁺ тп : зи	9,10 0,10		22
$n_2 + n_1 : Sn_1$	9,10	$n_2 + n$; n_2	20
$n_3 + n_1 \cdot 2n_2$	9,10 0,10	$2\pi_2$: π_2 + 2π	23 16
$n_3 + n : n_2 + 2n$	9,10		22
Colicional Ionization			22 11
	Q	$u_{2} + u_{2} + u_{1} + u_{1} + u_{2}$	11 7
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	9 11	$H^{+} + H_{0} I H + H_{0} + H^{+}$, 11
$2H + H + H + H + H^+ + 6$	11	$H^{+} + H_{0} + H^{+} + 2H_{0}$	11
2		··· 3 · ··· 2 · ··· · 2··· 2	

Table 1 Chemical Reactions and Physical Processes

References. (1) O sterbrook (1989); (2) Yan, Sadeghpour, & Dalgamo (1998, 2001); (3) de Jong (1972); (4) Pequignot, Petitjean, & Boisson (1991); (5) Schneider et al. (1994, 1997); (6) M cC allet al. (2003), with branching ratios given by Datz et al. (1995); (7) G alli & Palla (1998); (8) Theard & Huntress (1974); (9) Lepp, Stancil, & Dalgamo (2002); (10) following Dalgamo & M c-C ray (1973) we assume this rate equal to that of the reaction $H^+ + H + 2H$; (11) Hollenbach & M cK ee (1989); (12) A llison & Dalgamo (1969), with LTE distribution for the H₂ energy levels; (13) A bgrall et al. (1994) and A bgrall, Roue, & Drira (2000), with LTE distribution for the H₂ energy levels; (14) Stancil (1994), with LTE tting to T = 8400 K; (15) Bates & Opick (1968); (16) A bel et al. (1997); (17) K arpas, A nicich, & Huntress (1979); (18) Jura & Dalgamo (1971); (19) A rthurs & Hyslop (1957); (20) see text; (21) D onahue & Shull (1991); (22) Izotov & K olesnik (1984); (23) Shapiro & K ang (1987).