The cosm ic infrared background (C IR B) and the role of the \local environm ent of galaxies" in the origin of present-day stars

David E lbaz¹, D elphine M arcillac¹, and Emm anuel M oy¹

CEA Saclay=D SM =D A P N IA = Service d'A strophysique, O m e des M erisiers, F -91191 G if-sur-Y vette C edex, France

A bstract. A combination of evidence is presented suggesting that the majority of the stars in today's galaxies were born during a lum inous infrared phase (LIRP) triggered by the local environment of galaxies. The CIRB is a fossil record of these LIRPs and therefore rects the in uence of triggered star formation through galaxy-galaxy interactions, including non merging tidal encounters. This scenario, in which galaxies experienced several LIRPs in their history, is consistent with the measured redshift evolution of the cosm ic density of star formation rates and of stellar masses of galaxies.

1 Introduction

Stars represent only 15% of the cosm ic baryonic density, itself only equal to about 4% of the critical density ($_{\rm b}$ ' 0.04), and are unequally distributed into spheroids (10% of b, including spiral bulges) and disks (5% of b, Fukugita, Hogan & Peebles). It is usually assumed that disk stars formed quiescently while bulge stars form ed m ore e ciently and rapidly, as suggested by their redder colors and overabundance in -elem ent over iron ratio, typical of a SN II origin. However recent studies of the history of the star form ation of the disk of the M ilky W ay indicate that during the last 2 Gyr it has experienced about ve m a prevents of star form ation, starbursts, whose signatures can be found in the peaked ages of these open clusters (de la Fuente M arcos & de la Fuente M arcos 2004, and references therein). As a result we may wonder whether quiescent star form ation did play a major role in the form ation of present-day stars at all. There are some evidence that star form ation mostly takes place in globular clusters and is rarely isolated. These clusters are thought to evolve into unbound stellar associations, which evolve and dissolve in a time-scale of about 50 M yr. This tim escale is longer than the lifetime of massive stars which dom inate the lum inosity of starbursting galaxies or regions of galaxies. Hence, it is logical to expect that if star formation occured mainly in starburst episodes within galaxies, then the bulk of galaxies lum inosity will be absorbed by dust in the giant molecular clouds, while if star form ation is quiescent then only a minor fraction of a galaxy's lum inosity will be a ected by dust extinction. We will argue in the following that there is presently a solid collection of evidence suggesting that most stars that we see in the local universe form ed during starburst episodes.

A major peace of evidence for that comes from the detection of a strong di use cosm ic infrared background (CIRB, Puget et al. 1996, Hauser & Dwek and

2 E lbaz, M arcillac, M oy

references therein) which is majoritarily produced by lum inous infrared phases (LIRP) within galaxies located around z = 0.7, for the peak of the CIRB at 240 m is due to galaxies at z 2 and above (Elbaz 140 m, while the et al. 2002, Chary & Elbaz 2001). We have introduced the term LIRP instead of the classical one, lum inous and ultra-lum inous infrared galaxies, i.e. LIRGs and ULIRGs, because we wish to emphasize the idea that the scenario that is emerging from the study of distant galaxies is that LIRGs do not represent a population of galaxies that would require to be studied independently in order to determ ine which present-day galaxies are the rem nants of these LIRGs, but what is suggested instead is that they illustrate the om nipresence of rapid and e cient star form ation as a leading process in shaping the present-day universe, i.e. that any galaxy that we see today must have experienced a phase when it radiated the bulk of its light in the infrared (see also E lbaz & Cesarsky 2003). This phase should not be restricted to LIRGs and ULIRGs, i.e. galaxies with infrared (IR) lum inosities larger than 10^{11} L $\,$ or star form ation rates (SFR) larger than 20 yr¹, since the closest starburst M 82, for example, presents a spectral energy М distribution typical of m ost LIRGs, with the bulk of its lum inosity radiated in the IR although its lum inosity is only 4 $\,10^{10}\,{\rm L}\,$.

We present evidence suggesting that the bulk of local stars form ed during a LIRP. A spectroscopic diagnostic is used to quantify the typical duration of this phase and the am ount of stars that form ed during it. From the combination of both we will advocate that not only did all galaxies experience a LIRP but that they must have experienced several of them. Finally we will discuss the physical origin of the LIRP and present evidence that a major event in the lifetime of galaxies was probably underestim ated: the e ect of the \local environment of galaxies" (LEG) and its impact in terms of driving the conversion of gas into stars through passing-by galaxies.

2 Lum inous IR phases and the origin of present-day stars

The detection of a CIRB came as a surprise since local galaxies only radiate 30% of their bolom etric lum inosity in the m id to far IR range, i.e. sharing as a com m on origin stellar photons reprocessed by dust above 3 m.W ith about half of the di use background light radiated above and below this wavelength cuto, the extragalactic background light tells us that in the past, mapr star form ation events were strongly a ected by dust extinction even when galaxies were less metal rich. Deep surveys in the mid infrared (15 m with ISOCAM onboard ISO, E lbaz et al. 1999) brought independant evidence that infrared was m ore ubiquitous in the past. These surveys detected ten times m ore objects at faint ux levels than expected from the extrapolation of the local universe (no evolution models). These galaxies turned out to belong to the class of LIRGs and ULIRGs discovered by IRAS in the local universe but located at a median redshift of z 0.7. They do not exhibit any optical signature of such strong SFRs neither in their optical colors nor in their em ission lines, except if careful correction for extinction is applyied using the Balm er decrem ent (Cardiel et al. 2003,

Cosm ic infrared background 3

F bres et al. 2004). Only less than 20% of them were found to harbor or be dom inated by an active galactic nucleus (AGN; see Fadda et al. 2002). Unexpectedly, because of the complex and multiple physicalorigins of the m id and far IR photons (PAHs, Very Sm all G rains, B ig G rains), local galaxies do exhibit a strong correlation between their m id and far IR lum inosities over three decades in lum inosity, including the extrem e LIRGs and ULIRGs (E lbaz et al. 2002). W hen applyied to galaxies up to z 1 these correlations can be used to derive far IR lum inosities, i.e. SFRs, which are consistent with those derived from the radio, using the radio-far IR correlation, suggesting that these correlations are still valid at these redshifts (E lbaz et al. 2002).



F ig.1.a) cosm ic star form ation rate (C SFR) as a function of redshift and universe age, in a H₀ = 50 km s⁻¹ M pc⁻¹ and $q_0 = 0.5$ cosm ology (Fig.15 of C hary & E lbaz 2001). The data represent the SFR density derived from UV or H uncorrected for extinction (the various authors are quoted on the plot and the references can be found in C hary & E lbaz 2001).b) cosm ic stellarm ass history (C SM H) or redshift evolution of _2, the cosm ic stellarm ass density over critical density (cosm ology: = 0.7, _m = 0.3, H₀ = 70 km s⁻¹ M pc⁻¹). The data are from D ickinson et al. (2003). See text for description.

The median SFR of the galaxies responsible for the evolution of the mid IR 50 M yr ¹ and their contribution to the CIRB from 100 to 1000 m counts is is derived to be as large as two thirds of its measured intensity. Hence the bulk of the CIRB results from LIRPs at redshifts of the order of z 1, but due to cosmological dimming the in uence of more distant galaxies is more limited to the large wavelength tailof the CIRB. Som em odels have been designed to reproduce number counts in the mid IR (ISOCAM, 15 m), far IR (ISOPHOT-90, 175 m) and sub-mm (SCUBA, 850 m) together with the CIRB which can be used to derive the cosm ic star form ation history of the universe (CSFH) una ected by dust extinction and to disentangle the relative roles of rapid (LIRPs) and quiescent star form ation. In Chary & Elbaz (2001), we suggested to de ne a region delimiting all possible histories of star formation (see Fig. 1a), instead of a single line for any favorite m odel that would not represent the uncertainties of the m odel and observational constraints. One way to check the robustness of such m odels consists in comparing to direct measurements the resulting cosm ic stellar

4 Elbaz, Marcillac, Moy

m asses history (C SM H), i.e. the redshift evolution of 2, the m ass density of stars per com oving volum e over the critical density of the universe. The hatched area of F ig. 1a which represents the range of possible C SFH was converted into a range of possible C SM H in F ig. 1b, with the thin plain line showing the m ean value and the range of possible histories delimited by the two dot-dashed thin lines. The m odel to the data collected in D ickinson et al. (2003), where stellar m asses were directly m easured from optical-near IR m agnitudes.

Before deriving any conclusion, we wish to remind our assumptions: the CSFH was derived assuming mid to far IR correlations similar to locally (in agreement with the radio), the AGN fraction was supposed to make a minor contribution (see above), we assumed a universal IMF (we used the one of G ould et al. 1996, which combines a Salpeter IMF with the now standard in exion of the IMF below 1 M). If these assumptions are indeed justimed, then the t of the data in Fig. 1b illustrates the fact that the photons emitted by star forming regions do relect the stellar mass building of galaxies and as a consequence, it is now possible to derive which fraction of present-day stars were formed in a LIRP. The model CSFH was separated into three components shown as thick grey lines in the Fig. 1b, with 63% of present-day stars born during an IR phase of the LIRG type, which would be the dominant mode of star form ation in the universe. The shape of the redshift evolution of $\frac{1}{2}$ in plies that 80% of present-day stars were born below z= 2, and 50% below z= 1, most of which during a LIRP, leaving little room for quiescent star form ation.

If m ost of today's stars form ed during a LIRP and if this phase re ects e - cient and rapid star form ation then this suggests that som e \positive feedback", i.e. triggering, m ight be at play. This idea is com forted by the m orphology and local environm ent (LEG) of distant LIRGs. It is well-known that galaxies lie in large-scale structures m ade of walls, lam ents and clusters but LIRGs tend to appear exclusively in high density environm ents around z 0.7. The deepest ISO CAM survey was perform ed in a region of 27^{02} centered on the Hubble D eep Field N orth (HDFN), detecting 95 galaxies am ong which 47 lie above the com pleteness lim it of 0.1 m Jy.

The histogram of eld galaxies presents several redshift peaks. The location and the extent of these peaks can be quanti ed by adopting a tresholding S/N ratio of 3, where the "background" is sim ply the redshift distribution sm oothed with a gaussian with = 15,000 km s¹. M onte-C arlo simulations perform ed by extracting 100 random sam ples from the real distribution of eld galaxies show that the probability of reproducing by chance the level of clustering of the ISO CAM galaxies is much less than one chance over ten (largest error bars). These results are illustrated on Fig. 2a, where we plot the fraction of ISO CAM galaxies included in redshift peaks above a given S/N ratio as a function of this S/N. The corresponding curves for the eld galaxies and the simulated sam ples are also indicated. The thin (resp. thick) error bars show the 68 % (resp. 90 %) con dence level. The strong clustering of ISO CAM galaxies illustrates that at z 0.7, galaxies experiencing a LIRP are more clustered on average than eld galaxies. On the contrary, the study of LIRPs in the local universe using the



F ig.2. a) Cumulative fraction of galaxies located in a redshift peak of a given S/N, i.e. num ber of galaxies in all redshift peaks above a given S/N (see text).b) Equivalent width of the high-order Balm er absorption line, H 8, as a function of the 4000 A break. D ata with error bars are a sub-sam ple of z $0.7 \ LIRGs$. D ots were generated by a M onte C arlo simulation of 200,000 galaxies. The darkest region is populated by galaxies dom inated by continuous star form ation, or bursts older than 2 G yr.

shallower ELAIS survey (O liver et al., in these proceedings) shows that they are less clustered than eld galaxies locally. The natural explanation for this behavior is that galaxies m ight experience a LIRP when located in a region which is collapsing over a large scale, i.e. star form ation would be triggered by large-scale structures in the process of form ation.

How much stellar mass does a LIRG form ? We addressed this question using an optical spectroscopic diagnostic combining the equivalent width of the high order Balmer absorption line H8 to the 4000 A break to caracterize these starburst events (M arcillac et al., in prep.). The advantage of using H 8 instead of H is that it is in a bluer side of the spectrum, hence less a ected by sky lines, and that its underlying nebular emission line can be neglected. Fig. 2b shows a sub-sample of 22 LIRGs, at z 0.7, selected in three di erent locations of the sky compared to a Monte Carlo simulation of 200,000 galaxies using the code of Bruzual & Charlot (2004) such as those used by Kau mann et al. (2003) to reproduce the behavior of local galaxies in the Sloane survey. This simulation can be used to determ ine which histories of star form ation would reproduce these galaxies. It is found that only galaxies presently experiencing a burst of star form ation can fall in this region of the diagram and that this burst lasts approximately 10⁸ years and produce about 10% of the stars of the galaxies. These numbers were both derived by the simulation but they perfectly agree with the measured median mass of the ISOCAM galaxies of $5 \ 10^{10}$ (from Dickinson et al. 2003). Indeed in 10^8 years and with their median М 50 M yr ¹, they produce 5 10⁹ M of stars, i.e. 10% of the galaxy SFR mass. The mass of new ly form ed stars is also consistent with the typical mass of molecular gas found in local LIRGs. The model used in the Fig. 1 predicts that nearly 50% of present-day stars were form ed in a LIRP below z 1, hence

6 E lbaz, M arcillac, M oy

if this phase only makes up 10% of a galaxy's stars then the majority if not all of today's galaxies experienced up to ve or even more LRPs. This suggests that the dense surrounding of galaxies can trigger successive lum incus R phases, LIRPs, in galaxies. Finally, the optical morphology of z $0.7 \ \text{LRG}$ s derived from HST-ACS observations (E baz et al., in prep, see also Hammer et al. in these proceedings) shows that less than half of them look like the major mergers that we see locally in LIRGs. Major mergers might not be num erous enough to explain such a behavior and passing-by galaxies might play an important role by triggering strong star form ation events through tidal e ects. Hence the local environment of galaxies, or LEG, might be considered as a better candidate to understand the origin of distant LIRGs. The cosm ic star form ation history is therefore probably strongly dependent on the local density of galaxies which will also possibly determine their nalm orphology, i.e. spirals versus ellipticals, instead of the standard picture of the merger of two massive disks.

The recently launched Spitzer satellite will provide ideal observations to check the robustness of this scenario by observing larger patches of the sky (in particular the SW IRE legacy program), low ering the e ect of cosm ic variance, and to extend the study of lum inous IR phases to higher redshifts and low er lum inosities (with the M IPS G T O and G O O D S legacy program).

References

- 1. G.Bruzual, S.Charlot: MNRAS 344, 1000 (2003)
- 2. N.Cardielet al: ApJ 584, 76 (2003)
- 3. R.R. Chary, D. Elbaz: ApJ 556, 562 (2001)
- 4. R. de la Fuente M arcos, C. de la Fuente M arcos: to appear in New A stronom y (astro ph=0401360)
- 5. M .D ickinson et al: A pJ 587, 25 (2003)
- 6. D.Elbaz et al: A & A, 351, L37 (1999)
- 7. D.Elbaz et al: A & A, 384, 848 (2002)
- 8. D. Elbaz, C.J. Cesarsky: Science 300, 270 (2003)
- 9. D.Fadda et al: A & A 383, 838 (2002)
- 10. H.Flores et al.: A & A 415, 885 (2004)
- 11. M. Fukugita, C. J. Hogan, P. J. E. Peebles: ApJ 503, 518 (21998)
- 12. A.Gould, JN.Bahcall, C.Flynn: ApJ 465, 759 (2003)
- 13. M .Hauser, E.Dwek: ARAA 39, 249 (2001)
- 14. G.Kau mann et al: MNRAS 341, 33 (2003)
- 15. J.-L. Puget et al.: A & A 308, L5 (1996)