A utom ated C lassi cation of 2000 B right IR A S Sources

Ranjan Gupta

IUCAA, Post Bag 4, Ganeshkhind, Pune-411007, India

rag@iucaa.ernet.in

Harinder P. Singh

Department of Physics & Astrophysics, University of Delhi, Delhi-110007, India

hpsingh@physics.du.ac.in

K.Volk and S.Kwok

Department of Physics & Astronom y, University of Calgary, 500 University Dr., N.W., Calgary, Alberta, Canada T2N 1N4

volk@iras.ucalgary.ca ; kwok@iras.ucalgary.ca

ABSTRACT

An Arti cialNeuralNetwork (ANN) scheme has been employed that uses a supervised back-propagation algorithm to classify 2000 bright sources from the Calgary database of IRAS (Infrared A stronom icalSatellite) spectra in the region 8 m to 23 m. The database has been classified into 17 prede ned classes based on the spectralm orphology. We have been able to classify over 80 percent of the sources correctly in the rst instance. The speed and robustness of the scheme e will allow us to classify the whole of the LRS database, containing more that 50,000 sources, in the near future.

Subject headings: infrared: galaxies | m ethods: data analysis

1. Introduction

Infrared A stronom ical Satellite Low Resolution Spectrom eter (LRS) recorded spectra of som e 50,000 sources in blue (8 15 m) with = 40 and in red (13 23 m) with a resolution 20. A total of 5425 objects with better quality spectra were included in the A tlas of low-resolution IRAS spectra (1986, hereafter the A tlas). Volk & Cohen (1989a) published spectra of 356 RAS point sources with F (12 m) > 40Jy that were not included in the Atlas. These brighter sources were classified into nine classes based upon the spectral morphology. Sixty percent of the sources have silicate emission and red-continuum spectra associated with H II region sources. No emission-line sources formed part of the set of 356 spectra. This sample was also used to test the classification scheme of IRAS sources based on broad-band colors. Classificate extreme found for 338 of the sources in the sample. The remaining 18 sources had either extremely noisy or incomplete spectra. It was found that some class of sources overlapped on the color-color diagram s and, therefore, the nature of some of the IRAS sources could not be determined from the IRAS photom etry.

Volk et al. (1991) published an additional 486 spectra belonging to sources with 12 m uxes between 20 and 40Jy that were also not in the Atlas. Classi able spectra were found for 424 sources. The spectra were classi ed into nine groups as in Volk & Cohen (1989a) that describe the astrophysical nature of these sources. K wok, Volk & Bidelm an (1997) processed 11,224 spectra (including sources in the Atlas), corresponding to a ux limit of 7Jy at 12 m. These spectra were also classi ed by-eye and put into nine classes based on the presence of em ission and absorption features and on the shape of the continuum. They identi ed optical counterparts of these IRAS sources in the existing optical and infrared catalogs and listed the optical spectral types if they were known.

It is evident that large databases like the one referred to above require autom ated schem es for any analysis. A rti cialN euralN etw orks (ANN) have been em ployed extensively in several branches of A stronom y for autom ated data analysis (Lahav & Storrie-Lom bardi, 1994). ANN have been used previously by the IUCAA group in three distinct areas of stellar A stronom y. They have been applied to classify digitized optical and ultraviolet spectra (G ulatiet. al. 1994a,b; Singh et al. 1998); to compare a set of observed spectra of F & G dwarfs with a library of synthetic spectra (G ulatiet al. 1997a, G upta et al. 2001); and to determ ine the reddening properties of hot stars from the low-dispersion ultraviolet spectra (G ulatiet al. 1997b).

We have attempted to classify 2000 brightest sources from the Atlas into 17 classes by means of Arti cial Neural Networks. In the next section we describe features of the 17 new classes. In Section 3, we present details of the ANN scheme. Results are discussed in Section 4 and important conclusions of the study are presented in the last section. Further, an appendix has been added just before the reference section, to explain the general ANN architecture for the bene t of the readers.

2. Spectral C lasses

The set of 2000 spectra were classi ed into the 17 classes by-eye by one of us (K.Volk). These classi cations are assumed to from the reference set of \correct" classi cations. Such by-eye classi cation has an element of subjectivity in it, and is more apt to have problem s when the spectra are noisy. This set of 2000 sources are the brightest sources in the LRS atlas and have the least problem s with noise or spectral peculiarities. This makes the by-eye classi cation as accurate as possible despite the element of subjectivity.

The groups used for spectral classic cation are an extension of the nine classes described in Volk & Cohen (1989a). The current scheme introduces more classes, for three reasons:

(a) The $H \amalg$ region sources" group shows a variety of features, and so separate classes have been m ade for the m ore obvious feature types so that these classes are, one hopes, m ore uniform;

(b) A few new classes have been created within the 10 m silicate feature and 11.3 m SiC feature groups to divide the objects with weaker features from those with stronger features; and

(c) Som e sm all groups of unusual sources can be identied from the \usual/unknown" group of Volk & Cohen (1989a), hence these were given their own classes here such as class 14 for 21 m feature sources.

The division between the spectra with \stronger" and \weaker" features was made at LRS type 25 or 45 (for silicate em ission sources and SiC em ission feature sources respectively) since the original LRS classes are based upon the feature strength. Thus the spectra with strong features, LRS types 25 to 29 or 45 to 49, are separated from the spectra with weaker features, LRS types 21 to 24 or 41 to 44. All objects with LRS classes near the dividing line were exam ined by-eye to make sure that the LRS type accurately relative ects the feature strength. In some cases this led to spectra being placed into the \stronger" or \weaker" feature groups even when the LRS type is not as expected. In the large majority of cases no problem with the LRS type was found.

A short sum m ary of the 17 classes is given in Table 1, with representative spectra plotted in Figure 1. A detailed description of the di erent classes and their relation to the original LRS types now follows:

(0) Line em ission sources

This class is for spectra with strong emission lines (line peak to continuum ratio of 1 or larger) and no obvious dust emission or absorption features. These are nearly all planetary nebula sources. A few cases of H II region sources that may show the 12.8 m [N eII] line could have been included here, but were left in the other \H II region" groups since the 12.8 m emission line seen in LRS spectra is sometime due to an instrum ental problem rather than being an actual emission line, and since the possible [N eII] lines were relatively weak. C om paratively few H II region spectra have strong enough emission lines as observed by the LRS instrum ent to be potentially included in this group. In the original LRS classi cation, these are generally classi ed as types 81 to 96.

The prototype spectrum in Figure 1 shows several emission lines: 9.0 m [A rIII], 12.8 m [N eII], 15.5 m [N eV], and 18.7 m [SIII]. Other objects in this class tend to have fewer emission lines than this example.

(1) Sources with stronger 11.3 m SiC em ission features

In this group one has evolved carbon stars undergoing mass loss and showing a strong 11.3 m emission feature. In the original LRS classication these would be type 45 to 49 sources, implying the feature peak to estimated continuum ratio is larger than 1.648.

(2) Stellar continuum sources

Here is the class for stars without circum stellar dust shells that radiate signi cantly in the 8 to 23 m region, including Vega, Sirius, A kdebaran, and Antares. Most of these sources are of K-type to early M-type. Only a few have spectral types earlier than K 0. In the original LRS classic cation these would be type 16 to 19 sources

(3) Featureless sources with cooler colour tem peratures

This group is as in Volk & Cohen (1989a). The continuum colour temperature is lower than expected for the photospheric emission of any normal star, including late M -type stars. A side from the unusually low colour temperature for a stellar photosphere there is no overt sign of any dust emission. M any of these sources have optical counterparts among known M ira or irregular variables of M -type. In the original LRS classic cation these would be type 12 to 15 sources. All featureless continuum sources with F ratios of less than 3 between 7.9 and 11 m but which still have declining continua over the LRS wavelength range are placed in this class.

(4) Sources with UIR features at 7.6/8.5/11.3/12.5 m

This group includes sources with the UIR features, except for those with very steep (\red" according to the LRS atlas de nition) continua which form class 16. These would ideally be type 80 sources in the original LRS classic cation, but there was considerable confusion with other types of objects.

As long as the spectrum signal to noise ratio is reasonable it is possible to objectively distinguish between UIR feature spectra and silicate absorption, as discussed in Volk & Cohen (1989b).

(5) Sources with 10 m silicate absorption

This class includes objects with silicate absorption features, but we attempt to exclude cases of compact H II regions with foreground silicate absorption which have much lower continuum colour temperature than the \norm al" silicate absorption feature sources such as IRAS 01304+ 6211 (see Figure 1). If the spectral continuum is rising from 7.6 to 23 m and there is a silicate absorption feature at 10 m the spectrum is put in class 9. In the original LRS classic cation class 5 sources correspond to types 31 to 39. An additional rement is that sources that are in transition between silicate em ission and silicate absorption at 10 m are grouped in class 13.

(6) Sources with strong silicate em ission at 10 m

Spectra with a falling continuum over the LRS wavelength range and having a silicate em ission feature with a peak greater than 1.628 times the continuum are in this class. These would be types 25 to 29 in the original LRS classi cation. Sim ilar spectra but with weaker features are put into class 12.

(7) Sources with low temperature (100 K) dust continuum em ission

In this group are mostly HII region spectra which do not show very strong features due to silicates or the UIR features. The continuum must rise over the LRS wavelength range from 7.6 to 23 m for a spectrum to be included in this class, corresponding roughly to a colour temperature of less than 100 K, but generally the continuum shape cannot be described by a single colour temperature. In the original LRS classic cation these would have been given some type in the 60's, 70's, or 80's. In a few unusual cases there can be other types of spectra mixed into this group. When there are accompanying strong features such as the 10 m feature in emission or absorption the spectra are put into classes 8, 9, or 12.

(8) Sources with silicate em ission and a low tem perature dust continuum

Here are spectra with continuum shape sim ilar to that of class 7 but with a clear 10 m silicate em ission feature. In principle this corresponds to types 61 to 69 in the original LRS classi cation. Some of the sources are H II regions, but others are young planetary nebulae such as H b 12, V y 2–2, and Sw St–1; others are their im mediate progenitors caught in the phase between the asym ptotic giant branch phase and the planetary nebula phase (\protoplanetary nebulae" or PPN s) such as H D 161796 and IRAS 18095+ 2704. The spectrum of 18095+ 2704 is shown in F igure 1; this has unusually strong features and the continuum is not as steeply rising as is typically the case for spectra in this class.

(9) Sources with silicate em ission and a low tem perature dust continuum

This is another class where the continuum is required to rise over the LRS wavelength range from 7.6 to 23 m. Here we include spectra showing the 10 and 18 m silicate features in absorption. These are nearly all compact H II region sources with (presum ably) foreground cold dust in the associate m olecular cloud. In the original LRS classic cation these would be type 71 to 79 sources.

(10) Extrem e carbon stars

The prototype for this group is AFGL 3068 (IRAS 23166+1655), which is known to be a carbon star with an extrem ely optically thick dust shell. The spectra are selected for this group on the basis of having low colour tem peratures (of order 300 K) in the 10 M region of the spectrum and of having no strong emission or absorption features. In some cases there is some type of weak feature around 11 m which m ight be due to the 11.3 M SiC feature in emission or weak absorption. These objects are discussed in Volk, Kwok, & Langill (1992). (11) Low colour tem perature featureless spectra

The group of spectra have low colour temperature (about 150 to 200 K) featureless continua. They have noticeably lower colour temperatures than the extrem e carbon star spectra see F igure 1) but peak som ewhere in the LRS wavelength range and decline thereafter, unlike the H II region spectra in classes 7, 9, and 16. A few of these sources are known to be either young planetary nebulae or PPN s, and they all seem to have carbon-based dust shells. These objects are discussed by Volk, Kwok & W oodsworth (1993). In the original LRS classic cation they were generally assigned type 05 or 50.

(12) Sources with weaker silicate em ission features

This group of spectra is the analogue of class 6, but they have feature peak to continuum ratios of less than 1.628. In the original LRS at las classic cation these would be type 21 to 24 spectra. In various cases the feature was not detected at all in the original LRS at las classic cation and these were generally classic ed as type 15 to 18.

(13) Sources with interm ediate optical depth silicate dust shells

Prototype spectra for this class include IRAS 19192+0922, IRAS 17125 4814, IRAS 15119 6453, and IRAS 16546 4047 (see Figure 1). All these spectra have continua which fall between 7.6 and 23 m. The optical depth of the dust shells in these objects is such that the 10 m feature is in transition between emission and absorption while the 18 m feature is still in emission. In the original LRS classi cation these were often confused with SiC emission spectra, and so given erroneous types of 41 to 45.

(14) Source with the 21 m feature

A small number of sources are known to have a dust feature at 21 m, one possible identication of which is TiC grains (von Helden et al., 2000). The feature was discovered in the LRS spectra. Less than 20 such sources are known, the brighter of which are used to de ne the group here. All sources in this group are carbon-rich PPNs. A few extreme carbon stars and the planetary nebula IC 418 appear to have very weak 21 m features, but these require the Infrared Space Observatory to detect and so they are not included here.

(15) Sources with weaker 11.3 m SiC em ission

All spectra from carbon stars with the 11.3 m feature but with a feature to continuum ratio of less and 1.628 are put in this group, which is closely analogous to class 1. In the original LRS classic cation these would be of type 41 to 44.

(16) Cool continuum sources with strong UIR features

Any spectrum which has a continuum that rises from 7.6 to 23 m (allowing for the possibility of a strong 7.6 m feature at the short wavelength end of the spectrum) and strong U IR features at 7.6 and 11.3 m is placed in this class. These are H II region spectra with U IR features. In the original LRS classic cation these would be of type 80 or 81. In some cases is is possible that there is an underlying silicate absorption component as well as the U IR features, causing a very large rise at the short wavelength end of the LRS spectrum. How ever these spectra lack any 18 m absorption feature which makes the presence of strong 10 m silicate absorption less likely.

3. ANN Scheme

In this analysis, we have used the multi-layer back-propagation (M BPN) neural network schemewith supervised learning as described in Gulatietal. (1994a,b). This scheme requires a set of spectra to be predened as the training set and should include all the classes of the unknown set that the network is supposed to classify. The algorithm trains on this training set and subsequently in the test phase, classing as the unknown test set into the predened classes.

A training set of 170 spectra having 10 sources for each training class was set aside for training purposes. Fig. 1 shows a representative sample of each of these 17 classes. Each spectrum consists of 93 ux values in the range of 8 23 m. We may mention that 148 num bers of the 170 training spectra were included in the 2000 test spectra for validity checks.

The ANN scheme involved using this set of 170 input spectra with 17 assigned classes (0 to 16) to train the ANN algorithm. We refer to these classes as the catalog classes. The ANN con guration used was 93-16-16-17, im plying 93 data points for each of the 17 output spectral classes with two hidden layers of 16 nodes each (a detailed explanation on the ANN architecture is given in the Appendix). The training session involved an iterative procedure with the network weights getting modil ed at each iteration. During an iteration,

the computed output and the desired output were compared and the resultant error was then utilized to modify the network weights for the next iteration using a back propagation algorithm. The learning or training was stopped once the error wasm in in ized to a prede ned level and the network weights were considered to be frozen (Gulatiet al. 1994a). A learning curve for 10000 iterations is given in Fig. 2 (left panel).

Subsequent to the above training session, the test session uses the frozen weights determ ined above to perform the classic cation of the 2000 test set of spectra into 17 catalog classes.

4. Results and Discussion

A general picture of the result of our ANN classi cation scheme is given in Fig. 2 (right panel) in the form of a histogram. A total of 1,618 spectra have been classi ed correctly out of the total sample of 2000 source spectra indicating a success rate of 80% at the rst instance. Table 2 provides a sum mary of the ANN classi cation of the 2000 test sources into the 17 groups or classes. Table 3 lists 382 of the misclassi ed sources with their catalog class and the corresponding ANN class. In the following, we describe some ner details of the classi cation accuracy for individual classes 0 to 16.

Out of a total 23 spectra for class 0, 19 were classi ed correctly. From the four that were m isclassi ed, one each was classi ed as class 7, 8, 13, and 16. Table 3 lists the 4 spectra of class 0 that were m isclassi ed. Source 22036+ 5306 has been put in class 16 by the ANN probably because of the strong feature at around 8 m, typical of class 16.

Out of a total 136 spectra for class 1, 113 were classi ed correctly. From the 23 that were m isclassi ed, 11 were m isclassi ed into class 13; 6 were m isclassi ed into class 15, 3 were m isclassi ed into class 4, 2 were m isclassi ed into class 6 and 1 was m isclassi ed into class 14. The m isclassi ed sources of class 1 are listed in Table 3. Out of these, 18 spectra seem to belong to the class assigned by the ANN and not the catalog class, specially those put into classes 13 and 15. ANN C lassi cation accuracy of class 2 has the best agreem ent with the catalog class. Out of 89 spectra of class 2, 88 spectra were classi ed correctly while 1 spectrum was m isclassi ed into class 3.

O fthe 94 spectra for class 3,85 were classi ed correctly. O fthe 9 that were m isclassi ed, 5 were m isclassi ed into class 2 and 1 each into classes 1,12,13 and 15. At least 1 source (20056+1834) has a spectrum which does not seem to belong to class 3 and has been put in class 13 by the ANN. O fa total of 87 sources of class 4, 34 were correctly classi ed while 53 were m is classi ed; 22 into class 5, 11 into class 16, 10 into class 13, 4 into class 15, 2 each into lass 9 and 14 and 1 each into class 2 and 10. A s m entioned in Sec. 2, class 4 corresponds to type 80 sources in the original LRS classi cation in which there was a lot of confusion with the other types of sources. ANN put 22 sources into class 5, which has sources with 10 m silicate absorption feature. We note that in the training sample of class 4, there are at least two sources (16367 4701 & 19327+3024) which m ight be resulting in contam ination of this class with class 5 and hence less than e cient training for class 4.

Out of a total of 103 spectra for class 5, 88 were correctly classi ed. 7 spectra were m isclassi ed into class 16, 4 into class 4, 2 into class 9 and 1 each into class 13 and 14. ANN has correctly picked up at least 3 spectra from the catalog that should be put in class 4 (Table 3).

C lass 6 (strong 10 m feature) has the largest sample of 735 sources. Out of these 679 were classi ed correctly and 33 were m isinterpreted as class 12 (weak 10 m feature). Sim ilarly, class 12 has 198 spectra out of which 15 were m isclassi ed as class 6, while 27 were m isclassi ed into class 3. As discussed later, if one were to treat a weak and a strong 10 m feature sources as belonging to a single class, percentage of correctly classi ed sources will be in excess of 90% for these classes.

For class 7, out of the 53 sources, 45 have been classi ed correctly. 3 sources were incorrectly classi ed into class 0 while 2 sources each were incorrectly classi ed into classes 9 and 11. Class 8 has 13 sources out of which only one has been wrongly classi ed into class 0.

C lass 9 has 40 sources in all. 26 have been correctly classi ed, 4 each have been m isclassi ed into classes 7 and 14 and 3 each into classes 5 and 16. C lass 10 has 44 sources in alland 38 have been correctly classi ed but 3 have been w rongly classi ed to class 13, 2 into class 6 and 1 into class 1. C lass 11 has 12 sources out of which 8 were correctly classi ed. 3 sources were incorrectly classi ed into class 14 and 1 into class 0.

A sm entioned in Section 2, in the original LRS classi cation, class 13 spectra were often confused with SiC feature sources. In the ANN classi cation also, out of the 200 class 13 sources, 104 were wrongly classi ed (85 into class 6) while 96 were correctly classi ed. The wrongly classi ed sources have been listed in Table 3 for the purpose of review by the hum an classi ers.

C lass 14 has 3 sources out of which one was m isclassi ed into class 10. C lass 15 had 140 sources out of which 111 have been correctly classi ed. But 18 have been classi ed into class 1 and 3 each into classes 13 and 4; 2 each into class 2 and 4, and 1 into class 3. Lastly, 25 sources out of a total of 30 of class 16 were classi ed correctly. 2 sources were wrongly classi ed into class 7 while 1 each were classi ed into classes 0, 1 and 5.

If we assume classes 6 and 12 to be the same, (class 12 trained as class 6) and redo the classi cation exercise, we obtain the overall classi cation histogram as shown in Fig. 3. A total of 1,675 spectra have now been classi ed correctly which is about 84% of the total test data set. This process also improved the classi cation of the class 6 sources which is shown in Fig. 5.

We split o class 12 from class 6 because it in proved the classi cation of silicate em ission objects. W ith the full range of silicate em ission objects in one class there was more dispersion of values and the ANN system had more chance of misclassi cation. However, it does not matter that sources are misclassi ed from class 6 to class 12 or class 12 to class 6 because the dividing line between these two groups is not sharp an there is a continuous range of silicate feature strengths, and som e objects must fall at the boundary.

Fig. 6 shows the improvement in the class 13 classication, namely earlier 85 sources of this class were wrongly classications of which has now reduced to 50 wrong classications. This process also improved the classication of class 4 as seen in Fig. 4. There were such improvements noticed in general and Fig. 3 is really a combined elect of all these.

5. Conclusions

We have demonstrated in this paper the application of ANN scheme to a large database of 2000 IRAS spectra and have been able to correctly classify more than 80% of the data-set. The misclassi ed spectra were looked in detail and most of them could have been wrongly catalogued or had features which would have been confusing for even a hum an classi er.

W e stress here that the speed and robustness of this scheme can be very useful for classifying the whole of LRS database containing over 50,000 sources.

RG and HPS are grateful to SK and KV for kind hosp itality while on a visit to Calgary. We thank the two anonymous referees for very useful comments.

{ 12 {

A. ANN architecture

In a feedforward neural network there are several inputs, a few hidden layers, and several outputs (Bailer-Jones et. al., 2002). See the Figure 7 for a block view of this architecture. Each node in the input layer holds a value, x_i . In our example application, the input vector, $(x_1;x_2;:::;x_i;::)$, is the spectrum with 93 ux values, and the output vector, $(y_1;y_2;:::;y_1;::)$, has 17 nodes. Each of the input nodes connects to every node in the next layer of nodes, the rst hidden' layer, and each of these connections has a weight, $w_{i;j}$, associated with it. A node in the hidden layer form s a weighted sum of its inputs, and passes this through a nonlinear transfer function, such that the output from the jth hidden node is

$$p_{j} = \tanh \qquad \begin{array}{c} X \\ w_{i;j}x_{i} \\ i \end{array} \qquad (A:1)$$

These values are passed to a second hidden layer which performs a similar processing, the output from that layer being the vector q

$$q_{k} = \tanh \qquad w_{j,k}p_{j} : \qquad (A : 2)$$

The output layer then perform s a simple sum of its inputs, so that the network output, y_1 , is

$$y_1 = \bigvee_{k \in \mathcal{A}} w_{k;1} q_k : \qquad (A:3)$$

The tanh function in the hidden layers provides the nonlinear capability of the network. O ther nonlinear functions are possible; the sign oidal function $(1=(1 \exp [wx]))$ is used here. Both functions map an in nite possible input range onto a nite output range, 1 to +1 in the case of tanh. This in itates the transfer function of neurons.

REFERENCES

Atlas of Low Resolution IRAS Spectra. 1986. IRAS Science Team, prepared by F.M.O. Inon & E.Raim ond (A&AS, 65, 607) (Atlas)

- Bailer-Jones, C A L., Gupta, R. and Singh, H P. in \Autom ated D ata Analysis in Astronom y", Eds., Gupta, R., Singh, H P. and Bailer-Jones, C A L., Narosa Publishing House, N. Delhi (2002) pp. 51
- Gulati, R.K., Gupta, R., Gothoskar, P. & Khobragade, S., 1994a, ApJ, 426, 340
- Gulati, R.K., Gupta, R., Gothoskar, P. & Khobragade, S., 1994b, Vistas in Astronomy, 38(3), 293
- Gulati, R.K., Gupta, R., & Rao, N.K. 1997a, A&A, 322, 933

Gulati, R.K., Gupta, R., & Singh, H.P., 1997b, PASP, 109, 843

- Gupta, R., Gulati, R.K., & Singh, H.P., 2001, in Proc. 11th Cambridge Workshop \Cool Star, Stellar Systems, and the Sun { Challenges for the New Millennium ".Eds.R.J. Garcia-Lopez, R.Rebelo, & M.R.Zapatero Osorio, ASP Conf. Ser., 223, CDROM p.791
- Kwok, S., Volk, K., & Bidelman, W. P., 1997, ApJ, 112, 557
- Lahav, O., Storrie-Lom bardi, M.C. 1994, Vistas in Astronom y, 38 (Special issue on Neural Networks in Astronom y)
- Singh, H.P., Gulati, R.K., & Gupta, R., 1998, MNRAS, 295, 312
- Volk, K.& Cohen, M. 1989a, AJ, 98, 931
- Volk, K.& Cohen, M. 1989b, AJ, 98, 1563
- Volk, K., Kwok, S., Stencel, R. E., Brugel, E., 1991, ApJS, 77, 607
- Volk, K., Kwok, S., & Langill, P.P., 1992, ApJ, 391, 285
- Volk, K., Kwok, S., & Woodsworth, A.W. 1993, ApJ, 402, 292
- von Helden, G., Tielens, A.G.G.M., van Heijnsbergen, D., Duncan, M.A., Hony, S., Waters, L.B.F.M., Meijer, G. 2000, Science, 288, 313

This preprint was prepared with the AAS $\mathbb{P}T_EX$ m acros v5.0.

Table 1:A	short description	of the 17 ANN	training classes
-----------	-------------------	---------------	------------------

Source type	D escription	ANN Class
Stellar	Stellar photospheric spectra; B -type to early M -type	2
	Lowertemp.stellarcontinuum spectra with no	3
	features: m id to late M -type, little circum stellar dust	
Carbon stars	Strong 11.3 m em ission	1
	Lowertemp.continuum (250 K),	10
	e.g., AFGL 3068 & no strong features	
	Weaker 11.3 m em ission	15
0 xygen-rich AGB stars	Silicate absorption on an intermediate temp.	5
	continuum (400 200 K)	
	Stronger 10 m em ission on a	6
	higher tem p. continuum (600 K)	
	Weaker 10 m em ission on a	12
	higher tem p. continuum (600 K)	
	10 m features in transition from	13
	em ission to absorption	
P lanetary nebulae	Emission lines (PN)	0
(PN) & post-AGB	Silicate em ission on a low tem p. continuum	8
sources (PPN)	(oxygen-rich PN & PPN)	
	Lowertemp. (150 K) featureless	11
	(carbon-rich PN and PPN)	
	21 m em ission feature (carbon-rich PPN)	14
ISM spectra:	UIR or PAH features on a at continuum	4
m ostly H II regions;	(HII regions, galaxies, a few PN & PPN)	
includes a few PN,	Low tem p. continuum (100 K)	7
PPN,& galaxies	without strong features	
	10 m absorption on a low temp.continuum	9
	(100 K), includes a few PPN	
	UIR or PAH features on a low temp. continuum	16
	(100 K), includes a few PN	

							2		-		-) -							
Catalog								ΑN	NC	lass								Total
C lass	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	
0	19	•••	•••	•••	•••	•••	•••	1	1	•••	•••	•••		1	•••		1	23
1		113	•••	•••	3	•••	2	•••	•••	•••	•••	•••		11	1	6	•••	136
2	••••	•••	88	1	•••	•••	•••	•••	•••	•••	•••	•••		•••	•••	••••	•••	89
3		1	5	85	•••	•••	•••	•••	•••	•••	•••	•••	1	1	••••	1	•••	94
4		•••	1	•••	34	22	•••	•••	•••	2	1	•••		10	2	4	11	87
5		•••		•••	4	88	•••	•••		2	•••			1	1	•••	7	103
6		1		•••	•••	•••	679	•••	8	•••	9		33	5		•••		735
7	3	•••	•••	•••	•••	•••	•••	45	•••	2	•••	2		•••		•••	1	53
8	1	•••	•••	•••	•••	•••	•••	•••	12	•••	•••	•••		•••		•••		13
9	•••	•••	•••	•••	•••	3	•••	4	•••	26	•••	•••		•••	4	•••	3	40
10	•••	1	•••	•••	•••	•••	2	•••	•••	•••	38	•••		3		•••		44
11	1	•••	•••	•••	•••	•••	•••	•••	•••	•••	•••	8		•••	3			12
12		6	•••	27	•••	•••	15	•••	•••	•••	•••	•••	149	1				198
13	••••	•••		1	3	4	85	•••	3	•••	8			96				200

•••

•••

•••

•••

•••

2

•••

•••

•••

•••

•••

•••

1

•••

•••

•••

•••

•••

2

2

•••

•••

111

•••

•••

•••

25

•••

3

•••

•••

•••

•••

3

140

30

Table 2:2000 Test sources classi ed by ANN into 17 catalog classes.

•••

•••

1

14

15

16

•••

18

1

•••

2

•••

•••

1

•••

•••

3

•••

•••

•••

1

IRAS	Catalog	ANN	IRAS	Catalog	ANN	IRAS	C atalog	ANN
designation	class	class	designation	class	class	designation	class	class
04395+ 3601	0	13	15380 6545	3	12	14198 6115	4	5
17069 4149	0	7	17282 5102	3	2	14206 6151	4	16
21014 1133	0	8	17504 0234	3	2	15246 5612	4	16
22036+ 5306	0	16	19369+ 2823	3	1	15535 5328	4	5
03293+ 6038	1	15	20056+ 1834	3	13	16041 4912	4	5
04340+ 4623	1	13	22476+ 4047	3	2	16204 4717	4	5
05405+ 3240	1	15	23092+ 5236	3	15	16225 4844	4	14
08011 3627	1	6	23095+ 5925	3	2	16331 4637	4	16
08086 3905	1	13	00450 2533	4	16	16467 4255	4	13
08534 5055	1	13	01056+ 6251	4	5	16573 4619	4	5
08544 4431	1	13	02401 0013	4	10	17076 4702	4	13
10098 5742	1	14	04064+ 5052	4	13	17324 3152	4	16
12195 6830	1	13	05044 0325	4	5	17355 3241	4	2
13045 6404	1	13	05345+ 3157	4	5	17486 2345	4	5
13064 6433	1	13	06114+ 1745	4	16	18077 2614	4	5
15084 5702	1	13	06319 0501	4	5	18092 1742	4	9
15261 5702	1	15	07013 1128	4	13	18162 0246	4	13
15469 5311	1	4	07017 1114	4	5	18162 1612	4	16
16093 4808	1	15	08247 4223	4	5	18254 1149	4	5
16192 4900	1	15	08438 4340	4	9	18310 2834	4	5
16455 4349	1	4	09014 4736	4	13	18312 1209	4	5
17199 3512	1	4	10105 5719	4	16	18320 0352	4	15
17289 3106	1	13	10267 5658	4	13	18357 0604	4	5
18301 0656	1	6	10366 5931	4	13	18463 0052	4	16
18367 0452	1	13	11356 6144	4	13	19097+ 0847	4	16
18551+ 0323	1	15	11418 6706	4	13	19156 0935	4	15
21223+ 5114	1	13	12421 6217	4	15	19193+ 1504	4	5
16063 4906	2	3	13065 6354	4	5	19343+ 2026	4	5
00192 2020	3	2	14092 6506	4	5	21282+ 5050	4	14
21345+ 5410	4	15	05390+1448	6	12	19039+ 0809	6	12

Table 3. List of LRS sources m isclassi ed by ANN

Table 3 | Continued

IR A S designation	Catalog class	ANN class	IR A S designation	Catalog class	ANN class	IR A S designation	Catalog class	ANN class
22308+ 5812	4	16	06434 3628	6	12	19089+1542	6	8
23541+ 7031	4	5	06546 2353	6	12	19231 2717	6	13
09199 5447	5	14	08189+ 0507	6	12	19291+2012	6	10
17317 3331	5	16	10174 5704	6	8	19354+ 5005	6	12
17443 2949	5	16	10360 5633	6	12	19420+ 3318	6	10
18168 1520	5	13	11113 5949	6	12	20004+ 2955	6	8
18257 1000	5	16	11492 6052	6	12	20273+ 3932	6	1
18316 0746	5	16	12188 6246	6	10	20526 5431	6	12
18407 0619	5	4	12230 5943	6	12	21044 1637	6	12
18491 0207	5	16	14297 6010	6	10	21389+ 5405	6	12
19065+ 0832	5	16	16235+1900	6	12	21419+ 5832	6	12
19352+ 2030	5	9	16275 2638	6	13	22000+ 5643	6	12
20110+ 3321	5	9	16316 5026	6	12	22196 4612	6	12
20187+ 4111	5	4	16367 2046	6	12	22480+ 6002	6	8
20318+ 3829	5	4	16434 4545	6	10	23416+ 6130	6	8
20547+ 0247	5	4	17189 6501	6	10	05305+ 3029	7	16
23151+ 5912	5	16	17237 3102	6	12	09032 3953	7	11
00001+ 4826	6	12	17269 2625	6	12	11143 6113	7	0
01010+ 7434	6	12	17450 2724	6	10	14394 6004	7	9
01251+1626	6	12	17515 2407	6	13	16268 4556	7	0
01597+ 5459	6	12	17538 3728	6	12	16342 3814	7	9
02360+ 5922	6	12	18038 1614	6	13	17078 3927	7	0
03489 0131	6	12	18207 1029	6	10	20028+ 3910	7	11
03503+ 6925	6	12	18243+ 0352	6	12	21078+ 5211	8	0
04137+ 3114	6	12	18303 0519	6	13	06053 0622	9	7
04188+ 2819	6	8	18309 6955	6	12	07399 1435	9	7
04292+ 3100	6	12	18363 0523	6	8	10019 5712	9	5
04525+ 3028	6	8	18518+ 0358	6	10	10460 5811	9	7
13416 6243	9	14	07418 2850	12	6	18586 1249	12	3
16313 4840	9	14	07434 3750	12	6	19055+0613	12	3

Table 3 | Continued

IRAS	Catalog	ANN	IRAS	Catalog	ANN	IRAS	Catalog	ANN
designation	class	class	designation	class	class	designation	class	class
18032 2032	9	16	09411 5933	12	1	19111+ 2555	12	1
18100 1915	9	5	09508 4345	12	6	19328+ 3039	12	6
18316 0602	9	16	09564 5837	12	3	19510 5919	12	3
18317 0757	9	7	10383 7741	12	3	19585+ 5200	12	3
18379 0500	9	5	11202 5305	12	13	20111 4708	12	3
18566+ 0408	9	16	13248 7851	12	3	20416+ 1903	12	3
19310+ 1745	9	14	13548 3049	12	1	20438 0415	12	1
19566+ 3423	9	14	14129 5940	12	6	20507+2310	12	6
02152+ 2822	10	1	14234 5359	12	3	23212+ 3927	12	1
06505 0450	10	13	15410 0133	12	6	05073+ 5248	13	6
13522 5619	10	13	15492+ 4837	12	3	06259 1301	13	6
14318 5937	10	6	16052 2339	12	3	06308+ 0402	13	5
16279 4709	10	13	16306+ 7223	12	3	06364+ 0846	13	8
17056 3959	10	6	16328 4656	12	1	06491 0654	13	6
05573+ 3156	11	14	16334 3107	12	3	07180 1314	13	6
09370 4826	11	14	16383 1952	12	6	07376 2827	13	6
13308 6209	11	0	16387 2700	12	3	08357 1013	13	6
19500 1709	11	14	16438 1133	12	6	09354 5627	13	10
00007+ 5524	12	6	16494 1252	12	6	10028 5825	13	8
00245 0652	12	6	16534 3030	12	3	10077 5304	13	6
01150+ 5732	12	3	17201 4613	12	3	10277 5730	13	4
03082+1436	12	3	17297+ 1747	12	3	10287 5733	13	6
05132+ 5331	12	6	17318 2342	12	6	10323 4611	13	6
05450 3142	12	6	17505 7021	12	3	10481 6930	13	6
06210+ 4918	12	3	17578 1700	12	3	11214 6448	13	6
06261+1637	12	3	18347 0241	12	3	11296 4431	13	6
07091 2902	12	3	18378 3731	12	3	11467 6234	13	6
11528 5902	13	6	16320 4419	13	10	18061 1743	13	6
11575 7754	13	6	16335 4707	13	6	18083 2630	13	6
12043 6225	13	6	16350 4754	13	6	18085 1832	13	6

Table 3 | Continued

	IRA , .	S	Catalog	ANN	IRA , .	S	Catalog	ANN	IRAS	Catalog	ANN
_	designa	ation	CLASS	CLASS	aesigna	ation	CLASS	CLASS	designation	CLASS	CLASS
	12222	4652	13	6	16365	4717	13	6	18089 2952	13	3
	12384	4536	13	6	16399	3548	13	10	18222 1544A	13	6
	13243	6159	13	6	16409	5128	13	6	18373 0021	13	6
	13366	6222	13	6	16414	4941	13	6	18373 0922	13	6
	13517	6515	13	6	16490	4618	13	6	18381+ 0020	13	6
	13527	6117	13	10	16538	4135	13	6	18409+ 0431	13	6
	14273	6153	13	6	16567	4659	13	6	18471 0259	13	6
	14591	4438	13	6	16580	4424	13	6	18530+ 0817	13	6
	15008	5808	13	6	16598	4117	13	6	18588 1915	13	6
	15044	5822	13	6	17030	4246	13	10	19007 3826	13	6
	15054	5458	13	10	17109	3243	13	6	19043+ 1009	13	6
	15152	6241	13	6	17132	5003	13	6	19135+ 0931	13	6
	15163	5525	13	8	17186	4208	13	6	19171+ 1119	13	5
	15174	4821	13	6	17239	2812	13	6	19229+ 1708	13	6
	15236	5556	13	6	17362	3322	13	6	19244+ 1809	13	6
	15408	5413	13	4	17368	3000	13	6	19493+ 2905	13	10
	15500	5135	13	6	17382	1704	13	6	19520+ 2759	13	4
	15527	6041	13	6	17401	5730	13	6	20095+ 2726	13	6
	15532	4802	13	6	17479	2927	13	6	20171+ 3519	13	5
	16031	4856	13	6	17507	1122	13	6	20217+ 3330	13	6
	16038	5008	13	6	17559	2848	13	6	20267+ 2105	13	6
	16055	4621	13	5	18006	3213	13	6	20365+ 1154	13	6
	16061	4555	13	10	18022	1432	13	6	20509+ 4212	13	6
	16077	5830	13	6	18027	2314	13	6	21453+ 5959	13	6
	16109	4651	13	6	18046	2322	13	6	22525+ 6033	13	6
	16146	5257	13	6	18050	0518	13	6	16235 4832	14	10
	03112	5730	15	1	10329	3918	15	2	18400 0704	15	13
	05418	3224	15	1	12226+	0102	15	1	18551+ 1345	15	4
	05426+	2040	15	1	12544+	6615	15	2	19314 1629	15	1
	05440+	4311	15	1	13136	4426	15	1	19537+ 2212	15	1

ANN	IRAS	Catalog	ANN

Table 3 | Continued

IRAS	Catalog	ANN	IRA	S	Catalog	ANN	IRAS	Catalog	ANN
designation	class	class	designa	ation	class	class	designation	class	class
06183+ 1135	15	1	14232	6106	15	14	20014+ 2830	15	13
06230 0930	15	1	15193	5656	15	4	15357 5239	16	5
07098 2012	15	1	16296	4417	15	13	20068+ 3328	16	7
07538 3928	15	3	17044	3722	15	1	20286+ 4105	16	14
08304 4313	15	14	17047	2848	15	1	21334+ 5039	16	7
08439 2734	15	1	17048	1601	15	1	22566+ 5830	16	0
09521 7508	15	1	17544	2951	15	1			
10239 5818	15	4	18234	2206	15	1			



Fig. 1. The LRS spectra for the 17 training classes. There are ten training spectra each for the 17 C atalog classes but only one representative spectrum is shown. For each spectrum the IRAS name and the C atalog class (in parenthesis) are given above the plot. P lease note that there exists a crossover region between the "blue" and the "red" spectra (due to the instrum ent's settings) ranging between the 10–14 m and this is seen in these plots.



Fig. 2. ANN learning curve for 10000 iterations (left panel). The RMS error refers to the RMS di erence between the computed output and the desired output for each iteration. The right panel shows the histogram of classic cation accuracy for the 2000 test patterns with 17 training classes.



Fig. 3. The histogram of classic cation accuracy for the 2000 test patterns with 16 training classes (class 12 trained as class 6)



Fig. 4. | Histogram of classi cation accuracy for the 87 test patterns of class 4 with 17 training classes (left panel) and with combined training class 6 and class 12 (right panel)



Fig. 5. Histogram of classic cation accuracy for the 735 test patterns of class 6 with 17 training classes (left panel) and with combined training class 6 and class 12 (right panel)



Fig. 6. Histogram of classic cation accuracy for the 200 test patterns of class 13 with 17 training classes (left panel) and with combined training class 6 and class 12 (right panel)



Fig. 7. Feedforward articial neural network architecture for the problem under investigation