LARGE COSMIC VARIANCE IN THE CLUSTERING PROPERTIES OF LYMAN EMITTERS AT Z ' 5^1

K. SHIMASAKU^{2,3}, T. HAYASHINO⁴, Y. MATSUDA⁴, M. OUCHI², K. OHTA⁵, S. OKAMURA^{2,3}, H. TAMURA⁴, T. YAMADA⁶, AND R. YAMAUCHI⁴ shimasaku@astron.s.u-tokyo.ac.jp

ApJ, accepted

ABSTRACT

We reported in a previous paper the discovery of large-scale structure of Lyman emitters (LAEs) at z =4 86 0 03 with a projected size of $20h_{70}^{-1}$ Mpc $50h_{70}^{-1}$ Mpc in narrow-band data of a 25° 45° area of the Subaru Deep Field ($_0 = 0.3$; $_0 = 0.7$; $H_0 = 70h_{70}$ km s⁻¹ Mpc⁻¹). However, the surveyed area, which corresponds to $55h_{70}^{-1}$ Mpc $100h_{70}^{-1}$ Mpc, was not large enough that we can conclude that we are seeing a typical distribution of z' 5 LAEs. In this Letter, we report the results of follow-up imaging of the same sky area using a new narrowband filter (NB704, c = 7046Å and FWHM= 100Å) to detect LAEs at z = 4.79, i.e., LAEs lying closer to us by $39h_{70}^{-1}$ Mpc on average than the z = 4.86 objects. We detect 51 LAEs at z = 4.79 0.04 down to NB704 = 25.7, and find that their sky distribution is quite different from the z = 4.86 LAEs'. The clustering of z = 4.79 LAEs is very weak on any scales and there is no large-scale high-contrast structure. The shape and the amplitude of the angular correlation function are thus largely different between the two samples. These results demonstrate a large cosmic variance in the clustering properties of LAEs on scales of $50h_{70}^{-1}$ Mpc.

Subject headings: cosmology: observations — cosmology: early universe — cosmology: large-scale structure of universe — galaxies: high-redshift — galaxies: evolution — galaxies: photometry

1. INTRODUCTION

Search for Lyman emission of galaxies using a narrowband filter is a powerful tool to detect high-z faint galaxies. Indeed, many observations have successfully detected such Lyman emitters (LAEs) from z = 2 up to z' = 6.6 (e.g., Hu, Cowie, & McMahon 1998; Pascarelle, Windhorst, & Keel 1998; Campos et al. 1999; Hu, McMahon, & Cowie 1999; Kudritzki et al. 2000; Rhoads et al. 2000; Stiavelli et al. 2001; Ajiki et al. 2002; Hu et al. 2002; Venemans et al. 2002; Fynbo et al. 2003; Kodaira et al. 2003; Ouchi et al. 2003a; Shimasaku et al. 2003; see also Maier et al. 2003). Narrow-band surveys can also map effectively large-scale distributions of LAEs (e.g., Steidel et al. 2000; Venemans et al. 2002; Hu et al. 2004).

We recently made a survey of LAEs at z = 4.86 using the narrow-band filter NB711 ($_c = 7126\text{\AA}$, FWHM=73Å) in an area of 25[°] 45°, and found a large-scale structure of LAEs of $20h_{70}^{-1}$ Mpc $50h_{70}^{-1}$ Mpc size (Shimasaku et al. 2003). This is the first discovery of large-scale structure in young universes, suggesting that the birth of large-scale structure is very early in the history of the universe and that LAEs are strongly biased against dark matter, since Cold Dark Matter (CDM) models predict that the density fluctuations of dark matter are very small on such large scales. However, the size of the large-scale structure is nearly comparable to the size of the survey region $(55h_{70}^{-1} \text{ Mpc})$ $100h_{70}^{-1}$ Mpc), and thus we cannot safely conclude that we are seeing a typical distribution of LAEs at z' 5. To address this issue, it is strongly needed to enlarge the survey volume.

Motivated by this, we made a followup imaging survey of LAEs at z = 4.79, i.e., LAEs located closer to us by $39h_{70}^{-1}$ Mpc than those at z = 4.86, in exactly the same sky area. Using these data, we examine differences in the sky distribution of LAEs between the two redshifts. We adopt $_0 = 0.3$, $_0 = 0.7$, and $H_0 = 70h_{70} \text{ km s}^{-1} \text{ Mpc}^{-1}$.

2. DATA

We carried out deep imaging in the sky area of the Subaru Deep Field (SDF; centered at $(13^{h}24^{m}21;4;+27;29^{\circ}23^{\circ})$) [J2000.0]; Maihara et al. 2001) in a narrow-band filter centered at 7046 Å, NB704, with the prime-focus camera (Suprime-Cam; Miyazaki et al. 2002) on Subaru on 2003 February 2 and 3. The FWHM of the NB704 filter is 100 Å, giving a survey depth for LAEs along the sightline of z = 0.08 or, equivalently, $45h_{70}^{-1}$ Mpc. We observed two field-of-views (FoVs) of Suprime-Cam with a large overlap, the central FoV and the northern FoV, to obtain data for the same region $(25^{\circ} 45^{\circ})$ as imaged in the NB711 in search for z = 4.86 LAEs⁷. For both filters, the variation in the response profile is small across the field of view even through the fast (F/1.86) prime-focus optics, with a full variation in _c of 10Å (NB704) and 12Å (NB711). This ensures a considerably uniform selection of LAEs with respect to redshift over the survey field.

Individual CCD data are reduced and combined using IRAF

¹ Based on data collected at Subaru Telescope, which is operated by the National Astronomical Observatory of Japan.

² Department of Astronomy, School of Science, University of Tokyo, Tokyo 113-0033, Japan; shimasaku@astron.s.u-tokyo.ac.jp, ouchi@astron.s.u-tokyo.ac.jp, okamura@astron.s.u-tokyo.ac.jp.

³ Research Center for the Early Universe, School of Science, University of Tokyo, Tokyo 113-0033, Japan.

⁴ Research Center for Neutrino Science, Graduate School of Science, Tohoku University, Aramaki, Aoba, Sendai 980-8578, Japan; haya@awa.tohoku.ac.jp, matsuda@awa.tohoku.ac.jp, tamura@awa.tohoku.ac.jp, yamauchi@awa.tohoku.ac.jp.

⁵ Department of Astronomy, Kyoto University, Sakyo-ku, Kyoto 606-8502, Japan; ohta@kusastro.kyoto-u.ac.jp.

⁶ National Astronomical Observatory of Japan, Mitaka, Tokyo 181-8588, Japan; yamada@optik.mtk.nao.ac.jp.

⁷ The original 25[°] 45[°] area was determined as follows. We first observed the central FoV in NB711, finding in a northern part a large-scale overdense region of LAE candidates. We then imaged the northern FoV with an overlap of 15° to trace a northern extension of the overdense region.

and our own data reduction software (Yagi et al. 2002). The seeing sizes of the final images are 0.2090. To detect LAEs, we combine broad-band *R* and i^0 data with the NB704 data. These *R* and i^0 data, taken in the spring of 2001, have been used to detect LAEs at z = 4.86 combined with the NB711 data (Ouchi et al. 2003a; Shimasaku et al. 2003). The total exposure time and the limiting magnitude (3 on a 1.208 aperture) are: 90 min and 27.1 mag (*R*), 138 min and 26.9 mag (i^0), and 198 min and 26.7 mag (NB704) for the central FoV; 120 min and 27.5 mag (*R*), 110 min and 27.3 mag (i^0), and 216 min and 26.8 mag (NB704) for the northern FoV. All magnitudes are AB magnitudes.

Object detection and photometry are made using SExtractor version 2.1.6 (Bertin & Arnouts 1996). The NB704-band image is chosen to detect objects. We define the NB704-band limiting magnitude and the selection criteria for z = 4.79 LAEs so that the lower limit to the Lyman flux and the lower limit to the observed equivalent width of Lyman emission (EW_{obs}) be nearly equal to those for the z = 4.86 LAE sample in Shimasaku et al. (2003). We set the limiting magnitude to be NB704=25.7, corresponding to $f(Ly) = 1.1 \quad 10^{-17}$ erg s⁻¹ cm⁻² when the contribution from the continuum emission is negligible. The total number of objects down to NB704=25.7 is 42,440.



FIG. 1. — Distribution of all the detected objects in the Ri - NB704 vs. NB704 plane, where Ri (R + i)=2. The green lines indicate the distribution of 3 errors in brightness for a source with a flat (f = const) spectrum. The blue vertical line shows the faint limit to our catalog, and the blue diagonal line is for the detection limit of Ri (average of the two FoV data). The red line indicates our selection criterion on Ri - NB704 color. The red filled circles are our photometrically selected z = 4.79 LAEs that satisfy all the criteria (see text). The small inserted panel plots the response functions of the three filters (solid: NB704, dotted: R, dashed: i°).

We apply the following three criteria to the detected objects to isolate LAEs at z = 4.79: Ri - NB704 > 0.6, R - i > 0.5, and $i^0 - NB704 > 0$, where Ri $(R + i^0)=2$. Since the NB704 band measures fluxes between the *R* and i^0 bands, we define the offband continuum flux of objects as Ri (R+i)=2. The first criterion, Ri - NB704 > 0.6, is set to select LAEs with EW_{obs} 80 Å. The second and the third criteria are the same as those used to construct the z = 4.86 LAE sample. The second criterion reduces contamination by foreground galaxies whose emission lines other than Lyman happen to enter the NB704 band. Figure 1 plots all objects with NB704 26 in the Ri – NB704 vs. NB704 plane.

The number of objects passing the above criteria is 51. We have found that the contamination to the z = 4.86 LAE sample from low-*z* interlopers is about 20% on the basis of spectroscopic observations and Monte Carlo simulations (Shimasaku et al. 2003). We expect a similar contamination rate for the z = 4.79 LAE sample ⁸. We also expect that the completeness of the z = 4.79 sample is close to that of the z = 4.86 sample.



FIG. 2. — Sky distribution of z = 4.79 LAEs. North is up and east is to the left. Areas of relatively poor quality have been trimmed. A bright star in the upper left corner, around which detection of LAEs is impossible, has been masked. LAE candidates are shown by circles. Candidates common to the z = 4.86 sample are marked by open circles. Objects with -0.3 < NB704 - NB711 < 0.5 are plotted in cyan. The dotted, dashed, and solid lines correspond to contours of = 0;1, and 2, respectively (A top-hat smoothing of 8 Mpc radius is made over the LAE distribution to compute the local overdensity).

3. RESULTS

3.1. Sky Distribution

Figure 2 shows the sky distribution of photometrically selected 51 LAE candidates at z = 4.79, while Figure 3 is a similar plot for 43 z = 4.86 LAE candidates. For each sample, we estimate the local surface density of LAEs, (x_iy) , and compute the surface overdensity, $(x_iy) [(x_iy) -] =$, where

is the mean surface density of LAEs (is defined for each sample). The overdensity contours thus derived are drawn in each figure. Three objects are found to be common to the two samples (open circles in Figs. 2 and 3).

We point out two large differences in the distribution of

⁸ Although the NB704 data are deeper than the NB711 data, we have adopted considerably bright magnitudes for the limiting magnitudes of our samples. In this case, contamination is determined mainly from the depths of R and i^0 data, which are common in the two LAE samples.

LAEs in the two samples. First, the clustering of z = 4.79 LAEs is much weaker than that of z = 4.86 LAEs. In Fig. 2, no clear large-scale inhomogeneity is found, although the overdensity seems to become higher with decreasing Y. In contrast, a large-scale concentration of LAEs and large 'void' regions are seen in Fig.3.



FIG. 3. — Same as Fig.2, but for z = 4.86 LAEs. Candidates common to the z = 4.79 sample are marked by open circles.

Second, no concentration of z = 4.79 LAEs is found in the large-scale structure of z = 4.86 LAEs at Y' 20-30. We identify objects which are likely to be at z = 4.83 from NB704–NB711 color (-0.3 < NB704–NB711 < 0.5; 12 objects in total from the two samples), and plot them in cyan in Figs.2 and 3. We find in Fig.3 that the objects plotted in cyan are less concentrated on the sky than the others. We also note that the sky distribution of the 12 cyan objects seems to be in the middle of the distributions of the z = 4.86 LAEs and the z = 4.79LAEs. These findings indicate that the line-of-sight extension of the z = 4.86 large-scale structure toward lower redshift is not larger than $20h_{70}^{-1}$ Mpc.

3.2. Angular Correlation Functions

We calculate the angular correlation function (ACF), ! (), for the two samples, using the estimator defined by Landy & Szalay (1993), where is the angular separation. The results are plotted in Figure 4. The ACF for z = 4.86 LAEs has large positive values at < 8°, while the amplitude of the ACF for z = 4.79 LAEs is ' 0 at any angular separation.



FIG. 4. — Angular correlation functions (ACFs) of LAEs at z = 4.79 (filled circles) and z = 4.86 (open circles). Errorbars correspond to the 1 errors. The solid line indicates the best fit of A_1 ($^{-0.8} - C$) to the z = 4.79 data. The dotted and dashed lines represent, respectively, the best fit of A_1 ($^{-0.8} - C$) and A_1 ($^{-0.8} - C$) to the z = 4.86 data; we find the best-fit to be 0.2. The crosses indicate the ACF obtained if all LAEs lie in the large-scale structure and their distribution in the large-scale structure is uniform.

The ACF of galaxies, including Lyman-break galaxies (LBGs), is approximated fairly well by a power-law function, $^{-0.8}$ (for LBGs, see, e.g., Foucaud et al. 2003; Ouchi et al. 2003b). We thus fit ! () = $A_!$ ($^{-0.8} - C$) to our LAE data to examine whether this function is also a good fit to the LAE data and to compare $A_!$ between the two samples, where *C* is the integral constraint (*C* = 0:15 arcmin^{-0.8}). We obtain $A_! = 0:10$ 0.21 arcmin^{0.8} and $^{2}=N=1:1$ for z = 4:79 LAEs. For z = 4.86 LAEs, we obtain $A_! = 3:7$ 0.3 arcmin^{0.8}, about 40 times higher than that for z = 4:79 LAEs⁹, and find that the fit to the z = 4.86 data is extremely poor ($^{2}=N=14$). If we fit $A_!$ ($^{-}-C$) to the z = 4.86 data, with being another free parameter, then we find a very flat power-law slope, = 0.2, although the fit with this slope is still unsatisfactory ($^{2}=N=10$). For z = 4:79 LAEs we obtain = 0.9.

The ACF of z = 4.86 LAEs does not increase like -, but remains more or less flat around ! 1-2 at $< 8^{\circ}$ except for the data point at 0.5. Fig.3 shows that most of the z = 4.86 LAEs belong to the large-scale structure which occupies about 30% of the survey area. For such an extreme distribution, it is shown that the ACF has high values, ! 1-2, at $< 8^{\circ}$, even when we artificially re-distribute LAEs uniformly in the large-scale structure (crosses in Fig.4). Thus, the high, relatively constant amplitudes of the observed ACF are largely due to the presence of the large-scale structure and the large void regions.

3.3. *Number Density*

We estimate the spatial number density of LAEs to be (2:7 0:4) $10^{-4}h_{70}^3$ Mpc⁻³ for the z = 4.79 sample and (3:1 0:5) $10^{-4}h_{70}^3$ Mpc⁻³ for the z = 4.86 sample. In these calculations, we adopt a survey volume of $1.9 - 10^5h_{70}^{-3}$ Mpc³ (z = 4.79 sample) and $1.4 - 10^5h_{70}^{-3}$ Mpc³ (z = 4.86). We have not applied completeness correction here. The difference in the number density between the two samples is found to be within the 1 levels, although the z = 4.86 LAEs might be slightly more numerous.

⁹ Since the FWHM of the NB704 filter is wider than that of NB711 by about 30%, the ACF will be lower for z = 4.79 LAEs even when the intrinsic, spatial correlation is the same. However, the difference in A_{\perp} due to this effect is only about 15%.

The rms fluctuation of mass overdensity at z = 4 8 on a sphere with a volume equivalent to the average of our two survey volumes is calculated to be ' 0.08 for $_8 = 0.9$, where $_8$ is the present-day rms fluctuation of mass overdensity on a sphere of $8h_{100}^{-1}$ Mpc radius ($H_0 = 100h_{100}$ km s⁻¹ Mpc⁻¹). This implies that the number density of LAEs found in a volume similar to our average survey volume will fluctuate roughly by $0.08b_{g}$ (if Poisson noise is not included), where b_g is the bias parameter. The observed small fluctuation would prefer relatively small b_g values.

3.4. Possible Effects of Velocity Structures

In the analyses above, we have not considered the possibility that our LAEs form structures along redshift ('velocity structures'). A recent spectroscopic observation of a wide-field LAE sample at z' 5 7 has revealed distinct velocity structures with a depth of z = 0.05 (Hu et al. 2004; see also Fynbo et al. 2003 for a similar finding on QSO-absorber fields at z' 3). LAEs in our samples might have such structures. Multiple velocity structures within the filter bandpass make clustering analysis complicated. If, for instance, our z = 4.79 LAEs are divided into a few velocity structures, the true clustering amplitude will be higher than inferred from the apparent sky distribution (and from the ACF measurement). On the other hand, the largescale structure found at z = 4.86, elongated along right ascension, may be an edge-on view of a sheet-like structure extending along the line-of-sight. Velocity structures can also give an uncertainty to the estimates of number density made in §3.3. Spectroscopic data are needed for a significant fraction of our samples to examine such possibilities.

4. DISCUSSION

Hamana et al. (2004) found that the observed ACF of z = 4.86LAEs given in Ouchi et al. (2003a) cannot be reproduced by a simple halo model which assumes LAEs to be associated with dark haloes, because of too strong the observed correlation on scales > 2° . We find that our z = 4.86 LAEs have

large-scale structure and that their ACF is high and not fit by a power law, while the clustering of z = 4.79 LAEs is very weak. These results may suggest that the clustering of LAEs is typically weak, possibly tracing the dark-matter distribution, and that we happened to observe an unusual region in the z = 4.86universe where LAEs form a large, coherent structure of a size $50h_{70}^{-1}$ Mpc. Conversely, if it turns out, from a larger survey, that the z = 4.86 region we observed is relatively common in high-z universes, this will suggest that LAEs and LBGs are separate populations in terms of clustering properties, since the clustering of LBGs has been found to be approximated well by halo models (Hamana et al. 2004). Detailed modeling of the clustering of LAEs based on an enlarged sample will give important hints on the nature of LAEs.

Although we did not find in our two LAE samples a clear difference in the number density, a large field-to-field variance in the clustering of LAEs, including velocity structures, can influence measurements of the number density of LAEs based on narrow-band surveys, especially if the survey volumes are smaller than ours and if b_g is much larger than unity; the b_g value derived from the clustering of the z = 4.86 LAEs is as 10. Shallower surveys will suffer from larger varilarge as ances, since LAEs with brighter Lyman luminosities tend to be clustered more strongly (Ouchi et al. 2003a). For instance, Ajiki et al. (2003) found that the number density of z = 5.7LAEs in their sample of 2 $10^5 h_{70}^{-3}$ Mpc³ is three times higher than that estimated by Rhoads & Malhotra (2001) based on a similar survey volume.

To summarize, our observations show that it is necessary to survey a much larger volume than ours in order to derive the average clustering properties of LAEs.

We thank the anonymous referee for the useful comments. M. O. acknowledges support from the Japan Society for the Promotion of Science (JSPS) through JSPS Research Fellowships for Young Scientists.

REFERENCES

- Ajiki, M. et al. 2002, ApJ, 576, L25
- Ajiki, M. et al. 2003, AJ, 126, 2091
- Bertin, E., Arnouts, S. 1996, A&AS, 117, 393 Campos, A., Yahil, A., Windhorst, R. A., Richards, E. A., Pascarelle, S., Impey, C., & Petry, C., 1999, ApJ, 511, L1 Foucaud, S. et al. 2003, A&A, 409, 835
- Fynbo, J. P. U., Ledoux, C., Möller, P., Thomsen, B., Burud, I., 2003, A&A, 407, 147
- Hamana, T., Ouchi, M., Shimasaku, K., Kayo, I., Suto Y. 2004, MNRAS, 347, 813
- Hu, E. M., Cowie, L. L., & McMahon, R. G., 1998, ApJ, 502, L99 Hu, E. M., McMahon, R. G., & Cowie, L. L., 1999, ApJ, 522, L9 Hu, E. M. et al., 2002, ApJ, 568, L75 Hu, E. M. et al., 2004, AJ, 127, 563

- Kodaira, K. et al. 2003, PASJ, 55, L17 Kudritzki, R.-P., et al., 2000, ApJ, 536, 19
- Landy, S. D., Szalay, A. S. 1993, ApJ, 412, 64

Maier, C. et al. 2003, A&A, 402, 79 Miyazaki, S. et al. 2002, PASJ, 544, 833 Ouchi, M. et al., 2003a, ApJ, 582, 60

Maihara, T. et al., 2001, PASJ, 53, 25

- Ouchi, M. et al., 2003b, ApJ, submitted (astro-ph/0309657) Pascarelle, S. M., Windhorst, R. A., Keel, W. C., 1998, AJ, 116, 2659
- Rhoads, J. E., Malhotra, S., Dey, A., Stern, D., Spinrad, H., & Jannuzi, B. T., 2000, ApJ, 545, L85
- Rhoads, J. E., Malhotra, S. 2001, ApJ, 563, L5
- Shimasaku, K. et al. 2003, ApJ, 586, L111
- Steidel, C. C., Adelberger, K. L., Shapley, A. E., Pettini, M., Dickinson, M., Giavalisco, M. 2000, ApJ, 532, 170
- Stiavelli, M., Scarlata, C., Panagia, N., Treu, T., Bertin, G., Bertola, F., 2001, ApJ, 561, L37
- Venemans, B. P. et al. 2002, ApJ, 569, L11
- Yagi, M., Kashikawa, N., Sekiguchi, M., Doi, M., Yasuda, N., Shimasaku, K., Okamura, S. 2002, AJ, 123, 66