Sim ulating G alaxy C lusters - III: P roperties of the Intracluster Stars

Jesper Som m er-Larsen^{1;2;3?}, A lessio D. R om eo^{1;2;4}y and Laura Portinari^{1;5}z

¹Theoretical A strophysics C enter, Juliane M aries Vej 30, DK-2100 C openhagen, D enm ark

²N ordita, B legdam svej 17, D K -2100 C openhagen, D enm ark

 3 A stronom ical O bservatory, U niversity of C openhagen, Juliane M aries V ej 30, 2100 C openhagen , D enm ark

⁴D ipartim ento di Fisica e Astronom ia, Universita di Catania, via S.So a 64, 95123 Catania, Italy

⁵ Tuorla Observatory, Vaisalantie 20, FIN -21500 Piikkio, Finland

Accepted | .Received | ; in original form 2004 M arch 11

ABSTRACT

Cosm ological (CDM) TreeSPH simulations of the form ation and evolution of galaxy groups and clusters have been perform ed. The simulations invoke star form ation, chem ical evolution with non-instantaneous recycling, m etallicity dependent radiative cooling, strong star-burst, and (optionally) AGN, driven galactic super-winds, e ects of a meta-galactic UV eld and therm al conduction. Results for two clusters, one Virgolike (T ' 3 keV) and one (sub) C om a-like (T ' 6 keV), are presented. At z=0 the stellar contents of both clusters consist of a central dom inant (cD) galaxy surrounded by cluster galaxies and intracluster stars. The intracluster (IC) stars are found to contribute 20-40% of the total cluster B-band lum inosity at z=0 and to form at a mean redshift z_{i} 3 being on average about 0.5 G yr older than the stars in cluster galaxies. UBVR IJHK surface brightness pro les of the IC star populations are presented; the pro le of the larger cluster matches the observed V-band pro le of the cD in A bell 1413 (T ' 8 keV) quite well. The typical colour of the IC stellar population is B-R=1.4-1.5, com parable to the colour of sub-L E and SO galaxies. The mean Iron abundance of the IC stars is about solar in the central part of the cluster (r 100 kpc) decreasing to about half solar at the virial radius. The IC stars are -element enhanced with a weak trend of [0 /Fe] increasing with r and an overall [0 /Fe] 0.4 dex, indicative of dom inant enrichm ent from type II supernov .

The IC stars are kinem atically signi cantly colder than the cluster galaxies: The velocity dispersions of the IC stars are in the inner parts of the clusters (r 100-500 kpc) only about half of those of the cluster galaxies increasing slightly outward to about 70% at r= 1-2 M pc. The typical projected velocity dispersion in the V irgo-like cluster at R > 50 kpc is 300-600 km /s depending on orientation and projected distance from the cluster center. R otation is found to be dynam ically insigni cant for the IC stars. The velocity distributions of IC stars and clusters galaxies are in one cluster highly radially anisotropic, in the other close to isotropic.

Key words: cosmology: theory | cosmology: numerical simulations | galaxies: clusters | galaxies: formation | galaxies: evolution

1 IN TRODUCTION

Current hierarchical or \bottom -up" structure form ation theories predict that the eld star populations of habes of galaxies like the M ilky W ay should consist partly of stars originally born in sm all proto-galaxies and later tidally stripped from these by the main galaxy or through interaction with other proto-galaxies. The fraction of halo eld stars of such origin depends (apart from obvious cosm ic variance e ects) on the properties of the dark matter (e.g., cold dark matter, CDM, vs.warm dark matter, WDM : for WDM the hierarchy is broken at the free-stream ing mass scale, e.g., Som mer-Larsen & Dolgov 2001) and the nature of the star-form ation process. For exam ple, som e halo stars may be born in rapidly expanding, supernova driven proto-galactic supershells and hence be detached from their parent proto-

[?] E-m ail: jslarsen@ tac.dk

y E-m ail: aro@ ct.astro.it

z E-m ail: porti@ utu.

galaxies right from the outset in space, as well as phase-space (e.g., M ori et al. 1997; Som m er-Larsen, G otz & P ortinari 2003). If dark m atter is cold and star-form ation in expanding supershells is not a com m on phenom enon the above fraction could be quite large, in principle at least, approaching unity (e.g., H elm i et al. 2003). The halo stars resulting from tidal stripping or disruption of a proto-galaxy will stay localized in phase-space for a long period and several such \stream s" of halo stars have been detected in the haloes of the M ilky W ay and M 31 (e.g., H elm i et al. 1999; Ferguson et al. 2002).

From the point of view of structure form ation, galaxy clusters can be seen as scaled up versions of galaxies in hierarchical scenarios. In particular, gravity is expected to strip or disrupt cluster galaxies in a sim ilar way as satellite galaxies in galaxy haloes and a population of intracluster (or \ eld") stars should thus reside between the cluster galaxies. It has been known for decades that cD galaxies often are embedded in extended envelopes presum ed to consist of stars tidally stripped o galaxies in the process of being engulfed by the dD (e.g., 0 em ler 1976; D ressler 1979). In recent years it has been possible to perform quantitative studies of these stellar envelopes through ultra-deep surface photom etry of the general stellar population (e.g., Gonzalez et al. 2000; Gonzalez, Zabludo & Zaritsky 2004; Feldmeier et al. 2002, 2004a), or im aging/spectroscopy of individual planetary nebulae (e.g., A maboldiet al. 2002, 2003; Feldm eier et al. 2004b) or supernovae Type Ia (Gal-Yam et al. 2003). The potential in portance of intracluster stars in relation to the chem ical enrichm ent of the intracluster m edium (ICM) was recently discussed by Zaritsky, Gonzalez & Zabludo (2004) and Lin & Mohr (2004).

On the theoretical front N apolitano et al. (2003) have used an N-body dark matter only fully cosmological sim ulation of the formation of a Virgo-like cluster to make predictions about intracluster stars. They nd that unrelaxed velocity distributions and (bulk) streaming motions of the IC stars should be common due to the large dynamical timescales in clusters. Hence the evolution towards a relaxed coarse-grained phase-space distribution function through phase-mixing has proceeded much less far¹ than, e.g., in the G alactic halo, where dynamical timescales are much shorter.

The dark matter only simulations are complemented by various N-body simulations which invoke a more realistic modelling of the stellar properties of galaxies in order to study the fate of galaxies in a cluster environment (e.g., \galaxy harassment", Moore et al. 1996, \tidal stirring", Mayer et al. 2001 and the formation of the central cluster galaxy, Dubinski 1998). The advantage of this approach is that a very high resolution in the stellar component can be obtained. In relation to the properties of the IC stars, recent progress has been made by Dubinski et al. (2003), and Feldmeier et al. (2004a).

In general the properties of the system of IC stars are set by two main e ects: a) the cool-out of gas and subsequent conversion of cold, high-density gas to stars in individual galaxies and b) the stripping/disruption of the galaxies through interactions with other galaxies and the main clusterpotential. Since such interactions will generally a ect the star-form ation rate (as long as a reservoir of gas is available) the form er process is intim ately coupled to the latter.

Only fairly recently has it been possible to carry out fully cosm ological gas-dynam ical/N-body simulations of the form ation and evolution of galaxy clusters at a su cient levelofnum erical resolution and physical sophistication that the cool-out of gas, star-form ation, chem ical evolution and gas in ows and out ows related to individual cluster galaxies can be modelled to, at least som e, degree of realism (e.g., Valdamini2003; Tomatore et al. 2004). Recently M urante et al. (2004) analyzed 117 clusters form ed in a large cosm ological TreeSPH simulation (Borgani et al. 2004) to study the properties of the IC stars. They determ ined average density pro les of the IC stars as well as stars in galaxies for their large sample of clusters. One of the interesting conclusions reached is that only at fairly large cluster-centric distances $(r^{>} 0.4-0.5r_{vir})$ does the (spherically averaged) density of stars in galaxies (excluding the cD) exceed that of IC stars. Com plem entary work has recently been perform ed by W illm an et al. (2004), who sim ulated a C om a-like cluster at very high num erical resolution. They nd an IC star fraction of 10-20%, in good agreement with observations, and a tendency for the fraction to increase with time. In relation to on-going observationale orts, they moreover nd that lineof-sight velocities of intra-cluster planetary nebulae can be used for fairly accurate cluster m assestim ation, even for dynam ically quite unrelaxed clusters, if more than 5 elds at a range of projected radii are covered.

We have recently undertaken similar simulations of galaxy groups and clusters. W e are building on our TreeSPH code used for simulating galaxy formation (e.g., Sommer-Larsen, Gotz & Portinari 2003), in proved to include modelling of non-instantaneous chem ical evolution (Lia, Portinari & Carraro 2002a), m etallicity-dependent, atom ic radiative cooling, strong supernova, and (optionally) AGN, driven galactic winds and therm al conduction. The main results of the simulations concerning the properties of the ICM and cluster scaling relations are described in Paper I (Rom eo et al. 2004), while the characteristics of the population of the simulated cluster galaxies are presented in Paper II (Romeo, Portinari & Sommer-Larsen 2004) in this Paper III of the series, we discuss our ist results on the properties of the IC stars in two simulated clusters, with masses and temperatures sim ilar to the V irgo and C om a clusters, respectively. Our results com plan ent those of M urante et al. (2004) and W illm an et al. (2004), since we present for the IC stars also results on self-consistently calculated multiband surface brightness pro les, colours, abundances and detailed kinem atics (W illm an et al. 2004 also present a kinem atics study, which we compare to in this paper).

In section 2 we brie y describe the code and the numerical simulations, in section 3 we present and discuss the results obtained, and nally section 4 constitutes our conclusions.

2 THE CODE AND SIMULATIONS

The code used for the simulations is a signi cantly im proved version of the TreeSPH code we have used previously for galaxy form ation simulations (Sommer-Larsen, Gotz &

 $^{^1}$ A visualized example from one of our simulations is given at http://www.tac.dk/~jslarsen/IC stars

Portinari 2003): Full details on the code are given in Paper I, here we recall the main in provem ents over the previous version. (1) In lower resolution regions (which will always be present in cosm ological CDM simulations) an im provem ent in the num erical accuracy of the integration of the basic equations is obtained by solving the entropy equation rather than the therm all energy equation we have adopted the \conservative" entropy equation solving scheme suggested by Springel & Hemquist (2002). (2) Cold highdensity gas is turned into stars in a probabilistic way as described in Sommer-Larsen, Gotz & Portinari (2003). In a star-form ation event an SPH particle is converted fully into a starparticle. Non-instantaneous recycling of gas and heavy elements is described through probabilistic \decay" of star particles back to SPH particles as discussed by Lia, Portinari & Carraro (2002a). In a decay event a star particle is converted fully into an SPH particle. (3) Non-instantaneous chem ical evolution tracing 10 elements (H, He, C, N, O, Mg, Si, S, Ca and Fe) has been incorporated in the code following Lia et al. (2002a,b); the algorithm includes supernov of type II and type Ia, and m ass loss from stars of all masses. Metal di usion in Lia et al. was included with a diusion coe cient derived from models of the expansion of individual supernova rem nants. A much more im portant e ect in the present simulations, however, is the redistribution of m etals (and gas) by m eans of star-burst driven \galactic super{winds" (see point 5). This is handled selfconsistently by the code, so we set = 0 in the present sim ulations. (4) A tom ic radiative cooling depending both on the m etal abundance of the gas and on the m eta{galactic UV

eld, m odelled after H aardt & M adau (1996) is invoked.W e also include a simpli ed treatm ent of radiative transfer, by switching o the UV eld where the gas becomes optically think to Lyman limit photons on scales of 1 kpc. (5) Star-burst driven, galactic super-winds are incorporated in the simulations. This is required to expelm etals from the galaxies and get the abundance of the ICM to the observed level of about 1/3 solar in iron. A burst of star form ation is m odelled in the sam e way as the \early bursts" of Som m er-Larsen, Gotz & Portinari (2003), i.e. by halting cooling in the surrounding gas particles, to m in ic the initial heating and subsequent adiabatic expansion phase of the supershell powered by the star-burst; this scheme ensures e ective energy coupling and feedback between the bursting star particle and the surrounding gas. The strength of the super-winds ism odelled through a free param eter $f_{w ind}$ which determ ines how large a fraction of the new {born stars partake in such bursting, super-wind driving star form ation. We nd that in order to get an iron abundance in the ICM comparable to observations, $f_{w ind} > 0.5$ and at the same time a fairly top-heavy Initial M ass Function (IM F) has to be used. (6) Therm al conduction was im plem ented in the code following Cleary & Monaghan (1999).

W e ran simulations of two sm all and two larger galaxy groups as well as a V irgo-like and a (m ini) C om a-like cluster | we shall denote the latter two clusters C1 and C2 in the follow ing. A ll6 system s were chosen to be fairly relaxed (no > 12 m erging at z < 1). V irial m asses at z= 0 were approximately 3.4×10^{13} , 7.9×10^{13} , 2.0×10^{14} and 8.4×10^{14} h 1 M and X -ray em ission weighted tem peratures 1.1, 1.6, 3.0 and 6.0 keV, respectively. The groups and clusters were selected from a cosm ological, DM -only simulation of a at CDM

Properties of Intracluster Stars 3

model, with $_{M} = 0.3$, $_{b} = 0.036$, h = 0.7 and $_{8} = 0.9$ and a box-length of 150 h 1 M pc. M ass and force resolution was increased in Lagrangian regions enclosing the groups and clusters, such that $m_{gas} = m = 2.5 \times 10^8$ and $m_{DM} = 1.8 \times 10^9$ h¹M for the high resolution gas, star and dark matter particles. Particle numbers are in the range 100000-500000 baryonic+ DM particles. G ravitational (spline) softening lengths of g_{as} = 2.8 and $_{DM}$ = 5.4 h 1 kpc, respectively, were adopted. The gravity softening lengths were xed in physical coordinates from z=6 to z=0 and in comoving coordinates at earlier times. To test for numerical resolution e ects one simulation was run with eight times higher mass resolution and two times higher force resolution, yielding $m_{gas} = m = 3.1 \times 10^7$ and $m_{DM} = 2.3 \times 10^8$ h⁻¹M and gravity softening lengths of 1.4 and 2.7 h⁻¹ kpc, respectively.

For the simulation presented in this paper $f_{wind} = 0.8$, and an Arim oto-Yoshii MF (of slope x= {1, shallower than the Salpeter slope x= -1.35) with mass limits [0.1{100] M was adopted. AGN driven winds were not invoked. Moreover, the thermal conductivity was set to zero assuming that thermal conduction in the ICM is highly suppressed by magnetic elds (e.g., Ettori & Fabian 2000); notice how ever that, in relation to the cluster galaxy populations and hence also IC stars, no signi cant di erence is found between simulations with zero thermal conductivity and simulations invoking therm al conduction at 1/3 of the Spitzer level (Paper II).

3 RESULTS AND DISCUSSION

In this section we present and discuss results for clusters C1 and C2 run at our \standard" resolution. The resolution test discussed in Section 3.7 indicates that none of the following results concerning m ass and lum inosity fractions, m ean ages, surface brightness pro les, colours, abundances, kinem atics and density pro les of the IC stars change in any signi cant way with increased num erical resolution.

All results presented in the following refer to z=0, unless another epoch is explicitly specied. At z=0 1.2×10^4 and 2.3×10^4 star particles are located inside of $r_{\rm vir}=1.3$ and 2.0 h 1 M pc of the clusters C1 and C2². The corresponding mass in stars is 2.9×10^{12} and 5.6×10^{12} h 1 M or a fraction of about 1% of the total virial mass (the stellar mass fraction is smaller for C2 than for C1 – we shall return to this below). In both clusters the stellar mass constitutes 14% of the total baryonic mass inside of the virial radius. In the following we shall refer to star particles simply as \stars".

W hen preparing the initial conditions for C2, due to computational limitations we sampled with split DM + SPH particles a region som ewhat sm aller than the virial volum e". The volum e outside this region, but still inside of the virial volum e, is represented by high resolution, DM -only particles (with a mass scaled by $1=(1 f_b)$ with respect to the DM particles in the inner DM + SPH region, since they are not

 $^{^2}$ In this paper the virial radius $r_{\rm vir}$ is the radius corresponding to, at z=0, an over-density 337 times the mean density of the U niverse, as appropriate for a top{hat collapse in the adopted CDM cosm ology (e.g., B ryan & N orm an 1998). The virial mass M $_{\rm vir}$ of the cluster is the mass enclosed within the region of radius $r_{\rm vir}$.

split into baryonic and DM component; f_b is the universal baryon fraction, assumed to be 0.12 in this work); nally, the region outside the virial volum e is, as usual, sam pled by lower resolution dark matter particles with masses increasing with distance from the virial volume. As a consequence of this baryonic under-sam pling, about half of the mass inside of the virial radius of the nal cluster is in the form of (high { resolution) DM -only particles, i.e. not split into SPH + DM . The relative percentage of these DM -only particles increases with radius, hence the amount of late infall of gas and galaxies is reduced compared to the fully resolved case. This explains why the stellar mass fraction in C2 is sm aller than that in C1 as mentioned above (the C2 star m ass is only about twice the C1 starm ass, while the respective virialm asses di er by a factor of about four). The ratio of the respective number of galaxies N gal (C2)/N gal (C1) is also sm aller than the ratio of the corresponding virialm asses (again, a factor of 2 vs. a factor of 4, see below) and the average ages of the IC stars and galaxies in C2 are slightly larger than that in C1, due to the under-sam pling of late infall of galaxies at the outskirts (Section 3.3). Since the results we discuss refer to the inner 1{2 M pc, the baryonic undersam pling of the outskirts is not a crucial problem . M oreover, we have initiated a simulation of C2 with full baryonic sam pling of the virial volume; at a redshift of z = 0:8 this shows no signi cant di erence in relation to the results on IC stars presented here.

The stellar contents of both clusters are characterized by a massive, central dom inant (cD) galaxy surrounded by galaxies and intracluster stars. The e ective radii, calculated from the B-band surface brightness pro les of the two cD galaxies, are R_{eff} ' 5 kpc, taking the inner 80 kpc of the cluster as the extent of the cD s see below . A fter correction for an excess of central, young stars (section 3.2) the absolute m agnitudes of the cD s are M $_{\rm B}$ -23.Extrapolating the observational \K orm endy relation" (K orm endy 1977) such bright cD s have R_{eff} 20h⁻¹ kpc.Part of this discrepancy is due to an excess of central, young stars, as we shall discuss in section 3.2. W e also note that in a study of the form ation of the brightest cluster galaxy (BCG) based on a pure N-body simulation (but of higher resolution than ours) Dubinski (1998) found an e ective radius of 20 kpc for the central BCG.

3.1 Identifying C luster G alaxies and Intracluster Stars

The rst step in identifying the IC stars is to nd all galaxies in the clusters. To this end we proceed in the following simple way: by visual inspection of the z=0 fram es the stars in all galaxies except the cD are located within 10–15 kpc from the centers of the galaxies. Guided by this typical size of the individual cluster galaxies, a cubic grid of cube-length l=10 kpc is overlaid the cluster, and all cubes containing at least N th=2 stars are identi ed. Subsequently, each selected cube is embedded in a larger cube of cube-length 3 l. If this larger cube contains at least N m in=7 star particles, which are gravitationally bound by its content of gas, stars and dark matter the system is identi ed as a potential galaxy. Since the method can return several, alm ost identical versions of the same galaxy only the one containing the largest num ber of star particles is kept and classi ed as a galaxy. W e tested the galaxy identi cation algorithm by varying 1, N $_{\rm th}$, N $_{\rm m}$ in, and also the num erical resolution of the simulations (section 3.7) and found it to be adequately robust for the purposes of this paper.

The algorithm identi es N $_{gal}$ = 42 and 94 galaxies inside of the virial radii of C1 and C2, respectively, plus a cD at the center of each cluster. The ratio N $_{gal}$ (C2)/N $_{gal}$ (C1) is sm aller than the ratio of the corresponding virialm asses due to the baryonic under-sam pling in cluster C2, as discussed above. A part from the cD there are no galaxies inside of r_{cD} = 80 kpc in either of the z=0 fram es (this is not true in all z 0 fram es for clusters C1 and C2, though, but we tested that the results we present in the following did not depend on which z 0 fram e was chosen to represent the present epoch | see also below).

We de ne the system of IC stars as the stars located at cluster-centric distances r_{cD} r r_{vir} and not inside of the tidal radius of any galaxy in the cluster. The tidal radius for each galaxy is taken to be the Jacobi lim it

$$r_{\rm J} = \frac{m}{3M} D;$$
 (1)

where m is the mass of stars, gas and dark matter in the galaxy (inside of r_J), D is the distance from the cluster center to the galaxy and M is the total m ass of the cluster inside of D (e.g., Binney & Trem aine 1987). The cD itself is e ectively just the inward continuation of the system of IC stars and the division between IC stars and cD is som ewhat arbitrary (we hence below quote intracluster star fractions for $r_{cD} = 80$ as well as 40 kpc). The above de nition of IC stars is conservative, since we calculate the tidal radii on the basis of the z=0 fram e cluster-centric distances of the galaxies. For any galaxy which has been through at least one peri-center passage, this tidal radius should be taken as a mupper limit. Moreover, som e IC stars will be inside of the tidal radius of one of the cluster galaxies just by chance. As the total \tidal volum e" of all the cluster galaxies is found to be a few tenth of a percent of the virial volume of the cluster this e ect can be neglected, how ever.

3.2 The Intracluster Star Fraction

U sing the above de nition we nd that the intracluster stars constitute 21 and 28% of the stellarm ass inside of the virial radius for clusters C1 and C2, respectively (adopting $r_{cD} = 40$ kpc these numbers increase slightly to 28 and 34%). For com parison with observations it is more relevant to look at stellar lum inosities. Since ages and m etallicities are available for all star particles, the photom etric properties are straightforward to calculate treating each star particle as a Single Stellar Population (SSP; see Paper II for details). SSP lum inosities are computed by mass-weighted integration of the Padova isochrones (Girardi et al. 2002), according to the Arim oto-Yoshii IMF.We nd that the IC stars contribute only 9 and 11% of the cluster B-band lum inosities for C1 and C2 respectively. This is signi cantly less than the observationalestim ate of $L_{B;IC} = L_{B;tot} > 0.2$ (e.g., A maboldi 2004). The likely reason for this discrepancy is an excess of stars form ed fairly recently at the center of the cD and dom inating the B-band lum inosity: A fter a period of major merging at z 1-2 strong, quasi-stationary cooling ows develop at the centers of the clusters despite the strong, super-nova



F igure 1.M ean redshift of form ation of the cD + IC stars (solid lines) and stars in galaxies (dashed lines) at z= 0.R esults for cluster C1 (\V irgo-like") are shown by thin lines and for C2 (\m ini-C om a") by heavy lines. Statistical uncertainties are z $_{\rm f}$ < 0.05.

driven energy feedback to the IGM /ICM through galactic super-winds and the use of a fairly top-heavy IMF.As a consequence, stars continue to form steadily at the centers of the cD s (r < 10 kpc) at rates of 350 and 600 M /yr at z=0 for C1 and C2, respectively. At z=0 the cD s are too blue compared to observed dDs with central B-R colours of 1.0 (see Fig. 4) rather than 1.4-1.5 (note though that some star-formation is observed in many dDs at the base of the cooling ow e.g., McNamara 2004). To make a (albeit crude) correction for this excess of young stars at the centers of the simulated cD swe tem porarily discard the hum inosity contribution of all stars in the inner $r_c = 10 \text{ kpc}$ with form ation redshifts $z_{\rm f}$ < $z_{\rm c}$, where $z_{\rm c}$ is determ ined by requiring that the mean form ation redshift of the remaining stars inside of r_c should be sim ilar to the mean form ation redshift of stars at $r^{>} r_{c}$. We nd that this is the case for z_c 1 and shall in the following in various cases present results with and without this crude correction, adopting $z_c = 1$ for the form er.W ith this correction ${\rm L}_{\,{\rm B}\,;{\rm IC}}\,{=}{\rm L}_{\,{\rm B}\,;{\rm tot}}{=}\,0.29$ and 0.42 for the two clusters in better agreement with observations (for $r_{cD} = 40$ kpc we obtain $L_{B;IC} = L_{B;tot} = 0.42$ and 0.50, respectively), though som ew hat high com pared to the values found by Feldmeier et al. (2004a) for non-cD clusters. M oreover, the e ective radii of the cD s increase to 9 and 11 kpc, respectively. These values are in better agreement with observations (Korm endy 1977) and the nding ofD ubinski (1998), but still som ew hat sm aller. Part of this remaining discrepancy may be due to the e ect of gravitational \pinching" of our cD s by the excess, young central stars.

R egarding the intracluster star fraction we show in Paper II that clusters C1 and C2 are decient in bright galaxies apart from the cD s.A s discussed in Paper II, this is probably related to the fact that C1 and C2 are fairly relaxed system s,

Properties of Intracluster Stars 5

for which the last m a jorm erger took place at $z^> 1.H$ ence it is possible that the galaxies in these clusters su ered m ore tidal stripping and m erging than in an average cluster. To be conservative the above IC star fractions should then be considered upper limits to what would be found for a statistically representative sam ple of cluster m erging histories. Finally, we note that the observed IC star fraction of 20% has also been reproduced in the simulations of M urante et al. (2004) and W illm an et al. (2004).

3.3 M ean Form ation R edshifts, Surface B rightness P ro les and C olours of the Intracluster Stars

In Figure 1 we show the mean (spherically averaged) redshift of form ation, z_f , of the cD + IC stars (solid lines) and stars in galaxies except the cD (dashed lines) as a function of radial distance from the center of the cD for clusters C1 and C2, respectively. For both clusters the average form ation redshift of the IC stars is $z_{f;IC} = 3 \text{ at } r^{>} 100-200 \text{ kpc}$. The stars in galaxies (except the cD) are on average som ewhat younger with $z_{\rm f;gal}$ $2.5.\,T\,his$ is to be expected, since the bulk of the IC stars originate in (proto) galaxies, which have been partly or fully disrupted through tidal stripping in the main cluster potential or by galaxy-galaxy interactions. In contrast, the galaxies still remaining at z=0 have potentially been able to continue form ing stars out of rem aining cold gas or gas recycled by evolved stars and subsequently cooled to star-form ing tem peratures and densities. Still, due to ram -pressure stripping of the hot and dilute gas reservoir in galactic haloes and other e ects, the star-form ation rate in the galaxies decreases signi cantly from z=2 to 0, considerably more so than in eld galaxies cf. Paper II. Murante et al. (2004) nd a sim ilar trend that the IC stars are on average older than the stars still in cluster galaxies, but they nd som ew hat low er m ean redshifts of form ation, $z_f = 1.9$ and 1.7 respectively. The reason for this discrepancy is not clear: it m ay be related to di erent galactic super-w ind prescriptions or that we incorporate m etal-dependent radiative cooling in the simulations, whereas Borgani et al. (2004) use a prim ordial cooling function, but is probably not due to num erical resolution di erences (M urante 2004, private com m unication).

From Figure 1 it can be seen that both IC stars and galactic stars in cluster C2 are on average som ewhat older than those in C1. The reason for this is numerical, rather than physical, related to the under-sam pling of baryons in the outskirts of the virial volum e of the cluster discussed at the beginning of this section: the late infall of gas and galaxies onto the cluster is reduced with respect to a simulation with full baryonic sam pling of also the outer parts of the virial volum e.

In Figure 2 we show for cluster C 2 the azim uthally averaged UBVR IJK surface brightness proles of the dD + IC stars, both with and without the correction for central, young cD stars discussed above (the H-band results are om itted for clarity). The orientation is such that the cD is viewed along the m inor axis, but other orientations give very sim ilar results. The light proles are approxim ately described by $r^{1=4}$ laws. In reality the slope attens with increasing r, so the system can be seen as a central dom inant elliptical galaxy surrounded by an extended envelope, with



Figure 2. Multiband (UBVRIJK going bottom up) surface brightness pro les of cD + IC stars for cluster C2, shown with (thin lines) and without (thicker lines) the correction for the excess young, central stars, discussed in the text. A lso shown (thick dotted line) is V-band surface photom etry for the cD in the rich cluster A 1413 (T ' 8 keV) obtained by Feldm eier et al. 2002

the slope of the surface brightness pro le becom ing close to constant beyond r 40 kpc. These are characteristics of observed cD galaxies (e.g., Feldm eier et al. 2002). O bservationally, cD + IC stars can be traced by surface photom etry to V_{lim} '28.3 m ag/arcsec² (e.g., Feldm eier et al. 2002), which for cluster C2 corresponds to r 350 kpc. In Figure 2 we also show V band surface photom etry of the cD + envelope in the rich cluster A 1413 (T ' 8 keV) obtained by Feldm eier et al. (2002). Though the observational V band pro le is slightly atter than what we predict for the V band, overall agreem ent is quite good, especially when it is taken into account that the sam pling of the baryons becom es increasingly incomplete with radius in cluster C2 (see above). In Figure 3 we show the surface brightness pro les of C1 (\V irgo"). A lso shown is two B-band data points derived for V irgo IC stars (indirectly) using planetary nebulae by A maboldi et al. (2002). Our predicted B-band pro le tsthe inner point well, but not the outer one, which is too high (in fact slightly higher than the inner one). The V irgo cluster is know n, how ever, to have an unusually at galaxy surface brightness prole and possibly be dynam ically unrelaxed (Binggeli et al. 1987).

Shown in Figure 4 are the azim uthally averaged B - V, B-R and I-K colours of the cD + IC stars, again with and without the connection discussed above. At about 10 kpc the colours (e.g., B V 0.90-0.95) are typical of the stellar populations in the inner parts of cD s (e.g., M ackie 1992). W ithin 10 kpc this is also the case when correcting for the excess of central, young stars | if not, the core of the cD gets too blue with central colours approaching those of spiralgalaxies. O utside of 10 kpc the colour of the cD + IC stars gets bluer with increasing R, but the gradient is very shal-



F igure 3. Sam e as in Figure 2, but now for cluster C1. D ata points are B-band IC star estim ates based on planetary nebulae observations in the V irgo cluster by A maboldi et al. 2002



F igure 4. A zim uthally averaged B-V (solid lines), B-R (dashed lines) and I-K (dot-dashed lines) colours of cD + IC stars for clusters C2 (thick lines) and C1 (thinner lines), respectively. The results of applying the correction for the excess, central young stars is shown with thin dotted lines.

low: (B-V) per kpc of -0.0003 As the mean age of the stars is approximately constant with R (see Fig. 1), this is mainly due to a decrease in metal abundance with R (see below). At 100 kpc B R ' 1.45, typical of sub-L E and S0 galaxies (e.g., G ladders et al. 1998). We shall discuss this in a forthcoming paper, but note that cD s are found in general



F igure 5. Spherically averaged iron abundance of cD + IC stars (solid lines) and stars in cluster galaxies (dashed lines) for clusters C1 (thin lines) and C2 (thick lines), respectively (results for r < 20 kpc are not shown for clarity).

to have quite at colour pro les, in some cases getting redder with R (e.g., M ackie 1992; G arilli et al. 1997; G onzalez et al. 2000).

3.4 A bundance P roperties of the Intracluster Stars and C luster G alaxies

In Figure 5 we show the spherically averaged iron abundance of the cD + IC stars as well as of the stars in cluster galaxies as a function of cluster-centric distance. Iron is supersolar in the cD + IC stars at $r^{<}$ 100 kpc and decreases to about half solar at large r; the stars in cluster galaxies follow a sim ilar trend. The fairly large overall iron abundance of the IC stars, as compared to, e.g., stars in the halo of the M ilky W ay, re ects that it is the galaxies (past and present) which have to enrich the 5-10 times more massive hot intracluster medium (ICM) to an iron abundance of about 1/3 solar (Paper I). Durrell et al. (2002) carried out HST observations of an IC eld in the Virgo cluster at an average projected distance of 150 kpc from M 87 (which for the purposes here can be assumed coincident with cluster center). They con m an excess of red number counts, which they interpret as IC RGB stars. By comparison with observations of a dwarf irregular, they conclude that these stars have -0.8 < Fe/H > -0.2. Though this is som ewhat less than we predict at r 150 kpc, the discrepancy is only a factor of about two, and the observational abundances are clearly signi cantly larger than those of the halo stars of the M ilky W ay.

Figure 6 shows the corresponding oxygen-to-iron ratios as a function of r. [D /Fe] is super-solar everywhere. This is in agreement with present estimates for the lum inous elliptical galaxies that contain most of the stellar mass in cluster galaxies. For the IC stars no observational information



F igure 6. Spherically averaged oxygen-to-iron abundance ratio of cD + IC stars (solid lines) and stars in cluster galaxies (dashed lines) for clusters C1 (thin lines) and C2 (thick lines), respectively. The lim iting case for pure SN II enrichment [O /Fe]= 0.55 is shown by the thin, dotted line (results for r < 20 kpc are not shown for clarity).

is currently available (A maboldi 2004, private com m unication).

For pure type II supernovae enrichment and with the Arim oto-Yoshii IM F, one expects $[D/Fe]_{SN II}=0.55$ (e.g., Lia, Portinari & Carraro 2002a), so it follows from Fig. 6 that SN e Ia do contribute somewhat to the enrichment of the cD + IC stellar populations, and (not supprisingly) even more so for the stars still in galaxies at z=0 (in fact for an Arim oto-Yoshii IM F the global (t ! 1) value of the SN II+ SN Ia enrichment is [D/Fe]=0.18).

3.5 K inem atics of the Intracluster Stars and C luster G alaxies

Using observed velocities of planetary nebulae it will ultim ately be possible to kinem atically \dissect" the system s of cD + IC stars in nearby galaxy clusters, such as V irgo. It is hence of considerable interest to determ ine for our simulations the velocity distribution of the cD + IC stars and com pare it to that of cluster galaxies. To this end we proceed as follows: At z=0 cluster C1 and the cD at the center are som ew hat attened (ellipticity = 1 b=a ' 0.4) with sim ilar m inor axis orientations (we shall denote the m inor axis of the dD the z-axis" in the following). Cluster C2 is only slightly attened ('0.2), and so is the cD at the center, with approximately perpendicular minor axis orientations. For each cD + IC star and each cluster galaxy we determ ine three perpendicular velocity components: The radial component $v_r = v$ e, where e_r is the unit vector pointing radially away from the center of the cluster, the perpendicular (tane, where e is the unit vector gential) com ponent v = vperpendicular to er and aligned with the x-y plane and the



Figure 7. For cD + IC stars in cluster C1 we show the velocity dispersions r (solid lines), (dotted line) and (dashed line). The statistical uncertainties are marked with sm allerrorbars. The sim ilar quantities are shown for the cluster galaxies by led circles, crosses and led triangles, with larger errorbars marking the statistical uncertainties.



Figure 8. Sam e as Figure 7, but for cluster C2.

third (tangential) component v=v e, where e is the unit vector $e = e_{\rm r} \ e$.

W e calculate the mean rotation v and velocity dispersions $_{\rm r}$, and of cD + IC stars and galaxies in spherical shells. Outside r 10 kpc rotation is found to be dynam - ically insigni cant with v < 20-40 km/s in both clusters. W hen correcting for the excess young, central stars rota-



F igure 9.Velocity an isotropy parameter for cD + IC stars (solid line) and galaxies (solid circles) for cluster C1.The same is shown for cluster C2 by dashed line and solid triangles. An isotropic velocity distribution has = 1 indicated by a thin, dotted line.

tion is found to be also dynam ically unimportant at r < 10kpc. In Figures 7 and 8 we show the velocity dispersions of the dD + IC stars and cluster galaxies in clusters C1 and C 2, respectively, versus cluster-centric distance. The cD + IC stars are kinem atically signi cantly colder than the cluster galaxies: The velocity dispersions of the cD + IC stars are in the inner parts of the clusters (r 100-500 kpc) only about half of those of the cluster galaxies, increasing slightly to about 70% at r 1-2 M pc. A s stars and galaxies are m oving in the same gravitational potential this implies by Jean's theorem (e.g., Binney & Trem aine 1987) that the (num ber) density distribution of cluster galaxies is signi cantly atter than that of cD + IC stars, in particular in the inner parts of the clusters | see below . W illm an et al. (2004) also nd that the IC stars are kinem atically colder than the cluster galaxies, except for the innerm ost part of the cluster. The di erence here is probably due to the di erent de nitions of IC stars used: W illm an et al. (2004) require the IC stars to be unbound from the \cD " also | this results in large IC star velocity dispersions in the inner parts of the cluster.

The velocity distribution of the cD + IC stars in cluster C1 is highly radially anisotropic at r > 100 kpc, and, within the statistical uncertainties, this is also the case for the cluster qalaxies. We quantify this in Figure 9, where we show $= (2 \frac{2}{r} = (2 + 2))$, where 2 is the ratio of the kinetic energy in radial and m ean tangential (1D) motions, respectively. For an isotropic velocity distribution = 1. In cluster C1 there is alm ost tw ice as much kinetic energy in the radial direction than in the sum of the two tangential directions at r > 100 kpc. Interestingly, within the statistical uncertainties the same appears to hold for the galaxies, which at rst seem s surprising, since one would expect that it is predom – inantly galaxies on radial orbits which are tidally disrupted and transform ed into cD + IC stars. The explanation for this



Figure 10. Projected velocity dispersions for cluster C1 of cD + IC stars (thick lines) and dark matter (thinner lines) along the m inor axis (z-axis; dashed lines) and averaged along the x and y-axis (solid lines) of the cD /cluster. Thin dotted lines m ark the e ect of making the correction for the excess young central stars. Statistical uncertainties are <20 km/s.

lack of di erence in the orbital characteristics of the two populations is likely the combined e ect of radial infall of and dynam ical friction on cluster galaxies, but we defer a detailed discussion of this to a forthcom ing paper. W e note that W illm an et al. (2004) nd the velocity distribution of the IC stars to be more radially an isotropic than that of the galaxies in the outer parts of their sim ulated cluster. In cluster C2 the velocity distribution of the dD + IC stars is only mildly radially anisotropic and the velocity distribution of the cluster galaxies is, within the statistical uncertainties, isotropic. W e also note that Sommer-Larsen et al. (1997) found that the velocity distribution of Galactic halo stars becom es tangentially an isotropic in the outer halo the di erences probably relate to di erent accretion kinem atics during the build-up of the various haloes this will also be discussed in a forthcom ing paper.

Finally, concerning the attening of the systems for cluster C1 is about 50% larger than for cD + IC stars as well as galaxies, indicating that these systems are attened by anisotropic velocity distributions | see below.For cluster C2 ', consistent with this cluster being only mildly attened.

O bservationally, for the cD + IC stars one will only be able to determ ine line-of-sight velocities using planetary nebulae, not full 3D velocities. For direct comparison with observations we show in Figure 10 the projected velocity dispersions of the cD + IC stars and (for comparison) of the dark m atter in cluster C 1 versus projected distance from the center of the cD. By dashed lines we show the azim uthally averaged velocity dispersions along them inor axis of the cD. The projected stellar velocity dispersion is 350-400 km /s at the center of the cD, increases to 500 km /s at R 10-

Properties of Intracluster Stars 9

30 kpc and then decreases gradually with increasing R to about 300 km /s at R_{vir}. An increase of the stellar velocity dispersion with R has been observed in som e cD s, such as A 2029 (D ressler 1979), and has been interpreted as marking the transition from galactic to intracluster stars. The e ect is less pronounced after correction for the excess young central stars, but we note that these stars still contribute to the gravitational potential and therefore boost the central velocity dispersions. The projected velocity dispersion of the dark m atter follows a sim ilar trend with R as that of the stars, but is signi cantly larger. A s the stars and dark m atter are m oving in the sam e gravitational potential this im plies that the density distribution of dark m atter is signi cantly atter than that of the cD + IC stars | see below.

The projected velocity dispersions along the major axis of the cD are shown by solid lines. They have been calculated as the average of the projected velocity dispersions along the x-axis and y-axis. A long the x-axis the velocity dispersions are calculated for stars and dark m atter particles projecting onto 30 deg. wedges along the positive and negative y directions, and vice versa for dispersions along the y-axis. The m a praxis velocity dispersions follow similar trends as the m inor axis ones. However, for the cD + IC stars it is signi cantly larger and for the dark m atter halo som ew hat larger than along the minor axis. This shows that both systems are attened by an isotropic velocity dispersions and, as stated above, oriented in sim ilar ways. For cluster C2 the ndings are sim ilar (not shown) though the cD is not aligned with the cluster in this case and both systems are only mildly attened.

Sum marizing our ndings for the V irgo-like cluster, the typical projected velocity dispersion for the cD + IC stars at R > 50 kpc is 300-600 km/s depending on orientation and projected distance from the cluster center. Freem an et al. (2000) nd a (projected) velocity dispersion of 752 110 km/s for 23 IC planetary nebulae in the V irgo-cluster at a projected distance of about 150 kpc from M 87. This value is som ew hat larger than our prediction, but it m ay be due to the \real" V irgo-cluster being kinem atically unrelaxed (see also section 3.3); the observational sam ple is too sm all to test for velocity substructure (Freem an 2004, private com - m unication).

Since clusters are dynam ically fairly young it is important to establish that the results on kinem atics etc. are robust, i.e. they do not depend on the particular fram e chosen for the analysis. To this end we analyzed for each cluster two fram es 1.5 and 3 G yr before the present tim e, corresponding to redshifts 0.12 and 0.26, respectively. We indicate the none of the results presented in this paper change in any signi cant way when going to these earlier fram es.

3.6 Density Distributions of the Intracluster Stars, Cluster G alaxies and the D ark M atter

In Figure 11 we show for cluster C1 the spherically averaged density distributions of the cD + IC stars (with and without the correction for the excess, young central stars), the stars in cluster galaxies, the dark matter and, with an arbitrary normalization, the number density of cluster galaxies. As suggested previously, the number density pro le of cluster galaxies in the inner parts of the cluster as well as the dark matter density pro le in generalare signi cantly atter than





F igure 11. Spherically averaged density pro les for cluster C1 of cD + IC stars (solid line), dark m atter (dot-dashed line), stars in cluster galaxies (dashed line) and, with arbitrary norm alization, num ber density of cluster galaxies (dotted line). The result of applying the correction for the excess, young central stars is shown by the thin, dotted line.

the density pro le of cD + IC stars. For the cluster galaxies this is a manifestation of the transformation of galaxies to cD + IC stars by tidal disruption in the inner parts of the cluster.

M oreover, it follows from F ig.11 that the average (m ass) density of stars in cluster galaxies only exceeds that of the IC stars beyond about half the virial radius - this is in good agreem ent with the ndings of M urante et al. (2004).

3.7 A Numerical Resolution Test

It is in portant to test whether the properties of the IC star population presented in this paper depend much on the num erical resolution of the simulations. In the present paper, we address resolution e ects by discussing a smaller system, a group of virial mass 3:4 10^{13} h ¹M and em ission { weighted tem perature 1.1 keV. The group was run at the sam e \standard" resolution as clusters C1 and C2, as well as with eight times higher mass and two times higher force resolution. This results in particle m asses of $m_{qas} = m = 3.1 \times 10^{7}$ and $m_{DM} = 2.3 \times 10^8$ h⁻¹M and gravity softening lengths of 1.4, 1.4 and 2.7 h⁻¹kpc, respectively. In order to enable an optim alcom parison between the norm aland high resolution runs only Fourier m odes up the N yquist wavenum ber of the norm al resolution simulation were used to prepare the initial conditions for the high resolution run (i.e., additional high-wavenum berm odes up to the Nyquist wavenum ber of the high resolution simulation were not added to the Fourier m odes).

The resulting group looks like a scaled down" version of clusters C1 and C2. In particular, it has a prominent central galaxy, which we denote as the dD, like in the

F igure 12. For a M $_{\rm vir}$ = 3.4x10¹³ h ¹M (T = 1.1 keV) group is shown at z= 0.1 the cumulative mass in cD + IC stars (solid lines) and in stars in cluster galaxies (dashed lines) outside of r= 30 kpc for the norm al resolution (thin lines) and high resolution (thick lines) simulations, respectively.

cluster simulations | in this respect it is similar to the socalled \fossil" groups (e.g., Jones et al. 2003).

In Figure 12 we show the cumulative m ass of cD + IC stars and stars in galaxies outside of r = 30 kpc for the normal and high resolution runs at z = 0.1 (at z = 0 one large group galaxy is merging with the cD). There is reasonable agreement between the runs | the somewhat larger m ass in IC stars in the high resolution simulation at r > 200-300 kpc is likely due to the better resolution of star-forming gas in the lower over-density regions which come to populate the outer parts of the group.

As one goes to higher resolution one m ight naively expect that the fraction of IC stars in the simulations decreases as the cluster galaxies are increasingly well resolved. The interesting indication from F igure 12 is, however, that if anything the fraction of IC stars slightly increases with increasing num erical resolution.

Shown in F igure 13 at z= 0.1 are the azim uthally averaged velocity dispersions of cD + IC stars and dark m atter projected along the m inor axis of the cD for the norm al and high resolution simulations, and with the correction for excess young central stars discussed previously applied. The agreem ent is overall reasonable, especially when it is taken into account that there are at z= 0.1 just 1235 cD + IC stars inside of the virial radius in the norm al resolution simulation of this group. This is a factor 10-20 less than the corresponding num bers for clusters C1 and C2. The increase in central velocity dispersion (R < 10 kpc) for the high resolution sim – ulation, in particular for the stars, is likely an e ect of the increased force resolution. Note also the qualitative similarity between the velocity dispersion pro les shown in Fig. 13, and the results for cluster C1 (Fig. 10).

Finally, as a test of resolution dependence of chem ical





F igure 13. Velocity dispersions projected along the m inor axis of the $\$ D " for the cD + IC stars (low er curves and sym bols) and dark m atter (upper curves and sym bols) in the group. Results for the norm al resolution simulation are connected by dashed lines and are shown as solid circles with errorbars (stars) and solid triangles with errorbars (dark m atter). Results for the high resolution simulation are shown by solid lines.

evolution and m etal abundances we show in Figure 14 the cum ulative m ass of Iron in cD + IC stars and stars in galaxies outside of r= 30 kpc for the norm aland high resolution runs. There is reasonable agreem ent between the runs and we note that in relation to Figure 12 this is a nontrivial result, since a signi cant fraction of the iron produced by the stars ends up in the hot ICM (Paper I).

In general, as mentioned at the beginning of section 3, we carry out all the analysis of the (norm al resolution) cluster simulations C1 and C2 presented in this paper for the norm al and high resolution group simulations also. On this basis we nd that all results given in this paper appear largely robust to resolution changes. Ultimately, how ever, this can only be properly checked by running the \production" simulations C1 and C2 at higher num erical resolution. To this end we have initiated a high resolution simulation of C1 (with similar num erical characteristics as the above high-resolution group simulation) | at z=1.3 we nd no signi - cant di erences between the \standard" and high-resolution simulations of C1 in relation to the results presented in this paper.

4 CONCLUSIONS

In this paper we have discussed the properties of the intracluster (IC) stellar populations. Our results are based on cosm ological simulations of galaxy clusters including self-consistently m etal-dependent atom ic radiative cooling, star-form ation, supernova driven galactic super-winds, noninstantaneous chem ical evolution and the e ects of a m etagalactic, redshift dependent UV eld. In relation to m od-

F igure 14. Sam e as F igure 12, except that cum ulative m asses of Iron are shown.

elling the properties of the IC stars this is an important step forward with respect to previous theoretical works on the subject which were based on purely N-body (collisionless) simulations, apart from the recent, interesting work by M urante et al. (2004).

The main results from our simulations of a Virgo-like (C1) and a (sub) C om a-like (C2) galaxy cluster regarding the IC stellar populations are as follows:

The intracluster (IC) stars are found to contribute 20-40% of the total cluster B-band lum inosity at z=0 and to form at a mean redshift z_f 3, som ewhat larger than the mean form ation redshift of the stars in cluster galaxies which is about 2.5. This di erence corresponds to a time span of about 0.5 Gyr. M urante et al. (2004) nd som ewhat lower mean form ation redshifts of 1.9 and 1.7, respectively | the reason for this discrepancy between the two works remains to be identied.

W e calculate UBVR IJHK surface brightness pro les of the IC starpopulations and nd that the pro le of the larger cluster m atches the observed V-band pro le of A bell 1413 (T ' 8 keV) quite well. For the Virgo-like cluster we fail, however, to match the at pro le between 200 and 400 kpc projected cluster-centric distance observationally (but indirectly) inferred for the Virgo cluster by A maboldi et al. (2002) from planetary nebulae counts. W e note though that the galaxy distribution in Virgo is also unusually at (Binggeli et al. 1987). The V-band surface brightness is found to reach V_{lim} '28.3 mag/arcsec² (Feldmeier et al. 2002) in cluster C2 at r 350 kpc and in cluster C1 at 250 kpc. The typical colour of the IC stellar population r is B-R = 1.4-1.5, com parable to the colour of sub-L E and SO galaxies. The mean Iron abundance of the IC stars is about solar in the central part of the cluster (r 100 kpc) gently decreasing to about half solar at the virial radius. The IC stars are -element enhanced with, e.g, [0/Fe] increasing

slightly with r and characterized by a typical [O/Fe] 0.4 dex.

The IC stars are kinem atically signi cantly colder than the cluster galaxies: The velocity dispersions of the IC stars are in the inner parts of the cluster (r 100-500 kpc) only about half of those of the cluster galaxies increasing slightly to about 70% at r= 1-2 M pc. The typical projected velocity dispersion in the V irgo-like cluster at R > 50 kpc is 300-600 km /s depending on orientation and projected distance from the cluster center. R otation is found to be dynam ically insigni cant for the IC stars. The velocity distributions of IC stars and clusters galaxies are in one cluster highly radially anisotropic, in the other close to isotropic.

A test simulation of a T ' 1.1 keV group at higher numerical resolution indicates that the results presented are largely robust to resolution changes. W ork is in progress to simulate also at higher resolution the two clusters discussed in this paper. M oreover, we are in the process of enlarging signi cantly our sam ple of galaxy clusters.

ACKNOW LEDGEMENTS

We have bene ted considerably from discussions with M agda A maboldi, John Feldm eier, K en Freem an, A nthony G onzalez, G iuseppe M urante and B eth W illm an. M oreover, the comments by the anonym ous referee signi cantly im - proved the presentation of our results.

A llcom putations reported in this paper were perform ed on the IBM SP4 facility provided by D anish C enter for Scienti c C om puting (D C SC). We gratefully acknow ledge the abundant access to com puting time on this system. This work was supported by D anm arks G rundforskningsfond through its support for the establishm ent of the T heoretical A strophysics C enter (TAC), and the V illum K ann R asm ussen Foundation.

REFERENCES

- A maboldi, M., 2004, IAU Symp., 217, Recycling intergalactic and interstellarm atter, eds.P.-A.Duc, J.Braine, and E.Brinks., p.54
- A maboldi, M .et al., 2002, AJ, 123, 760
- A maboldi, M . et al., 2003, A J, 125, 514
- Binggeli, B., Tammann, G.A., & Sandage, A., 1987, AJ, 94,251
- Binney, J., & Tremaine, S. 1987, Galactic Dynamics. Princeton Univ. Press, Princeton
- Borgani, S., 2004, M NRAS, 348, 1078
- Bryan G L., Norm an M L., 1998, ApJ 495, 80
- Cleary, P.W., & Monaghan, J.J., 1999, Journal of Computational Physics, 148, 227
- D ressler, A., 1979, ApJ, 231, 659
- Dubinski, J., 1998, ApJ, 502, 141
- Dubinski, J., Koranyi, D., & Geller, M., 2003, IAU symposium, 208, 237
- Durnell, P.R., Ciardullo, R., Feldmeier, J.J., Jacoby, G.H., & Sigurdsson, S., 2002, ApJ, 570, 119
- Ettori, S., & Fabian, A.C., 2000, MNRAS, 317, 57
- Feldm eier, J.J., et al., 2002, ApJ, 575, 779
- Feldm eier, J.J., et al., 2004a, ApJ, 609, 617

- Feldmeier, J.J., Ciardullo, R., Jacoby, G.H., & Durell, P.R., 2004b, A.p.J. in press (astro-ph/0407274)
- Ferguson, A M N., Irwin, M J., Ibata, R A., Lewis, G F., & Tanvir, N R., 2002, AJ, 124, 1452
- Freeman, K.C., et al., 2000, ASP Conf. Series, 197, 389
- Gal-Yam, A., et al., 2003, AJ, 125, 1087
- Garilli, B., et al., 1997, AJ 113, 1973
- G irardi, L., et al., 2002, A & A , 391, 191
- G ladders, M D ., et al., 1998, ApJ, 501, 571
- Gonzalez, A.H., et al., 2000, ApJ, 536, 561
- Gonzalez, A.H., Zabludo, A.J., & Zaritsky, D.2004, ApJ, submitted (astro-ph/0406244)
- Haardt, F., & Madau, P. 1996, ApJ, 461, 20
- Helm i, A., W hite, SD M., de Zeeuw, P.T., & Zhao, H., 1999, Nature, 402, 53
- Helm i, A., W hite, S.D. M., & Springel, V., 2003, MNRAS, 339, 834
- Jones, L.R., Ponman, T.J., Horton, A., Babul, A., Ebeling, H.& Burke, D.J., 2003, MNRAS, 343, 627
- Kommendy, J., 1977, ApJ, 218, 333
- Lia, C., Portinari, L., & Carraro, G. 2002a, MNRAS, 330, 821
- Lia, C., Portinari, L., & Carraro, G. 2002b, MNRAS, 335, 864
- Lin Y.-T., Mohr J.J., 2004, ApJ in press (astroph/0408557)
- Mackie, G. 1992, ApJ 400, 65
- M cN am ara, B.R. 2004, in \The R iddle of Cooling F lows in G alaxies and C lusters of G alaxies", Charlottesville, V irginia, USA, Eds. T H Ruprecht, JC K em pner & N Soker M ori, M., Yoshii, Y., T su jim oto, T., & N om oto, K. 1997,
- ApJ, 478, L21
- M urante, G ., et al., 2004, ApJ, 607, L83
- Napolitano, N.R., et al., 2003, ApJ, 594, 172
- 0 em ler, A . Jr., 1976, ApJ, 209, 693
- Romeo, A D., Portinari, L., & Sommer-Larsen, J., 2004, MNRAS, submitted (astro-ph/0404445, Paper II)
- Romeo, AD., Sommer-Larsen, J., Portinari, L., & Antonuccio, V., 2004, MNRAS, in preparation (Paper I).
- Som m er-Larsen J., Dolgov A., 2001, ApJ 551, 608
- Som mer-Larsen, J., Beers, T.C., Flynn, C., W ilhelm, R., & Christensen, P.R. 1997, ApJ, 481, 775
- Som m er-Larsen J., Gotz M., Portinari L., 2003, ApJ, 596, 46
- Springel, V., & Hemquist, L., 2002, MNRAS, 333, 649
- Tomatore, L., Borgani, S., M atteucci, F., Recchi, S., & Tozzi, P., 2004, MNRAS, 349, L19
- Valdamini, R., 2003, MNRAS, 339, 1117
- W illman, B., Governato, F., W adsley, J., & Quinn, T., 2004, MNRAS, submitted (astro-ph/0405094)
- Zaritsky, D., Gonzalez, A.H., & Zabludo, A.I., 2004, ApJ, 613, L93