# H igh speed photom etry of faint C ataclysm ic Variables: IV . V 356 A ql, A qr1, F IR ST J1023+ 0038, H 0242{2802, G I M on, A O Oct, V 972 O ph, SD SS 0155+ 00, SD SS 0233+ 00, SD SS 1240-01, SD SS 1556-00, SD SS 2050-05, FH Ser

Patrick A.W oudt<sup>?</sup>, Brian W amery, M agaretha L.P retoriusz Departm ent of A stronom y, University of Cape Town, Private Bag, Rondebosch 7700, South A frica

# ABSTRACT

W e present results from high speed photom etry of a further thirteen faint cataclysm ic variables. V 356 A ql (N ova A ql 1936) shows are-like outbursts with recurrence time 3000 s, but no coherent periodicities. A qrl is an interm ediate polar with a scales spin period of 6.7284 m in and a probable orbital period  $P_{orb} = 3.226$  h derived from orbital sidebands. Its orbital sideband frequencies show very variable am plitudes. The published spectroscopic period of 2.0 h suggests that A qr1 is similar to GW Lib and FS Aur in having an additional periodicity of unknown origin.FIRST J1023+0038 has  $P_{orb} = 4.7548$  h with an orbital modulation of range 0.45 m ag, probably caused by re ection e ect from a hot white dwarf primary; as such it may have been a nova som et in e in the past few decades. H 0242 is a deeply eclipsing very low mass-transfer rate system, probably a dwarfnova of very long outburst interval, with  $P_{orb} = 1.792$  h. G IM on, an old nova, has optical modulations at 4.325 h and possibly also at 48.6 m in and is thus a candidate interm ediate polar. AO Oct, an SU UM a type dwarf nova, shows orbital modulation at quiescence with  $P_{orb} = 94.097 \text{ min} \cdot \text{V} 972 \text{ Oph}$  (Nova Ophiuchi 1957) showed no ickering activity during one set of observations, but did so at a later time, con ming the correctness of the identication of this object, but it shows no orbitalm odulation. SD SS 0155 is a deeply eclipsing polar with an orbital period of 87.13 m in. SD SS 0233 shows aring activity but no discernible periodicity. No periodicity was found in the new AM CVn candidate SDSS 1240.SDSS 1556 shows a periodic modulation at 1.778 h which is possibly due to orbital motion. SD SS 2050 is an eclipsing polar with Porb = 1.5702 h.FH Ser (Nova Serpentis 1970) has strong ickering activity but no detectable orbitalm odulation.

K ey words: techniques:photom etric { binaries:eclipsing { close { novae, cataclysm ic variables

#### 1 IN TRODUCTION

C ontinuing the University of Cape Town survey of faint cataclysm ic variable stars (CVs) that have not previously been studied with high time resolution photom etry, we present results for a further 13 stars. The observational technique is the same as that employed previously: the UCT CCD photom eter (O 'D onoghue 1995) used in frame transfer mode and white light, attached to the 1.9-m (74-in) or 1.0-m (40in) re ectors at the Sutherland site of the South A frican A stronom ical Observatory. In addition a few of our runs were m ade with the SAAO CCD photom eter on the 1.0-m re ector.W ith white light, and with the very non-standard ux distributions of CVs, no transform ations onto recognised photom etric systems are possible { but in any case we are largely interested in the brightness variations that occur on short time scales, and which are measured di erentially with respect to the mean brightness of each variable star. Nevertheless, by observing hot white dwarf standards we are able to provide magnitudes that are roughly on the V scale, and are good to 0.1 mag.

In the previous papers (W oudt & W amer 2001, 2002a, 2003a) we have concentrated on faint nova rem nants. Here we also give results for som e old novae, but we have extended our coverage to include a num ber of faint CV s that have been found in recent large-scale surveys.

<sup>&</sup>lt;sup>?</sup> E-m ail: pw oudt@ circinus.ast.uct.ac.za

y E-m ail:warner@physci.uct.ac.za

z E-m ail: retha@ m ensa.ast.uct.ac.za



F igure 1. The light curves of V 356 A ql. Runs S6258, S6262, S6266, S6279 and S7061 have been displaced vertically down-wards by 0.35, 0.7, 1.2, 1.7 and 2.3 m ag, respectively, for display purposes only.

In Section 2 we give the detailed results of our observations. Table 1 contains the observing log for all of the runs on the various observed stars. Section 3 sum m arises the results.

#### 2 OBSERVATIONS

## 2.1 V 356 A quilae

V 356 A ql was N ova A quilae 1936, N o. 1, discovered by Tam m (1936), reaching maximum brightness at  $m_{pg} = 7.7$ and receiving classi cation as a slow nova. It settled down at a post-emption brightness of V = 18.3 (Szkody 1994). It has no determ ination of orbital period, and has not been studied with time resolved photom etry. The spectrum in the quiescent state (R ingwald, N aylor & M ukai 1996) show s weak H em ission on a continuum that rises rapidly at short wavelengths probably indicative of a high mass transfer accretion disc of m odest inclination. W ith the measured H equivalent width of 15 A and the assumption of an accretion disc background lum inosity, an inclination 50 is deduced (see Figure 2.20 of W amer 1995).

O ur observational runs on V 356 A qlare listed in Table 1; we observed it on 6 nights in Septem ber/O ctober 2001 and then revisited in August 2003 to check whether it showed the same behaviour. The light curves are given in Fig. 1. W e have been unable to interpret these in any useful way. The light curves of V 356 A qlduring runs S6258, S6262 and S6279 (see Fig. 1) show are-like outbursts of 0.3 m ag recurring on time scales 3000 s, but there is no de nite periodicity, and the other nights show di erent behaviours. The FT s quite naturally show excess power at 3000 s and harm onics, but there is no coherent period. There are no periodicities at shorter tim e scales.

There are some similarities of the light curves to those of large amplitude ZZ C et stars (e.g.W armer & N ather 1972),



Figure 2. The light curves of A qr1 (SD SS 2238). Runs S6602, S6604 and S6608 have been displaced vertically downwards by 0.5, 1.0 and 1.5 m ag, respectively.

but the time scale is much longer and the surface tem perature of a white dwarf that erupted only 70 years ago should be much hotter than the ZZ Cet instability strip. Because of accretion of hydrogen, the hotter helium -rich pulsational instability strip should not be relevant to a nova rem nant like V 356 A ql.

T in e-resolved spectroscopy used to investigate the changes that take place during the are-like outbursts could help to interpret this unusual object.

#### 2.2 A qr1 (= SD SS J223843.84+ 010820.7)

Listed in the Downes et al. (2001) catalogue as Aqr1, this star was identi ed as a CV by Berg et al. (1992) in a survey for bright Q SO s, where it was designated 2236+0052. It was rediscovered in the Sloan D igital Sky Survey (Szkody et al. 2003), where it is listed as SD SS 2238. The spectrum has strong Balm er em ission on a blue continuum and show s H eII of m oderate strength.

Szkody et al. (2003) obtained 7 spectra in a 1.7 h nun and deduced  $P_{orb} = 2.0$  h but this must be som ew hat uncertain because of the undersam pling; they also obtained 3.75 h of photom etry, with 10 m in integrations, and described the result as random variability at the 0.02 m ag level.

Our observations of A qrl are listed in Table 1 and the light curves shown in Figs. 2 and 3. The light curves show that A qrl has a brightness modulation with a time scale of several hours and variable range up to 0.2 mag, upon which is superimposed a regular periodicity near 6.7 m in and maximum range 0.10 mag. Fig. 3 shows a blow up of part of the light curve of run S6600 { the 6.7-m in periodicity is clearly visible in the light curve. This modulation is the signature of an intermediate polar (IP). However, there is some am biguity in the interpretation of this star, so we give m ore details of the analysis.

The shorter period appears not to be signi cantly

# Table 1.0 bserving log.

0 b ject	Туре	Run No.	Date of obs.	HJD of rstobs.	Length	t <sub>in</sub>	Tel.	V
			(start of hight)	(+ 2452000.0)	(n)	(S)		(mag)
V 356 A ql	NR	S6251	2001 Sep 24	2177,23744	2.43	20,40	40-in	17.7
		S6254	2001 O ct 09	2192.23163	3.56	20	40-in	17.8
		S6258	2001 O ct 10	2193.22754	2.80	20,40	40-in	17.8
		S6262	2001 O ct 11	2194,22808	3.34	30,40	40-in	17.9
		S6266	2001 O ct 12	2195,22790	2.32	20,40	40-in	17.8
		S6279	2001 O ct 19	2202 23856	2.20	8	74-in	17.8
		S7061	2003 A ug 28	2880,21490	2.40	45	40-in	17 <b>.</b> 8 <sup>b</sup>
A qr1	P	S6600	2002 O ct 31	2579.32931	2.30	20	74-in	18.4
		S6602	2002 N ov 01	2580.25848	4.98	20	74-in	18.3
		S6604	2002 N ov 02	2581,25284	4.98	20	74-in	18.3
		S6608	2002 N ov 04	2583,25411	4.76	20	74-in	18.2
F irst	P?	S6708	2002 D ec 30	2639.47195	3.30	10,15	74-in	17.6
		S6711	2003 Jan 28	2668.39976	5.01	20	40-in	17.5
		S6714	2003 Jan 29	2669.45782	3.90	20	40in	17.5
		S6716	2003 Jan 30	2670.40913	4.91	20	40-in	17.5
		S6719	2003 Jan 31	2671.38651	4.70	20	40-in	17.5
		S6736	2003 Feb 03	2674.56766	1.26	20	40-in	17.6
		S6744	2003 Feb 05	2676.52060	2.76	25	40-in	17.6
		S6785	2003 Feb 21	2692.30455	4.08	25	40-in	17.5
		S6803	2003 Feb 24	2695.45347	3.23	25	40-in	17.6
H 0242{2802	DN	S6591	2002 O ct 29	2577.37719	3.69	45,90	74-in	18.7:
		S6595	2002 O ct 30	2578.46831	0.56	25	74-in	18.7:
		S6597	2002 O ct 30	2578.53360	0.95	45	74-in	18.7:
GIM on	NR	S6142	2000 D ec 21	1900.45337	1,21	5	40in	16.3
		S6155	2000 D ec 25	1904.36811	5.43	10	40-in	16.4
		S7227	2003 D ec 30	3004.40587	4.60	30	40-in	16 <b>.</b> 5ª
		S7228	2003 D ec 31	3005.35415	4.91	30	40-in	16 <b>.</b> 5ª
		S7230	2004 Jan 01	3006.35296	5.98	30	40-in	16 <b>.</b> 4ª
AO Oct	DN	S6102	2000 Jun 05	1701.51362	4.02	60,120	74-in	20.5
		S6121	2000 A ug 22	1779.39636	6.62	45,60,120	74-in	20.1
		S6125	2000 A ug 24	1781.50967	3.90	45,60	74-in	20.2
		S6131	2000 Aug 27	1784.47835	3.85	60	74-in	20.0
V 972 Oph	NR	S6342	2002 Apr 02	2367.45468	1.29	6,15	74-in	15.9
-		S7064	2003 A ug 29	2881.21042	4.66	30	40-in	16 <b>.</b> 1ª
		S7071	2003 A ug 30	2882,21084	2.56	30	40-in	16 <b>.</b> 2ª
SD SS 0155+ 00	Р	S7141	2003 Oct. 04	2917,39123	0.85	45	74-in	18.0
		S7143	2003 O ct 05	2918.36033	1.76	10,45	74-in	18.0
SD SS 0233+ 00	DN?	S6519	2002 Aug 31	2518.52361	1.65	90	74-in	19.9
		S6525	2002 Sep 01	2519.48797	1.65	90	74-in	19.9
		S6527	2002 Sep 01	2519.64325	0.60	120	74-in	19.8
		S6594	2002 O ct 30	2578.30153	3.83	100	74-in	19.8
		S7116	2003 Sep 23	2906.54136	2.10	60	74-in	19.9
		S7119	2003 Sep 24	2907.50635	2.03	100	74-in	19.9
SD SS 1240-01	AM CVn	S7240	2004 Feb 14	3050.50814	1.60	90	40-in	19.6
		S7246	2004 Feb 16	3052.45836	0.83	120	40-in	19.6
		S7254	2004 Feb 19	3055.50047	2.13	90	74-in	19.6
		S7257	2004 Feb 20	3056.50522	1.22	90	74-in	19.6
		S7260	2004 Feb 21	3057.55213	2.20	100	74-in	19.7
SD SS 1556-00	DN	S6812	2003 Feb 26	2697.51362	0.70	60	40-in	17.9
		S6870	2003 M ar 26	2725.49996	3.08	45	74-in	18.0
		S6872	2003 M ar 27	2726.48824	2.09	45	74-in	18.0
		S6881	2003 M ar 29	2728.46586	2.06	45	74-in	18.0
		S6887	2003 M ar 30	2729.43842	2.16	45	74-in	18.0

Notes: NR = N ova R em nant; IP = Interm ediate Polar; P = Polar; DN = D warf Nova;  $t_{in}$  is the integration time;  $\forall$  denotes an uncertain value; mean magnitude out of eclipse; a taken with the SAAO CCD; b taken with a V lter.

Table 1.Continued: observing log.

0 b ject	Т уре	Run No.	D ate of obs.	HJD of rstobs.	Length	t <sub>in</sub>	Tel.	V
			(start of night)	(+ 2452000.0)	(h)	(s)		(m ag)
SD SS 2050-05	Р	S7062	2003 Aug 28	2880.32638	3.31	30	40-in	18.1 <sup>;a</sup>
		S7065	2003 Aug 29	2881.41126	0.88	30	40-in	18.2 <sup>;a</sup>
		S7080	2003 Sep 03	2886.31274	1.95	30	40-in	17.7
		S7091	2003 Sep 16	2899,23195	1.13	60	40-in	17.8
		S7093	2003 Sep 17	2900,26348	0.62	60	40-in	17.8
		S7099	2003 Sep 20	2903,22917	1.60	60	40-in	18.0
		S7105	2003 Sep 21	2904.31274	1.05	60	40-in	18.0
		S7128	2003 Sep 29	2912,23631	0.73	30	74-in	18.0
FH Ser	NR	S7045	2003 Aug 22	2874.23498	3.49	10	40in	17.2:
		S7050	2003 Aug 23	2875,21135	3.85	10	40-in	17.0:
		S7056	2003 Aug 27	2879,23047	4.46	30	40-in	17.0ª;b

Notes: NR = Nova Rem nant; P = Polar;  $t_{in}$  is the integration time; Y denotes an uncertain value; mean magnitude out of eclipse; a taken with the SAAO CCD; b taken with a V lter.



Figure 3.A detailed view of part of the light curve of run S6600 of A qr1 (SD SS 2238).



F igure 4. The Fourier transform of the four long runs on Aqrl. The top panel shows a detailed section of the Fourier transform around the high frequency m odulation.

aliased in the FT of the combined four runs (upper panel of Fig. 4), and is at P = 6.72841(16) m in with a mean am - plitude 0.028 m ag. There are evident sidebands to the main peak, both at lower and higher frequencies. K eeping in m ind that in other IP s the sideband frequencies are displaced from the main peak by multiples of the orbital frequency (e.g. W arner 1986), we looked for such a suite of frequencies in A qrl and met with an unexpected problem : in the total FT there is no set of frequencies that produces a convincing

t. There is a strong peak at low frequency (lower panel of Fig. 4) corresponding to 2.295 h, with approximately oneday aliases at 2.08 h and 2.55 h. These are near to the spectroscopic period, but are primarily due to the two strong ares seen in run S6602 (Fig. 2). If we exclude run S6602 from the FT, the peak at 2.295 h disappears. In the FT with run S6602 excluded, a low frequency peak with a fair num ber of possible aliases is seen around 86 Hz, m atching the orbital sidebands noted previously.

Denoting the rotation frequency of the prim ary as !, these sidebands qualitatively t the scheme (W amer 1986) ! ,!,!+ . There is some indication in the FT of run S6604 (Fig. 5) for a sideband at ! 2 , which suggests the principal peak as the rotational frequency ! (in some IPs the strongest m odulation in the optical is at ! ).

We nd from a non-linear least squares t of sinusoids that ! = 2391.05 Hz, and ! + = 2563.19 Hz. The m ean value of from the orbital sidebands is 86.1 Hz. These peaks are m arked by vertical bars in the upper panel of Fig. 4. A lthough far less probable, the one-day alias of the orbital frequency at 74.5 Hz cannot be excluded from these data.

Furtherm ore, we noticed that adding the fourth night of data to the set three produced confusion (in the sense of extra structure within the sideband envelopes) in the FT. This led us to suspect that the sideband amplitudes vary from night to night, which the FT will interpret as amplitude modulation and generate appropriate extra sidebands, typically with 1/(2 d) spacings, which are what we see within the sideband patterns.

That this is the case is seen in Fig. 5, where we show FTs for the individual nights. A lthough the frequency resolution is low the relative am plitudes of the sidebands vary greatly: in S6600 the high frequency sideband is not resolved but





F igure 5. The Fourier transforms of the four individual runs on A qr1 (SD SS 2238), centred on the high frequency m odulation. The Fourier transform of runs S6600, S6602 and S6604 have been displaced vertically by 60, 40 and 20 mm ag, respectively, for display purposes.

has about one third of the am plitude of the m ain peak and the low frequency sideband is not clearly present; in S6602 both the high and the low frequency sidebands are of low am plitude; in S6604 the high frequency sideband is at about half the am plitude of the m ain peak, and there are two bands on the low frequency side of which the furthest is not seen in the other runs and appears at about twice the spacing of the usual sideband; in S6608 the high frequency sideband is at about 80% of the height of the m ain peak and the low frequency sideband is about 55% of the m ain peak.

W e are now confronted by a contradiction: in standard IP s the sideband frequency splitting agrees exactly with the orbital frequency (), but in A qrl our deduced orbital period from the 86.1 Hz splitting is  $P_{\rm orb} = 3.226$  h, which is quite di erent from the spectroscopic period  $P_{\rm sp} = 2.0$  h m easured by Szkody et al. (2003). This appears to be a further case of a m easured  $P_{\rm sp}$  not agreeing with  $P_{\rm orb}$ : the previous examples are (a) GW Lib, in which the photom etric period  $P_{\rm phot} = 2.08$  h (W oudt & W amer 2002b) bears no obvious relation to the measured  $P_{\rm sp} = 1.28$  h (Thorstensen et al. 2002), (b) FS Aur in which  $P_{\rm sp} = 1.428$  h and  $P_{\rm phot} = 3.4247$  h (Tovm assian et al. 2003), and (c) HS 2331+ 3905 where  $P_{\rm sp}$  is not stable but is always near to 3.5 h and  $P_{\rm phot} = 1.351$  (A rau p-Betancor et al. 2004). In the last case there is no doubt, from the presence of eclipse, that  $P_{\rm phot} = P_{\rm orb}$ .

In A qrl we deduce the presence of a magnetic prim ary rotating with a period of 6.7 m in, which elim inates any interpretation of  $P_{\rm sp}$  as arising in a slow ly rotating prim ary.

# 2.3 FIR ST J102347.8+ 003841 in Sextans

This star was discovered as a radio source with the use of the Very Large A rray and was identied in the optical as a 17th m agnitude star (B ond et al. 2002). It was later independently discovered in the Sloan D igital Sky Survey, where it is given the designation SD SS 1023 (Szkody et al. 2003). CVs are not strong radio sources, and the few detections are of system sm any magnitudes brighter. The spectrum has strong em ission lines of H e II 4686 A and of H e I, as well as the usual B alm er lines, which B ond et al. suggest could be the signature of a magnetic prim ary, which m ight go som e way to explaining the radio are. Spectra and radial velocities obtained by Szkody et al. (2003) over on ly 1.7 h suggest a  $P_{\rm orb}$  3 h and variations that they consider to be m ore typical of an interm ediate polar than a polar.

A CV discovered by such a unique m ethod invites further study. O ur photom etric runs are listed in Table 1, the individual light curves are shown in Fig. 6 and the binned m ean light curve, phased according to the ephem eris derived below, is given in Fig. 7.

F IR ST J1023+ 0038 has a strong repetitive m odulation with a range of 0.45 m ag about a mean of V 17.5 and a period of 4.7548 h. Low amplitude ickering is evident, showing the presence of mass transfer. Fourier transforms of our light curves show only the fundamental and rst harmonic of the principalm odulation { there is no evidence for any other periods in the system. The absence of a subharmonic shows that the observed period is the actual orbital period. The ephemeris for the times of minima is

# $H JD_{m in} = 2452668.428 + 0^{d}.198115(2) E:$ (1)

The prole of the brightness modulation resembles that of a relection elect, rather than an eclipse. The implication is that there is a hot white dwarf in the system, possibly indicative of a nova eruption, overlooked some time in the not too distant past (e.g. a few decades ago). A list of relection elects in known recent novae is given in W oudt & W arner (2003b) in connection with the CV RX J1039.7{ 0507. FIRST J1023+0038 is only 10 degrees away in the same constellation, with G alactic coordinates '= 243 and b = +46 and is therefore far from the direction of the G alactic bulge where m ost searches for novae tend to concentrate.

The optimal condition for a large re ection e ect is reached for short orbital period systems, where the separation of the components is small, and/or in absence of screening of the secondary by an optically thick accretion disc { which can only happen if there is a strongly m agnetic prim ary (Prialnik 1986; Kovetz, Prialnik & Shara 1988). At P<sub>orb</sub> = 4.75 h F IR ST J1023+ 0038 does not have the required sm all Porb and therefore is probably strongly magnetic, i.e. it is likely to be a polar. If so, it could have been desynchronised by the putative nova eruption { detection of m odulated polarisation at a period slightly di erent from that of Porb would con mour suggestion that it is a nova remnant. The pole switching in a desynchronised polar (e.g. M ason et al. 1998) and/or reconnection events in the magnetic eld lines connecting the primary and the secondary could produce just the conditions to explain the unusual radio are in FIRST J1023+ 0038; but in any case, the independent deduction of the presence of a strong magnetic eld is compatible both with the spectrum and the existence of radio em ission.

#### 2.4 H 0242{2802 in Fornax

The star known currently as H  $0242\{2802, which is in the constellation Formax and will be abbreviated here to H <math>0242$ , was discovered in an H emission line survey



F igure 6. The light curves of FIR ST J1023+0038, obtained in 2003 January and February, plotted according to the ephem eris given in Eq. 1. Light curves for runs S6708, S6711 and S6714 are at the correct brightness, light curves for runs S6716, S6719 and S6736 have been displaced vertically downwards by 0.6 m ag, and light curves for runs S6744, S6785 and S6803 have been shifted downwards by 1.2 m ag for display purposes.



F igure 7. The average light curve of FIRST J1023+ 0038 (plotted tw ice), folded on the ephem eris given in Eq.1.

(C lowes et al. 2002) and found to have a spectrum resem – bling W Z Sge, with double peaked emission lines and the underlying white dwarf absorption spectrum, by H owell et al. (2002), who measured its brightness at B = 19.0. It is clearly a CV with a very low rate of mass transfer.

O ur photom etric nuns on H 0242 are listed in Table 1. H 0242 is a deeply eclipsing system with a period of 1.790 h, showing an orbital hump lasting about half the orbital period, interrupted by an eclipse, very much in the classical style of Z C ha and O Y C ar. O ur photom etric period agrees very well with the recently obtained spectroscopic period of 108 5 m in of H 0242 (M ason & H ow ell 2004). We acquired observations of four eclipses on two consecutive nights. The m ean light curve is shown in Fig. 8 and the ephem eris of m id-eclipse is

$$H JD_{m in} = 2452577:4376 + 0^{d}.07460(1)E:$$
 (2)

H 0242 is a candidate for searching for non-radial pulsations in its white dwarf prim ary. The spectrum and the



Figure 8. The average light curve of H 0242 (plotted twice), folded on the ephemeris given in Eq. 2.

eclipse pro le indicate that the prim ary contributes at least half of the light in the system; such objects are com m only found to have ZZ C et prim aries (W arner & W oudt 2003).

#### 2.5 GIM onoceros

N ova M onocerotis was discovered at 8.5 m ag by M .W olf in February 1918 som e weeks after m axim um, which probably was V 5.2 at the beginning of the year. It was a fast nova, later designated as G I M on .M odem spectra show weak H em ission on a very blue continuum (Liu & Hu 2000), typical of a high M- optically thick accretion disc.

P rior to eruption a single photographic detection shows it to have been at m agnitude 15.1, a brightness to which it returned by 1975 (R obinson 1975). How ever, there has been



F igure 9. The light curves of GIM on, phased according to the ephem eris given in Eq. 3. Run S7227 is displayed at the correct brightness; runs S6155, S7228 and S7230 are displaced vertically by -0.1, 0.2 and 0.35 m ag, respectively.

a decline since then: Szkody (1994) m easured V = 16.2 in 1989, which agrees with our current estimate (Table 1). This is a greater range than could be attributed to the slow (60 d) oscillations in brightness with a range of 0.6 m ag, typical of high M- discs, found to be present in GIM on (Honeycutt 2001).

Our observations are listed in Table 1 and consist of a two runs obtained in 2000 (of which only S6155 is long enough to be useful) and three more three years later. The latter were obtained with the SAAO CCD photom eterm ounted on the 1-m re ector, which we had access to at the same time that we were using the UCT CCD photom eter on the 1.9-m telescope. The longer read-out time for the SAAO CCD resulted in lower time resolution than what we were able to achieve with the UCT CCD.

The light curves are shown in Fig. 9, with the 2003/4 group aligned according to the period given below. The mean light curve for the latter is shown in Fig. 10.G I M on shows a brightness variation with a range of 0.10 m ag and a period of 4.325 h, which is indicative of orbital or superhum p modulation in a system of moderate inclination. The ephemeris for minimum brightness of these observations is:

$$H JD_{m in} = 245 3004 4634 + 0^{4} 1802 (3) E:$$
 (3)

The FT for the 2003/4 light curves, after prew hitening with the 4.325 h m odulation and its rst harm onic, shows a peak at 48.6 m in (Fig. 11). W ithin the errors of the least squares t, a peak at its rst harm onic (at 24.2 m in) is identie ed in the FT; both the fundam ental and rst harm onic of this m odulation are marked in Fig. 11. The mean range of the 48.6 m in m odulation, as seen in the FT, is 16 m m ag but individual cycles have m uch large am plitudes.

In the FT of run S6155 (taken three years earlier) excess power is seen around 52 m in; given the relative short data length (7 cycles of the putative 48.6 m in modulation), this peak is not well resolved in this individual run. In Fig. 9 we have marked the consecutive m inim a of the 48.6 m in modulation in run S6155 by vertical bars. There are no shorter period coherent modulations in the light curves.



F igure 10. The average light curve of GIM on (plotted twice), folded on the ephemeris given in Eq. 3.



Figure 11. The Fourier transform of GIM on, prewhitened at the orbital frequency and its rst harmonic. The peak at 48.6 m in and its rst harmonic are marked by vertical bars.

Our conclusion is that G IM on is a candidate interm ediate polar with an orbital period of 4.33 h and a spin-related period of 48.6 m in . Our observations are not su cient to detect possible orbital side bands to the 48.6 m in signal, so we do not know whether it is the white dwarf rotation period or a reprocessed signal.

# 2.6 A 0 0 ctantis

A0 Oct is a dwarf nova of the SU UM a type, showing superoutbursts approximately once per year. At maximum of outburst it reaches V  $15.3^1$ ; at minimum it can be as faint as V = 21 (Vogt & Bateson 1982; Howell et al. 1991). Superhum ps with a period of 96.70 min are observed during superoutburst (Patterson et al. 2003). Early photom etric observations of A0 Oct at quiescence failed to show any orbital modulation (Howell et al. 1991), but those data were very noisy and at the lim it of the instrum ent available (V = 20.9, observed with a 1-m telescope and a V liter). More recent photom etry (Septem ber 2000) has been successful (Patterson et al. 2003).

<sup>1</sup> The G eneralC atalogue of Variable Stars (K holopov et al. 1985) states that m axim um is at V = 13.5, and this has been perpetuated in the literature. However, we believe this to be a transcriptional error as the original discovery paper by von G essner & M einunger (1974) reports a maxim um of 15.3.



8 Patrick A.W oudt, Brian W amer and M agaretha L.P retorius

Figure 12. The light curves of AO Oct, obtained in 2000 June and August.



Figure 13. The mean light curve of AO Oct in August 2000 (plotted twice), folded on the ephemeris given in Eq. 4.

O ur photom etric observations are listed in Table 1.W e detected orbitalm odulation in our rst observation in June 2000 and later extended coverage with three nights in August 2000, when it happened that AO O ct was about 0.5 m ag brighter than in June. The light curves are shown in Fig. 12; an orbitalm odulation with a peak-to-peak range of 0.2 m ag is clearly seen, but there was also a great deal of ickering during run S6121. The FT of the August light curves has a peak at 94.10 m in, which we increased in accuracy by performing a non-linear least squares t to nd a period of 94.097 (0.022) m in. This is in reasonable agreement with the value 94.43 (0.19) m in obtained from photom etry at quiescence by Patterson et al. (2003). The ephemeris for maximum light is given by

$$H JD_{max} = 245 1781 :5261 + 0^{d} :065345 (15) E :$$
 (4)

Fig. 13 shows the m ean light curve obtained by folding our light curves at the orbital period. The superhum p period is 2.8% longer than the orbital period, giving a beat period between the two of 2.4 d, which is typical of SU UM a stars of this orbital period (W amer 1995).

## 2.7 V 972 Ophiuchi

V 972 O ph was N ova O phiuchi 1957, discovered by H aro at m agnitude 9.8 (H aro 1957), but reaching 8.0 at m axim um . It



F igure 14. The light curves of V 972 O ph. The light curve of run S7071 is displaced vertically dow nw ards by 0.25 m ag for display purposes.

is classi ed as a slow nova. The spectrum obtained by R ingwald et al. (1996) shows a red continuum with no distinct em ission lines; they measured its brightness as V = 16.7. However, Zwitter & M unari (1996) measured B = 17.5 and although they also saw a red continuum, they found double-peaked H e II em ission, strong H em ission, and weak H em ission superim posed on wide absorption, typical of a high M- disc. The absence of H e I lines led them to conclude that the white dwarf in V 972 O ph is still extrem ely hot. The continuum slope is due to reddening in this low galactic latitude (b = +2) nova.

The spectra con m that the nova remnant has been correctly identied (Duerbeck 1987) but our rst photom etric observation, made in A pril2002, showed no sensible short time scale variations. Revisiting in A ugust 2003 revealed the expected activity. It is possible that the changes in level of activity are correlated with the strength of the emission lines, and that the spectrum obtained by Ringwald et al. is characteristic of the quieter moments.

Our runs are listed in Table 1 and the light curves are shown in Fig. 14. The 2003 runs show considerable activity, but there is no sign of any orbital modulation on time scales



F igure 15. The light curves of SD SS 0155 (plotted in intensity), aligned according to the ephem eris given in Eq. 5.

of hours and the FTs do not show any persistent periodicities.

## 2.8 SD SS J015543.40+ 002807.2 in Cetus

SD SS 0155 is a CV announced in the rst Sloan D igital Sky Survey release (Szkody et al. 2002), where its strong H e II em ission and its identi cation as a RO SAT source clearly pointed to its being a polar. C om parison of SD SS photom etry and earlier photographic im ages showed that SD SS 0155 has high and low states of lum inosity, over the range 14.7 { 17.6 m ag, com patible with a polar classi cation. R adial velocity m easurem ents of the H eII lines obtained when the system was in a high state gave an orbital period of 88 2 m in.

O ur photom etric observations are listed in Table 1 and were obtained when SD SS 0155 was in a low state. The light curves are given in Fig. 15 and show the star to be a deeply eclipsing system with an eclipse lasting 320 s and a period of 87.13 0.02 m in. Because it is too faint to detect when in eclipse we have plotted the light curve on an intensity instead of a magnitude scale. We have used zero intensities (rather than noise-determ ined upper lim its) for the integrations obtained in eclipse. The ephem eris for the middle of eclipse of the prim ary is

$$H JD_{m in} = 2452918.42776 + 0.06051 (1) E$$
 (5)

The light curve shows that SD SS 0155 is probably a single pole accretor, with the accretion region hidden behind the prim ary for roughly half an orbital period.

#### 2.9 SD SS J023322.61+ 005059.5 in Cetus

SD SS 0233 is another CV in the rst release of the SD SS (Szkody et al. 2002), where its spectrum was found to be dom inated by narrow Balmer em ission and weak He I on a blue continuum. In appearance it possibly resembles a low inclination dwarfnova in quiescence, but Szkody et al. found



High-speed photom etry of faint Cataclysm ic Variables: IV. 9

Figure 16. The light curves of SD SS 0233.

it to be a highly variable X -R ay source, which is uncharacteristic of a CV with such a low energy optical spectrum .

Our observations are listed in Table 1 and the light curves are displayed in Fig. 16. The star shows slow aring activity of considerable amplitude, and faster ickering, but despite our having devoted a moderate amount of photom etric time to it there are no hopeful indications of any persistent periodicity, so we have relegated it.

# 2.10 SD SS J124058.03-015919.2 in V irgo

SD SS 1240 was discovered recently as a candidate AM CVn star following a system atic search for AM CVn systems in the nst data release of the Sloan Digital Sky Survey (Roelofs et al. 2004). SD SS 1240 was previously identi ed as a DB white dwarf following the 2dF quasar redshift survey (Croom et al. 2001) { the 2dF spectrum shows broad absorption lines in the blue part of the spectrum. Roelofs et al. (2004) have obtained spectra with the 6.5-m M agellan telescope and clear double-peaked H e I em ission lines are present, as well as H e II (4686 A) and N III (4634 and 4640 A) em ission lines.

Ourphotom etry is listed in Table 1 and the longer light curves (runs S7240, S7254 and S7260) are shown in Fig. 17. Surprisingly, no variations are seen in the light curves and com bined FT. There is little or no ickering, and no orbital m odulation is detectable. The orbital period will have to be determ ined from spectroscopy.

#### 2.11 SD SS J155644.24-000950.2 in Serpens

SD SS 1556 em erged as a V 18.1 m ag star in the rst release of S ban D igital Sky Survey C V s (Szkody et al. 2002) where the spectrum is seen to have B alm er em ission lines, with no H e II em ission, and the H em ission is centred in the absorption trough of the underlying white dwarf spectrum.



F igure 17.The light curves of SD SS 1240.R un S7240 is displayed at the correct brightness; runs S7254 and S7260 are displaced vertically downwards by 0.3 and 0.6 m ag, respectively.

These are the characteristics of a dwarf nova with low  $M_{-}$ , probably of short orbital period.

Our observations are listed in Table 1 and the light curves are shown in Fig. 18. There is a clear non-sinusoidal modulation with a period of 1.78 h, a peak-to-peak range of 0.4 m ag. and sm all variations in prole from cycle to cycle. The mean light curve, sum med modulo the measured period, is shown in Fig. 19. The FT of the light curves, seen in Fig. 20, has an aliasing am biguity at the fundam ental frequency, but the presence of the rst (and fourth) harm onic gives further estimates of the period, and there is only one alias at the fundam ental that furnishes ts (to within 1

Hz) to the aliases in the harm onic patterns. This gives a period of 1.778 h (106.7 m in). The ephem eris for m axim um light is:

$$H JD_{max} = 2452725 \cdot 5160 + 0^{4} \cdot 07408 (1) E:$$
 (6)

It is probable that this period is P<sub>orb</sub> for SD SS 1556, but the pro le of the m ean light curve (Fig.19) is similar to that of the long outburst interval dwarf nova GW Lib (W oudt & W armer 2002b) which we have discussed in Sect. 2.2, so we advise caution in interpretation of the photom etric period.

The FT prew hitened at the fundam ental and rst and fourth harm onic shows low am plitude but signi cant structure, with peaks near 1558 s, 1137 s, and 834 s (the low er panel of Fig. 20); the low frequency peak o the scale in the prew hitened FT is due to the data length of som e observing runs. A s SD SS 1556 is a low M- system it could be expected to show non-radial oscillations of the ZZ C et type, as detected in a num ber of other low M- CV s (W armer & W oudt 2003). O ur observed periods are som ew hat beyond the observed long end of non-accreting ZZ C et stars, but are theoretically possible (see discussion by M ukadam et al. 2002). W e need to obtain further observations before drawing a conclusion on the nature of these periodicities.

# 2.12 SD SS J205017.84-053626.8 in A quarius

SD SS 2050 is another product of the second release of CV s in the Sloan D igital Sky Survey (Szkody et al. 2003). The spectrum has very prom inent H e II 4686 A, alm ost as strong as neighbouring H , which is a frequent characteristic of m agnetic CV s. Szkody et al. (2003) nd a tentative  $P_{\rm orb}$  of



Figure 18. The light curves of SDSS1556, folded on the ephem eris given in Eq. 6. Run S6870 is displayed at the correct brightness. Runs S6872, S6881 and S6887 are vertically displaced by 0.4, 0.8 and 1.2 m ag, respectively. Run S6812 is displaced upwards by -0.4 m ag and arbitrarily shifted in phase to m atch the tim e of m axim um.



F igure 19. The average light curve of SD SS 1556, folded on the ephem eris given in Eq. 6.

2 h from only 1.8 h of spectral coverage. O f considerable signi cance is their observation of a low state in July 2002, compared with a month later when it was bright enough to obtain the spectra; again this is characteristic of magnetic CVs { in particular, polars.

O ur observations of SD SS 2050 are listed in Table 1 and the longest light curve (S7062) is shown in Fig.21. The other runs were obtained around the times of eclipses in order to improve the period determ ination. The light curve is clearly that of a high inclination polar, with eclipses 1.5 m ag deep of 260 s duration recurring with a period  $P_{\rm orb} = 1.5702$  h. The ephemeris for m id eclipse is given by

 $H JD_{m in} = 245 2880 : 32762 + 0^{d} : 0654246 (5) E :$  (7)



High-speed photom etry of faint Cataclysm ic Variables: IV. 11



Figure 22. The light curves of FH Ser. Runs S7045 and S7056 are displaced vertically by -0.7 m ag and +0.5 m ag, respectively. Run S7050 is displayed at the corrected brightness.

tions of SD SS 1556. The upper panel shows the total FT with the orbital frequency marked (and the rst and fourth harm onic). The lower panel shows the total FT after prew hitening at the orbital frequency and the two harm onics.



F igure 21. The light curve of SD SS 2050, phased according to the ephem eris given in Eq. 7.

#### 2.13 FH Serpentis

FH Ser was Nova Serpentis 1970, discovered by Hirose & Honda (1970) and, during eruption, was one the best-studied novae of the twentieth century, showing a great increase in infrared lum inosity at the time that the optical brightness dim inished through the form ation of dust in the ejecta. It reached V = 4.4 at maximum and has declined to postoutburst brightness of V 17.6. Its modern classi cation is a slow nova of the Fe II class (W illiam s 1992). It has an observed ejecta shell (Gill & O'Brien 2000). Despite its in portance in the history of understanding the evolution of nova light curves, it does not have a known orbital period, and no high speed photom etry for it has been published. The spectrum in 1991 was still dom inated by em ission lines from the ejecta (Ringwald et al. 1996). We observed FH Ser in the hope of detecting an orbital brightness modulation. Our observations are listed in Table 1 and the light curves are displayed in Fig. 22.

Table 2. Stars showing the GW Lib phenom enon.

Star	P <sub>phot</sub> (h)	P <sub>sp</sub> (h)	P <sub>phot</sub> =P sp
GW Lib	2.09	1.280	1.63
A qrl	3.226	2.0	1.6
FS Aur	3.425	1.428	2.40
H S 2331	1.351	3.5	1/2.6

FH Sershows rapid ickering superim posed on slow excursions of 0.5 m ag on time scales 2.5 h. The FTs do not show evidence for any coherent periodicity on time scales up to a few hours. The large amplitude of ickering will mask orbital modulation; to obtain the orbital period by a photom etric method will probably require a multi-site collaborative cam paign.

# 3 SUMMARY AND DISCUSSION

O ur observations are of a m ixed group of CVs { four known old novae (G IM on, V 356 A ql, V 972 O ph and FH Ser), in the latter three we found no orbital m odulation, and a possibly overlooked nova (FIR ST J1023). We have found two new polars with deep eclipses (SD SS 0155 and SD SS 2050) and a new interm ediate polar (A qr1). We have determ ined orbital periods for two dwarf novae in quiescence (A O O ct and H 2042) and observed periodic m odulations in one other dwarf novae (SD SS 1556) that m ight be orbital but could also be a possible further manifestation of the GW Lib phenom enon of a non-orbital m odulation of unknown origin. We nd no periodic photom etric signal in SD SS 0233, nor in the new AM CVn star SD SS 1240.

A m ong our results is one CV, A qrl, for which the spectroscopic period di ers considerably from the photom etric period.W e sum m arise this in Table 2 and draw attention to the clustering of period ratios, which is perhaps indicative of som e resonance e ect in operation.

# 12 Patrick A.W oudt, Brian W amer and M agaretha L.P retorius

#### ACKNOW LEDGMENTS

W e thank Dr.D.O'D onoghue for the use of his EAGLE program for Fourier analysis of the light curves. PAW is supported by funds m ade available from the N ational R esearch Foundation and by strategic funds m ade available to BW from the University of Cape Town.BW 's research is supported by the University.M LP is supported by a bursaries from the D epartm ent of Labour and the N ational R esearch Foundation.

## REFERENCES

- A rau jo-B etancor, S., G ansicke, B.T., H agen, H.-J., M arsh, T., Thorstensen, J., et al., 2004, P roc. IAU Coll. 194, R evM exA A (SC), in press
- Berg C ., W egner G ., Foltz C B ., Cha ee Jr. F H ., Hewett P.C ., 1992, A pJS, 78, 409
- Bond H E ., W hite R L ., Becker R H ., O 'Brien S ., 2002, PASP, 114, 1359
- $C \mbox{ low es } R$  .,  $H \mbox{ uber } M$  .,  $H \mbox{ ow ell } S \mbox{ B}$  .,  $M \mbox{ ason } E$  ., 2003, in preparation
- Croom, SM., Smith, RJ., Boyle, BJ., Shanks, T., Loaring, N.S., et al., 2001, MNRAS, 322, 29
- Downes R A., Webbink R F., Shara M M., Ritter H., Kolb U., Duerbeck H W., 2001, PASP, 113, 764
- Duerbeck H W ., 1987, Sp.SciRev., 45, 1
- GillC D ., O 'B rien T .J., 2000, M N R A S, 314, 175
- Haro G., 1957, HAC No. 1405
- Hirose H., Honda M., 1970, IAUC, 2212
- Honeycutt R K ., 2001, PASP, 113, 473
- HowellSB., DobrzyckaD., Szkody P., KreidlT.J., 1991, PASP, 103, 300
- Howell S.B., Mason E., Huber M., Clowes R., 2002, A&A, 395, L47
- K holopov P N ., 1985, (ed.) G eneral C atalogue of Variable Stars, M oscow
- Kovetz A., Prialnik D., Shara M. M., 1988, ApJ, 325, 828
- LiuW., Hu J.Y., 2000, ApJS, 128, 387
- M ason E , Howell S B , 2004, Proc. IAU Coll. 194, RevM exAA (SC), in press
- Mason PA., Ramsay G., Andronov I., Kolesnikov S., Shakhovskoy N., Pavlenko E., 1998, MNRAS, 295, 511
- M ukadam, A S., K epler, S.O., W inget, D E., Bergeron, P., 2002, A pJ, 580, 429
- 0 'D onoghue D ., 1995, Baltic A str, 4, 517

Patterson J., Thorstensen J., K em p J., et al., 2003, PASP, in press

- Prialnik D., 1986, ApJ, 310, 222
- Ringwald F A ., Naylor T ., M ukai T ., 1996, M N R A S, 281, 192
- Robinson E L., 1975, AJ, 80, 515
- Roelofs, G H A ., G root, P J., Steeghs, D ., Nelemans, G ., 2004, Proc. IAU Coll. 194, RevM exAA (SC), in press
- Szkody P., 1994, AJ, 108, 639
- Szkody P., Anderson S.F., Agueros M., Covarrubias R., Bentz M., et al., 2002, AJ, 123, 430
- Szkody P., Fraser O., Silvestri N., Henden A., Anderson S.F., et al., 2003, A.J., 126, 1499
- $\texttt{Tam}\,\texttt{m}$  N ,1936,AN,260,375
- Thorstensen J.R., Patterson J., K em p. J., Vennes S., 2002, PASP, 114, 1108
- Tovm assian G ., Zharikov, S., M ichel, R ., Neustroev, V ., G reiner, J., et al., 2003, PASP, 115, 725
- Vogt N ., B ateson F M ., 1982, A & A S, 48, 383
- von Gessner H., Meinunger I., 1974, Verand. Sterne 6, 249
- W arner B ., 1986, M NRAS, 219, 347
- W armer B ., 1995, C ataclysm ic V ariable Stars. C U P, C am bridge
- W amer B., Nather R.E., 1972, MNRAS, 156, 1

W amerB., W oudtPA., 2003, Proc.IAU Colloq.No.193, in press
W illiam sR E., 1992, AJ, 104, 725
W oudtPA., W amerB., 2001, M NRAS, 328, 159
W oudtPA., W amerB., 2002a, M NRAS, 335, 44
W oudtPA., W amerB., 2002b, Ap& SS, 282, 433
W oudtPA., W amerB., 2003a, M NRAS, 340, 1011
W oudtPA., W amerB., 2003b, M NRAS, 339, 731
Zw itter T., M unariU., 1996, A&AS, 117, 449