Evidence of P lanetesim al infall on to the very young H erbig B e star LkH 234 $^{\rm 1}$

Abhijit Chakraborty, Jian Ge and

Suvrath M ahadevan

525 Davey Laboratory, Pennsylvania State University, University Park, PA 16802 abhijit@astro.psu.edu, jian@astro.psu.edu, suvrath@astro.psu.edu

ABSTRACT

We report here the stervidence for planetesim al infall onto the very young Herbig Be star LkH 234. These results are based on observations acquired over 31 days using spectroscopy of the sodium D lines, the He I 5876A, and hydrogen H lines. We nd Redshifted Absorption C omponents (RAC) with velocities up to 200 km /s and very mild B lueshifted Absorption C omponents (BEC) up to 100 km /s in the Na I lines. No correlation is observed between the appearance of the Na I RAC & BEC and the H and He I line variability, which suggests that these (Na I RAC & BEC) are form ed in a process unrelated to the circum stellar gas accretion. We interpret the Na I RAC as evidence for an infalling evaporating body, greater than 100 km in diam eter, which is able to survive at distances between 2.0 to 0.1 AU from the star. The dram atic appearance of the sodium RAC and mild BEC is readily explained by the dynam ics of this infalling body making LkH 234 the youngest (age 0.1 M yr) system with evidence for the presence of planetesim als.

Subject headings: stars: form ation | stars: pre{main-sequence | circum stellar matter | stars: individual (LkH 234)

1. Introduction

LkH 234 (m $_{\rm v}$ = 11.9m ag) is a star em bedded in a nebula associated with the NGC 7129 star form ing region and is at a distance of about 1250 parsecs. This is one of the youngest known Herbig Be objects and has an age 0.1 m illion years and spectral type of B5 (Fuente et al. 2001). Herbig Ae/Be stars are intermediate mass (2 to 10M) pre-main-sequence stars (Herbig 1960). A number of Herbig Ae stars have been found to possess circum stellar disks and show both photom etric and spectroscopic variability (H erbst et al. 1999). LkH 234 though much hotter and more massive also exhibits these properties (Fuente et al. 2001, Herbstetal. 1999, Polom skietal. 2002). The star show sphotom etric variability of 12 m agnitude in the V band and is known posses a circum stellar disk with the disk mass to the stellar mass ratio (M $_{\rm D}$ /M) 0.02. A detailed description of LkH 234 and its circum stellar region can be found in Fuente et al. (2001, and references therein).

Spectroscopic variations in m etallic lines (like Na I, Ca II, etc.) in young A type stars like P ictoris (20 to 100 m illion years) are offen interpreted as the signature of star grazing com ets or falling evaporating bodies with sizes oftens of kilom eters (G rinin et al. 1996, Sm ith & Terrile 1984, Thebault & Beust 2001). The only Herbig Be star where grazing com etary transient events have been inferred so far is HD 100546 (V iera et al. 1999). How ever, HD 100546 ism uch older (10M yrs) and cooler (spectral type B9) com pared to LkH 234. H igh resolution spectroscopic monitoring of very young (0.1 to 1 M yrs) Herbig Be stars are scanty

 $^{^1\}mathrm{B}\,\mathrm{ased}$ on observations from the 9.2m Hobby-Eberly Telescope

and uncharted.

Here we report dram atic variations in the spectroscopic lines of N a I5890A (D 2), and 5896A (D 1), He I 5876A, and H of LkH 234 over a period of 31 days during O ct. N ov. 2003. The observations consists of high resolution spectra (R = 30,000) from the H obby-E berly Telescope (HET) using the High Resolution Spectrograph (HRS). Section 2 describes observations and in section 3 we discuss the results and section 4 gives the conclusion.

2. Observations

We have obtained high signal to noise ratio and high resolution spectra (R = 30,000, 0.2A) of LkHa234 covering HeI5876A, NaID2 & D1 lines and the Balmer H at 6563A. The observations were taken with the 9.2m Hobby-Eberly Telescope (HET) (Ram sey et al. 1998) using its High Resolution Spectrograph (HRS) (Tull 1998). The echelle spectral orders were optim ized to cover the wavelength range between 5700 to 6750A. This allowed the spectral lines of interest to be recorded by the \blue" CCD of the spectrograph that has fewer num ber of bad pixels than the \red" CCD . The results com prise of ve data sets over a tim e period of 31 days. The observation dates were: 7th, 13th, 23rd, 27th O ctober and 8th N ovem ber 2003. O n each night of observation, three ten m inutes exposures on LkH 234 were taken which were com bined to get high signal to noise ratio (S/N = 70 to)100). All data were reduced using standard IRAF routines under ECHELLE tasks. Every spectrum wasbias subtracted, and at-elded. SpectralCalibrations were done using Thorium -Argon lamp spectra taken either immediately before or after the observations. W e found that the wavelength calibration is accurate up to 0.05A.

H (6563A), He I 5876A, and Na I D 2 & D 1 spectral lines are shown in gures 1 and 2 respectively and the dates of observations are noted along side of the spectra. The x-axis is in term s of relative velocity (km / s) with respect to the center of the lines in the stellar rest fram e and the y-axis shows line intensities normalized with respect to the stellar continuum. In gure 2 the zero of the relative velocity scale is with respect to the stellar rest fram e of the sodium D 2 line.

3. Results & Discussion

3.1. G aseous accretion

We illustrate in gure 1 the variability in H and He I lines. These are seen as stellar photospheric absorption lines in the spectra observed on 7th O ct. 2003. However, in all the later days of observations the H appeared to be in em ission consisting of two components, the Red shifted Em ission C om ponent (REC), and the B lue shifted Em ission C om ponent (BEC). The H equivalent width varied from + 3A (on 7th O ct.) to -50A (13th O ct.) in just 6 days and it continued to rise up to -100A (27th O ct.) before it decayed to -74A (8th N ov.). The relative velocity between the REC and BEC varied between 150 km /s to 225 km /s with the REC stronger than the BEC by a factor of 2.5 to 3.5.

The spectra obtained on 7th Oct. 2003 do not show any sign of emission or activity in H or He I. This observation shows the star in a quiescent state, with neither accretion nor stellar wind a ecting the absorption line pro le. The variations in the He I line are found to correlate with the variations in the H . W henever the H BEC and REC are stronger the He I either lls up due to mild emission or shows a very large absorption width (alm ost 1.7 tim es the photospheric line width). The line width varied from 150 km/s to 260 km /s. A B5 star does not have su ciently energetic photons to ionize H e I (Jaschek & Jaschek 1995), so the lling up of the He I absorption line due to mild emission is an indicator of accretion phenomena (de Winter et al. 1999). We nd the disk lum inosity to be about 1% of the stellar lum inosity (L 10³L) with maximum disk temperature of 21000K, if we consider an accretion rate of 10 ⁷M per year (N atta et al. 2000).

Therefore, the double peak H (REC & BEC) proles and the He I mild emission are because of orbiting accreting gas on to the star and as well as out ow of circum stellar gas (BEC in H due to accretion heat). However, when the accretion lum inosity is not strong enough to produce enough ionizing photons, the He I line appears as a very broad absorption feature. The broad absorption feature is the e ect of the com bination of the stellar photospheric line and various unresolved circum stellar gas com ponents corresponding to REC and BEC of H .

The e ect of gaseous accretion are probably also seen on the nebular em ission lines. W e found a correlation between the total equivalent width in H and the peak nebular emission in H and Na I. Figures 1 & 2 show that the nebular H and Na I lines were of minimal intensity (1.06 above the continuum) on 7th 0 ct. The same lines are strongest (H 7.9, on the blue wing of the REC, and NaID2 line 3.0) on 27th Oct. when the H equivalent width is maximum (-100A). We plan to model the impact of the episodic accretion on the nebula using a photo-ionizing code (for eq. the Cloudy code) which will be published along with the entire spectra of LkH 234 elsewhere in the near future.

3.2. Planetesim al infall

Figure 2 shows the variability in the Na ID2 and D1 lines. Redshifted Absorption Components (RAC) were observed in Na I lines on only one night (13 Oct. 2003) with a maximum redshifted velocity of 200 km /s. The RAC's associated with both the Na I lines are of similar depth 100km/s) which is an indicator (for redshifts of saturation (unshielded Na I column density = 10^{12} /cm²) (de W inter et al. 1999). However, the maximum depth of these components is about 30% of the stellar continuum im plying partial coverage of the stellar photosphere. In addition to Na I RAC, very mild BEC (60 to 100 km/s and 1.05 to 1.07 above the continuum) are also seen on the 13th Oct. spectra. The variations in He I and H are observed even when no Na IRAC & BEC are seen (see qures 1 & 2). Thus the appearance and disappearance of the NaIRAC & BEC are uncorrelated with the variations in H and He I line pro les. We conclude that we witnessed a transient phenom enon on 13th Oct. 2003 whose e ects lasted a few days at most.

From the K eplerian dynamics of an infalling object (Beust et al. 2001) the most redshifted N a I absorption component velocity (200 km/s) should correspond to a distance of 0.1 AU from the star. At this distance both the number of ionizing photons and intensity of the stellar wind will be too high for any unshielded N a I to survive (Sorelli et al. 1996). Their calculations using photo-ionizing codes under local therm odynam ical equilibrium (LTE) and non-LTE show that only a solid body like a com et or asteroid can approach a hot star this close before it disintegrates and evaporates. Further, theoreticalm odel calculations for stars up to B9 show that the size of solid bodies which are able to survive up to a distance of 0.1 AU should be at least 100 km in diam eter (Beust et al. 2001). The mild Na I BEC could then be part of the falling evaporating body being blown away by the stellar wind or radiation pressure. How ever, this will also m ean that the falling solid body m ay have a high eccentric orbit so that the blown away m aterial is not projected against the stellar surface. Our Na I RAC and BEC observations are consistent with such an object dissociating between 2 AU and 0.1 AU from the B5 star LkH 234.

In many Herbig Ae stars, variations in Na I RACs and BAC's usually correlates with H and He Ivariations (deW interet al. 1999, Grinin et al. 1996). Such N a I variations are now thought to be generated by m agneto-hydrodynam ic funnelling of neutral gas on to the star rather than solid body infall (Beust et al. 2001, M ora et al. 2002, N atta et al. 2000). No such correlation is present in the current LkH 234 data set. Of particular signi cance is the observation on 27th Oct. 2003 when the REC and BEC of H are the strongest. The absence of any NaIRAC and BEC in this observation is a strong argum ent against the N a I line variations being caused by neutral matter funnelled onto the star by magnetic elds. The variability of H and He I in LkH 234 are due to episodic gas accretion, while the observed variations in N a I lines is a transient event that exhibits the dynam ical signature of a body (100 km in diam eter) falling onto the star. It is worth mentioning here that recent studies on m eteors from com etary origin by Trigo-Rodriguez et al. (2004) have shown greater sodium abundances than those expected for interplanetary dust particles and chondritic m eteorites and Potter et al. (2002) have detected com et like sodium tail from planet M ercury, and also the discovery of presence of sodium tail in com ets (C rem onese et al. 1997). Thus an infalling solid body of asteroid size and de cient in H can produce the observed Na IRAC and BEC and no correlation with the H and He I lines.

The dusty circum stellar disk of the young star LkH 234 could be a new ly form ed protoplanetary disk consisting of solid bodies of di erent sizes (Fuente et al. 2001). Theoretical models of form ation of planetary bodies in circum stellar disks around a 1M mass star show that planetesim als of sizes of up to a few hundred kilom eters can quickly form within the rst 10^5 years and the typical cum ulative number of such bodies can be 10^4 to 10^6 between 0.5 to 1.5 AU from the star (Lissauer 1993, W eidenschilling et al. 1997). W hile such numbers are not known for B type stars, LkH 234 (age 0.1 M yrs) may have high frequency of infall events (G rady et al. 2000).

4. Conclusion

W e have presented here high resolution spectroscopic monitoring of a Herbig Bestar LkH 234 in the Na I (D 2& D 1), He I (5876A) and H lines. W e found no correlations between the Na I and the He I and H line variability in the ve data sets over a period of 31 days (7th Oct. 2003 to 8th Nov. 2003) though we do nd a correlation between He I and H . Thus the origin of variation in the Na I line (RAC up to 200 km /s and BEC up to 100 km /s seen only on 13th Oct. 2003) is di erent from those of H and He I.W hile the H and He I line variations are due to episodic gaseous accretion, the Na IRAC indicate a dram atic transient event of solid body infall on to the star and the BEC as the blown away parts of the solid body not projected against the stellar surface. Considering the Keplerian dynamics and the harsh environm ent of a B5 star, we estim ate that a solid body of size 100 km broke up and disintegrated at a distance between 0.1 to 2.0 AU from the star. This makes LkH 234 the youngest system (0.1 M yrs) with evidence for protoplanetary bodies of asteroidal size. W e plan to pursue further spectroscopic monitoring of the starwith the HET to determ ine the frequency of such events and further understand the com plicated dynam ics of this very young circum stellar environm ent.

The authors would like to thank Larry R am sey for additional discretionary time on the Hobby-Eberly Telescope and Eric Feigelson and M ike Eracleous for important discussions on the subject. The authors would also like to thank the referee M ike Sitko for very valuable suggestions which m ade the content of the paper strong. The Hobby-Eberly Telescope (HET) is a joint project of the University of Texas at Austin, the Pennsylvania State University, Stanford University, Ludwig-Maxim ilians-Universitat Munchen, and Georg-August-Universitat Gottingen. The HET is named in honor of its principal benefactors, W illiam P. Hobby and Robert E. Eberly. A \mathcal{L} . and J \mathcal{G} .would like to acknow ledge National Science Foundation for grant AST-01-38235, AST-02-43090 and NASA for grants NAG 5-12115 and NAG 5-11427. SM.would like to acknow ledge the support of the Michelson Fellow ship Program (JPL, NASA).

REFERENCES

- Beust, H., Karmann, C. & Lagrange, A.M. 2001, A & A, 366, 945
- C rem onese, G ., B oehnhardt, H ., C rovisier, J., et al. 1997, ApJ, 490, L199
- de W inter, D , G rady, C A , van den Ancker, M E . et al. 1999, A & A , 343, 137
- Fuente, A., Neri, R., Martin-Pintado, J., et al. 2001, A&A, 366, 873
- G rady, C A., Sitko, M L., Russel, R W., et al. 2000, in Protostars & Planets IV, page 613, The University of Arizona Press, editors: V.Mannings, A P.Boss and S.S.Russel
- G rinin, V P., K ozlova, O .V., The, P.S., et al. 1996, A & A , 309, 474
- Henbig, G.H. 1960, ApJS, 4, 337
- Herbst, W . & Shevchenko, V S. 1999, AJ, 118, 1043
- Jaschek, C. & Jaschek, M. 1995, The behavoir of chem ical elements in stars, Cambridge University Press

Lissauer, J. 1993, ARA & A, 31, 129

- Mora, A., Natta, A., Eiora, C., et al. 2002, A&A, 393, 259
- Natta, A., Grinin, V.P. & Tambovtseva, L.V. 2000, ApJ, 542, 421
- Natta, A., Grinin, V. & Mannings, V. 2000, in Protostars & Planets IV, page 559, The University of Arizona Press, editors: V. Mannings, A P. Boss and S.S. Russel

- Polom ski, E F., Telesco, C M., Pina, R., et al. 2002, AJ, 124, 2207
- Potter, A.E., Killen, R.M. & Morgan, T.H. 2002, Meteoritics & Planetary Science, 37, 1165
- Ram sey, L W ., A dam s, M .T ., Barnes, T G ., et al. 1998, Proc. SP E, 3352, 34
- Sm ith, B.& Terrile, R. 1984, Science, 226, 1421
- Sorelli, C., Grinin, V P.& Natta, A. 1996, A&A, 309, 155
- Thebault, P.& Beust, H. 2001, A&A, 376, 621
- Trigo-Rodriguez, JM ., Llorca, J. & Fabregat, J. 2004, MNRAS, 348, 802
- Tull, R.G. 1998, Proc. SP E, 3355, 387
- Viera, SLA., Pogodin, MA. & Franco, GAP. 1999, A&A, 345, 559
- W eidenschilling, S.J., Spaute, D., Davis, D.R., et al. 1997, Icanus, 128, 429

T h is 2-column preprint was prepared with the AAS ${\rm IAT}_E X$ m acros v5.0.



Fig. 1. The left panel shows the spectra of LkH 234 around the H (6563A) emission line and the right panel around HeI (5876A). The x-axis is in relative velocity with respect to the rest wavelengths of H and HeI respectively. The y-axis is the ux normalised with respect to the stellar continuum. The dates of observations are shown with each spectrum. The narrow 20 km /s width nebular H emission line is seen near the core of the stellar photospheric absorption line on 10-07-03 and so are m any nebular absorption lines. The nebular H line can also be seen on the blue wing of the REC on the other dates of observations.



Fig. 2. | The spectra of LkH 234 in the region of Na ID2(5890A) and D1(5896A) absorption features. The x-axis has been converted into relative velocity (km/s) with respect to the Na ID2 line at the stellar rest frame. The y-axis is same as in gure 1. The Red shifted Absorption Components (RAC), the nebular emission and the interstellar absorption lines are marked. The dates of observations are shown along with each spectrum. A lso seen are the very mild B lueshifted Absorption Components (BEC; on 10-13-03) on the blue side of the Na I (D2&D1) interstellar absorption lines. The interstellar Na I lines are o set by 27 km/s from the rest frame of the star.