# GMRT Discovery of A M illisecond Pulsar in a Very Eccentric Binary System

Paulo C. Freire<sup>1</sup>, Yashwant Gupta<sup>2</sup>, Scott M. Ransom<sup>3;4</sup>, Ishwara-Chandra C. H<sup>2</sup>.

# ABSTRACT

We report the discovery of the binary millisecond pulsar J0514 4002A, which is the rst known pulsar in the globular cluster NGC 1851 and the rst pulsar discovered using the G iant M etrew ave R adio Telescope (GMRT). The pulsar has a rotational period of 4.99m s, an orbital period of 18.8 days, and the m ost eccentric pulsar orbit yet m easured (e = 0.89). The companion has a minimum m ass of 0.9M and its nature is presently unclear. A fler accreting matter from a low m ass companion star which spun it up to a (few) millisecond spin period, the pulsar eventually exchanged the low m ass star for its more massive present companion. This is exactly the same process that could form a system containing a millisecond pulsar and a black hole; the discovery of NGC 1851A demonstrates that such systems might exist in the Universe, provided that stellar m ass black holes exist in globular clusters.

Subject headings: binaries: general | pulsars: general | pulsars: individual (PSR J0514-4002A) | globular clusters : general | globular clusters: individual (NGC 1851)

### 1. Introduction

Since 1987, several globular cluster surveys at Jodrell Bank, A recibo, Parkes and more recently the G reen Bank telescope have found 79 pulsars in a total of 23 globular clusters

<sup>&</sup>lt;sup>1</sup>NAIC, A recibo O bærvatory, HC3 B ox 53995, PR 00612, USA; pfreire@naic.edu

<sup>&</sup>lt;sup>2</sup>National Centre for Radio Astrophysics, P.O. Bag 3, Ganeshkhind, Pune 411007, India; ygupta@ncra.tifr.res.in, ishwar@ncra.tifr.res.in

<sup>&</sup>lt;sup>3</sup>D epartm ent of P hysics, R utherford P hysics B uilding, M cG ill U niversity, 3600 U niversity Street, M ontreal, Q uebec, H 3A 2T 8, C anada; ransom@physics.mcgill.ca

<sup>&</sup>lt;sup>4</sup>Department of Physics and Center for Space Research, Massachusetts Institute of Technology, Cambridge, MA 02139

 $(G C s)^5$ . These notings have con m ed that most of the binary millisecond pulsars (M SP s) in G C s have low -m ass white dwarf (W D) companions and nearly circular orbits, as observed in the G alactic disk. This is an important con mation of the evolutionary scenarios proposed by A par et al. (1982) for the form ation of M SP s.

In GCs, exchange encounters, which only have a signi cant probability of occurring in dense stellar environments, occasionally exchange one of the components of a binary system with a typically more massive star. The exchanges may occur during encounters with either other binaries or with isolated stars. In GCs such encounters can place isolated neutron stars into binaries with a main sequence (MS) star which eventually evolves, \recycles" the neutron star, and nally forms a MSP WD binary system. Such a process explains the anomalously large numbers of MSPs in GCs (by mass) when compared to the Galaxy. If the neutron star placed in orbit around a low-mass MS star is already a pulsar, we observe \irregular" eclipsing binary pulsars, such as PSR J0024 7204W in 47 Tucanae (Edmonds et al. 2002), PSR B1718 19 in NGC 6342 (Lyne et al. 1993), and PSR J1740 5340 in NGC 6397 (Ferraro et al. 2001), which are peculiar to GCs.

In this paper we report and discuss the discovery of a unique binary m illisecond pulsar, PSR J0514 4002A (henceforth NGC 1851A) in the GC NGC 1851. We discovered this system in a new 327-M Hz survey of GCs carried out using the G iant M etrew ave R adio Telescope (GMRT), at K hodad near Pune, India.

# 2. The GMRT survey observations and the discovery of NGC 1851A

O ur 327-M Hz survey aims to nd faint pulsars with steep radio spectra which would be missed by the high-frequency searches. It bene ts from the large gain of the central array of the GM RT of 4.6 K /Jy when used in the phased array mode of the A rray C om biner (G upta et al. 2000; P rabu 1997). Thism ode produces a beam on the sky with a diam eter of about 3 arcm inutes. The number of spectral channels across the available 16-M Hz band is 256 and the sam pling intervalused is 258 s. Each observation consists of a pair of 72-m inute scans containing  $2^{24}$  sam ples. Between scans, the 14-antenna central array was re-phased using a reference source.

We observed a set of 16 GCs in 2003 February. The data were written to tape and taken to McG ill University, where we processed them using the BORG (a 104-processor Beowulf cluster dedicated for pulsar processing) running the PRESTO software package (R ansom 2001).

<sup>&</sup>lt;sup>5</sup>See http://www2naic.edu/ pfreire/GCpsr.htm land references therein

O ne of the GCs observed was NGC 1851. Its distance (D) from the Sun is about 12.6 kpc (Cassisi, De Santis, & Piersim oni 2001), and its Galactic coordinates are l = 244:51; b = 35:04 (Harris 1996)<sup>6</sup>. It is a relatively bright GC (M<sub>v</sub> = 8:33) with a very condensed core (c = log ( $r_t=r_c$ ) = 2:32, where  $r_t$  and  $r_c$  are the tidal and core radii). It is am ong the ten clusters in the Galaxy with the highest central lum inosity density ( $_0$  ' 2 10<sup>5</sup> L pc<sup>3</sup>).

Neither of the prominent electron density models of the Galaxy make reliable predictions of the electron column density (normally expressed as the Dispersion Measure, or DM) to NGC 1851. The Taylor & Cordes (1993) model predicts a DM 28 pccm<sup>3</sup> for D > 3 kpc, while the Cordes & Lazio (2002; NE2001) model predicts DM 45 pccm<sup>3</sup> for D > 7 kpc. We therefore dedispersed the raw data at 500 trial DMs, from 20 to 70 pccm<sup>3</sup> and spaced by 0.1 cm<sup>3</sup> pc in order to minimize dispersive smearing and keep the total time resolution of our data between 0.5 and 1.5 m s. For each of these DM trials, a full Fourier-domain matchedlter-based acceleration search (Ransom, Eikenberry, & Middleditch 2002) was carried out, enabling the detection of binary systems with relatively short orbital periods. This is only possible because of the computing power of the BORG.

In the rst scan for NGC 1851, taken 2003 February 10, we detected a clear pulsed signal with a period of 4.991 m s (see Figure 1). The signal was present at all frequency channels within the observing band and the arrival times versus frequency are well described by a DM of 52.15(10) pc cm<sup>3</sup>. This value will help to re ne electron density models of the G alaxy in the direction of NGC 1851. The observed pulsations had a width of 0.8 m s, which is exactly the time resolution of the system after accounting for the dispersive sm earing within each of the 62.5-kH z channels and the sam pling interval. The intrinsic pulse is therefore essentially unresolved, in plying that future observations with better time and frequency resolution will signi cantly improve the signal-to-noise and timing precision. We con immed this signal in NGC 1851 data taken on the February 10, 11 and 20, the signal is not present in pointings m ade towards other clusters. This is therefore the rst pulsar discovered in NGC 1851 and, incidentally, the stGMRT pulsar discovery. Highly signi cant changes of the barycentric pulsed period both within and between observations of the cluster immediately identi ed the pulsar as a member of a binary system. Further observations of NGC 1851 were made in 2003 December in order to determ ine the orbit of the pulsar, using the same observing system and parameters as for the original search. A nearly analysis of the orbital trajectory of the pulsar in the P-P-diagram (Freire, K ram er, & Lyne 2001) suggested that the orbit is not circular.

<sup>&</sup>lt;sup>6</sup>See http://physwww.physicsmcmaster.ca/harris/mwgc.dat for the updated version of the table of GC parameters presented in this paper

A nalysis of the rotational periods using TEMPO<sup>7</sup> proved most surprising: the best-t model (see Figure 2) indicates e = 0.389(2), the most eccentric orbit of any known binary pulsar, and several orders of magnitude more eccentric than the typical orbits of fast M SP s<sup>8</sup>. The orbital period P<sub>b</sub> is 18.7850(8) days, and the sem im a praxis of the orbit projected along the line-of-sight (x) is 36.4(2) light seconds. This implies a minimum companion mass of 0.9M , assuming a pulsar mass of 1.35M (for the median of expected inclinations, 60, the companion mass is 1.1M ). The epoch of periastron T<sub>0</sub> is M JD = 52984.46(2), the longitude of periastron ! is 82(1) and the barycentric rotational period P is 4.990576(5) m s.

## 3. Flux density and position

W hen the pulsar is observed, the cross-correlations between the voltages of all active antennae (including those outside the central square, which are not directly used to search for the pulsar) are also recorded. This allows for simultaneous high-resolution imaging of the cluster. We found a faint object at  $= 05^{h}14^{m}06.74 \quad 0.06^{s}$ ;  $= 40.02^{0}50.0 \quad 1.3^{00}$  (see Figure 3), which is only 0.1 arcm inutes from the center of the cluster ( $= 05^{h}14^{m}062^{s}$ ; =

40  $02^{0}50^{0}$ , Harris 1996). This places the source, as projected in the plane of the sky, just outside the cluster's 0.06-arcm inute core. Such a position is common among the pulsars with known positions within GCs as is expected due to the elects of dynamical friction and m ass segregation (Kulkarni & Anderson 1996; Freire et al. 2001).

The ux density S for this object is variable. For example, on the 5D ecem ber in age, the source is not detectable (4 lim it of 2.4 m Jy), while on 6, 12 and 30D ecem ber it is detectable with ux densities of 5.5 0.5, 3.5 0.5 and 4.2 0.5 m Jy, respectively. This correlates well with the relative intensities of the 4.99 m s pulsed signal for those days. The brighter source to the north-east, on the other hand, has a constant ux density (within 10%, which is the typical accuracy of the ux calibration schem e). There is therefore little doubt that the faint source near the center is NGC 1851A. The in age in F igure 3 is a combination of these four epochs and gives a nalestim ate of the mean ux density towards the cluster, indicating that we are not m issing the detection of bright pulsars because of selection e ects such as very short rotational or orbital periods. We also observed the cluster at 610M Hz on 7 and 23

<sup>&</sup>lt;sup>7</sup>http://pulsar.princeton.edu/tem.po

<sup>&</sup>lt;sup>8</sup> for this and following quantities, the number in parenthesis indicates the 1- uncertainty, which we conservatively estimate to be ten times the form alvalue computed by TEMPO based on the fact that there is still no phase-connected timing solution

December, as well as 1 January. We did not detect the pulsar in any of the individual epochs nor in a combined map made from all three observations at a 4-level or better. From the rms noise of 0.1 m Jy, we estimate the 610-M Hz ux density to be < 0.4 m Jy. This implies a very steep spectrum, with < 3.4. This underlines the relevance of low-frequency surveys; it is unlikely that this pulsar could have been found at any other frequency.

#### 4. Form ation and nature

All known eccentric (e > 0:1) binary pulsars in the disk of the G alaxy have relatively m assive companions. This varied set of companions includes blue giants, other neutron stars and heavy W D s. Blue giants live only a few M yr which is probably not long enough to allow the sustained m ass accretion required to spin up a pulsar companion to m illisecond spin periods. This is in accordance with the observations, these pulsars have rotational periods of tens or hundreds of m s. The second supernova event is likely to m ake the orbit signi cantly eccentric, presum ing the binary survives. Since both stars are now compact, tidal circularization is henceforth in possible. The prolonged episode of stable m ass accretion needed to spin up a neutron star to m illisecond periods is only possible from evolved low erm ass M S stars. Such large tim escales allow e ective tidal orbit circularization as well. This process likely created the M SP currently in NGC 1851A, although with a presently unknown low -m ass W D companion.

SomeMSP + low-mass white dwarf systems in GCs, like PSR B1802 07 (P = 23:1ms, P<sub>b</sub> = 2:62 days, e = 0:212, (D 'Am ico et al. 1993)) can becom e m ildly eccentric due to interactions with other objects in the cluster (Rasio & Heggie 1995). This is alm ost certainly not the origin of the present NGC 1851A binary system since its eccentricity is probably too high to be explained by this mechanism and its companion too massive. We are therefore lead to the conclusion that after recycling, NGC 1851A exchanged its form er low -m ass com panion with a more massive object. This is the rst system presenting clear evidence of such a process. A massive star (the pulsar's present companion) passed within a distance smaller than about four times the separation of the components of the previous binary system. The most likely outcome from such an event is the form ation of an eccentric binary system containing the two more massive objects (Hut 1995). The low-mass component of the previous binary is ejected, causing the new binary system to recoil in the opposite direction. The probability for encounters is obviously larger for wider binaries, but the resulting recoil velocities are smaller. These events lead to relatively wide binaries (as observed for NGC 1851A) which are likely to remain near the center of the GCs. This is consistent with the projected position of NGC 1851A relative to the center of the cluster. If the original binary system is tightly

bound, the resulting new binary system is also likely to be very tight. Such an encounter would produce a large recoil velocity and could make the resulting system escape the cluster, or send it to the cluster's outer regions.

There are three previously known eccentric pulsar binaries with massive companions in GCs, but none of them presents such strong evidence of this kind of exchange encounter. The rst is PSR B2127+11C (Anderson et al. 1990), a double neutron star in M 15 (P = 30.53 m s, e = 0.6813, P<sub>b</sub> = 0.33 days). This system is remarkably similar to the rst known Galactic binary pulsar, PSR B1913+16 (Weisberg & Taylor 2003); therefore no mechanisms specic to GCs are needed to explain its formation. However, the fact that this pulsar is much more distant from the center of M 15 than any of the other 7 pulsars known in that cluster (Anderson 1992) hints at an exchange encounter with a powerful recoil from the ejected object. The second system is PSR J1750 37, (Possentiet al. 2001). Its main characteristics (P = 111:6 ms, e = 0.71,  $P_{b} = 17.3$  days) are very similar to another Galactic binary pulsar, PSR J1811 1736 (Lyne et al. 2000). An exchange encounter is, again, not the only possible form ation mechanism, but de nitely a possibility. The third system is PSR J2140 2310B, in M 30 (Ransom et al. 2004), which contains a 13-m s pulsar and e > 0.5. Its rotational period is about half of the sm allest rotational period found for the eccentric Galactic systems. The formation of such an object is very likely to require an exchange encounter, but there is still a small probability that this is a M SP m assive W D system that became eccentric through distant encounters with other stars, like PSR B1802 07.

The nature of the companion of NGC 1851A is as yet unclear, it could be either a compact or extended object. In NGC 1851, a cluster with an age of 9 Gyr (Salaris & W eiss 2002) 1-M stars are now leaving the main sequence. Because of the lengthy episode of M SP recycling that preceded its formation, the present NGC 1851A binary system is very likely to be a few Gyr younger than the cluster in which it lies. M athieu, M eibom and D olan (2004) have determined that, for the open cluster NGC 188 (with an age of 7 Gyr and a stellar population similar to that of GC s), binary system s containing M S stars with orbital periods larger than 15 days have not yet had time to circularize. Therefore, the observed eccentricity of the NGC 1851A system does not rule out the possibility of the companion being an extended object. In fact, partial and/or irregular \eclipses" from an extended object such as a M S starm ay explain the apparent ux variability from NGC 1851A.

### 5. Conclusion

We have discovered a remarkable 5-m s binary pulsar, the rst to be found in the GC NGC 1851 and the rst pulsar to be discovered with the GMRT. Its orbit is them ost eccentric known for any system containing a pulsar, while its rotational period is much shorter than that of any other pulsar in an eccentric binary system. This indicates that, after becoming an MSP by accreting matter from a low-mass companion star, this neutron star has alm ost certainly exchanged it for its present, signi cantly more massive companion. If black holes exist in GCs, a MSP-black hole binary could be formed in exactly the same way.

If the companion is a compact object, then two relativistic elects will be measurable, namely the rate of advance of periastron and the Einstein delay. The measurement of both elects would lead to the determination of the masses of the two components of this system. A third relativistic elect, the orbital decay due to emission of gravitational waves, will be masked by the unpredictable acceleration of the binary in the gravitational eld of the cluster, so no tests of general relativity are likely to be possible. If the companion is a M S star, we might measure the orbital precession caused by the quadrupole moment of the companion star and possibly a rotation of the orbital plane around the axis of rotation of the companion (W ex et al. 1998). In addition, variations in DM with orbital phase may be detectable (Johnston et al. 1996), particularly if the companion is an active star. For NGC 1851A none of these elects have been conclusively measured to date.

We thank J. Coppens, B. Bhattacharya and the sta of the GMRT for help with the observations. The G iant M etrew ave R adio Telescope is run by the N ational C enter for R adio A strophysics of the T ata Institute of Fundam ental research. SMR acknow ledges the support of a Tom linson Fellow ship awarded by M of ill University. The computing facility used for this research was funded via a N ew O pportunities R essearch G rant from the C anada Foundation for Innovation. We also thank C. Salter and A. D eshpande for their suggestions concerning the present paper.

#### REFERENCES

Alpar, M.A., Cheng, A.F., Ruderman, M.A., & Shaham, J. 1982, Nature, 300, 728

Anderson, S.B. 1992. PhD thesis, California Institute of Technology

Anderson, S.B., Gorham, P.W., T.A. Prince, S.R.K., & Wolszczan, A. 1990, Nature, 346, 42

Cassisi, S., De Santis, R., & Piersim oni, A.M. 2001, MNRAS, 326, 342

- Cordes, J.M. & Lazio, T.J.W. 2002, ApJ. submitted, http://xxx.lanlgov/abs/astroph/0007310
- D'Am ico, N., Bailes, M., Lyne, A.G., Manchester, R.N., Johnston, S., Fruchter, A.S., & Goss, W.M. 1993, MNRAS, 260, L7
- Edmonds, P.D., Gilliland, R.L., Camilo, F., Heinke, C.O., & Grindlay, J.E. 2002, ApJ, 579, 741
- Ferraro, F.R., Possenti, A., D'Amico, N., & Sabbi, E. 2001, ApJ, 561, L93
- Freire, P.C., Camilo, F., Lorimer, D.R., Lyne, A.G., Manchester, R.N., & D'Amico, N. 2001, MNRAS, 326, 901
- Freire, P.C., Kramer, M., & Lyne, A.G. 2001, MNRAS, 322, 885
- Gupta, Y., Gothoskar, P., Joshi, B.C., Vivekanand, M., Swain, R., Sirothia, S., & Bhat, N.D.R.2000, in ASP Conf. Ser. 202: IAU Colloq. 177: Pulsar Astronomy - 2000 and Beyond, 277
- Harris, W. E. 1996, Astron. J., 112, 1487
- Hut, P.1995, in ASP Conf. Ser. 72: Millisecond Pulsars. A Decade of Surprise, 46
- Johnston, S., Manchester, R. N., Lyne, A. G., D'Amico, N., Bailes, M., Gaensler, B. M., & Nicastro, L. 1996, MNRAS, 279, 1026
- Kulkami, S.R. & Anderson, S.B. 1996, in Dynamical Evolution of Star Clusters Confrontation of Theory and Observations, ed.P.Hut & J.Makino, (Dordrecht: Kluwer), 181
- Lyne, A.G., Biggs, J.D., Harrison, P.A., & Bailes, M. 1993, Nature, 361, 47
- Lyne, A.G. et al. 2000, MNRAS, 312, 698
- Mathieu, R.D., Meibom, S., & Dolan, C.J. 2004, ApJ, 602, L121
- Possenti, A., D'Amico, N., Manchester, R. N., Sarkissian, J., Lyne, A. G., & Camilo, F. 2001, in Talk presented at the ACP Summer 2001 Workshop \Compact Objects in Dense Star Clusters", Aspen, 26 June 2001, http://arxiv.org/abs/astro-ph/0108343.
- Prabu, T. 1997. Master's thesis, Indian Institute of Science. see also http://www.rri.res.in/dsp\_ral/gac/gaomain.htm

Ransom, S.M. 2001. PhD thesis, Harvard University

Ransom, S.M., Eikenberry, S.S., & Middleditch, J. 2002, Astron. J., 124, 1788

Ransom, S.M., Stairs, I.H., Backer, D.C., Greenhill, L.J., Bassa, C.G., Hessels, J.W. T., & Kaspi, V.M. 2004, ApJ. submitted, http://xxx.lanl.gov/abs/astro-ph/0310347

Rasio, F.R. & Heggie, D.C. 1995, ApJ, 445, L133

Salaris, M. & Weiss, A. 2002, A&A, 388, 492

Taylor, J.H. & Cordes, J.M. 1993, ApJ, 411, 674

- Weisberg, J. M. & Taylor, J. H. 2003, in Radio Pulsars, ASP Conference Proceedings, Vol. 302. Held 26-29 August 2002 at Mediterranean Agronom ic Institute of Chania, Crete, Greece. Edited by Matthew Bailes, David J. Nice and Stephen E. Thorsett. San Francisco: Astronom ical Society of the Paci c, 2003. ISBN : 1-58381-151-6, p.93, 93
- Wex, N., Johnston, S., Manchester, R.N., Lyne, A.G., Stappers, B.W., & Bailes, M. 1998, MNRAS, 298, 997

This preprint was prepared with the AAS  $IPT_EX$  m acros v5.0.



Fig. 1. Discovery observation for NGC 1851A, in the globular cluster NGC 1851. Pulsed em ission is persistent throughout the scans. The pulse prole is rather narrow (top), corresponding to the time resolution of the system for this DM.



Fig. 2. The measured rotational periods of NGC 1851A, as would be observed at the barycenter of the solar system, as a function of the pulsar's orbital phase. The solid lines indicate the prediction of the best-tm odel.



Fig. 3. A radio in age of NGC 1851 at 327 MHz, made by combining the data from 4 di erent epochs of observations in December 2003. The pulsar is the faint source near the center of the cluster, which is indicated with a cross. The two circles indicate the core and half-mass radii (0.06 and 0.52 arcm inutes respectively). The brighter source to the north-east is not the pulsar; when the interferom eter was pointed at it no pulsed emission was detected. The sm all ellipse in the lower right corner indicates the dimensions of the synthesized beam of the interferom eter, which includes all the active antennas. The diam eter of the beam of the central square, used in the pulsar search, is about 3 arcm inutes.