

Far-Ultraviolet Spectroscopy of Star-Forming Regions in Nearby Galaxies: Stellar Populations and Abundance Indicators ¹

William C. Keel

Department of Physics and Astronomy, University of Alabama, Box 870324, Tuscaloosa,
AL 35487; keel@birmingham.ua.edu

Jay B. Holberg

Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ 85721;
holberg@argus.lpl.arizona.edu

Patrick M. Treuthardt

Department of Physics and Astronomy, University of Alabama, Box 870324, Tuscaloosa,
AL 35487

Received _____; accepted _____

Submitted to the Astronomical Journal

¹Based on observations made with the NASA-CNES-CSA Far Ultraviolet Spectroscopic Explorer. FUSE is operated for NASA by the Johns Hopkins University under NASA contract NAS5-32985.

A B S T R A C T

We present FUSE spectroscopy and supporting data for star-forming regions in nearby galaxies, to examine their massive-star content and explore the use of abundance and population indicators in this spectral range for high-redshift galaxies. New far-ultraviolet spectra are shown for four bright H II regions in M 33 (NGC 588, 592, 595, and 604), the H II region NGC 5461 in M 101, and the starburst nucleus of NGC 7714, supplemented by the very-low-metallicity galaxy I Zw 18. In each case, we see strong Milky Way absorption systems from H₂, but intrinsic absorption within each galaxy is weak or undetectable, perhaps because of the "UV bias" in which reddened stars which lie behind molecular-rich areas are also heavily reddened. We see striking changes in the stellar-wind lines from these populations with metallicity, suggesting that C II, C III, C IV, N II, N III, and P V lines are potential tracers of stellar metallicity in star-forming galaxies. Three of these relations – involving N IV, C III, and P V – are nearly linear over the range from O/H = 0.05–0.8 solar. The major difference in continuum shapes among these systems is that the giant H II complex NGC 604 has a stronger continuum shortward of 950 Å than any other object in this sample. Small-number statistics would likely go in the other direction; we favor this as the result of a discrete star-forming event ~3 Myr ago, as suggested by previous studies of its stellar population.

Subject headings: galaxies: individual (M 33, NGC 7714, M 101, I Zw 18) {
galaxies: stellar content | galaxies: abundances | ultraviolet: galaxies

1. Introduction

The history of galaxies is, in large part, the history of star formation. Massive stars play key roles both as highly visible tracers of star formation and as players in altering surrounding star formation and both energy and chemistry of the interstellar medium. It is these stars which dominate the observed properties of actively star-forming galaxies. The massive part of stellar populations is most clearly observed in the ultraviolet, where their energy distributions peak and competing light from cooler stars is minimal, as long as the foreground extinction allows escape of enough of this radiation. In such environments, ultraviolet studies of star-forming regions have proven fruitful in understanding these populations. The recent opening of the far-ultraviolet window, between Lyman and the Lyman limit, for deep observations, allows study of massive hot stars in a range where they fully dominate the spectrum. This relatively narrow band contains an embarrassment of spectral riches, with numerous lines from stellar winds as well as interstellar material both atomic and molecular. These include the strong and highly-ionized lines of O VI and the unique ability to measure cold H_2 . In addition, this piece of the spectrum is accessible for high-redshift galaxies, at least for composite samples where the Lyman forest can be averaged adequately, allowing the possibility of direct comparisons of stellar populations over a large span of cosmic time.

The very sensitivity of the far-UV light to star formation and reddening makes it a purer probe of some properties of star-forming regions than observations at longer wavelengths. Since only short-lived stars contribute significantly in the far-UV range, the details of star-forming history should matter only for very brief bursts (such as might be found in individual H II regions, but are less likely on galaxy scales). This makes the far-UV spectrum more sensitive to the stellar population itself than to its history. Furthermore, although the extinction is high, its differential effect across the far-UV band is modest, and

paradoxically the effective reddening to the stars we see is smaller than found at longer wavelengths. In observing stars intermingled with highly structured dust distributions, the "picket-fence" effect (Heisler & Ostriker 1988) means that most of the stars are so reddened as to make no significant contribution in the deep ultraviolet; all the stars we see are only lightly reddened. This also reduces the effects of the forest of H_2 absorption features because of the mixing of molecular gas and dust.

Observational data on nearby galaxies, before the Far-Ultraviolet Spectroscopic Explorer (FUSE), were limited to a handful of star-forming systems. Four starburst galaxies were observed using HUT on Astro-2 (Leitherer et al. 1995, 2002), largely to measure escaping radiation in the Lyman continuum, which provided initial data for comparison with models based on stellar spectra dating back to Copernicus. The strongest features fall into two blends near 970 ($Ly + C\ III$) and 1030 Å ($O\ VI + Ly + C\ II$). A Voyager 2 observation of M 33, with some spatial resolution in one direction, was analyzed by Keel (1998), showing that its far-UV continuum is virtually identical to those of the powerful starbursts, and that NGC 604 is bluer in this range than the overall disk. These data also suggested significant Lyman emission, with spatial profile suggesting an origin in the diffuse ISM rather than giant H II regions. Ironically, until the availability of FUSE observations, the richest information on galaxies shortward of Lyman came from objects at high redshift, particularly composite spectra of Lyman-break galaxies (Steidel et al. 2001, Shapley et al. 2003). To enable comparisons between local, well-studied star-forming systems and these powerful, young objects, we have undertaken a series of FUSE observations of star-forming regions in nearby galaxies. We present here the analysis of these spectra in the context of their stellar populations and systematic changes with metallicity. A companion paper (Keel, Shapley, & Steidel 2004) considers the comparison with composite spectra of Lyman-break galaxies.

2. Observations

2.1. Object selection

The FUSE targets were selected for known high UV flux and hence low reddening, and to span a range of metallicity. The regions targeted in M 33 include the brightest H II regions in the mid-UV, and span a radial range within the disk from 1.6–4.5 kpc (see Fig. 1). They also include examples with a single dominant star cluster and with multiple clusters or more diffuse associations (also illustrated in Fig. 1 using HST data at 1700 Å). None of the objects in M 33 or M 101 achieves "super star cluster" (SSC) status, if we follow common usage in requiring not only high luminosity and stellar mass (Melnick, Moles & Terlevich 1985) but that the stars be concentrated into a single clump on 10-pc scales as noted by such studies as that of Meurer et al. (1995). While NGC 604 has the right luminosity, its stars are widely spread throughout a 100-pc region including multiple clumps (Hunter et al. 1996). In the more distant systems, the FUSE aperture samples multiple regions; Fig. 2 shows the bluest archival HST imagery with the aperture superimposed. In NGC 7714, there are ~10 luminous SSCs, comparably luminous in the near-UV, within the aperture, more clearly shown in the inset to Fig. 2. In each of these cases, essentially the entire star-forming region fits within the aperture, an important desideratum in comparing these observations with global measurements of distant galaxies.

For each of these objects, there are abundance measurements from the traditional optical emission lines, which we use as an widely-applicable tracer of the abundances of recently-formed stars as well. The M 33 objects span much of its disk's abundance gradient, while NGC 5461 and the nucleus of NGC 7714 were selected to sample lower and higher metallicity. We also analyze the summed archival FUSE spectrum of I Zw 18 from Aloisi et al. (2003), as a comparison with the lowest known gas-phase abundances.

Table 1 includes our adopted values from the literature for $[O/H]$, the best-measured of the abundances due to the optical lines from multiple ionization stages of oxygen and its importance as a coolant.

2.2. FUSE Spectroscopy

The FUSE optical system and detectors are described by Moos et al. (2000). Four primary mirrors are used to feed independent detectors optimized for subsections of the far-UV band; maintaining the coalignment of these optical systems is an important operational issue. Each detector has a distinct wavelength calibration, but for our purposes, we are not pushing the resolution limit of FUSE and can combine the various data segments in 0.1-Å pixels.

As shown in Figs. 1 and 2, the FUSE pointing positions were set at the midpoint of the stellar distributions from available UV images. The large 30" aperture was specified for the M 33 and M 101 H II regions at the outset, both for flux integrity in alignment of the four instrumental channels and to include essentially all their starlight, and adopted for NGC 7714 as well when the performance penalty for using a smaller aperture became clear during the mission. The requested center coordinates and total exposure times are listed in Table 1.

Of these objects, NGC 604 is the brightest in the far-UV range by a factor 10, and will thus play a continued role in our knowledge of stellar populations in this spectral range. In each of these spectra, the useful resolution is limited by the signal-to-noise ratio, dictating the wavelength binning to detect features of interest. For NGC 604, we can work at 0.1-Å binning, for a velocity spacing typically 30 km s^{-1} . Only in this object does it (marginally) matter that it is not a point source, with starlight coming from most of the aperture area.

This smearing along the dispersion axis contributes a broadening of about 60 km s^{-1} .

The spectra were processed through the FUSE pipeline; we inspected the two-dimensional spectra to verify data quality and background subtraction, and to note such effects as the "worm", a shadowing of the detector by a repeller wire, near the red end of some spectra. Each of our targets was observed during two orbits, so we could compare each orbit's summed spectra as a check on the errors at each wavelength. Loss of flux due to the "worm" does occur in some of the spectra around from $1170\text{--}1180 \text{ \AA}$, a range which does not affect any of the features we analyze. Similarly, there are spurious emission features near 1044 and $1169 \text{ \AA}$ which are second-order scattered solar features, which do not seriously confuse features of interest. For clarity of illustration, we interpolate across terrestrial airglow features in Ly γ , Ly β and O I at the highest resolution before further averaging and display.

We also use the summed archival spectrum of I Zw 18 described by Aloisi et al. (2003), who kindly provided it in electronic form. We compare the spectra of NGC 604 and I Zw 18 at $0.1\text{-\AA}$ resolution in Fig. 3, showing foreground absorption from H_2 , atomic absorption lines in the foreground and in these galaxies, and prominent stellar-wind lines.

2.3. Supporting Data

We draw on additional data from a variety of sources for these well-observed objects. IUE spectra in the large $10\text{--}20$ -arcsecond apertures give mid-UV spectra in roughly matched apertures; we have collected spectra from the archive, rejecting those with obviously discrepant pointings or flux levels from the combined spectra.

HST imaging is invaluable in examining the stellar populations, resolving the brightest stars in each of these associations. Archival WFC2 data exist for each of the M 33 regions

in F170W plus longer-wavelength filters, so we can assess the stellar statistics at least into the mid-ultraviolet. The best UV images of NGC 604 were taken in support of a slitless-spectroscopy program, using the STIS NUV-MAMA detector working at 1800–2700 Å.

Each of these stellar collections ionizes a substantial surrounding region. H α data (as in Bosch et al. 2002) show structure extending 200–800 pc. All four of the M 33 regions fit their description of an evolved H II region, one in which star formation has been long enough to generate extensive filaments and loops driven by stellar energy input; they estimate ages > 4 M yr for all except NGC 604, at 3 M yr, for the current episode of star formation (matching the conclusions of Gonzalez Delgado & Perez 2000).

Stars which contribute strongly in the far-UV have spectral types B0 and earlier, for ages up to 15 M yr. However, the continuum slopes of the shorter-lived O stars vary only slightly across the far-UV range, so the continuum offers little signature of the recent star-forming history (Robert et al. 2001). To this point, the far-UV line behavior is not well enough calibrated to infer the star-forming history independently.

3. Interstellar Absorption Features

The far-UV range is rich in narrow absorption lines from the ISM, both ionic and molecular. The spectrum of NGC 604 is especially complex, with over 100 detected and identified transitions. Foreground H $_2$ in the Milky Way is so prominent in all these objects that it must be carefully accounted for in identifying intrinsic absorption features. For the strongest Fe II lines, we detect not only the M 33 ISM (at about -200 km s $^{-1}$), but the foreground high-velocity cloud structure near -370 km s $^{-1}$ described by Wakker et al. (2003).

3.1. Atomic features

Numerous atomic absorption features are seen in these spectra, with stronger absorption in most cases from foreground Milky Way gas. Their strengths in the integrated spectra will reflect the distribution of gas toward the stellar associations weighted by contribution to the far-UV flux; for these extensive distributions of stars, derived column densities should be regarded as characteristic values.

3.2. Molecular hydrogen

The Lyman and Werner band systems of H_2 are at once an advantage and a nuisance in far-UV spectroscopy, detected in almost every line of sight crossing a significant distance in our own galaxy's ISM. A total of 56 such lines are individually detected in NGC 604. The H_2 features in these M 33 data have also been analyzed by Bluhm et al. (2003), who found that H_2 absorption at the velocities appropriate for disk gas in M 33 is weakly detected in NGC 588, 592, and 595, at column densities $N(H_2) = 10^{16} - 3 \times 10^{17} \text{ cm}^{-2}$, but only an upper limit 10^{15} could be derived for NGC 604, despite the higher data quality and comparable column densities from atomic lines, with all the values approximate due to the lines being too weak for independent measurement of the Doppler-width parameter. Bluhm et al. also present instructive simulations on the difficulty of deriving unique column densities when the background source consists of stars spanning a range of column density and position within the spectroscopic aperture. For example, the nondetection in NGC 604 might result from patchy extinction, correlated with the H_2 distribution, and the far-UV background flux being dominated by those stars with the shallowest H_2 absorption. This will act in addition to any physical effects related to the radiation field and shocks in the vicinity of such an active star-forming region. These regions are certainly associated with large concentrations of molecular gas; all three of them within the survey area of Engargiola

et al. (2003) are associated with molecular clouds (NGC 592, 595, and 604 with their clouds 40, 32, and 8, respectively). Their H_2 masses are estimated in the range 1.2×10^5 solar masses.

In spite of these issues in physical interpretation of molecular absorption, the empirical results are directly comparable to what we may see in high-redshift systems, as this spectral region becomes accessible for at least composite study in Lyman-break galaxies (Shapley et al. 2003). We can use both the sets of H_2 line strengths seen from Galactic gas in these spectra, and synthetic absorption spectra, to ask what signatures of molecular absorption remain at modest spectral resolution. The least confusion with other spectral features for modest H_2 column densities occurs for blends of features near 1005 and 1072 Å. The wavelengths and shapes of these blends depend on the spin-level populations in the gas, generally shifting to longer wavelengths for warmer gas (McCandliss 2003).

We have re-examined the NGC 604 spectrum for evidence of associated H_2 absorption. The ability to see such features at low redshift is limited largely by blending with foreground Galactic features, since the spacing between multiplet members provides ample opportunity for overlap. For atomic absorption species, we find a difference between foreground and M33 absorption typically 220 km s^{-1} , which gives wavelength shifts very close to the spacing of low-order members of the H_2 multiplets near 963, 982, 1002, and 1014 Å. As a result, only the bluest members of these multiplets could be securely detected from gas near the velocity of NGC 604 in the presence of the much stronger foreground gas. To guard against unrecognized atomic features, we also require that a putative NGC 604 feature not be seen in the I Zw 18 spectrum at the corresponding emitted wavelength. Compared to 56 H_2 lines seen from foreground Milky Way gas, only two potential detections resulted for NGC 604. These are the transitions at rest wavelengths 1001.69 (5;4 ! 13;1) and 1008.38 (2;3 ! 8;0) at equivalent widths of roughly 0.02 and 0.08 Å, respectively. The 1008 line,

with better S/N ratio, appears substantially broader, with a profile compatible with that of the other narrow features. If accurate, these detections are insufficient to derive the absorbing column density, lacking information on the intrinsic Doppler broadening and level populations. If we adopt the "optimum" b -value 10 km s^{-1} for the NGC 588 and 592 detections from Bluhm et al., and incorporating the H_2O OLS template models described by McCandliss 2003, we derive $N(\text{H}_2) = 3 \times 10^{17} \text{ cm}^{-2}$ from this single transition and assuming the same mix of level populations as in the foreground gas. Hotter molecular gas could give comparable absorption at column densities several times lower, close to the Bluhm et al. limit.

Lines arising in the states $J = 0-3$ are seen in the foreground Milky Way gas, well below the saturated level that drove Rachford et al. (2002) to use detailed profile fitting. As in their work, we also find significant changes in derived $N(\text{H}_2)$ between various absorption bands. We follow them in obtaining a characteristic kinetic temperature from the level populations in $J = 0;1$ as

$$T_{01} = \frac{74 \text{ K}}{\log N(0) - \log N(1) + 0.954} : \quad (1)$$

For the NGC 604 foreground H_2 spectrum, we obtained $T_{01} = 113 \text{ K}$, at the high end of values seen by Rachford et al. for dense and self-shielded environments but reasonable for the more diffuse "intercloud" ISM. The populations in $J = 0;1$ are taken to represent a thermal (collisional) temperature, with "excess" absorption from higher J representing radiative excitation, so that only the lowest levels are useful in deriving the thermal environment of the molecular gas.

Similar issues affect the strongest H_2 lines in I Zw 18, albeit with less overlap from Milky Way features because of the larger velocity shift. From the stacked spectrum, Aloisi et al. (2003) set an upper limit of $N(\text{H}_2) = 5 \times 10^{14} \text{ cm}^{-2}$ against the emerging continuum.

4. Stellar Content

4.1. Stellar winds

Most of the stellar signature in far-UV spectra occurs in the wind lines, which dominate this spectral range thanks to the numerous resonance transitions of metals. Both the wind lines and the evolutionary tracks when winds are important should have strong metallicity dependences, since the radiation pressure driving the winds acts largely through the opacity of heavy elements. This behavior has been shown, albeit sometimes in complex ways, upon comparison of stellar spectra from the Milky Way and Magellanic Clouds. Leitherer et al. (2001) examined wind lines in the mid-UV, notably C IV, from SMC to Galactic abundances, confirming a significant metallicity dependence in its strength but noting that ionization states other than the dominant ones can behave differently as secondary abundance effects shift the ionization balance in the winds (as seen in Si IV, which is stronger in LMC stars than the Milky Way). These changes are most pronounced for the brighter luminosity classes. Among the most luminous stars, those in the Magellanic Clouds have smaller wind velocities even for comparable depth (e.g. Kudritzki & Puls 2000).

While photospheric lines in the far-UV range are weak enough to scale directly with abundances, they are either extremely weak or overlain by wind lines. Lamers et al. (1999) suggest that line blanketing is a more secure route to the stellar abundances. From the comparison by Robert et al. (2001), Si IV $\lambda 1122=8$ are high-excitation photospheric lines that should serve as useful indicators of the stellar population, independent of winds.

In interpreting the O VI profiles from galaxy-scale systems such as NGC 7714, Gonzalez Delgado et al. (1998) caution from HUT spectra that interstellar absorption from large-scale outflows can significantly overlap wind lines, specifically between Ly α and O VI $\lambda 1032$.

Even FUSE-based spectral syntheses in the far-ultraviolet extend only to 1000 Å (Robert et al. 2003), so we continue our empirical approach in comparing systems of various properties to seek differences linked to composition or history. We can, however, be guided by available syntheses of some of the prominent blends of features, such as the Ly γ -O VI-C II regions from 1025-1038 Å (Gonzalez Delgado et al. 1997). They find that Ly γ and C II absorption arise in B stars, at the cool end of far-UV contributors, while O VI P Cygni profiles come from the most massive stars. High spectral resolution is crucial to separating the interstellar absorption contributions in each case; for Ly γ , hydrogen column densities $N_H > 10^{21} \text{ cm}^{-2}$ lead to blending of stellar and interstellar components. Robert et al. (2001) show that O VI is very weak at SMC metallicities, and that the youngest population can be diagnosed from the presence of C II 1176 and C IV/N IV 1169 which are specific to hot O stars. At Magellanic Cloud abundances, the S IV 1063=73=74 lines have a wind contribution only from supergiants, and P V 1118=25 exhibit similar behavior.

To show some of the spectral differences seen with metallicity, Fig. 4 compares the spectra of NGC 604 and I Zw 18, now with the strongest Galactic H₂ features removed by fitting Gaussians or Voigt profiles (for the stronger lines), and plotted in the emitted wavelength frame. Fig. 5 is a similar comparison of NGC 7714 with NGC 604, illustrating the yet stronger features seen at near-solar metallicity.

The wind lines seen in these systems may be described as follows. For the fainter objects with short exposures (NGC 588, 592, 595, 5461, 7714) the data have been smoothed by typically 0.7 Å in making these assessments.

NGC 604 (Fig. 4): Combining both members of the O VI doublet to reduce confusion, there is a broad wind absorption reaching 2800 km/s and probably blending with Ly γ . O VI 1037, C III 977, and N II 1083 have nearly black cores. A distinct detached absorption may be present in C III from 1600-2000 km/s. There may be a broad wind

feature in N III to ~ 3000 km/s. P V is weak except for possibly photospheric components near zero velocity. The emission sections of P Cygni profiles are strong in O VI 1037, N II 1082, and N III 991.

NGC 588/592/595: These spectra have wind features as much like one another as their S/N ratio can tell, both individually and when averaged. Compared to NGC 604, the cores of C III, O VI, and N II are shallower, with residual intensities 10-40%. As noted below, the continuum level in NGC 604 is higher than in any of the other systems below 950 Å.

NGC 5461: a trough occurs in N IV to about 1000 km s^{-1} . There is a single well-defined trough in C IV to 600 km s^{-1} . No feature is obvious in N III or S III. Broad absorption is seen in both O VI lines to about 1000 km s^{-1} .

I Zw 18: There is at most weak O VI absorption between $1700\text{--}2800 \text{ km s}^{-1}$, seen only in the 1037 line. P Cygni emission is absent in O VI, N II, and N III by comparison with NGC 604. Essentially no blueshifted troughs occur for N III. Two features may be present for C III from $0\text{--}1200$ and $1700\text{--}2000 \text{ km/s}$, but blending with saturated interstellar C II is an issue. P V is very weak.

NGC 7714: This nucleus has the strongest wind features in our study. Both O VI lines are strong, extending to 900 km s^{-1} . The P V lines at 1118;1125 are broad and blueshifted suggesting a wind contribution, which is plausible for non-supergiants at its metallicity. N III 991 may have a broad wind component blending with the interstellar O I lines at about 1000 km s^{-1} . The core of C III is blended with O I but shows a wind trough to beyond 1000 km s^{-1} .

For high-redshift objects, it will be difficult or impossible to separate wind and interstellar contributions to some of these lines. For purposes of comparison, we have generated simple equivalent-width values with respect to the adjacent (pseudo)continuum

for lines in clean parts of the spectrum, typically spanning a 5-Å line region, and tabulate these in Table 2. A irglow emission contaminated the fainter M 33 regions (combined in the table as "M 33 avg") too strongly for a reliable measurement of N III 991. For the N II and N III lines we list both the absorption and net equivalent widths (in parentheses), for use when the components are not resolved. We include values for C IV 1548,1550 from summed IUE low-dispersion spectra, as these values were taken with a large enough aperture to sample most of the stellar population in a way much like the FUSE aperture. Errors on the FUSE value are typically 0.2 Å for such broad features, as reflected in the limits for some lines in I Zw 18, while the C IV errors are closer to 1 Å. Each of these lines shows a strong trend with emission-line metallicity (Fig. 6), showing that they do in fact have potential use as abundance indicators. The wind lines measure stellar values directly, in contrast to the large regions of the interstellar medium sampled by emission-line techniques. These relations are quite sensitive in the sub-solar regime of particular interest for the evolution of galaxies at $z > 3$. The equivalent widths measured for N IV, C III, and P V vary almost linearly with abundances. Our results indicate that these strong features can be used as metallicity indices for high-redshift stellar populations, in regimes for which the optical emission lines lie in infrared bands of high atmospheric emission and absorption, and can be a valuable tool for approaching the chemical evolution of galaxies in the range $z = 3 - 4$.

For ease of use, we present least-square quadratic fits to the data in Fig. 6, along with the derived constants needed to invert these fits with line equivalent widths as the independent variables. These interpolation curves are shown in Fig. 6, and the numerical values are found in Table 3. The O/H ratio is in solar units as found from emission-line analysis (as cited in Table 1, updated to the "new" solar abundance scale with $12 + \log(\text{O}/\text{H}) = 8.69$), and all equivalent widths are in Å in the emitted frame. For each line, the data have been approximately fitted in the form $EW = a_1 + a_2(\text{O}=\text{H}) + a_3(\text{O}=\text{H})^2$.

Similarly, an inverse fitting function for metallicity derived from each line is obtained from the quadratic formula, with constants tabulated as $(O/H) = c_1 + c_2 (c_3 + (c_4 EW))^{\frac{1}{2}}$. The values are based on least-squares fits with equal weights, slightly modified in two cases to keep the fit monotonic across the metallicity range of our sample. For [N II], we adopt a flat value $EW = 1 \text{ \AA}$ for $(O/H) < 0.5$ solar. These forms are for interpolation purposes, and their limitations appear from the fact that they do not all approach zero line strength at zero metallicity. Crudely speaking, wind-dominated lines would have a quadratic dependence on metallicity until saturation sets in, since the mass-loss rate and fraction of the mass in the right ionic state each depend on metallicity, while mostly photospheric lines (such as P V) are more nearly linear in strength with the abundances.

Strictly speaking, these relations apply only to galaxies with a long (even if episodic or weak) history of star formation, since the elements involved in these lines come from very different stellar processes. For example, oxygen should be enriched rather quickly, coming from massive stars, while carbon will take longer coming from intermediate masses, with the dominant sources nitrogen still somewhat ambiguous. At best, these relations could be taken seriously for single elements in high-redshift galaxies, and can in fact be used to test for the differential enrichment history of such systems, in much the same way that the behavior of Si IV and C IV lines with redshifts has been used to infer such differences by Mehlert et al. (2002), who derive a relation between C IV EW and metallicity consistent with our IUE analysis included in Fig. 6.

4.2. Far-UV spectra and the stellar mix

Beyond the stellar-wind lines, the greatest difference among all these far-UV spectra is the flux excess in NGC 604 from about 912–940 Å, above what is seen in any of the other objects either of higher or lower metallicity. Small-number statistics in the massive stars

might be expected to be more important for the less rich systems, but in M 33 it is these which have the same spectral shape as the more luminous and distant systems, leaving NGC 604 as the odd one out. Previous work on its stellar content indicates that there was a strong peak in its star-forming history about 3 Myr ago, recent enough to affect the mix of stars contributing in this spectral range. We have examined additional UV data on the stellar content of these M 33 H II regions to see how NGC 604 might be different.

The stellar population in NGC 604 has drawn considerable attention, in being one of the brightest star-forming complexes in the Local Group while offering a dramatic structural contrast to the dominant, compact stellar cluster in 30 Doradus. Bruhweiler, Miskey, & Smith Neubig (2003) combined mid-UV WFC2 images with a wide-slit STIS spectrum to extract individual spectra of the brightest members. They note that the ten most UV-luminous stars will dominate the integrated spectral features from NGC 604, and that two of these are located ~ 0.3 magnitude above the usual 120-solar-mass limit. Specifically, the "top ten" stars contribute 46% of the total measured flux just longward of Lyman α . Mid-UV STIS images (central wavelengths 1820–2700 Å) obtained in support of a slitless spectroscopy program by J. M. Ais-Apellaniz show, via the nearest pixel sampling, that there are additional spatially resolved companions to each of the brightest stars identified by Bruhweiler et al., although none so bright as to bias the measured colors or magnitudes. NGC 604 also contains significant numbers of WR stars, although with a WR/O star ratio near 0.1 rather than the 0.3 seen in, for example, NGC 595 (Driszen, Moutat, & Shara 1993).

To compare the populations in these H II regions as they affect the integrated far-UV spectra, we produced color-magnitude arrays for each based on the archival WFC2 data in F170W and F555W (as well as intermediate-wavelength data when available). Lacking imagery in the far-UV band itself, we use the mid-UV properties as proxies to at least identify the hottest luminous stars. Reddening corrections do not enter for our immediate

purposes, since we need to know only how many stars contribute to the UV flux and extinction does not vary greatly between 1700 and 1100 Å. Bruhweiler et al. find, for stars in NGC 604, that the 1100-Å extinction is about 1.5 times that at 1700 Å. Likewise, we are interested here only in stars selected from UV flux, so the diagrams sample only those stars well detected in the mid-UV data. These observational HR diagrams are shown in Fig. 7. We include only stars without serious crowding issues, measured within 0.3" radii. The samples include stars to about $m_{170} = 19$ on the STMAG scale (based on flux per unit wavelength), although the completeness varies considerably even at $m_{170} = 18$ because of differences in crowding; for NGC 604, the cumulative counts with flux suggest statistical completeness only above $m_{170} = 17.0$. The brightest stars in the rich population of NGC 604 are 1.5–2 magnitudes brighter than found in any of the other associations. The color range among UV-bright stars in NGC 588 is smallest, extending from the blue envelope near $m_{170} - m_{555} = -3.5$ redward only to -2.0 , while all the other regions have the diagram populated to $m_{170} - m_{555} = 0$, and in NGC 604 to 2.4. This may in part be a reddening issue, since the stellar distribution in NGC 588 is more compact, with less scope for differential reddening across the association, than the others. Indeed, dust lanes are prominent in the continuum images of NGC 604 in both the optical continuum and UV/optical colors; it is clear observationally that the UV reflects only the least-extinguished stars in this object. The observed color distribution in NGC 604 is rich in the bluest stars, but these stars are also represented with similar color in NGC 588 and 595. A more important difference is that the richer population in NGC 604 includes several stars with F170W magnitude brighter than seen in any of the other clusters. While richness effects mean that the brighter star-forming regions will have brighter first-ranked stars, the studies referenced above suggest that NGC 604 has undergone a distinct burst of star formation about 3 Myr ago, younger than the other regions, an event which is recent enough to leave its mark on the massive-star population. We now focus on these brightest stars.

These data allow us to address how much the brightest stars dominate the mid-UV flux in each case, incorporating the total F170W flux within the FUSE aperture. Fluctuations in the bright population could affect the integrated spectra of the lower-luminosity, sparser regions more strongly simply from statistics, but the detailed recent history of star formation will enter through aging as well. Given the temperature range of these stars ($3-4 \times 10^4$ K following Bruhweiler et al.), the hottest stars should be more dominant at the shorter wavelengths of the FUSE data. It is not clear that all the far-UV flux is from direct starlight; Hill et al. (1995a,b) and Malamuth et al. (1996) showed evidence that a significant fraction of mid-UV light from similar systems is scattered. The HST F300W image of NGC 604 shows reflection nebulosity, and smoothed versions of the mid-UV STIS images match its morphology, indicating that scattering is important at shorter wavelengths as well, as would be expected for a roughly λ^{-4} Rayleigh behavior. Since the FUSE aperture is not much larger than the stellar distribution in NGC 604, the widths of narrow absorption features are not a sensitive test of whether scattering is important at these wavelengths. We do see a role for scattering around some of the brightest stars in the HST STIS image at 2400 Å, from analysis of the point-spread functions of stars. This is easier to interpret than the distinct emission and absorption structures, since reflected continuum can be confused with emission from the weak [O II] doublet near 2471 Å. Some of the stars match the nominal PSF closely, while two of the brightest ones exhibit excess light from 0.3-1.0" from the core. This excess contains as much as 28% of the mid-UV light in the brightest case. These data leave open the possibility that scattered light is important at the shorter far-UV wavelengths.

We therefore bracket the total UV flux between the sum of detected stars and the large-aperture sum. For larger fractions of scattered light, the brightest individual stars are more dominant, since the total number of stars producing the observed light is smaller. Cumulative distributions of F170W magnitude are shown in Fig. 8, along with simple

geometric estimates of corrections for crowding (which are small, but underestimate the actual effect where stars are more clumped than random within the H II region). Accumulating the flux from the bright stars, we find that only about 20% of the total flux within the FUSE aperture at 1700 Å comes directly from the stars, less than half of what Bruhweiler et al. (2003) find at 1200 Å within a narrow slit. This may be a sign that scattering is important, since much of the scattered light we see at mid-UV wavelengths in the HST imagery is on larger scales than this. In contrast, as a fraction of the light from detected stars, the brightest ten (a good approximation to the stars brighter than any found in the other H II regions) contribute about 40%, more in line with the Bruhweiler et al. results. This also makes sense for these stars being able to affect the overall spectrum. However, the high temperature needed for the excess component in NGC 604 means that we are seeing a difference in history rather than simply small-number statistics in what stars appear at a given time.

4.3. Stellar populations

In general, metallicity will be manifested in the composite spectra both directly, through photospheric and wind lines, and indirectly, as the evolutionary tracks of stars change with abundances. Effects on the initial-mass function are too small to see at the abundances found in the M 33 disk, as shown by Malamuth et al. (1996) for some of the same H II regions we observed. They suggest that the excitation trends seen in the associated ionized gas result from the different emergent ionizing fluxes for stars at various metallicities. However, different evolutionary histories are still implied by the strong changes in wind properties seen with metallicity.

Comparison of the our spectra suggests that very recent events in the star-formation history do have observable impact in the far-UV, as exemplified by NGC 604. Its mix

includes light from a greater fraction of hot stars than the other systems, which fits with the other properties of this region in suggesting a burst so recent that even in the far-UV, it does not look like constant star-formation (in this case, ages ~ 3 Myr). In general, the timescales may be metallicity-dependent; Robert et al. (2003) show that evolved O I/III and B I/III stars appear later at lower abundances.

5. Summary

We have used FUSE spectra of star-forming regions in nearby galaxies, whose gas-phase metallicities range from 0.05–0.8 solar, to explore the utility of far-ultraviolet spectra in measuring the abundances in star-forming galaxies, as well as to probe the massive-star populations in these galaxies. The absorption lines from radiatively-driven winds prove to be very sensitive to metal abundance; all six strong and unblended species (including C IV from archival IUE data) have a strong, monotonic metallicity dependence. For N IV, C III, and P V, the relation between straightforward equivalent-width values and oxygen abundance from emission-line spectra is closely linear, suggesting that these lines will be useful in tracing the chemical history of galaxies from $z = 3$ –4, beyond which the Lyman forest makes even composite spectra progressively less informative.

The continuum of NGC 604 departs from the uniform shape of the other objects below 950 Å. After considering the effects of small-number statistics among the massive stars in these objects, we conclude that this difference probably traces to a discrete burst of star-formation ~ 3 Myr ago in NGC 604. This region had been considered by several previous studies to have hosted such a burst, on grounds of both morphology of the gas and fitting of the H-R diagram.

B.-G. Andersson was helpful in understanding some of the issues in scheduling and

data analysis from FUSE. We thank Alessandra Abissi and her collaborators for providing their summed FUSE spectrum of I Zw 18. We also acknowledge the community service provided by Steven McCandless in making his H₂OOLS compilation of data and routines available. Dick Tipping patiently explained some of the intricacies of the H₂ spectrum on several occasions. This work was supported by NASA through FUSE GI grant NAG 5-8959. We also made use of the MAST archive system in retrieving data from HST, IUE, and UIT. We thank the referee, Claus Leitherer, for a detailed, expeditious, and helpful critique.

R E F E R E N C E S

- Allende Prieto, C ., Lambert, D .L ., & Asplund, M . 2001, *ApJ*, 556, L63
- Alisi, A ., Savaglio, S ., Heckman, T .M ., Hoopes, C .G ., Leitherer, C ., & Sembach, K .R .
2003, *ApJ*, 595, 760
- Bluhm , H ., de Boer, K .S ., Marggraf, O ., Richter, P , & Wakker, B .P . 2003, *A & A* , 398, 983
- Bosch, G ., Terlevich, E ., & Terlevich, R . 2002, *MNRAS*, 329, 481
- Bruhweiler, F .C ., Miskey, C .L ., & Smith Neubig, M . 2003, *AJ*, 125, 3082
- Corbelli, E . & Schneider, S .E . 1997, *ApJ*, 479, 244
- de Vaucouleurs, G . 1959, *ApJ*, 130, 728
- Drissen, L ., Moutat, A .F .J ., & Shara, M .M . 1993, *AJ*, 105, 1400
- Engargiola, G ., Plambeck, R .L ., Rosolowsky, E ., & Blitz, L . 2003, *ApJS*, 149, 343
- Gonzalez-Delgado, R .M ., Perez, E ., Diaz, A .I ., Garcia-Vargas, M .L ., Terlevich, E ., &
Vilchez, J .M . 1995, *ApJ*, 439, 604
- Gonzalez Delgado, R .M ., Leitherer, C ., & Heckman, T . 1997, *ApJ*, 489, 601
- Gonzalez Delgado, R .M ., Leitherer, C ., Heckman, T ., Lowenthal, J .D ., Ferguson, H .C ., &
Robert, C . 1998, *ApJ*, 495, 698
- Gonzalez Delgado, R .M . & Perez, E . 2000, *MNRAS*, 317, 64
- Heisler, J . & Ostriker, J .P . 1988, *ApJ*, 332, 543
- Hill, J .K . et al. 1995a, *ApJ*, 438, 181
- Hill, J .K . et al. 1995b, *ApJS*, 98, 595

- Hunter, D . A ., Baum , W . A ., O 'Neil, E . J., & Lynds, R . 1996, ApJ, 456, 174
- Izotov, Y . I., Cha ee, F . H ., Foltz, C . B ., Green, R . F ., Guseva, N . G ., & Thuan, T . X .
1999, ApJ, 527, 757
- Keel, W . C . 1998, ApJ, 506, 712
- Keel, W . C ., Shapley, A E ., & Steidel, C C . 2004, submitted
- Kim , M ., Kim , E ., Lee, M G ., Sarajedini, A ., & Geisler, D . 2002 AJ, 123, 244
- Kudritzki, R . & Puls, J. 2000, ARA & A , 38, 613
- Kuzio, R . E .; Ciardullo, R .; Feldmeier, J. J.; Jacoby, G . H . 1999 AAS 195, .1103
- Lee, M G ., Kim , M ., Sarajedini, A ., Geisler, D ., & G ieren, W . 2002, ApJ 565, 959
- Leitherer, C ., Ferguson, H . C ., Heckman, T . M ., & Lowenthal, J. D . 1995, ApJ, 454, L19
- Leitherer, C ., Leao, J. R . S ., Heckman, T . M ., Lennon, D . J., Pettini, M ., & Robert, C .
2001, ApJ, 550, 724
- Leitherer, C ., Li, I. H ., Calzetti, D ., & Heckman, T . M . 2002, ApJS, 140, 303
- Luridiana, V ., Esteban, C ., Peimbert, M ., & Peimbert, A . 2002, Revista M exicana de
Astronom ia y Astro sica, 38, 97
- Malum uth, E . M ., Waller, W . H ., & Parker, J. W . 1996, AJ, 111, 1128
- M cCall, M L ., Hill, R ., & English, J., 1990, AJ, 100, 193
- M cCandliss, S R . 2003, PASP, 115, 651
- M ehler, D ., et al. 2002, A & A , 393, 809
- M elnick, J., M oles, M ., & Terlevich, R . 1985, A & A , 149, L24

Meurer, G. R., Heckman, T. M., Leitherer, C., Kinney, A., Robert, C., & Garnett, D. R. 1995, *AJ*, 110, 2665

Moos, H. W. et al. 2000, *ApJL*, 538, L1

Quillen, A. C. & Yukita, M. 2001, *AJ*, 121, 2095

Rachford, B. L. et al. 2002, *ApJ*, 577, 221

Robert, C., Pellerin, A., Aloisi, A., Leitherer, C., Hoopes, C., & Heckman, T. M. 2003, *ApJS*, 144, 21

Shapley, A. E., Steidel, C. C., Pettini, M., & Adelberger, K. L. 2003, *ApJ*, 588, 65

Steidel, C. C., Pettini, M., & Adelberger, K. L. 2001, *ApJ*, 546, 665

Testor, G. & Lortet, M.-C. 1987, *A & A*, 178, 25

Vilchez, J. M., Pagel, B. E. J., Diaz, A. I., Terlevich, E., & Edmunds, M. G. 1988, *MNRAS*, 235, 633

Wakker, B. et al. 2003, *ApJS*, 146, 1

Waller, W. H., Parker, J. W., & Malumuth, E. M. 1996, *ASP Conf. Ser.* 99: Cosmic Abundances, 354

Fig. 1. | Locations and stellar distributions in the regions observed in M 33. The background image is U II image FUV-0496 (1500 Å), with ellipses corresponding to galactocentric radii in the inner disk of M 33. Each cutout from HST mid-UV image is 45 arcseconds square with north at the top, with an outline of the FUSE large science aperture at the recorded position for each observation. These regions range from single, compact associations (as in NGC 588) to the extended and multiple collections in NGC 592 and 604. The dispersion direction for these observations runs ENE-W SW; the limiting spectral resolution is set by the distribution of the far-UV starlight in this direction, which is a factor only for NGC 604. The images are displayed with an offset logarithmic intensity scale. The ellipses indicate distance from the nucleus in the disk plane, taking the geometric parameters for this part of the disk (within the inner non-warped region) from the optical fits by de Vaucouleurs (1959) and the H I fits of Corbelli & Schneider (1997). We adopt a distance of 850 kpc, in the middle of the range of distances from Cepheids (Lee et al. 2002), the tip of the red-giant branch (Kinn et al. 2002), and planetary nebulae (Kuzio et al. 1999).

Fig. 2. | FUSE aperture location and size for the NGC 5461 and 7714 observations, superimposed over the shortest-wavelength HST observation available. The ACS image of NGC 5461, from program 9490 led by K. Kuntz does not include the entire FUSE aperture. An inset shows the multiple luminous clusters in the starburst nucleus of NGC 7714, observed by Windhorst et al. under program 9124. As for the cutouts in Fig. 1, each image section spans 45 arcseconds with north at the top. The dispersion direction for NGC 5461 runs SSE-NNW, while for NGC 7714 it is ENE-W SW.

Fig. 3. | FUSE spectra of NGC 604 (heavy line) and I Zw 18, shown with 0.1-Å pixels in the heliocentric velocity frame, to reduce clutter from the H₂ absorption features. Fluxes have been scaled up by 10¹⁴ erg cm⁻² s⁻¹ Å⁻¹ for NGC 604, and by 2 × 10¹⁴ for I Zw 18. The numerous H₂ features (of which the most prominent are marked by ticks below the spectra)

are from Milky Way gas. Features in NGC 604 are slightly blueshifted (240 km s^{-1} , about $0.8 \text{ \AA}$ at $1000 \text{ \AA}$), while the redshift of I Zw 18 is $cz = 751 \text{ km s}^{-1}$ for a typical wavelength shift of $+2.4 \text{ \AA}$. Atomic interstellar absorption lines are marked at each redshift by vertical lines above the spectrum, where the shorter line is for I Zw 18. Angled symbols near the top indicate the zero-redshift locations of important stellar-wind lines. The continuum of NGC 604 is higher below about $940 \text{ \AA}$. While the interstellar lines are nearly as strong in I Zw 18 as the much higher-metallicity disk of M 33, the stellar-wind features are substantially weaker. The continuum level in NGC 604 is significantly higher shortward of about $955 \text{ \AA}$, and P Cygni emission redward of the 1037 line is prominent in NGC 604 but not in I Zw 18.

Fig. 4. | Comparison of the FUSE spectra of NGC 604 and I Zw 18, as in Fig. 3, now with Galactic molecular absorption removed and both spectra plotted in the emitted wavelength frame. Wind and interstellar lines are marked as before; some unpatched foreground absorption remains. Interstellar absorption features intrinsic to the surrounding galaxies stand out by matching in both spectra. This comparison shows the difference in both wind absorption and P Cygni between the abundances of I Zw 18 (O/H about 0.02 solar) and NGC 604 (0.4 solar). Both effects are clear for O VI, C III, and N II.

Fig. 5. | Comparison of the H_2 -corrected spectrum of NGC 604 to NGC 7714, in the emitted frame as in Fig. 4. Stronger absorption is prominent in the blue wings of O VI and C III, and in the overall profiles of N II and Si III/IV. The NGC 7714 data have been boxcar-smoothed by $0.7 \text{ \AA}$ and scaled by a factor 10^{14} .

Fig. 6. | Equivalent widths of stellar-wind lines in the spectra of star-forming regions. Each shows a strong metallicity dependence, here quantified using the traditional emission-line results for O/H. Typical errors for the far-UV lines are $0.2 \text{ \AA}$, with the IUE spectra used for C IV accurate to about $1 \text{ \AA}$. The three fainter M 33 regions are averaged into single

points for each transition. The quadratic interpolation functions with coefficients listed in Table 3 are overplotted as guides, where we take a fixed value $EW = 1 \text{ \AA}$ for $N_{\text{H II}}$ below $(O/H) = 0.5$.

Fig. 7. | Observed color-magnitude arrays for H II regions in M 33, derived from archival W FPC2 images. The two brightest objects in NGC 604 have less certain colors due to saturation in the F555W images. These are shown in the STMAG system, in which zero color index corresponds to constant F . Since we are interested in which stars contribute to the far-UV flux, no reddening corrections have been applied. The stars of interest are so blue that red-leak corrections in the F170W filter are negligible for our purposes.

Fig. 8. | Cumulative star counts as observed at $1700 \text{ \AA}$ for the M 33 H II regions, including stars within the FUSE apertures. In each region, the upper curve includes a simple correction for crowding, made by assuming that the fainter stars are uniformly distributed through the populated region in each association.

Object	RA	Dec	cz	Exposure	O/H	source
	2000	2000	km s ⁻¹	seconds	(solar units)	
NGC 588	01 32 45.50	+ 30 38 55	-174	5220	0.41	Vilchez et al. 1988
NGC 592	01 33 12.27	+ 30 38 49	-162	3965	0.48	Interpolated
NGC 595	01 33 33.60	+ 30 41 32	-178	7123	0.56	Vilchez et al. 1988
NGC 604	01 34 32.50	+ 30 47 04	-226	7151	0.66	Vilchez et al. 1988
NGC 5461	14 03 41.30	+ 54 19 05	298	5189	0.68	Luridiana et al. 2002
NGC 7714	23 36 14.0	+ 02 09 19	2798	6023	0.81	Gonzalez-Delgado et al. 1995
IZw 18	09 34 02.30	+ 55 14 25	751	95097	0.05	Izotov et al. 1999

Table 1: FUSE Targets and Properties

Note. | O/H is in solar units, converted when necessary assuming a solar value of $12 + \log \text{O/H} = 8.60$ following Allende Prieto, Lambert, & Asplund (2001)

Object	N IV 955	C III 977	N III 991	N II 1083	P V 1122	C IV 1549
IZw 18	0.27:	0.47	< 0.2 (< 0.2)	1.11 (1.02)	0.27	2.0:
M 33 avg	1.26	1.19	... (...)	0.86 (0.24)	0.66	8.4 (7.8)
NGC 604	2.17	1.83	1.77 (0.59)	1.50 (1.33)	0.89	9.49 (7.97)
NGC 5461	2.11	1.56	2.40 (1.87)	2.19 (1.92)	0.90	7.82 (6.03)
NGC 7714	3.26	2.56	3.68 (2.36)	2.52 (2.52)	1.01	10.1 (10.2)

Table 2: Equivalent Widths of Stellar Wind Lines

Note. | All values are in Å in the emitted frame. Parenthesized values include the emission component of a P Cygni profile.

Transition	a_1	a_2	a_3	c_1	c_2	c_3	c_4
N IV	0.30	-0.656	5.2201	0.063	0.0958	-5.8314	20.8804
N III	0.05	-1.079	6.5539	0.082	0.0763	-0.0794	26.2157
N II	1.28	-4.029	7.0447	0.286	0.0710	-19.7753	28.1790
C IV	1.14	19.210	-10.8494	0.885	-0.0461	418.6269	-43.3975
C III	0.50	-0.502	3.6120	0.070	0.1384	-6.9750	14.4481
P V	0.22	0.910	0.1037	-4.386	4.8224	0.7356	0.41473

Table 3: Quadratic Fits for Line Strength versus Metallicity

Note. | Entries are coefficients of forward and inverse quadratic fits as listed in the text, when equivalent widths are in Å and O/H is in solar units.

This figure "fig1.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/0403499v1>

This figure "fig2.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/0403499v1>

This figure "fig3.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/0403499v1>

This figure "fig4.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/0403499v1>

This figure "fig5.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/0403499v1>

This figure "fig6.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/0403499v1>

This figure "fig7.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/0403499v1>

This figure "fig8.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/0403499v1>