

Comment on “Strangelets as Cosmic Rays beyond the Greisen-Zatsepin-Cuzmin Cutoff”

Shmuel Balberg

Racah Institute of Physics, The Hebrew University, Jerusalem 91904, ISRAEL

(Dated: March 20, 2022)

PACS numbers: 98.70.Sa, 24.85.+p, 97.60.JD, 26.60.+c

In [1] Madsen and Larsen propose that strangelets - stable lumps of strange quark matter (SQM) - are candidates for the highest energy cosmic rays. They point out that the expected properties of strangelets make them easier to accelerate and less prone to energy losses than protons or ions. The purpose of this Comment is to point out that if ultra-high-energy cosmic rays (UHECRs) are indeed strangelets, their flux would guarantee that all neutron stars must be strange quark (SQ) stars, while no “conventional” (i.e., composed of nucleons and other baryons) neutron stars can coexist. This seems highly unlikely in view of various observed phenomena in neutron stars.

The hypothesis that SQM, composed of approximately equal abundances of up, down and strange quarks, is the true ground state of matter [2] has proved to be of special interest in the context of astrophysical compact objects. Self-bound SQM objects with masses of $M \lesssim 2 M_{\odot}$ would be gravitationally stable and, for the most part, resemble conventional neutron stars [3]. While there have been attempts to identify observations of some neutron stars as indication that they are SQ-stars, these were far from conclusive, and to date the existence of strange quark stars remains speculative.

If SQM is the ground state of matter, a strangelet will absorb free neutrons, converting them to additional SQM (ordinary matter is repelled by positively charged strangelets, and so is unaffected). Hence, if a strangelet penetrates a neutron star, the object converts to a SQ-star on a time scale of minutes [4]. Similarly, a massive stars which accumulates an SQM seed in its core will later give birth to a SQ-star, not a neutron star. The potential conversion of neutron stars into SQ-stars raises a key question: can both types of objects coexist in nature? If SQ-stars exist, then a fraction of them must participate in binary mergers with other compact objects, in which some SQM is injected into the galactic disk. The rate of binary mergers in the galaxy appears to be large enough so that if SQM is absolutely stable, the flux of strangelets would ensure that all objects suspected as neutron stars are actually SQ-stars. Correspondingly, if some neutron stars are identified as being conventional, the inevitable conclusion is that SQM is not the ground state of matter and there are only conventional neutron stars [5, 6].

This argument might be undermined if the magnitude of SQM contamination of the galaxy is suppressed, for example, if SQ-stars are disfavored in binaries (see [7] and references therein). However, there can be no ambi-

guity in the context of a “SQM as UHECRs” scenario, for which the flux of strangelets is predetermined by construction. At 10^{20} eV the flux is about one particle per 100 km^2 per year (see, e.g., [8]). If UHECRs are strangelets, an existing neutron star should absorb several of them per year, and a massive star in the red giant stage would be impacted by as many as 10^{16} yr^{-1} . Moreover, if the strangelets are accelerated in astrophysical phenomena (e.g., the Fermi mechanism in shocks), the flux at lower energies must be orders of magnitude larger (the differential spectrum will be no harder than $dN/dE \propto E^{-2}$).

In his reply [9], Madsen argues that if a significant fraction of the strangelet energy is deposited in the strangelet as it impacts particles in the target star, high energy strangelets will disintegrate. In particular, if the binding energy per baryon is E_B , the upper limit for the minimal strangelet baryon number that will survive disintegration is $A_{min} = E/E_B$, where E is the initial energy. The physics of strangelet fragmentation is essentially unknown, so the fraction of the initial energy which is deposited in the strangelet is unclear, as opposed to, e.g., pion production and acceleration of the target particles. A very high energy strangelet may also absorb baryons and grow in mass. But even if Madsen’s upper limit is realistic, lower energy strangelets can be stopped prior to disintegration with relatively small baryon numbers. The key is that the total flux of strangelets (at all energies) required to have an ultra-high energy component at the observed level is so large, that even a minute probability for strangelet survival guarantees complete conversion of neutron stars to SQ-stars. Coexistence of both types of objects is ruled out.

There do exist several observable indications that at least some neutron stars cannot be SQ-stars. In the more likely case of color-flavor-locked (CFL) [10] SQM, the crusts of SQ-stars would certainly too thin to be consistent with the thick crust required to explain Type I X-ray bursts, in which light elements fuse to heavier ones in explosive flashes [11]. Moreover, CFL SQ-stars probably cannot support millisecond rotation periods due to r -mode instabilities [12]. While non-paired SQM stars will have larger crusts that may be marginally consistent with these phenomena, they apparently cannot glitch (undergo a sudden spin up observed in several pulsars; this is the original argument of [5, 6]), and would also cool far too quickly to be compatible with observations [13]. While the absence of a satisfactory model for SQ-stars

which is consistent with all these observations cannot be considered as proof to the opposite, the more likely conclusion is that at least some neutron stars are, indeed, conventional. If so, the galaxy cannot be contaminated

with strangelets at the level required for an UHECR-model.

I wish to thank K. Rajagopal for valuable discussions and advice.

-
- [1] J. Madsen and J. M. Larsen, Phys. Rev. Lett. **90**, 121102 (2003).
 - [2] A. R. Bodmer, Phys. Rev. D **4**, 1601 (1971); E. Witten, Phys. Rev. D **30**, 272 (1984).
 - [3] C. Alcock, E. Farhi and A. V. Olinto, Astrophys. J. **310**, 261 (1986).
 - [4] A. V. Olinto, Phys. Lett. **B192**, 71 (1987).
 - [5] J. Madsen, Phys. Rev. Lett. **61**, 2909 (1988).
 - [6] R. R. Caldwell and J. L. Friedman, Phys. Lett. **B264**, 143 (1991)
 - [7] K. Belczynski, T. Bulik and W. Kluzniak, Astrophys. J. **567**, L63 (2002).
 - [8] L. Anchordoqui et al., Int. J. Mod. Phys. **A18** 2229 (2003).
 - [9] J. Madsen, Phys. Rev. Lett. **92**, 119002 (2004).
 - [10] M. Alford, Ann. Rev. Nucl. Part. Sci. **51**, 131 (2001).
 - [11] A. Cumming and L. Bildsten, Astrophys. J. **559**, L127 (2001); H. Shatz et al., Nucl. Phys. A. **718**, 247C (2003) and references therein.
 - [12] J. Madsen, Phys. Rev. Lett. **85**, 10 (2000).
 - [13] D. Page et al., Phys. Rev. Lett. **85**, 2048 (2000).