# Detection of $D_2H^+$ in the Dense Interstellar M edium

 $C.Vastel^1$  and  $T.G.Phillips^1$ 

California Institute of Technology, Downs Laboratory of Physics, MS 320-47, 1200 East California Boulevard, Pasadena, CA 91125, US

H.Yoshida<sup>2</sup>

Caltech Subm illim eter Observatory, 111 Nowelo Street, Hib, HI 96720, US

vastel@submm.caltech.edu

### ABSTRACT

The 692 GHz para ground-state line of  $D_2H^+$  has been detected at the Caltech Subm illimeter O bservatory towards the pre-stellar core 16293E. The derived  $D_2H^+$  abundance is comparable to that of  $H_2D^+$ , as determined by observations of the 372 GHz line of ortho  $H_2D^+$ . This is an observational veri cation of recent theoretical predictions (R observed in cold, high-density regions of the interstellar medium associated with low mass pre-stellar cores and protostars. This detection con m s expectations that the multiply deuterated form s of  $H_3^+$  were missing factors of earlier models. The inclusion of  $D_2H^+$  and  $D_3^+$  in the models leads to predictions of higher values of the D/H ratio in the gas phase.

Subject headings: ISM : abundances | ISM : clouds | ISM : molecules | molecular data | molecular processes | radio lines: ISM

## 1. Introduction

Recently, m illim eter and subm illim eter spectroscopy of the dense interstellar m edium has shown that, in cold dense regions, deuterated m olecular species are highly abundant, som etim es m ore than  $10^{-1}$  of the H version. A m azingly, doubly and triply deuterated species can be observed, e.g.  $D_2CO$  (C eccarelli et al. 1998), NHD<sub>2</sub> (R oue et al. 2000), CHD<sub>2</sub>OH (Parise et al. 2002),  $D_2S$  (Vastel et al. 2003), ND<sub>3</sub> (L is et al. 2002; van der Tak et al. 2002), CD<sub>3</sub>OH (Parise et al. 2004). Severalm odels have been developed to account for such high levels of deuteration (T ielens 1983; R oberts & M illar 2000a,b). Phillips & Vastel (2002)

have pointed out that the deuteration of  $H_3^+$  will be extended beyond  $H_2D^+$ , to  $D_2H^+$  and  $D_3^+$ , and that detection of  $D_2H^+$  might be possible. A calculation taking a high degree of deuteration into account has been carried out by Roberts, Herbst & M illar (2003) and W alm sley, F lower & P ineau des F ôrets (2004), con m ing the expectation that, in dense depleted regions, the abundance of  $D_2H^+$  will be similar to that of  $H_2D^+$ , and that  $D_3^+$  will be abundant.

The key enabling work in the astronom ical search for  $D_2H^+$  is the laboratory measurement of the para ground-state transition  $(1_{10}-1_{01})$  by H irao and Am ano (2003). We report here the rst astronom ical detection of that transition.

Chem ical reactions go in the direction to m inim ize energy. The chem ical fractionation process favors the production of the heavier m ore deuterated species, because of the m ass dependence of the zero-point vibration energies of the isotopic variants. G as phase species are expected to be depleted at the centers of cold, dark clouds, since they accrete on the dust grains (see, e.g., Chamley 1997). A series of observations has shown that the abundances of m olecules like CO decrease in m any pre-stellar cores (B acm ann et al. 2002). The rem oval of these reactive species a ects the gas-phase chem istry and particularly the deuterium fractionation within the cloud. Indeed, the rem oval of species that would norm ally destroy  $H_3^+$  (e.g. CO; R oberts & M illar 2000a) m eans that  $H_3^+$  is m ore likely to react with HD and produce  $H_2D^+$ . For example, if  $[CO/H_2] = 5 = 10^6$  (B acm ann et al. 2002), this leaves HD at  $[HD/H_2] = 5 = 10^5$  as the m ost abundant m olecule available for reaction with  $H_3^+$  and  $H_2D^+$ , and favors the production of high deuterium content m olecules:

$$H_3^+ + HD ! H_2D^+ + H_2 + E_a$$
 (1)

 $H_2D^+ + HD + D_2H^+ + H_2 + E_b$  (2)

$$D_2H^+ + HD ! D_3^+ + H_2 + E_c$$
 (3)

where  $E_a$ ,  $E_b$  and  $E_c$  are the released energies of the exotherm ic reactions. Using the zero-point energies computed by Ramanlal, Polyanski Tennyson (2003), and the energy of the rst allowed rotational state of the  $H_3^+$  molecule permitted by the Pauli exclusion principle (92 K), these values are:  $E_a = 230$  K,  $E_b = 180$  K and  $E_c = 230$  K.

A fter a long frustrating search (Phillips et al. 1985; Pagani et al. 1992; van D ishoeck et al. 1992; Boreiko & Betz 1993), and with the advent of new submillim eter receivers,  $H_2D^+$  was detected toward two young stellar objects, NGC 1333 IRAS 4A (Stark et al. 1999) and IRAS 16293-2422A (Stark et al. 2004), although with relatively low signal strength.

The  $H_2D^+$  search has now been extended to pre-stellar cores, and has been detected with relatively strong emission (Caselli et al. 2003; Caselli et al. 2004, in preparation; Vastel et al. 2004, in preparation) con rm ing that  $H_2D^+$  is dram atically enhanced in a gas depleted of m ost m olecules.

The ammonia and DCO<sup>+</sup> emission around the proto-binary system IRAS16293-2422 (W ootten & Loren 1987; M izuno et al. 1990; Lis et al. 2002) does not only peak on IRAS16293-2422 itself but shows a second peak, about 90<sup>th</sup> to the southeast, in a condensation called 16293E. Lis et al. (2002) found that CO in this region is depleted by a factor of 7. It is known to be a region where deuterium fractionation is strong and was chosen to be searched for  $D_2H^+$ .

## 2. Observations and results

The pure rotational transition  $(1_{10}-1_{01})$  of  $D_2H^+$  has been measured in the laboratory by Hirao and Am ano (2003). Spectroscopic observations of 16293E, presented here, were carried out in February 2004 using the facility receivers and spectrom eters of the Caltech Submillimeter Observatory (CSO) on Mauna Kea, Hawaii. The position chosen was the  $DCO^+$  peak em ission ( 2000 = 16h32m 29.4s, 2000 = -24°28°52.6°, Lis et al. 2002). We observed both the  $1_{10}$ - $1_{11}$  transition of ortho- $H_2D^+$  ( = 372.42134 (20) GHz, Bogey et al. 1984) and the  $1_{10}-1_{01}$  transition of para-D<sub>2</sub>H<sup>+</sup> ( = 691.660440 (19) G H z, H irao and Am ano 2003) (see Figures 1 and 2) which are the only lines currently available for these species. The data were taken under good weather conditions (225 GHz zenith opacity between 0.03 and 0.05). The CSO main beam e ciencies are 60% for the 345 G H z receiver and 40% for the 650 G H z receiver, determ ined from total power observations of M ars and Saturn. If the em ission is extended compared to the beam size of CSO, as appears to be the case for DCO  $^{+}$  (Lis et al. 2002), then the e ciency is about 70% at 372 GHz and 60% at 692 GHz. The FW HM beam size at 372 GHz is about  $20^{\circ}$ , compared to  $11^{\circ}$  at 692 GHz. Typical calibration 24%. The pointing of the telescope was determ ined from observations uncertainties are of Jupiter, and was stable about  $2^{00}$  (mm s). We used both the 50 MHz and 500 MHz bandwidth acousto-optical facility spectrom eters. The 500 MHz system was used to check system perform ance and calibration with the CO (6! 5) line, which is o set by 82 km s<sup>1</sup> from D<sub>2</sub>H<sup>+</sup>. There are no known lines of other interstellarm olecules within 50 M H z, likely to em it from such a cold region. In Table 1 we report the frequency, the antenna tem perature, the linewidth and the velocity relative to the local standard of rest, for the two lines. In Figure 3, we present 8.2 and 4.4 detections of H  $_2D^+$  and D  $_2H^+$  respectively, obtained in 23 m inutes and 103 m inutes on-source integration time, respectively.

Since the para ground-state transition for  $D_2H^+$  is the only line available to existing telescope facilities, it is not possible to obtain con m ation of the identi cation. However, the situation is very di erent from molecule detection in hot core regions, such as OM C-1, where line confusion is rampant. At the 692 GHz transition frequency, any heavy molecule would need to be in a high J state, but the 10 K excitation temperature m eans that any such state cannot be occupied. Of course, CO (6 ! 5) is observable, but only at  $T_a = 2.3$  K.NoU-lines have been observed in this source. The identi cation rests on the comparison of the deduced  $V_{LSR}$  for  $D_2H^+$  with that for other deuterated molecules, and also the line width and line strength. Note that the quoted uncertainty for the line frequency is about 0.16 km s<sup>1</sup> for  $H_2D^+$  and 0.008 km s<sup>1</sup> for  $D_2H^+$ . The very slight di erence in the observed  $V_{lsr}$  for these two lines (  $0.2 \text{ km s}^1$ ) could be due, in part, to the uncertainty in the line frequencies, particularly  $H_2D^+$ . A lso the accuracy of the astronom icalm easurem ent is limited by the resolution of the acousto-optic spectrom eter at about 0.1 km s<sup>1</sup>.

### 3. D iscussion

From the observed line strengths, given in column 3 of Table 1, we estimate the  $H_2D^+$ and  $D_2H^+$  column densities (see Table 2) for an excitation temperature  $T_{ex}$  of 10 K, assuming a 25% calibration uncertainty (3). The column density is given by:

$$N_{tot} = \frac{8^{-3} Q(T_{ex})}{c^{3} g_{u} A_{u1}} \frac{e^{E_{u} = T_{ex}}}{e^{h_{ex} - T_{ex}}} \frac{Z}{1} dv$$
(4)

where Q (T $_{\rm ex}$ ) is the partition function, A sum ing LTE conditions, we can estimate the optical depth from the observed line intensity:

$$T_{m b} = [J (T_{ex}) \ J (T_{bg})](1 \ e)$$
 (5)

where J (T) =  $(h = k)=(e^{h = kT})$  1) is the radiation temperature of a blackbody at a temperature T, and T<sub>bg</sub> is the cosm ic background temperature of 2.7 K. In the case of the H<sub>2</sub>D<sup>+</sup> transition,  $g_u = 9$ ,  $A_{u1} = 1.04 \ 10^4 \ s^1$ ,  $E_{u1} = 17.9 \ K$ ; in the case of the D<sub>2</sub>H<sup>+</sup> transition,  $g_u = 9$ ,  $A_{u1} = 4.55 \ 10^4 \ s^1$ ,  $E_{u1} = 33.2 \ K$ . The derived column densities depend on the assumed value of the excitation temperature. U sing N H<sub>3</sub>, M izuno et al. (1990) estimate the gas temperature to be 12 K. U sing D<sub>2</sub>CO line ratios, Loinard et al. (2001) obtained a rotational temperature of 8–10 K. Thus we quote, in Table 2, the values obtained for an excitation temperature of 10 K. Figure 4 presents the evolution of the ortho-H<sub>2</sub>D<sup>+</sup> and para-D<sub>2</sub>H<sup>+</sup> column densities as well as the para-D<sub>2</sub>H<sup>+</sup>/ortho-H<sub>2</sub>D<sup>+</sup> ratio, as a function of

tem perature between 9 and 15 K. Figure 4 and Table 2 represent the case where the source em ission is extended compared to the beam size. If the source em ission is comparable to or sm aller than the CSO beam size, the para- $D_2H^+$  /ortho- $H_2D^+$  ratio is then increased by a factor of 1.5 at the average excitation tem perature of 10 K. At therm al equilibrium, the ortho to para (respectively para to ortho) concentration ratio for  $H_2D^+$  (respectively  $D_2H^+$ ) exp(-86.4/T) (respectively 9/6  $\exp(-502/T)$ , so that at 10 K, this ratio is equal to 9 10<sup>3</sup> (respectively 10<sup>2</sup>). However, taking into account the limited rates would be 2 of the spin allowed collisions with H<sub>2</sub>, it is found that at these low tem peratures, the ortho to para  $H_2D^+$  concentration ratio is close to unity (G erlich, H erbst & R oue 2002). The para to ortho  $D_2H^+$  ratio is estimated by W alm sley, F lower & P ineau des Fôrets (2004) to be about one, for the same conditions. The para- $D_2H^+$  /ortho- $H_2D^+$  ratio presented in Figure 4 should then approximately represent the actual  $D_2H^+/H_2D^+$  ratio.

The 1.3 mm dust continuum strength (see Lis et al. 2002) is  $0.3 \text{ Jy in a } 11^{00} \text{ beam}$ and  $1.3 \text{ Jy in a } 20^{00} \text{ beam corresponding to the angular resolution of the } H_2D^+ \text{ and } D_2H^+$ data. A ssum ing a dust temperature of 12 K and a mass opacity coecient of 0.005 cm<sup>-2</sup> g<sup>-1</sup> (appropriate for pre-stellar cores), we derive an H<sub>2</sub> column density of  $5 \cdot 10^{23} \text{ cm}^{-2}$ . We then derive the  $H_2D^+$  and  $D_2H^+$  abundances to range between  $10^{-10}$  (at 10 K) and  $10^{-11}$  (at 15 K) compatible with abundances found by Roberts, Herbst & Millar (2003) for a cloud at 10 K and n (H<sub>2</sub>)=  $3 \cdot 10^6 \text{ cm}^{-3}$ .

The main result of this work is the detection of  $D_2H^+$ , with an abundance comparable to that of  $H_2D^+$ . This is a rem arkable veri cation of recent theoretical predictions, aim ed at explaining the large deuteration ratios observed in low mass pre-stellar cores and protostars. Two models, Roberts, Herbst & Millar (2003) and Walmsley, Flower & Pineau des Fôrets (2004), have recently considered the e ect of including all possible deuterated isotopom ers of H<sub>3</sub><sup>+</sup> in the chem ical networks, as suggested by Phillips & Vastel (2002). Roberts, Herbst & M illar (2003) studied the tem poral evolution of a cold and dense cloud, and found that at late times ( 10<sup>4</sup> yr), when CO is severely depleted in the gas phase (m ore than a factor of 1000), the  $D_2H^+/H_2D^+$  ratio reaches unity. W alm sley, F lower & P ineau des Fôrets (2004) studied the evolution of gas, depleted in CO, as a function of gas density and grain size distribution. For densities larger than  $10^6$  cm  $^3$ , they also found that D $_3^+$  can be the most abundant ion and that the  $D_2H^+/H_2D^+$  ratio reaches unity (see their gure 2). Both models need an extrem e CO depletion to account for such a ratio. As discussed in the Introduction, the measured CO depletion in 16293E is a factor 7 (Liset al. 2002), rather than the extrem e CO depletion needed by the models. However, the CO was measured in a large region (31") compared with that probed by the present observations:  $20^{0}$  for the H<sub>2</sub>D<sup>+</sup> and  $11^{0}$  for the  $D_2H^+$  observations. A lso, regions containing CO will not contain much  $H_2D^+$  and  $D_2H^+$ and vice-versa, so the inevitable inhom ogeneities in the region inhibit a clear result.

In sum mary, after some years of inconclusive results for theoretical models in understanding the observed high deuteration ratios of doubly and triply deuterated molecules, the present observation seems to suggest that the basic process is now almost completely understood: the large deuteration is due to extrem e CO depletion, and the factor that was previously missing in the models is the multiply deuterated forms of  $H_3^+$ . This is quite an achievement, and one remaining step will be to verify that the last prediction, a signi cant abundance of  $D_3^+$ , is also fulled. At present, the lines detected here are the only ones available for  $H_2D^+$  and  $D_2H^+$ . Knowledge of the ortho to para ratios and abundances of  $H_2D^+$  and  $D_2H^+$  would be considerably in proved if the ground state transitions of para- $H_2D^+$  (at 1370.15 G H z) and ortho- $D_2H^+$  (at 1476.60 G H z) were available. These lines could be detected from space telescopes such as Herschel with the Heterodyne Instrument for the Far-Infrared and also possibly from the stratospheric observatory SOF IA. However,  $D_3^+$ , like  $H_3^+$  has no perminanent dipole moment. Therefore this molecule probably can only be detected in absorption in the near-infrared.

The authors are grateful to John Pearson for providing an earlier spectroscopic com putation for  $D_2H^+$  and to Cecilia Ceccarelli for discussion. This research has been supported by NSF grant AST-9980846 to the CSO.

#### REFERENCES

- Baomann, A., Le och, B., Ceccarelli, C., Castets, A., et al., 2002, A&A 389, 6
- Bogey, M., Demuynck, C., Denis, M., Destombes, J.L., Lemoine, L., 1984, A&A 137, L15
- Boreiko, R.T., Betz, A.L., 1993, ApJ 405, L39
- Caselli, P., van der Tak, F., Ceccarelli, C., Bacmann, A., 2003, A&A 403, L37
- Caselli, P., et al., in preparation
- Ceccarelli, C., Castets, A., Loinard, L., Caux, E., Tielens, A.G. G. M., 1998, A&A 338, 43
- Chamley, S.B., 1997, MNRAS 291, 455
- Gerlich, D., Herbst, E., Roue, E., 2002, P&SS 50, 1275
- Hirao, T., Am ano, T, 2003, ApJ 597, 85
- Lis, D.C., Roue, E., Gerin, M., et al., 2002, ApJ571, L55

- Loinard, L., Castets, A., Ceccarelli, C., Caux, E., Tielens, A., 2001, ApJ 552, L163
- M izuno, A., Fukui, Y., Iwata, T., Nozawa, S., Takano, T., 1990, ApJ 356, 184
- Pagani, L., Wannier, P., Frerking, M., et al. 1992, A&A 258, 472
- Parise, B., Ceccarelli, C., Tielens, A.G.G.M., Hersbt, E., et al., 2002, A&A 393, L49
- Parise, B., Castets, A., Herbst, E., Caux, E., et al., 2004, A&A 416, 159
- Phillips, T.G. and Vastel, C., 2002, in Chemistry as a Diagnostic of Star Formation, ed.C.L. Curry & M.Fish, astro-ph/0211610
- Phillips, T.G., Blake, G., Keene, J., Woods, R., Churchwell, E., 1985, ApJ 294, L45
- Ramanlal, J., Polyansky, O L., Tennyson, J., 2003, A & A 406, 383
- Roberts, H., Herbst, E., Millar, T.G., 2003, ApJ 591, 41
- Roberts, H. & Millar, T. J., 2000a, A & A 361, 388
- Roberts, H. & Millar, T. J., 2000b, A&A 364, 780
- Roue, E., Tine, S., Coudert, L., Pineau des Fôrets, G., et al., 2000, A&A 354, 63
- Stark, R., van der Tak, F., van Dishoeck, E., 1999, ApJ 521, L67
- Stark, R., Sandell, G., Beck, S., Hogerheijde, M., et al., 2004, astro-ph/0402604
- Tielens, A.G.G.M., 1983, ApJ 119, 177
- van der Tak, F., Schike, P., Muller, H., Lis, D., et al., 2002, A&A 388, L53
- van Dishoeck, E., Phillips, T.G., Keene, J., Blake, G., 1992, A&A 261, L13
- Vastel, C., Phillips, T.G., Ceccarelli, C., Pearson, J., 2003, ApJ 593, L97
- Vastel, C., Phillips, T.G., et al., in preparation
- W alm sley, M ., Flower, D R ., Pineau des Fôrets, G ., 2004, accepted in A & A
- W ootten, A., Loren, R., 1984, ApJ 317, 220

This preprint was prepared with the AAS  $\square T_{E}X$  m acros v5.0.



Fig. 1. Diagram of the lowest energy levels of the  $D_2H^+$  molecule.







Fig. 3. Spectra of the ortho- $H_2D^+$   $1_{10}-1_{11}$  and para- $D_2H^+$   $1_{10}-1_{01}$  transitions towards 16293E.



Fig. 4. Column densities of the ortho- $H_2D^+$   $1_{10}-1_{11}$  (solid line) and para- $D_2H^+$   $1_{10}-1_{01}$  (dashed line) on the left side of the graph, and the para- $D_2H^+$ /ortho- $H_2D^+$  ratio (dotted line) on the right side, in the case where the source emission is extended compared to the beam size (see text).

Table 1. Results of Gaussian ts to the H  $_2D^+$  and D  $_2H^+$  spectra.

Line		T <sub>a</sub>		v	V L	SR
	(GHz)	(K)	(km s <sup>1</sup> )		(km s $^1$ )	
H <sub>2</sub> D <sup>+</sup> (1 <sub>10</sub> -1 <sub>11</sub> )	372 <b>.</b> 42134 (20) <sup>a</sup>	1.31	0.36	0.04	3.55	0.02
D <sub>2</sub> H <sup>+</sup> (1 <sub>10</sub> -1 <sub>01</sub> )	691 <b>.</b> 660440 (19) <sup>b</sup>	0.34	0.29	0.07	3.76	0.03

 $^{\mathrm{a}}\mathrm{M}$  easured frequency by Bogey et al. (1984).

 $^{\rm b}{\rm M}$  easured frequency by H irao and Am ano (2003).

Table 2. Ortho- $H_2D^+$  and para- $D_2H^+$  column densities at  $T_{ex} = 10 \text{ K}$ , in the case where the source emission is extended compared to the beam size (see text).

	T <sub>ex</sub> = 10 K				
	N (10 $^{13}$ cm $^2$ )				
o-H <sub>2</sub> D <sup>+</sup>	0.74	1.73	0.43		
р-Д <sub>2</sub> Н <sup>+</sup>	0.61	1.29	0.32		
$pD_2\text{H}^+/\text{o}H_2\text{D}^+$	0.75 0.37				