## IRAS 05436{0007 and the Emergence of M cN eil's N ebula

Bo Reipurth<sup>1</sup>

Institute for A stronom y, University of Hawaii, 2680 W oodlawn D rive, Honolulu, H I 96822

reipurth@ifa.hawaii.edu

and

Colin Aspin<sup>2</sup>

Gem iniObservatory, 670 North A bhoku P lace, University Park, Hib, HI 96720

caa@gemini.edu

#### ABSTRACT

We present a study of M cN eil's N ebula, a new ly appeared re ection nebula in the L1630 cloud, together with photom etry and spectroscopy of its source. New IR photom etry compared to earlier 2M ASS data shows that the star has brightened by about 3 m agnitudes in the near-infrared, changing its location in a J-H /H -K <sup>0</sup> diagram precisely along a reddening vector. A G em iniN IR IK-band spectrum shows strong CO-bandhead em ission and Br is in em ission, indicative of strong accretion. A G em iniG M O S optical spectrum shows only a red, heavily veiled continuum, with H strongly in em ission and displaying a pronounced P C ygnipro le, with an absorption trough reaching velocities up to 600 km s<sup>1</sup>. This im plies signi cant m ass loss in a powerful wind. However, no evidence is found for any shocks, as commonly seen in collim ated out ow s from young stars. Apparently the eruption has dispersed a layer of extinction and this, together with the intrinsic brightening of the IRAS source, has allowed an earlier out ow cavity to be ooded with light, thus creating M dN eil's N ebula.

Subject headings: stars: formation | stars: pre{main-sequence | Re ection nebulae | ISM : individual(M dN eil's N ebula) | stars: variables: other

## 1. Introduction

W ide-eld photographic and CCD in ages have docum ented the presence of num erous small compact relection nebulae in star-forming dark clouds. Such nebulae may possibly represent a transitional stage in which a young star goes from being an embedded infrared source to a visible T Tauri star or Herbig Ae/Be star. This process is widely believed to involve powerful bipolarout ow activity, as re ected in them orphology of the nebulae, which are often com etary, with the partially embedded star illum inating an out ow cavity. W hen observed over longer time-scales, such small rejection nebulae are often found to vary in brightness and occasionally in illum ination pattern. Perhaps the earliest known example is NGC 2261, which re ects light from the young starR M on and was found to vary by Hubble (1916). More recently, Cohen et al. (1981) demonstrated the variability of the fan-shaped nebula em anating from PV Cephei. At som e stage, the out ow activity of a star for the rst time punches a hole in its surroundings, suddenly allowing a beam of light to sweep across the surroundings. Subsequent eruptions in the source will again ood the out ow channel with light. The emergence of the nebula Re 50 in the Lynds 1641 cloud near the embedded source IRAS 05380 {0728 was docum ented by Rejourth & Bally (1986). Another case may have been the nebula IC 430 in L1641, which at present is quite sm all, but at the end of the 19th century appears to have been a very large and much brighter object (Pickering 1890; Strom & Strom 1993).

# 2. Observations

The observations presented below were taken on the "Frederick C.G illett" G em iniN orth Telescope<sup>1</sup>, on M auna K ea, H aw aiion UT dates February 03 (near-IR) and 14 (optical), 2004. The near-IR imaging and spectroscopy was acquired with the facility imager/spectrom eter, N IR I, using the f/6 cam era giving an image scale on the A laddin InSb 1024x1024 detector of 0.116"/pixel. Images in J, H and K<sup>0</sup> were obtained in total on-source integration times of 3.6 seconds (0.18 seconds, 10 coadds, two spatial positions) in each liter. The data were calibrated using similar observations of the UK IRT faint standard stars FS113, 119 and 135. The spectroscopic data were acquired using a 0.5" wide long-slit and K-band grism with a

<sup>&</sup>lt;sup>1</sup>Based on observations obtained at the G em ini O bservatory, which is operated by the A ssociation of U niversities for R essarch in A stronom y, Inc., under a cooperative agreem ent with the NSF on behalf of the G em inipartnership: the N ational Science Foundation (U nited States), the Particle Physics and A stronom y Research Council (U nited Kingdom), the N ational Research Council (C anada), CONICYT (Chile), the A ustralian Research C ouncil (A ustralia), CNPq (B razil), and CONICET (A rgentina).

total on-source integration time of 60 seconds (30 seconds, 1 coadd, two spatial positions) resulting in a spectrum with resolving power 780 covering the wavelength region 2.05-2.45 m icrons. Telluric feature correction was performed using observations of the A0 V star H IP 28056. The optical data were acquired using the facility optical multi-object spectrom eter, GM OS-N. Im ages in the GM OS Sloan g', r', i' and z' liters were obtained with exposure times of 60 seconds and 10 seconds, the shorter exposure time being used to give unsaturated in ages of the young star. The GM OS spectrum was a 5 m in exposure obtained using the R 831 grating, 0.5" wide long-slit, and a central wavelength of 5800 A resulting in a spectrum covering the wavelength range 4800-6800 A at a resolving power of R 4500. Infrared im ages were obtained of the eruptive star at the UH 2.2m telescope through an R-band liter (5 m in exposure) on Feb 1, 2004 and through a narrow band H<sub>2</sub> 2.12 m liter (20 m in exposure) on Feb 14, 2004 UT.

## 3. Pre-outburst Observations

0 n January 23, 2004, M cN eil (2004)<sup>2</sup> discovered the presence of a bright nebula m ore than an arm inute in extent south-west of M 78 in the L1630 cloud, a nebula that was not seen on any of the blue or red POSS-I, POSS-II and UKSTU sky at las plates taken between 1951 and 1991. M cN eil's Nebula is clearly com etary-shaped, with a highly reddened, dustobscured star at its apex. Subsequent exam ination of recent im ages obtained in the fall of 2003 suggests that M dN eil's N ebula rst appeared in late N ovember 2003 (B ricero et al. 2004; M cN eil, priv. comm.). However, Mr. John W elch of Phoenix, Arizona, kindly brought our attention to a photograph of the M 78 region obtained by E.K reimer on Oct 22, 1966 on which M dN eil's Nebula is seen at virtually the sam e brightness as at discovery (M allas & K reim er 1978). This o ers the important insight that the current eruption of the illum inating star is not the rst one. We here sum marize what is known about this star prior to the appearance of the nebula. The star lies within the error ellipse  $(45^{\circ} 7^{\circ})$  at PA 88°) of the IRAS source 05436{0007, which is detected only at 12 and 25 m with uxes of 0.53 and 1.19 Jy, respectively, with a ag that indicates an 80% probability that the uxes are variable. The star also coincides with the 2M ASS source 05461313{0006048, which is located at  $_{2000}$ :  $05^{h}46^{m}13.1^{s}$ , and  $_{2000}$ : { $00^{\circ}06^{0}05^{\circ\circ}$ . The 2M ASS photom etry is listed in Table 1. A faint optical counterpart to this 2M ASS source is visible on an I-band in age

 $<sup>^{2}</sup>$  This reference is listed as M cN eil, Reipurth, & M eech (2004) in the A strophysics D ata System. The second and third authors only provided a simple con m atory CCD in age, and full credit for the important discovery of the eruption in IRAS 05436{0007 should go solely to M r. M cN eil. Consequently, we suggest that future reference to the discovery announcement in this IAU C incular be given as M cN eil (2004).

obtained by E isb el & M undt (1997). In the sub-m illimeter, the source was detected as a compact, isolated dust continuum source at 350 and 1300 m by Lis, M enten, & Zylka (1999), their source LM Z 12, a designation we w ill use for the source in the following until a variable star name is assigned in the GCVS. Subsequently, the same source was detected at 850 m by M itchellet al. (2001) and Johnstone et al. (2001), their source 0 riB sm m 55. Lis et al. (1999) suggested a pre-outburst bolom etric luminosity for LM Z 12 of 2.7 L . These sub-m illimeter observations additionally detected extended, di use em ission about 40 arcsec north of LM Z 12, coinciding with part of M cN eil's N ebula. Limited m illimeter observations by Lis et al. (1999) did not reveal any m olecular out ow from the source, and only a sm all core in  $\rm HCO^+$ .

## 4. Observational Results

We present in Fig. 1 a color in age of M cN eil's Nebula based on g', r', and i' in ages taken with GMOS at the G em ini-N telescope in  $0.5^{\circ}$  seeing. The nebula has approximate dimensions of 30 60 arcsec, and shows considerable structure, with a very bright patch of illumination near the source. Photom etry of LMZ 12 at its apex is given in Table 1. Note that because of the bright nebulosity, the stellar magnitudes are very sensitive to the aperture-size employed, we used a small aperture of  $0.9^{\circ}$  radius.

The nebula encompasses the Herbig-Haro object HH 22 (Herbig 1974). However, the detection of a collimated jet associated with HH 22 emanating from a source further to the East suggested to E islo el & M undt (1997) that there is no connection between this HH object and LM Z 12. A nother HH object, HH 23, is located further to the N orth. It consists of two knots on an axis that passes close to LM Z 12, and E islo el & M undt (1997) suggested that the faint optical source coincident with LM Z 12 is the driving source. M dN eil's N ebula opens up in the approximate direction towards HH 23, and we suggest that it represents the illuminated out ow cavity carved at some time in the past by LM Z 12 into the L1630 cloud. P roper motion measurements of HH 23 are needed to establish a mer link with LM Z 12.

Infrared J, H, K<sup>0</sup> photom etry of the source obtained on Feb 03, 2004 is listed in Table 1. It is evident that the source is much brighter now than when the 2M ASS data were obtained on 0 ct 07, 1998. At the time of our observations, LM Z 12 had brightened by J=3.6, H=3.2, and K  $^{0}=2.9$  m agnitudes. W hen plotted in a J-H/H-K  $^{0}$  diagram, it is evident that it displays a substantial infrared excess (Fig. 2). It is noteworthy that in its current high state, LM Z 12 shows considerably less reddening than at the time of the 2M ASS observation in 1998.

In Fig. 3 we show a contour diagram of LM Z 12 as seen in our K  $^{0}$ -band images. It is immediately obvious that the star is surrounded by a compact rejection nebulosity and that, in particular, this nebula shows a curved tail characteristic of many stars undergoing highaccretion events (e.g., Herbig 1977). In Fig. 4, we show a K-band spectrum of LM Z 12, which displays a red continuum with strong CO-bandhead emission, and the Br and Na I lines are in emission. Our optical spectrum of LM Z 12 shows a red continuum with a prominent H line in emission, but no other emission or absorption lines, suggesting the presence of heavy veiling. The H line, shown in Fig. 5, has an equivalent width of 32 A and displays a characteristic P Cygnipro le. The absorption component has an equivalent width of 5.6 A.

### 5. D iscussion

The two major classes of eruptive variables among pre-main sequence stars are the FU ors and the EX ors (Herbig 1966, 1977, 1989). In the following we discuss whether LM Z 12 can be linked to one of these two categories.

FU ors are characterized by large-am plitude (V 5-6 m ag.) brightenings lasting several or m any decades, and in the optical they show F-G type spectra without emission lines. In the K-band region, FU ors display deep CO-bandhead absorption (e.g., Reipurth & A spin 1997). A lthough LM Z 12 has about the right am plitude, it remains to be seen how long it stays bright. Spectroscopically, however, LM Z 12 looks very di erent from m ature FU ors both in the optical and in the infrared. The prominent H emission and the strong CO emission that are presently seen do not suggest a classication as a FU or as we currently understand the phenom enon. On the other hand, Bricero et al. (2004) emphasize that the earliest optical spectrum after outburst of the FU or V1057 Cyg had some similarities to that of LM Z 12. Further photom etric and spectroscopic monitoring of the star is required to settle this.

EX or eruptions, which m ay occur repeatedly in a given star, have am plitudes that can be comparable to those of FU ors, but they have durations which are m uch shorter, from a few m onths to a few years. Spectroscopic studies of EX ors are limited, because of the rarity and shorter durations of these eruptions. The few such studies that have been m ade (e.g., Lehm ann, Reipurth, & Brandner 1995; Herbig et al. 2001; Parsam ian et al. 2002) all show that EX ors in eruption have at least the lower Balm er lines in emission, together with emission lines of He I and other lines characteristic of the m ost active T Tauri stars (Herbig 1962). A high-resolution spectrum of the H line of SVS 13 in outburst (E islo el et al. 1991) shows a P Cygnipro le rather sim ilar to the one seen in Fig. 5. How ever, we do not see any other emission lines than H in the spectral range observed (4800-6800 A) in LM Z 12. Infrared spectra of EX ors are even more limited, one case is the eruption of SVS 13, for which a K-band spectrum shows the CO bandheads strongly in emission as well as Br in emission (Carr & Tokunaga 1992). However, a similar spectrum of EX Lup shows Br in emission, but the CO bandheads and Na I in absorption (Herbig et al. 2001). It thus does not seem that EX ors have a unique infrared spectral signature. Based on a comparison with these som ewhat limited observations of EX ors, we conclude that LM Z 12 does have a certain resem blance to EX ors.

The spectroscopic data give us some insight into the physical processes behind the observed eruption. The detection of Br in emission testi es to a region of hot gas. Br emission has been shown to correlate tightly with accretion luminosity, so its presence indicates that the eruption is linked to an episode of accretion (N a jita et al. 1996; M uzerolle et al. 1998). The pronounced P Cygnipro le seen at H in the optical spectrum is likely to be form ed in a strong wind that has su cient optical depth to produce the deep blueshifted absorption trough (e.g., M uzerolle et al. 2001). Only few T Tauri stars show such well-developed P Cygnipro les at H . The well-de ned blue edge of the absorption trough indicates of up to 600 km s<sup>1</sup>, even m ore than the extrem e wind that em – anates from FU O rionis (e.g., Herbig et al. 2003).

It is noteworthy that we see no spectral features indicative of shocks: the infrared spectrum shows no evidence for  $H_2$  emission, and the optical spectrum does not display the [SII] 6717/6731 lines characteristic of Herbig-Haro jets. Our  $H_2$  2.122 m image of LM Z 12 shows no presence of any extended jet ow. It thus appears that LM Z 12, in common with most FU ors and EX ors, has powerfulm ass loss, but apparently not in the well-collim ated fashion that enables shocked HH jets.

O ne nal insight into the LM Z 12 EX or event comes from the changes of the infrared colors from before to after the eruption. The pre-and post-outburst colors show that LM Z 12 has moved precisely along a reddening vector, indicating that the star has brightened partly because its visual extinction diminished by about 4.5 m agnitudes. U sing the reddening curve of Rieke & Lebofsky (1985), we nd that this corresponds to  $A_J = 1.26$ ,  $A_H = 0.81$ , and  $A_K = 0.50$ . Since we know that the infrared colors changed by J = 3.64, H = 3.16, and K = 2.87, we see that the intrinsic brightening of the star corrected for the change in extinction is 2.4 m agnitudes in each of the J, H, and K Iters. Through a combination of this intrinsic brightening and the clearing away of an obscuring layer, the star has been able to illum inate its out ow channel, thus creating M dN eil's N ebula.

W e are grateful to Nuria Calvet for helpful comments, M ike Connelley for taking an infrared  $H_2$  in age, George Herbig for illuminating discussions, Inger Jorgensen and Kathy

Roth for a GMOS spectrum, Jay M dN eil for alerting us to the outburst, K aren M eech and Yan Fernandez for a CCD im age, Peter M ichaud and K irk Pu'uohau-Pum m ill for preparing the color-im age, Jean-R ene R oy for approving all the G em ini 8m ToO observations, B rian Ski for inform ation about both recent and pre-discovery im ages, John W elch for inform ation about the 1966 outburst, and an anonym ous referee for helpful com m ents.

#### REFERENCES

- Briceno, C. et al. 2004, ApJ Letters, in prep.
- Carr, J.S., & Tokunaga, A.T. 1992, ApJ, 393, L67
- Cohen, M., Kuhi, L.V., Harlan, E.A., Spinrad, H. 1981, ApJ, 245, 920
- E islo el, J., & M undt, R. 1997, AJ, 114, 280
- E isb el, J. et al. 1991, ApJ, 383, L19
- Herbig, G.H. 1962, Adv. Astr. Astrophys., 1, 47
- Herbig, G.H. 1966, Vistas in Astronomy, 8, 109
- Herbig, G.H. 1974, Lick Obs. Bull. No. 658
- Herbig, G.H. 1977, ApJ, 217, 693
- Herbig, G.H. 1989, in ESO W orkshop on Low Mass Star Formation and Pre-Main Sequence Objects, ed.B.Reipurth, p. 233
- Herbig, G.H., Aspin, C., Gilmore, A.C., Imho, C.L., Jones, A.F. 2001, PASP, 113, 1547
- Herbig, GH., Petrov, PP., & Duemmler, R. 2003, ApJ, 595, 384
- Hubble, E.P. 1916, ApJ, 44, 190
- Johnstone, D., Fich, M., Mitchell, G.F., Moriarty-Schieven, G. 2001, ApJ, 559, 307
- Lehm ann, T., Reipurth, B., Brandner, W. 1995, A&A, 300, L9
- Lis, D.C., Menten, K.M., & Zylka, R. 1999, ApJ, 527, 856
- Mallas, J.H., & Kreimer, E. 1978, The Messier Album, Sky Publishing Corporation
- M dN eil, J.W . 2004, IAUC 8284

- M itchell, G F., Johnstone, D., M oriarty-Schieven, G., Fich, M., Tothill, N F.H. 2001, ApJ, 556, 215
- M uzerolle, J., Calvet, N., & Hartmann, L. 1998, AJ, 116, 2965
- Muzerolle, J., Calvet, N., & Hartmann, L. 2001, ApJ, 550, 944
- Najita, J., Carr, J.S., & Tokunaga, A.T. 1996, ApJ, 456, 292
- Parsamian, E.S., Mujica, R., & Corral, L. 2002, A strophysics, 45, 393
- Pickering, E.C. 1890, Ann. Harvard College Obs., 18, 113
- Reipurth, B., & Aspin, C. 1997, AJ, 114, 2700
- Reipurth, B., & Bally, J. 1986, Nature, 320, 336
- Rieke, G.H., & Lebofsky, M.J. 1985, ApJ, 288, 618
- Strom, K.M., & Strom, S.E. 1993, ApJ, 412, L63

This preprint was prepared with the AAS  ${\mathbb I}{\mathbb P} T_E X$  m acros v5.0.

D ate	g <b>′</b>	r <b>′</b>	i'	J	Н	K <sup>0</sup>	Telescope
0 ct 7, 1998				14.74 .03	12.16 .03	10.27 .02	2M ASS
Feb 03 <b>,</b> 2004 <sup>a</sup>				11.1 .1	9.0 .1	7.4 .1	Gemini
Feb 14 <b>,</b> 2004 <sup>b</sup>	22.8	17.4	15.6				Gemini

Table 1. Photom etry of IRAS 05436{0007

<sup>a</sup>W e list large and conservative error estim ates for the G em ini IR photom etry, since this object is very bright for an 8m telescope, requiring very brief integrations.

 $^{b}$ GMOS photom etry is obtained in an aperture with radius 0.9<sup>00</sup> in order to avoid bright nearby nebulosity. It is quoted to one decimal place only since we use standard zeropoints from the GeminiGMOS web site.



Fig. 1. A color-in age of M dN eil's N ebula obtained by combining broadband g', r', and i' in ages obtained with G M O S at the G em ini-N 8m telescope. LM Z 12 is at the bottom of the nebula, and H H 22 is the bluish curved object on the northeastern edge of M dN eil's N ebula. The height of the gure is about 80 arcsec. N orth is up and East is left.



Fig. 2. | A J-H/H-K<sup>0</sup> diagram showing the location of LM Z 12 as observed with 2M ASS [1] on 0 ct 7, 1998 and with G em in i-N [2] on Feb 3, 2004. The dashed line is the T Tauri locus, and the solid straight lines and dot-dashed line are reddening vectors of  $A_V = 15$  m ag.



Fig. 3. A compact nebula is seen around the illum inating star of M dN eil's N ebula in this  $K^{0}$ -band in age obtained with N IR I at the G em in i-N 8m telescope.



Fig. 4. | A K-band spectrum of LM Z 12 obtained with N  ${\rm IR}$  I at the G em in i-N 8m telescope.



Fig. 5. | The H line in LMZ 12 shows a pronounced P Cygnipro le. The rest velocity of the star is indicated.