

Why a Single-Star Model Cannot Explain the Bipolar Nebula of Eta Carinae

Noam Soker

ABSTRACT

I examine the angular momentum evolution during the 1837–1856 Great Eruption of the massive star η Carinae. I find that the new estimate of the mass blown during that eruption implies that the envelope of η Car substantially spun-down during the 20 years eruption. Single-star models, most of which require the envelope to rotate close to the break-up velocity, cannot account for the bipolar nebula the Homunculus formed from matter expelled in that eruption. The kinetic energy and momentum of the Homunculus further constrains single-star models. I discuss how η Car can fit into a unified model for the formation of bipolar lobes where two oppositely ejected jets inflate two lobes (or bubbles). These jets are blown by an accretion disk, which requires stellar companions in the case of bipolar nebulae around stellar objects.

Key words: binaries: close circumstellar matter stars: individual: η Carinae stars: mass loss stars: winds

1. Introduction

The bipolar structure of the Eta Carinae (η Car) nebula the Homunculus is not unique. Its basic structure, that of bubble pair, where each bubble is bounded by a thin dense shell, filled with a low density interior, and having a narrow waist between them, is seen in many diverse objects. Examples include the Perseus cluster of galaxies (Fabian et al. 2002), the symbiotic nebula He 2-104 (Corradi & Schwarz 1995; Corradi et al. 2001; here the dense shell is not closed), and the planetary nebula (PN) NGC 3587 (PN G 148.4+57.0; e.g., Guerrero et al. 2003). The similarity between bubble pairs in clusters of galaxies and

¹Department of Physics, Technion – Israel Institute of Technology, Haifa 32000 Israel, and Department of Physics, Orenan, Israel; soker@physics.technion.ac.il

in PN s is discussed in Soker (2003a,b; 2004a), while in Soker (2004b) I discuss the sim ilarity of these system s w ith symbiotic nebulae and the bipolar nebula of Car. In clusters of galaxies such bubbles are know n to be form ed by oppositely ejected jets, which are detected by radio em ission (e.g., Hydra A ; M cN am ara et al. 2000). In a series of papers I argue that this suggests that bubble pairs in PN s (Soker 2003a,b; 2004a) and symbiotic nebulae (Soker 2004b) are also form ed by such jets (note that not all PN s are shaped by jets). In partic- ular, there are m ore and m ore observations show ing and hinting at the presence of jets in symbiotic system s (e.g., Kellogg, Pedelty, & Lyon 2001; Brocksopp et al. 2004), in PN s, and PN progenitors (e.g., Imai et al. 2002; Hirano et al. 2004; Sahai et al. 2003).

In previous papers I attribute the bipolar structure of Car (e.g., Ishibashi et al. 2003) and the presently blown fast polar wind found by Smith et al. (2003a) to interaction w ith the binary companion (Soker 2001, 2003c). In particular, in Soker (2001) I proposed that the bipolar nebula was shaped by jets blown by the companion, via an accretion disk, during the 20 years Great Eruption a century and a half ago. In that paper I listed som e argum ents in favor of such a m odel. However, the new nding of Smith et al. (2003b) of a m ore m assive Homunculus, and several new papers using a single star m odel for the shaping of the wind and circum stellar m atter of Car (e.g., D warkadas & O wocki 2002; Smith et al. 2003a; Gonzalez et al. 2004; van Boekel et al. 2003) m otivate m e to reconsider the single star m odel. Other papers in recent years study the role of rotation and/or binary companion in Car. Stothers (1999) nd that the rotation does not a ect m uch the instability of lum inous blue variables (LBV), which is in support of the view I take in the present paper. Davidson (1999; see also Davidson 2000), on the other hand, argues that a binary m odel for Car is not "a Panacea". I accept Davidson's view that som e crucial aspects of the Car eruptions are due solely to the behavior of the m assive star, and not to the companion. My m ain interest is in show ing that the bipolar structure of the nebula is due to a binary companion. In section 2 I study the m om entum and energy budget of Car. In section 3 I consider the required stellar angular m om entum in single-star m odels. I w ill not follow the evolution of rotating m assive stars w ith any speci c m odel (for these see, e.g., Heger & Langer 2000; M aeder, & Meynet 2003; Meynet & M aeder 2003). I w ill use analytical approach. In section 4 I sum m arize m y nding that single-star m odels encounter severe problem s.

2. Wind's energy and m om entum

Som e of the argum ents given here are not new . However, I put them in a broader context in light of the new estimated m ass that was blown during the 20 years Great Eruption $M_{GE} \approx 12M_{\odot}$ (Smith et al. 2003b). For a wind driven by radiation pressure the effective

number of times a photon is scattered by the wind material, i.e., in the positive radial direction, is given by

$$n_s = \frac{M_w v_w}{L=c} = 700 \frac{L}{10^{7.3} L} \frac{M}{0.5 M} \frac{v_w}{600 \text{ km s}^{-1}} ; \quad (1)$$

where M_w is the mass loss rate into the wind, v_w is the terminal wind speed, L is the stellar luminosity, and c the light speed. The different variables are scaled with the mass loss rate (Smith et al. 2003b) and luminosity (Davidson & Humphreys 1997) during the Great Eruption. This is a very large number, which is not encountered in other stars. For example, in most PNs and highly evolved red giant stars $n_s < 1$ (Knappe 1986). The PN with the highest value of $n_s = 600$ in the list given by Knappe (1986) is NGC 2346. The central star of this PN contains a close binary companion (Bond 2000), which went through a common envelope evolution and released large amount of orbital energy. Therefore, the binary companion is behind the large value of n_s in the PN NGC 2346. Smith et al. (2003b) notice that such a high value of n_s occurs in explosions. However, the mass loss rate during the Great Eruptions lasted for 20 years, a time longer by more than an order of magnitude than the dynamical time of Car, even if it swelled to 10 AU. Shaviv (2001) compares the super-Eddington wind of the Great Eruption to that of novae and obtains satisfactory fit to the Great Eruption outflow, assuming spherical wind. Although he assumes a steady state mass loss process, the photospheric radius of the nova ejected mass expands by several orders of magnitude during the explosion, and the mass involved occupies a thin layer on the white dwarf surface prior to the explosion. The $10 M_{\odot}$ ejected mass in the Great Eruption, on the other hand, originated in a thick envelope layer of Car. In addition, the formation of the bipolar structure in an explosive event requires different models than the one I criticize in the present paper. The results of Shaviv (2001), though, show that a single star can lose mass at a high rate. Such a high mass loss rate is needed in the presently proposed binary model, but here the wind speed, hence the momentum and kinetic energy supplied by the primary, can be much lower.

The new estimated energy of the Homunculus is $E_{\text{Hom}} = 10^{49.6} = 10^{50}$ erg (Smith et al. 2003b). The ratio of this energy to the energy radiated during the 20 years Great Eruption is

$$\frac{E_{\text{Hom}}}{L \cdot 20 \text{ yr}} = 1 \frac{L}{10^{7.3} L} \frac{E_{\text{Hom}}}{5 \cdot 10^{49} \text{ erg}} \quad (2)$$

This implies that half the energy liberated during the Great Eruption was radiated, and half was converted to the kinetic energy of the wind. The present mass loss rate of $2 \cdot 10^{-3} M_{\odot} \text{ yr}^{-1}$ (van Boekelet al. 2003) over the 150 years span since the Great Eruption contributes negligible amount to this energy.

The required transfer efficiency of momentum and energy from radiation to the wind in single star models of Car is much higher than in any other object blowing similar winds. These values are more typical for binary systems, as I now discuss. In principle there are two processes by which a binary companion can account for such a high kinetic energy of the wind. First, the binary system can release orbital energy $E_{orb} \approx 0.5GM_1M_2/a_f$ where M_1 and M_2 are the two masses inward to the final orbital separation of a_f , and assuming that the final orbital separation is much smaller than the initial orbital separation. If a fraction η of the released orbital energy is deposited into the expelled mass, then the final orbital separation for a companion to explain the Homunculus energy is

$$a_f \approx 70 \frac{E_{Hom}}{5 \times 10^9 \text{ erg}}^{1/2} \frac{1}{0.5} \frac{M_1}{120M} \frac{M_2}{30M} R \quad (3)$$

The presently observed orbital period is 5.5 yr, which for primary and companion masses of $M_1 = 120M_\odot$ and $M_2 = 30M_\odot$, respectively, implies an average orbital separation of $a \approx 16.5 \text{ AU}$ (for observational support for the presence of a binary and its properties see, e.g., Damiani 1996; Ishibashi et al. 1999; Damiani et al. 2000; Corcoran et al. 2001a,b; Pittard & Corcoran 2002; Duncan & White 2003; Fernandez-Lajus et al. 2003). Therefore, the orbital energy of the presently observed binary system cannot account for the kinetic energy of the Homunculus. However, Car could have swallowed a closer, lower mass third star during its Great Eruption. For example, a third star with $M_3 = 5M_\odot$ could have spiral down to the core of Car, and released more than the required energy. The Lesser Eruption of 1890 makes this scenario unlikely, although it can't be ruled out based on pure physical arguments.

In the second process, the companion is outside the primary envelope, it accretes mass via an accretion disk and blows two jets (or a collimated fast wind: CFW). In this scenario, which was proposed for the formation of the Homunculus in Soker (2001; where more supporting arguments are given), the primary in Car expelled its mass at low speeds. In principle, the companion may accrete half of the mass expelled by the primary and blow a fraction of 0.2 at a speed equals to its escape velocity $v_j \approx 2000 \text{ km s}^{-1}$. For an accreted mass of $10M_\odot$ the energy carried in the jets is then $8 \times 10^9 \text{ erg}$, enough to account for the energy of the Homunculus. In addition, in such a scenario a substantial fraction of the energy radiated by the system during the great eruption came from the accretion energy onto the companion. Recently, Smith & Morse (2004) reported the discovery of extremely fast material, $v > 3200 \text{ km s}^{-1}$, in Car. They attribute this material to mass loss from the primary star. In the binary model, on the other hand, this extremely fast material was first accreted onto an accretion disk around the companion, and then was ejected at this high speed.

3. The angular momentum evolution of Car

3.1. Moment of Inertia

Following Soker & Harpaz (1999)² I examine the ratio of the density at the photosphere, ρ_p , to the average envelope density ρ_a . As shown below, this ratio indicates the moment of inertia, a relevant quantity for the slowing down process. The photospheric density is given by (Kippenhahn & Weigert 1990)

$$\rho_p = \frac{2}{3} \frac{m_H}{k_B} \frac{GM_1}{R^2 T}; \quad (4)$$

where m_H is the mean mass per particle, k_B is the Boltzmann constant, and κ is the opacity. Substituting typical values for Car in the last equation gives

$$\rho_p = 1.5 \cdot 10^{13} \frac{1}{1 \text{ cm}^2 \text{ g}^{-1}} \frac{T}{10^4 \text{ K}}^3 \frac{L_1}{10^7 L} \frac{M_1}{120M} \text{ g cm}^{-3}; \quad (5)$$

where a black body luminosity was assumed to eliminate the stellar radius. Using the (solar composition) opacity as given by Alexander & Ferguson 1994; see also Rogers & Iglesias 1992), I find the following approximate, but adequate, fitting in the relevant photospheric densities

$$\rho_p = \begin{cases} 10 (T=10^4 \text{ K})^{14} \text{ cm}^2 \text{ g}^{-1} & \text{for } 5;000\text{K} \leq T \leq 8;500\text{K} \\ 1 \text{ cm}^2 \text{ g}^{-1} & \text{for } 8500\text{K} \leq T \end{cases} \quad (6)$$

I take the envelope mass M_{env} to contain about half of the Car stellar mass. For LBV as Car, the envelope mass is expected to be lower, hence increasing the slowing down process and strengthening the effect studied here. The average density in the envelope is given by

$$\rho_a = \frac{M_{con}}{4 R^3} \approx 7 \cdot 10^8 \frac{T}{10^4 \text{ K}}^6 \frac{L}{10^7 L} \frac{M_{env}}{60M} \text{ g cm}^{-3}; \quad (7)$$

As was shown for AGB stars (Soker & Harpaz 1999), and can be checked for more massive stars, when ρ_p is not much smaller than ρ_a , the envelope density profile is shallow. In low mass envelopes, the density profile is almost flat in the outer region (Soker & Harpaz 1999). For massive stars this can be seen in the model of an initial 120M star which was reduced

²Note that the density scale in Figs. 1-5 of SH 99 is too low by a factor of 10; the correct scale is displayed in their Fig. 6.

to $66:6M_{\odot}$ as calculated by Stothers & Chin (1993). In that model (their fig. 2), the effective temperature is $T_p = 8000 \text{ K}$, and luminosity $2 \times 10^6 L_{\odot}$. The radius is $700R_{\odot}$. The envelope mass is very low, because of the mass lost by the star. On average, the density profile from $0.05R$ to the surface is $\rho \propto r^{-1}$. The parameters in the evolved $120M_{\odot}$ model of Mader (1981) without mass loss are $L \approx 10^6 L_{\odot}$, $T_p \approx 4000 \text{ K}$, $R \approx 2000R_{\odot}$, and $M_{\text{env}} = 30M_{\odot}$. From the equations above, and using the correct opacity for low densities of $\kappa \approx 3 \times 10^4 \text{ cm}^2 \text{ g}^{-1}$, $I_{\text{nd}} = 10^{10} \text{ g cm}^{-3}$ and $\rho_a = 5 \times 10^9 \text{ g cm}^{-3}$. The ratio of 50 is moderate, and the density profile in the outer envelope is $\rho_e \propto r^{-3}$.

The structure of the envelope considered above determines the moment of inertia of the star (that of the core is negligible for giants)

$$I = M_{\text{env}} R^2 \quad (8)$$

For an envelope density profile of $\rho_e \propto r^{-2}$ one finds $\beta = 2.9$, while for $\rho_e \propto r^{-3}$ the value is $\beta = [\ln(R=r_{\text{in}})]^{-1}$, where r_{in} is the inner radius of the envelope. In the model described above of Mader (1981), $R=r_{\text{in}} \approx 10$, and $I_{\text{nd}} \approx 0.15$. For steeper density profiles the value of β is lower. In the sun, for example, equations (5)–(7) give a very large ratio of $\log(\rho_a = \rho_p) \approx 5$. The density profile in most of the solar interior (beside the very outer envelope) can be fitted by

$$\rho(r) = \rho_0 \exp(-K r); \quad (9)$$

with $K R = 10.54$ (Bahcall, Pinsonneault, & Basu 2001). For such a profile, and with $r_{\text{in}} \approx R$, the moment of inertia coefficient is $\beta \approx 8 = (K R)^2$. For the sun this gives $\beta = 0.07$.

This subsection shows that the ratio $\rho_a = \rho_p$ can be used as a crude indicator for the moment of inertia. The exact value of β can't be predicted from this ratio, but for the present goal it is enough to use the crude relation

$$\begin{aligned} \beta &< 0.2; && \text{for } \log(\rho_a = \rho_p) < 2 \\ &: 0.1 - 0.2 && \text{for } 2 < \log(\rho_a = \rho_p) < 4 \\ &: > 0.1 && \text{for } \log(\rho_a = \rho_p) > 4 \end{aligned} \quad (10)$$

3.2. Angular Momentum Loss

Consider an envelope of a giant star rotating as a solid body. The angular momentum loss rate from the envelope to the wind is

$$J_{\text{wind}} = \dot{M} R^2 \omega; \quad (11)$$

where ω is the stellar angular velocity, J is the stellar angular momentum, and \dot{M} depends on the mass loss geometry: for a constant mass loss rate per unit area on the surface $\dot{M} = 2\pi R^2 \dot{\sigma}$,

while for an equatorial mass loss $\beta = 1$. Smith et al. (2003b) argue for an enhanced polar mass loss rate during the Great Eruption, for which $\beta < 2=3$. However, still some extra mass resides in the equatorial plane even according to Smith et al. (2003b). Therefore, I will take $\beta = 0.5 - 0.7$ for the Great Eruption mass loss geometry. The change of the envelope's angular momentum with mass loss is given by (e.g., Soker & Harpaz 2000)

$$\frac{d \ln J_{\text{env}}}{d \ln M_{\text{env}}} = \frac{d \ln \Omega}{d \ln M_{\text{env}}} + \frac{d \ln I}{d \ln M_{\text{env}}} = \frac{1}{(M_{\text{env}})} \quad ; \quad (12)$$

If the structure of the atmosphere does not change much while mass loss occurs then $d \ln I = d \ln M_{\text{env}} = 1$ and Ω is constant. The solution of the last equation becomes

$$\frac{\Omega}{\Omega_0} = \frac{M_{\text{env}}}{M_{\text{env}0}} \quad ; \quad (13)$$

3.3. Angular Velocity Evolution During the Great Eruption

Most single-star models I am aware of for the formation of the Homunculus require Car to rotate at $\Omega = \beta \Omega_{\text{Kep}} \approx 0.7$, where Ω is the angular velocity of the stellar envelope and Ω_{Kep} is the Keplerian velocity on the equator (i.e., the break-up angular velocity). Dwarkadas & Owocki (2002) and Smith et al. (2003a), for example, take $\beta = 0.9$, while M aeder & Desjacques (2001) take $\beta = 0.8 - 0.9$. The model presented by Langer, Garcia-Segura, & Mac Low (1999) is different in that they consider the ratio of luminosity to the Eddington limit. Similar considerations are qualitatively presented by Zethson et al. (1999) to account for the slow ejecta in the equatorial plane of Car. They emphasize that even slowly rotating luminous stars can have highly non-spherical mass loss geometry. They also point out that their model cannot explain why the slow ejecta are in dense compact condensations. In the binary model, these condensations are assumed to be formed by the influence of the companion. The model of Langer et al. (1999) which gives two lobes similar to those in Car have $\beta \approx 0.2$. However, most of the mass in their model is being lost at low velocities, $\approx 400 \text{ km s}^{-1}$, and it resides in the equatorial plane rather than in the Homunculus. Therefore, the total kinetic energy in their model is much below that of the Homunculus. Aerts, Lamers, & Molenberghs (2004) study the influence of rotation on the mass loss from Car, and also find the wind velocity to be too low. Therefore, in what follows I will refer only to the models require $\beta \approx 0.7$.

To facilitate an analytical treatment that will demonstrate the problems of a rotating single star, I consider two extreme cases: an eruption accompanied by expansion of the envelope to $R \approx 10 \text{ AU}$ (e.g., Davidson & Humphreys 1997), and an eruption leaving the

Car primary star a hot star. The angular momentum of Car at the beginning of the Great Eruption is determined by both single star evolution and binary interaction. To possess fast rotation, though, the star must have interacted with a stellar companion, because single stars slow down with mass loss (Maeder & Meynet 2000), e.g., large mass loss events before the Great Eruption, such as the one proposed by Bohigas et al. (2000), would have substantially slowed down Car. As a star like Car expands to $R \approx 10 \text{ AU}$, its moment of inertia increases both because of the increase in size and the increase in moment of the inertia coefficient (eq. 8). Presently, Car has a radius of $100R_{\odot}$ and an effective temperature of $20,000 - 30,000 \text{ K}$ (Hillier et al. 2001; Smith et al. 2003a). For such a model $\beta = 0.1$ by equations (5), (7) and (10). For a Great Eruption luminosity of $10^{7.3} L_{\odot}$ and a radius of 10 AU , the effective temperature becomes $8,300 \text{ K}$, and $\beta \approx 0.1$. Such an expansion implies, because of conservation of angular momentum, that the envelope must rotate very slowly, even if before expansion the star was almost at break-up angular velocity. I take now the following parameters: total mass lost during the Great Eruption equal to $M_{GE} = 12M_{\odot}$; an envelope mass of about half the stellar mass, or $M_{env0} = 70M_{\odot}$ at the beginning of the great eruption (as stated above, this is an upper limit on the envelope mass; the envelope mass is likely to be lower, increasing the effect studied here); mass loss geometry with $\mu = 0.5$ (see discussion following eq. 11); and $\beta = 0.1 - 0.15$. For these parameters, $\beta_1 = (\beta = \mu) \beta_1 = 4$. By equation (13), the envelope angular velocity at the end of the Great Eruption is $\beta_f = \beta_0 (58=70)^4 = 0.5 \beta_0$, where β_0 is the (non-dimensional) angular velocity before mass loss starts but after expansion. At the middle of the Great Eruption, $\beta_m = \beta_0 (64=70)^4 = 0.7 \beta_0$. For $\beta = 0.15$, as in the extended $60M_{\odot}$ model of Maeder (1981), these values are $\beta_f = 0.65 \beta_0$ and $\beta_m = 0.8 \beta_0$. Since in this scenario the angular velocity at the beginning of the Great Eruption should be very low as well $\beta_0 \ll 1$, I conclude that in the case when Car expanded during the Great Eruption the angular velocity is very low, and a single star model cannot account for the bipolar structure.

Consider now the other extreme, where Car remained a hot star during the Great Eruption. Equations (5) (7) and (10) implies $\beta = 0.1$. Again, I take $\mu = 0.5$, i.e., a moderate polar enhanced mass loss rate. The value of β cannot be too low. This is because a low value of β implies highly enhanced mass loss rate along polar directions. This, however, will make the problem of the radiation momentum transfer to the wind (eq. 1), even more severe, as only radiation escaping along the polar directions accelerate the mass there. Therefore, a value of $\beta = 4$, as in the extended envelope case, may be an underestimate. In any case, for this value of β , and for a maximum, possible value of $\beta_0 = 1$, the angular velocity and middle and end of the Great Eruptions are $\beta_m = 0.7$ and $\beta_f = 0.5$, respectively.

The main conclusion of this section is that as a single star, Car could not have maintained a fast rotation during the 20 years Great Eruption.

4. Summary

Most single-star models for the formation of the bipolar nebula the Homunculus of Car are based on fast rotation of the progenitor during the Great Eruption, 1843–1856. These models do not consider the origin of the fast rotation, or the way the envelope maintains its fast rotation during the Great Eruption. A single-star spin-up mechanism as proposed by Heger & Langer (1998) requires the outer convective part of the envelope to lose mass to inner regions. This will decrease substantially the ratio $M_{\text{env}}/M_{\text{env}0}$ in equation (13), hence substantially reducing the angular velocity as a result of the mass loss. The single-star models for the bipolar Great Eruption, therefore, do not directly address the question of what is the main factor behind the bipolar structure. There is, however, more to the single star model. Smith et al. (2003a), for example, attribute the 5.5 year period to angular momentum redistribution in the envelope. The star eruptively expels large amount of mass and angular momentum, such that its surface slows down. The star then relaxes, and angular momentum diffuses outward. After a period of 5.5 years, the surface rotates fast enough for a new eruption to occur. Although I see some problems with this scenario, it does, however, show that the angular momentum evolution in a single star scenario is more complicated than what I described in previous sections.

The goal of the present paper was to show that the origin of the fast envelope rotation required by these models cannot be ignored. Even when fast rotation occurs, it is not clear how much it can affect instabilities and the mass loss process (Stothers 1999). I used general considerations, rather than a specific model. (Most papers dealing with a fast rotating progenitor of Car do not actually provide such a stellar model!) The first problem is the initial fast rotation required. This in principle can be provided by a relatively low mass companion which entered the envelope of Car in the past. The more difficult problem to overcome, as I showed in section 3 above, is to maintain a fast envelope rotation during the Great Eruption itself. On top of the angular momentum problem, the models are further constrained by the huge momentum and kinetic energy of the mass blown during the great eruption (section 2 above). These, for example, rule out a model where most of the mass is blown from the polar caps of the progenitor.

Binary companions, on the other hand, can relatively easily account for the shaping of the Homunculus (Soker 2001), and the energy and momentum of the Great Eruption (section 2 above). The basic process is that of a companion accreting from the primary wind, forming an accretion disk, and blowing two opposite jets. The energy source is the gravitational energy of the mass accreted on the companion. Comparing LBV's nebulae with other objects, e.g., PNs, strongly suggests that binary interaction is indeed behind the shaping of these non-spherical nebulae (e.g., O'Hara et al. 2003). The binary model has

another advantage: it incorporates *Car* into a unified model explaining all objects having two low density lobes with an equatorial waist between them; sometimes these are termed bubble pair. These objects include clusters of galaxies, where the bubbles are X-ray deficient bubbles, symbiotic nebulae, and planetary nebulae (see Soker 2003 a,b; 2004a). The shaping in all these objects is via supersonic jets. The jets are blown by accretion disks at a speed about equal to the escape velocity from the accreting object (Livio 2000).

Following the analytical/exploratory papers of the shaping problem from a binary-model perspective (Soker 2001; 2003c, and the present paper), the next step in understanding the shaping of the Homunculus is to conduct a 3D numerical simulation of the mass transfer from *Car* to its companion, including two oppositely ejected jets, with possibly wide opening angle (Soker 2004a), and including the orbital motion.

ACKNOWLEDGEMENTS

I thank an anonymous referee for useful comments. This research was supported in part by a grant from the Israel Science Foundation.

REFERENCES

- Aerts, C., Lamers, H. J. G. L. M., Molenberghs, G. 2004, *A & A*, 418, 639
- Alexander, D. R., & Ferguson, J. W. 1994, *ApJ*, 437, 879
- Bahcall, J. N., Pinsonneault, M. H., Basu, S. 2001, *ApJ*, 555, 990
- Bohigas, J., Tapia, M., Ruiz, M. T., & Roth, M. 2000, *MNRAS*, 312, 295
- Bond, H. 2000, *ASP Conf. Ser.* 199, *Asymmetrical Planetary Nebulae II: From Origins to Microstructures*, ed. J. H. Kastner, N. Soker, & S. Rappaport (San Francisco: ASP), 115 (astro-ph/9909516)
- Brockopp, C., Sokolowski, J. L., Kaiser, C., Richards, A. M., Muxlow, T. W. B., & Seymour, N. 2004 *MNRAS*, 347, 430
- Corcoran, M. F., Ishibashi, K., Swank, J. H., & Peter, R., 2001a, *ApJ*, 547, 1034
- Corcoran, M. F., et al. 2001b, *ApJ*, 562, 1031
- Corradi, R. L. M., Livio, M., Balick, B., Munari, U., & Schwarz, H. E. 2001, *ApJ*, 553, 211
- Corradi, R. L. M., & Schwarz, H. E. 1995, *A & A*, 293, 871
- Damiani, A. 1996, *ApJ*, 460, L49

- Dam ineli, A ., Kaufer, A ., Wolf, B ., Stahl, O ., Lopes, D .F ., & de Araujo, F .X . 2000, *ApJ*, 528, L101
- Davidson, K . 1999, in *Eta Carinae At The Millennium*, *ASP Conf. Ser.* 179, eds. J. A . Morse, R .M . Humphreys, and A .Dam ineli (*Astronomical Society of the Pacific*; San Francisco)
- Davidson, K . 2000, in *Cosmic Explosions*, *ASP Conf. Vol.* 522, Eds. S.S. Holt and W .W . Zhang, (*American Institute of Physics*) 421.
- Davidson, K ., & Humphreys, R .M . 1997, *ARA & A*, 35, 1
- Duncan, R .A ., & White, S .M . 2003, *MNRAS*, 338, 425
- Dwarkadas, V .V ., & Owocki, S .P . 2002, *ApJ*, 581, 1337
- Fabian, A .C ., Celotti, A ., Blundell, K .M ., Kassim, N .E ., & Perley, R .A . 2002, *MNRAS*, 331, 369
- Fernandez Lajus, E ., Gam en, R ., Schwartz, M ., Salerno, N ., Llinares, C ., Farina, C ., Am orin, R ., & Niem ela, V . 2003 (*astro-ph/0311437*)
- Gonzalez, R .F ., de Gouveia DalPino, E .M ., Raga, A .C ., Velazquez, P .F . 2004, *ApJ*, 600, L59
- Guerrero, M .A ., Chu, Y .H ., Manchado, A ., & Kwitter, K .B . 2003, *AJ*, 125, 3213
- Heger, A ., & Langer, N . 1998, *A & A*, 334, 210
- Heger, A ., & Langer, N . 2000, *ApJ*, 544, 1016
- Hillier, D .J ., Davidson, K ., Ishibashi, K ., & Gull, T . 2001, *ApJ*, 553, 837
- Hirano, N ., Shinnaga, H ., Zhao, J .H ., Young, K ., Fong, D ., & Dinh, V .-T . 2004, in *Asymmetrical Planetary Nebulae III*, eds. M .M eixner, J .H .K astner, B .B alick, & N .Soker, *ASP Conf. Series*, in press
- Im ai, H ., Obara, K ., Diamond, P .J ., Om odaka, T ., & Sasao, T . 2002, *Nature*, 417, 829
- Ishibashi, K ., Corcoran, M .F ., Davidson, K ., Swank, J .H ., Peter, R ., Drake, S .A ., Dam ineli, A . & White, S . 1999, *ApJ*, 524, 983
- Ishibashi, K . et al. 2003, *AJ*, 125, 3222
- Kellogg, E ., Pedelty, J .A . & Lyon, R .G . 2001, *ApJ*, 563, L151155
- Kippenhahn, R ., & Weigert, A . 1990, *Stellar Structure and Evolution*, Springer-Verlag, Berlin
- Knapp, G .R . 1986, *ApJ*, 311, 731
- Langer, N ., Garcia-Segura, G ., & Mac Low, M .-M 1999, *ApJ*, 520, L49

- Livio, M . 2000, in *Asymmetrical Planetary Nebulae II: From Origins to Microstructures*, ed. J.H. Kastner, N. Soker, & S. Rappaport, *ASP Conference Series*, Vol. 199, (San Francisco: ASP), 243
- M aeder, A . 1981, *A & A* , 99, 97
- M aeder, A ., & Desjacques, V . 2001, *A & A* , 372, L9
- M aeder, A ., & Meynet, G . 2000, *ARA & A* , 38, 143
- M aeder, A ., & Meynet, G . 2003, *A & A* , 411, 543
- M cNam ara, B . R ., et al. 2000, *ApJ*, 534, L135
- Meynet, G ., & M aeder, A . 2003, *A & A* , 404, 975
- O 'H ara, T . B ., M eixner, M ., Speck, A . K ., U eta, T ., Bobrowsky, M . 2003, *ApJ*, 598, 1255
- P ittard, J.M ., & Corcoran, M . F . 2002, *A & A* , 383, 636
- Rogers, F . J ., & Iglesias, C . A . 1992, *ApJS*, 79, 507
- Sahai, R ., Morris, M ., K napp, G . R ., Young, K ., & B ambaum , C . 2003, *Nature*, 426, 261
- Shaviv, N . J . 2001, *MNRAS*, 326, 126
- Sm ith, N ., Davidson, K ., G ull, T . R ., Ishibashi, K ., & Hillier, D . J . 2003a, *ApJ*, 586, 432
- Sm ith, N ., G ehrz, R ., D ., H inz, P . M ., H o m ann, W . F ., H ora, J . L ., M am a jek, E . E ., & M eyer, M . R . 2003b, *AJ*, 125, 1458
- Sm ith, N ., & M orse, J . A . 2004, *ApJ*, 605, 854
- Soker, N . 2001, *MNRAS*, 325, 584
- Soker, N . 2003a, *PASP*, 115, 1296
- Soker, N . 2003b, *Nature*, 426, 236
- Soker, N . 2003c, *ApJ*, 597, 513
- Soker, N . 2004a, *A & A* , 414, 943
- Soker, N . 2004b, in *Asymmetrical Planetary Nebulae III*, eds. M . M eixner, J . H . K astner, B . B alick, & N . Soker, *ASP Conf. Series*, in press (extended version in astro-ph/0309228)
- Soker, N ., & H arpaz, A . 1999, *MNRAS*, 310, 1158
- Soker, N ., & H arpaz, A . 2000, *MNRAS*, 317, 861
- Stothers, R . B . 1999, *ApJ*, 513, 460
- Stothers, R . B ., & Chin, C .-W . 1993, *ApJ*, 408, L85
- van Boekel, R . et al. 2003, *A & A* , 410, 37

Zethson, T., Johansson, S., Davidson, K., Humphreys, R.M., Ishibashi, K., & Ebbets, D.
1999, *A & A*, 344, 211