T IM M I2 and V LT-ISAAC Spectroscopy of C ircum stellar D ust D isks { A Spatially R esolved 3.3 µm PAH Feature A round H D 100546

V incent C.G eers¹, Jean-Charles Augereau¹, K laus M.Pontoppidan¹, Hans U.Kau², Anne-M arie Lagrange³, Gael Chauvin^{3;4}, and Ewine F.van D ishoeck¹

¹ Leiden Observatory, P.O. Box 9513, 2300 RA Leiden, The Netherlands

² ESO, Garching, Germany

³ Laboratoire d'A strophysique de G renoble, France

⁴ ESO, Santiago, Chile

A bstract. W e present prelim inary results of spectroscopic data obtained with the La Silla 3.6m T IM M I2 and the V LT -ISA AC instrum ents. W e have obtained 23 T IM M I2 N-band and 18 Q-band spectra for our total sample of 36 stars in M arch 2003 to investigate the evolution of the grain properties in circum stellar disks. O ur sample consists of H erbig stars, T Tauri stars and Vega-like stars, som e of which have recently been identied to be members of nearby (< 100 pc) stellar associations. Silicate and P olycyclic A rom atic H ydrocarbon (PAH) em ission features are detected and the spatial resolution of T IM M I2 allowed to place upper lim its on the spatial extent of the em ission. W e also obtained ISA AC L-band spectra of 4 well-studied disks. For the H erbig B e star HD 100546, we detect spatially extended PAH em ission at 3.3 µm which corresponds to a source with radius 11 3AU in this line.

1 Introduction

In the last decade, circum stellar disks at di erent evolutionary stages have been intensively studied from optical to m illim eter wavelengths. Low -m ass (T Tauri) and interm ediate-m ass (Herbig Ae/Be) young stars have been shown to have young dusty and gaseous protoplanetary disks for which in aging can provide constraints on the structure [6,9,17]. M ore evolved gas-free dust disks around main-sequence stars, studied at sub-millimeter and near-infrared wavelengths, show a \lumpy" structure with gaps, holes and/or asymmetries in the dust distribution, indicative of gravitational perturbations by stellar com panion (s) or unseen planet (s) [2,3,18]. Near- and m id-infrared spectroscopy of circum stellar disks around Pre M ain Sequence (PMS) stars at di erent evolutionary stages has shown a wealth of gas and dust features, enabling detailed studies of dust m ineralogy and its relation with grain processing and disk evolution [10]. The shape of the $10\,\mu\text{m}$ feature can serve as a tracer of grain growth in disks and the presence of crystalline silicates, not observed in the ISM, is interpreted as evidence that the dust composition evolves as the disk is dissipating and large protoplanetary bodies are form ing [13,15].

2 The Spatially Resolved ISAAC Spectrum of HD 100546

W ith ISAAC at VLT we obtained in m id-2000 low resolution (R 600) L-band (2.8{ $4.2\,\mu$ m}) spectra of four old PMS stars using a 0.6^{00} 120⁰⁰ slit w ith a spatial resolution of 0.146^{00} /pixel. For HD 100546 the long slit was aligned with the m a jor axis of the disk resolved in scattered light [1,7]. The reduction procedure is described in [12]. It includes the rem oval of bad pixels and jitter correction before co-adding 2-D spectra, spectrum extraction, wavelength and ux calibration.



Fig.1. (Upper panel) ISAAC L-band spectrum of HD 100546, scaled by a factor 1.25 (see discussion in the text) to match the ISO spectrum and then shifted by +4Jy for clarity. The grey band highlights the position of the 3.3 μm PAH fea-(Lower panel) ture. FW HM of the spatial pro le, as extracted from 2D spectral image, of HD 100546 (thick line) and its corresponding standard BS 4757 (thin line). The inset shows the FW HM of HD 100546 only around $3.3\,\mu\text{m}$

The L-band spectrum for HD 100546 is plotted in the top panel of Fig.1 together with the ISO spectrum. The two spectra agree reasonably well on absolute ux, although the ISAAC continuum ux is 20% lower than the ISO ux. This di erence cannot be explained by aim ass di erence between the science and standard target. The spectrum shows several hydrogen em ission lines, which are stronger in the smaller aperture ISAAC data. The Pf δ line falls on top of a broad spectral feature at 3.3 µm attributed to PAHs [10]. Tabulated HI recombination line ratios from [8] were used to estim ate and subtract the contribution of the HIPf δ line to the intensity of the PAH feature. The PAH em ission feature to continuum ratios are sim ilar for both ISAAC and ISO spectra.

In the bottom panel of Fig.1, the measured FW HM of the spatial pro k, assumed to be Gaussian, as a function of wavelength is plotted for HD 100546 and the standard star BS 4757, which was used for telluric correction and ux calibration. The wavy pattern of the FW HM pro less is currently not understood

but it is sim ilarly observed for the science and standard stars. D istortion correction of the 2D spectra before FW HM extraction does not in prove the results. The spike at $3.31\,\mu\text{m}$ is caused by almost complete atmospheric extinction. W hen rescaling the measured FW HM of BS 4757 according to the di erence in measured airmass and DIMM seeing and dividing by a factor 1.05, we nd that the rescaled FW HM of BS 4757 ts reasonably well with the FW HM of the continuum of HD 100546. A ssum ing BS 4757 is a point source, we conclude that the continuum of HD 100546 is unresolved. A signi cant bump is observed at $3.3 \mu m$ with a spatial extent (FW HM) of 0.22^{00} 0.064⁰⁰(3) after correcting for interpolated FW HM of the continuum . A dopting a H ipparcos distance of 103^{+7}_{6} pc [16], this corresponds to a radius of 11 3AU, where the uncertainty is largely due to the noise in the extracted FW HM. In conclusion we nd a signi cant bum p in the FW HM at the wavelength of 3.3μ m which indicates that the disk is spatially resolved in this PAH feature. The $3.3 \mu m$ PAH carriers are thus spread over a larger area than the therm alized dust grains responsible for the adjacent continuum or the ionized gas responsible for the H lines.

3 TIM M I2 N -band Spectra of C ircum stellar D isks

W ith T M M I2 at the ESO /3.6m telescope we obtained 23 low resolution N-band (7.5{ 13.9μ m; R 160; $1.2^{\circ0}$ 70° slit) and 18 low resolution Q-band (14.5{ 23.9 μ m; R 130; $3.0^{\circ0}$ 70° slit) spectra for our sam ple of 36 H erbig A e/Be and T Tauri stars with a spatial resolution of $0.45^{\circ0}$ /pixel in N-band and $0.6^{\circ0}$ /pixel in Q-band. Spectra for targets with IRAS 12 μ m uxes as faint as about 0.5{1Jy



Fig.2. Selection of T M M I2 N-band spectra; the grey band indicates the position of the 9.6 μ m atmospheric extinction band. (Left panel) our T IM M I2 spectrum of HD 100546 is scaled to m atch the overplotted ISO spectrum and then shifted + 20 Jy for clarity. (M iddle panel) Spectrum of K K O ph, normalized to its ux at 12 μ m; showing a broad em ission feature between 9{12 μ m and a sm all feature at 11.2 μ m. (R ight panel) Spectra of HD 34282 and TY C rA; for comparison our T IM M I2 spectrum of HD 169142 is added, for which ISO observations have shown [10] that the em ission features at 8.6 and 11{11.2 μ m can be attributed to the presence of PAHs

4 V.C.Geers, J.-C.Augereau, K.M. Pontoppidan et al.

could be observed in N-band. The reduction procedure is similar to that of the ISAAC observations. Absolute ux calibration is not perform ed; for comparison of features all the spectra are normalized to their ux at 12 μ m, except HD 100546 which is scaled to m atch the ISO spectrum. Preliminary m id-IR N-band spectra for a selection of our T IM M I2 sample are shown in Fig.2.

For HD 100546, the shape of our T IM M I2 spectrum agrees reasonably well with the ISO spectrum but we do observe some interesting di erences: the sharp feature at 11.0 μ m in the ISO spectrum does not show up in our T IM M I2 spectrum while we do see a more pronounced feature at 11.8 μ m. The HD 100546 spectrum has a clear feature at 11.2 μ m, which can be attributed to blended features from both forsterite and PAH [4]. Using the same approach as for our ISAAC spectrum, we do not indicate of an excess spatial extent at the 11.2 μ m feature compared to the extent of the continuum in our T IM M I2 data, with an upper limit of a radius of 50 11AU.

KK O ph, a Herbig Be star, shows a broad em ission feature peaking between 9 and 12 μm , which could be attributed to sm all am orphous silicate grains, plus additional sm aller features like e.g. that close to 11 $2\,\mu m$, possibly due to crystalline silicates. TY C rA is a triple, perhaps quadruple [5], system and shows clear em ission features at 8.5 and 11.1 μm of which the latter feature has, based on ISO-SW S data, been attributed to PAHs [14]. HD 34282 is another Herbig star for which a gas disk in K eplerian rotation has recently been resolved at m illim eter wavelengths [11]. We detect in its mid-IR spectrum two features at 8.6 (Δ = 0.28 μm) and 11.2 μm (Δ = 0.20 μm) again consistent with the presence of PAHs con rm ing the tentative detection by ISO .

4 Future W ork with VLT Facilities

In relation to the work presented here, we plan to use the high spatial and spectral resolution of V ISIR to obtain spatially resolved N - and Q-band spectra of young P re M ain Sequence stars with circum stellar disks. This will provide the unique opportunity to simultaneously obtain a clean spectral separation of am orphous and crystalline silicates and distinguish for example the 11.2 μ m PAH and 11.3 μ m forsterite features as well as constrain their spatial distribution. C om - pared to T IM M I2 on the La Silla 3.6m telescope, V ISIR at the 8.2m V LT will allow us to put m uch stronger constraints on the spatial extent of the 11.2 μ m PAH feature of HD 100546 because of the higher spatial resolution.

A cknow ledgem ents: A strochem istry in Leiden is supported by a NW O Spinoza grant. J.C. A ugereau is supported by a fellow ship from the European Research Training Network The O rigin of P lanetary System s" (PLANETS; contract num ber HPRN-CT-2002-00308) at Leiden O bservatory.

References

- 1. Augereau, J.C., Lagrange, A.M., Mouillet, D., et al. 2001, A&A, 365, 78
- 2. Augereau, J.C., Nelson, R.P., Lagrange, A.M., et al. 2001, A&A, 370, 447
- 3. Augereau, J.C. & Papaloizou, J.C.B. 2004, A & A, 414, 1153
- 4. Bouwm an, J., de Koter, A., Dom inik, C. et al. 2003, A&A, 401, 577
- 5. Chauvin, G. et al. 2003, A & A, 406, L51
- 6. Dartois, E., Dutrey, A., & Guilloteau, S. 2003, A & A, 399, 773
- 7. G rady, C.A. et al. 2001, AJ, 122, 3396
- 8. Hummer, D.G. & Storey, P.J. 1987, MNRAS, 224, 801
- 9. Mannings, V. & Sargent, A. I. 1997, ApJ, 490, 792
- 10. M eeus, G ., W aters, L.B.F.M ,Bouwm an, J. et al. 2001, A & A , 365, 476
- 11. Pietu, V., Dutrey, A., & Kahane, C. 2003, A & A, 398, 565
- 12. Pontoppidan, K.M., Fraser, H.J., Dartois, E. et al. 2003, A & A, 408, 981
- 13. Przygodda, F., van Boekel, R., Abraham, P. et al. 2003, A&A, 412, L43
- 14. Siebenm orgen, R., Prusti, T., Natta, A., et al. 2000, A&A, 361, 258
- 15. van Boekel, R., W aters, L.B.F.M., Dom inik, C. et al. 2003, A&A, 400, L21
- 16. van den Ancker, M E., The, P.S., T jin A D jie, H R E. et al. 1997, A&A, 324, L33
- 17. W olf, S., Padgett, D. L., & Stapelfeldt, K. R. 2003, ApJ, 588, 373
- 18. W yatt, M.C.2003, ApJ, 598, 1321