DENIS detections of highly obscured galaxies in the area around PKS 1343 { 601

Anja C. Schroder

Dept. of Physics & Astronomy, University of Leicester, University Road, Leicester LE17RH, UK

Renee C.K raan-Korteweg

Depto. de Astronom a, Universidad de Guanajuato, Apdo. Postal 144, Guanajuato GTO 36000, Mexico

Gary A.Mamon

IAP (CNRS UMR 7095), 98 bis B kmd A rago, 75014 Paris, France, and GEPI (CNRS UMR 8111), Observatoire de Paris, 92195 M eudon, France

Patrick A.W oudt

Dept. of Astronomy, University of Cape Town, Rondebosch 7700, South Africa

A bstract. We present results of a search for galaxies around the highly obscured giant radio galaxy PKS 1343 { 601 using the near-infrared DEN IS survey. We compare our notings with surveys in the B-band, at the 21 cm line emission, and with 2MASS. Recession velocities of galaxies in this area suggest a low-velocity-dispersion group or cluster of galaxies including an X-ray con med Seyfert 2. The colours of the galaxies have been used to exam ine the extinctions in this low-latitude area where IRAS/DIRBE estimates are unreliable. We not the true extinction to be roughly 15% lower than the IRAS/DIRBE extinctions.

#### 1. Introduction

PKS 1343 { 601 (Centaurus B) is a very strong radio source in the southern Zone of Avoidance (ZoA). Near-infrared (NIR) observations revealed it to be a giant elliptical galaxy as often found in the centre of galaxy clusters. At a distance of

10 from the G reat Attractor (GA) and with a recession velocity of 3872 km s <sup>1</sup> (W est & Tarenghi 1989) it lies within the GA overdensity. Finding a new cluster here will have considerable impact on the local velocity eld calculations. On the other hand, X-ray observations with ASCA have only revealed di use emission from PKS 1343 { 601 itself (Tashiro et al. 1998, see also the discussion in Ebeling et al. 2002) which would rule out a rich cluster. A more sensitive XMM—New ton observation does not indicate any obvious cluster emission either but will have to be analysed in detail for an improved estimation.

A detailed discussion of PKS 1343 { 601 and the potential for a cluster in this area can be found in K raan-K orteweg et al. (these proceedings), who present

I-band observations of a 2  $\,$  2  $\,$  area around PKS 1343 { 601. Nagayam a et al. (2004) have analysed deep HJK observations of the inner  $36^{0}$   $36^{0}$  (see also Nagayam a et al., these proceedings). A more detailed presentation of the following work will be given in Schroder et al. (2004).

To search for galaxies in this highly obscured area ( ${\rm fA_B~i~'}~12^{\rm m}$ ) we have used the N IR survey D E N IS (Epchtein 1997, 1998). The N IR has the advantage that it is less a ected by the foreground extinction as compared to the optical (the extinction in K is about 10% of the extinction in B), and that it is sensitive to early-type galaxies, tracers of massive groups and clusters which are neither uncovered in far infrared surveys nor in the 21 cm radiation where extinction is negligible (see M am on 1994).

# 2. The DEN IS survey

The DEN IS survey is a N IR survey of the southern sky in the Cousins-I (0.8 m), J (1.25 m) and K (2.15 m) passbands with a resolution of  $1^{0}$  in I and  $3^{0}$  in J and K (Epchtein 1997, 1998). The observations were carried out between 1996 and 2001 with a dedicated 1 m ESO telescope at La Silla (Chile). About 92% of the southern sky (+2 DEC 88) were covered.

DENIS in ages are  $12^0$   $12^0$  large with an exposure time of 9s. The observing mode consisted of step-and-stare scans of 180 in ages in declination, resulting in strips of  $12^0$  30. There is an overlap region of  $1^0$  on each side of every in age. The limiting magnitudes (at a sensitivity of about 3) are  $18^n$ . 5,  $16^n$ . 5,  $13^n$ . 5 for the I-, J-, and K-bands, respectively, while the limits for high completeness and reliability of the galaxy extraction at high galactic latitudes are  $16^n$ . 5,  $14^n$ . 8, and  $12^n$ . 0, respectively (M am on 1998).

Interpolating from C ardelli et al. (1989), the extinctions in the DEN IS N IR passbands are  $A_{\rm I}=0^{\rm o}.45$ ,  $A_{\rm J}=0^{\rm o}.21$ , and  $A_{\rm K}=0^{\rm o}.09$  for  $A_{\rm B}=1^{\rm o}.0$ . Thus the decrease in number counts of galaxies in the ZoA as a function of extinction is considerably slower in the N IR than in the optical (Schroder et al. 2004, Schroder, K raan-K orteweg, & M am on, 1999). While DEN IS easily detects galaxies up to extinctions of  $A_{\rm B}$  '  $10^{\rm m}$ , we can indiminishably bright and close by galaxies in the K-band at much higher extinctions.

# 3. DENIS detections

We have searched 29 DENIS slots each with 37 im ages around PKS1343 { 601. The total area covered amounts to 29:8 square degrees. Using the DENIS visualisation package Denis 3d by E.Copet we scanned each image simultaneously in the three passbands. That way the relative appearance of highly-obscured galaxies in the three bands, which is dierent to most stars, could be used to distinguish galaxies from blended stars and extended Galactic objects.

The automatic extraction package SExtractor (Bertin & Amouts 1996) was used to obtain IJK photometry for the visually detected galaxies. We have derived total magnitudes as well as colours within a 7-arcsecond aperture (to avoid strong contamination by superimposed stars). We have subsequently checked the magnitudes and colours of each galaxy and compared them with




Figure 1. The distribution of galaxies (circles) and Galactic objects (crosses) in the searched area (tilted rectangular) is shown. Filled circles are galaxies visible with DENIS, and open circles stand for uncertain DE-NIS galaxies. Extinction contours according to the IRAS/DIRBE maps are displayed as labelled. The galaxy PKS1343 { 601 (l =  $309:7;b = +1:8,A_B = 12\%:3$ ) is close to the centre in this im age.

the im age to see whether SExtractor has deblended all objects close by and whether all the param eters agree w ith each other.

On 1073 searched im ages we found 83 galaxies (plus 38 uncertain candidates). Of these, 79 (33) are visible in the I-band, 82 (35) in J, and 67 (25) in K (see Schroder et al. 2004 for details). Figure 1 shows the distribution of the detected galaxies: lled circles stand for de nite galaxies, and open circles show uncertain candidates. Crosses denote Galactic objects. The contours depict the extinctions at  $A_B=2^m$ ,  $3^m$ ,  $5^m$ ,  $10^m$ ,  $20^m$ ,  $30^m$ ,  $50^m$ , and  $60^m$ , as indicated.

W hile PKS 1343 { 601 is prom inent in all three passbands, a nearby equally large galaxy is not visible in the I-band and is probably an early-type spiral with an elongated halo. Two objects at very high extinctions (with  $A_B=39^m$  and  $24^m$ , for DZOA 4641-06 and DZOA 4645-13, respectively) are classified as uncertain galaxies; they are only visible in the K-band. With extinction-corrected (but not diam eter-corrected, cf. Cameron 1990) magnitudes of K°=8.9 and 8.3, respectively, they are only slightly fainter than the central elliptical galaxy at K°=8.0, which would indicate that these two galaxy candidates could lie at the distance of the cluster or closer.

At very high extinctions, we also detected extended G alactic objects like planetary nebulae, H  $\scriptstyle\rm II$  regions, and rejection nebulae, which are often associated with young stellar objects. Some of these objects are easily recognisable as non galaxian, but there are cases where a distinction could only be made on the basis of high foreground extinction (which is usually  $A_B > 50^m$ ). A careful comparison between unambiguous galaxies and Galactic objects was used to classify cases at intermediate extinctions.

## 4. Com parison with other catalogues

#### 4.1. B-band

W oudt & K raan-K orteweg (2001, hereafter W KK), who have conducted a deep search for galaxies using copies of the ESO-SERC B<sub>J</sub>-band photographic plates, list 35 galaxies within our search area. Only one of their galaxies (W KK 2589, D ZOA 4655-08) is not visible on the DEN IS im ages; it is a sm all, very low surface brightness galaxy, probably of type Sm or Irr. Five B-band galaxies, all classi ed as uncertain galaxies by W KK, were identified as (blended) stars with the higher spatial resolution of the DEN IS I-band im ages.

M ost B -band galaxies are found in the low extinction regions, that is, at  $A_B < 3^m$ . The highest extinction for a B -band galaxy in the searched area is  $A_B = 5^m$ . 2 (D ZOA 4641-07, W K K 2301). The completeness limit for B° = 15 $^m$ . 5 and D° = 60 $^m$  of the W K K -catalogue is  $A_B = 3^m$ .

## 4.2. 2M ASS

We have extracted 65 objects from the 2M ASS all-sky extended source catalogue (2M ASS, 2003) within the region  $13^h:45 < RA < 14^h:11$  and 63:8 < Dec < 57:4, 11 of which are just outside the searched DEN IS area. Forty nine objects are common in both data sets, nine of which are classified by us as Galactic objects and two as uncertain galaxies. Five 2M ASS objects were not found with DEN IS; one is a Galactic object, two are misidentications, and two are very small but bright galaxies that were not recognised as such in the DEN IS search. This gives a rough estimate of the reliability of 80% for the automatic 2M ASS extraction in this highly confused and obscured region, while the completeness of the 2M ASS galaxy extraction in this area is 50%. Many of the DEN IS galaxies which are not in the 2M ASS extended source catalogue were either not recognised as extended by the 2M ASS automatic search algorithm (they will be in the 2M ASS point source catalogue however), or they are very faint in J and K and were only found through the DEN IS I-band.




Figure 2. The di erence of DEN IS and 2M ASS m agnitudes versus DEN IS m agnitudes in J (upper panel), in K (m iddle panel) and for the 2M ASS isophotal K-band m agnitude in the bottom panel.

Figure 2 shows a comparison of the 2M ASS J- and K-band magnitudes with our magnitudes. The J-band shows a magnitude dierence of 0 $^{\circ}$  08 0 $^{\circ}$  07 and a standard deviation of 0 $^{\circ}$  41 (upper panel), while in the K-band (middle panel) the dierences is 0 $^{\circ}$  30 0 $^{\circ}$  06 with a standard deviation of 0 $^{\circ}$  33, that is, there is a substantial oset between the two catalogues in the K-band. This oset is due to the large dierence in magnitude limits of the two surveys: the high-latitude (bj> 20) nominal values for 2M ASS are 14 $^{\circ}$  3 and 15 $^{\circ}$  8 for K and J, respectively, while for DEN IS they are 12 $^{\circ}$  0 and 14 $^{\circ}$  8, respectively. If we use the 2M ASS isophotal K-band magnitudes (at 20 $^{\circ}$  per square arcsecond) instead the mean of the dierences becomes maller at 0 $^{\circ}$  18 0 $^{\circ}$  06 ( = 0 $^{\circ}$  32, bottom panel), indicating that the DEN IS isophotal limit out to which SExtractor determines the total K-band magnitudes is even brighter.

U sing the 2M ASS isophotal J-band the change in o set is negligible (the mean is now  $0^{n}$ , 03  $0^{n}$ , 07 w ith  $= 0^{n}$ , 34; not shown) as expected by the smaller dierence in limiting magnitudes.




Figure 3. The X-ray spectrum of the AGN DZOA4653-11 with a model of a cold absorber plus a power law tted, top panel, and the residuals of the t, bottom panel.

## 4.3. X -rays

One of the galaxies found in our DEN IS search, DZOA 4653-11, was discovered in the serendipitous 1XMM catalogue (XMM SSC, 2003) and was identified as a highly obscured AGN (Motch, priv.comm.) with high foreground extinction. A deep I-band in age obtained by Motch indicates an early type spiral galaxy. A prominent Holine in the optical spectrum obtained with EFOSC at the 3.6m ESO telescope by Motch leads to a heliocentric velocity of  $v=3643\,\mathrm{km~s^1}$ , which means this galaxy is a cluster member. The X-ray ux over the energy range 0.2 12.0 keV is slightly variable with time and gives (1:351 0:007) 10  $^{11}$  erg cm  $^2$  s  $^1$ , i.e.,  $L_{\rm X}=2.2$   $10^{42}h_{70}^{\,2}$  erg s  $^1$ .

Fitting a cold absorber plus a power law to the XMM-Newton spectrum (Fig. 3) we not a photon index of  $1.27\,$  0.02 and a hydrogen column density of N $_{\rm H}\,=\,(2.05\,$  0.04)  $10^{22}$  cm  $^2$ . The Galactic column density at this position from the IRAS/DIRBE estimate from Schlegel et al. is only N $_{\rm H}\,=\,1.07\,$   $10^{22}$  cm  $^2$ , indicating there is signicant intrinsic absorption (note, though, that the IRAS/DIRBE maps are uncertain at this latitude), suggesting it to be a Seyfert 2 galaxy. A faint iron line (44 eV equivalent width) is also visible.

## 4.4. 21 cm wavelength

At 21 cm, the Galactic plane is transparent and the search for galaxies here is limited because of the increased noise due to strong Galactic continuum sources. We have extracted H I velocities for galaxies in this area from the blind H I Parkes All Sky Survey ZoA (HIZOA) survey of the southern ZoA conducted with the multibeam receiver on the Parkes telescope (Kraan-Korteweg et al.




Figure 4. Radial velocities as a function of distance to PKS 1343 { 601. The led circles represent the low velocity dispersion sub-system while the open circles show the velocity outliers.

2004), which covers the entire southern ZoA (jbj 5) in the velocity range 1200 to 12700 km s<sup>1</sup> with an integration time of 25 m inutes. To these, we added 7 optical velocities of galaxies within the search area (Fairall et al. 1998; V isvanathan & Yam ada 1996; W est & Tarenghi 1989; and the AGN D ZOA 4653–11).

Figure 4 shows the velocities plotted as a function of the distance to the giant elliptical PKS 1343 { 601. The velocities in close multiplets and pairs were replaced by theirm edian values, so as to avoid double counting multiply-identi ed single objects, as well as to remove the binary contribution to the group velocity dispersion. The velocity dispersion out to 2 and for  $1850 < v < 6500 \,\mathrm{km \ s^{\ 1}}$  is  $907 \, \mathrm{km} \, \mathrm{s}^{\, 1}$  . However, Figure 4 shows wide gaps in the velocity distribution. In the range  $3000 < v < 4700 \, \mathrm{km \ s^{\, 1}}$  there are 18 galaxies with a velocity dispersion  $v = 136 \,\mathrm{km} \,\mathrm{s}^{\,1}$  , and the gaps outside of the extrem e velocities among these 18 galaxies correspond to over 5  $_{
m v}$ . Taking the velocity dispersion at face value, we estimate the virial radius of the system to be  $233 h_{70}^{1} \text{ kpc} = 0:5$ , but we only have two velocities within this distance to PKS 1343 { 601, and, iterating over assum ed values of the virial radius, one nds that the velocities within the virial radius are compatible with the idea that the system around PKS 1343 { 601 is richer with a larger true velocity dispersion, although it cannot be as rich as a rich cluster (see Schroder et al. 2004, for details). This is consistent with the upper lim it on the cluster X -ray em ission (Ebeling et al. 2002). However, more velocity measurements and deeper NIR imaging will be needed to con rm this result.

#### 5. Extinction and N IR colours

Colours of galaxies are independent of distance but sensitive to extinction. Hence, NIR colours can be used to derive extinctions in low latitude areas where the IRAS/DIRBE maps are not properly calibrated. Cameron (1990) has shown that one needs to apply both a correction to the isophotal magnitude of a galaxy as well as for the reduction of the major axis of an obscured galaxy. Since the

colours used for the following analysis were derived from 7-arcsecond apertures (where contamination by superimposed stars is small) we only applied the magnitude correction according to the extinction in the IRAS/DIRBE maps and looked for dependencies in the residuals.

The left-hand panels in Fig. 5 show the extinction-corrected colours as a function of extinction in the B -band (lled circles are galaxies, open circles uncertain candidates). Galaxies at higher extinctions are clearly too blue, i.e., the extinction correction according to the IRAS/DIRBE maps overestimates the true extinction.

W ith the least square to the led circles (i.e., excluding the uncertain candidates), written in the form  $C^{\circ} = aA_B + b$  where C is the colour, we nd:

$$(I \ J)^{\circ} = (0.038 \ 0.013) A_B + (1.071 \ 0.057); = 0.21$$

$$(J K)^{\circ} = (0.008 0.009) A_B + (0.940 0.049); = 0.18$$

$$(I K)^{\circ} = (0.061 0.018) A_B + (2.149 0.086); = 0.28$$

If we assume the true extinction to be a constant factor f of the IRAS/DIRBE value of Schlegel et al., i.e.,  $A_B^{true}=f\,A_B$ , we can derive the correction from the above equations as

$$f = 1 + \frac{a}{E = A_B}$$
;

where E is the reddening in the respective colour. Combining the (independent) estimates of f for I  $\,$  J and J  $\,$  K  $\,$ , we derive

$$A_{B}^{true} = (0.86 \ 0.04) A_{B}$$
:

To verify this result and to exclude the possibility that this particular region shows an unusual deviation, we have also derived DENIS colours of galaxies found in a di erent region of the ZoA. These are part of a project that looks for DENIS counterparts of galaxies found with the blind HIZOA shallow survey (HIZSS, Henning et al. 2002) within a 10-arcm inute search radius. Forty HI detections in the area 210  $\,<\,1<\,260$ , jbj  $\,<\,5:7$  have been searched so far (Schroder et al., in preparation). The right-hand panels in Fig. 5 show the results for all galaxies found in these regions and with good photom etry. The least square ts to the led circles give:

$$(I J)^{\circ} = (0.054 0.013) A_{B} + (1.199 0.037); = 0.22$$
  
 $(J K)^{\circ} = (0.014 0.019) A_{B} + (1.171 0.065); = 0.26$   
 $(I K)^{\circ} = (0.056 0.027) A_{B} + (2.365 0.092); = 0.36$ 

D espite the di erent coverage in extinction the slopes of the least square ts agree well within the errors. Combining now the estimates for both regions in the colours I  $\,$  J and J  $\,$  K  $\,$  we derive

$$A_{B}^{true} = (0.83 \ 0.03) A_{B}$$
:

In other words, we not the true extinction in the southern ZoA to be about 17% lower than the estimates of the maps by Schlegel et al. (1998).




Figure 5. The NIR colours, corrected for extinction according to the IRAS/DIRBE maps, are plotted versus extinction in the B-band for the galaxies around PKS 1343  $\{$  601 (left panels) and for galaxies around H I detections in a dierent region of the ZoA (right panels). Filled circles are galaxies, open circles are uncertain galaxies. Least square to the led circles are shown.

O ne has to keep in m ind that, apart from the di culties in interpreting high IRAS/DIRBE measurements as reddening values, the gas/dust distribution in the Galactic P lane shows variations on a small spatial scale. This patch in essmay lead to a bias in adding galaxies preferentially in the low extinction areas and therefore to underestimate the true extinction. However, the agreement between the two regions we have used is excellent despite the fact that one covers a range in extinction of  $2^{\rm m} < {\rm A_B} < 16^{\rm m}$  while the other is restricted to  $0^{\rm m} < {\rm A_B} < 8^{\rm m}$ . In other words, the fraction of galaxies missed in the range  $8^{\rm m} < {\rm A_B} < 16^{\rm m}$  due to much higher extinction patches is negligible and does not a ect our result.

We will continue with this work using the colours of galaxies found within the search radii of the other half of the 110 HIZSS detections at higher Galactic longitudes (and offen at higher extinctions) to improve the statistics. Future deeper NIR surveys, like UKIDSS (Warren, 2004) and VISTA (McPherson, Craig, & Sutherland 2003), can be used to go to higher extinction values. A comparison with the current work will be important to verify that no high-extinction patches have substantially in uenced the here presented results.

A cknow ledgm ents. This research has made use of the NASA/PAC Infrared Science Archive (2MASS) and the NASA/PAC Extragalactic Database (NED), which are operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. The authors are grateful to the HPASS-ZOA team and the DENIS teams in Chile and at PDAC for all the eorts in observing and reducing the data. We also thank Ch. Motch for making unpublished data available to us. ACS and PAW thank the National Science Foundation for nancial support; RCKK thanks CONACyT for their support (research grants 27602 and 40094-F).

#### R eferences

Bertin, E., & Amouts, S. 1996, A&A 117, 393

Cameron, L M . 1990, A & A 233, 16

Cardelli, JA., Clayton, G.C., & Mathis, J.S. 1989, ApJ 345, 245

Ebeling, H., Mullis, C.R., & Tully, R.B. 2002, ApJ 580, 774

Epchtein, N. 1997, in 2nd Euroconference, The Impact of Large Scale Near-Infrared Surveys, eds. F. Garzon et al., (Dordrecht: Kluwer), 15

Epchtein, N. 1998, in 179th Symposium of the IAU, New Horizons from Multi-Wavelength Sky Surveys, eds. B.J.McLean, D.A.Golombek, J.J.E. Hayes, H.E. Payne, (Dordrecht: Kluwer), 106

Fairall, A.P., Woudt, P.A., & Kraan-Korteweg, R.C. 1998, A&ASS 127, 463

Henning, P.A., Staveley-Sm ith, L., Ekers, R.D., Green, A.J., et al. 2000, A.J. 119, 2686 (H. IZSS)

K raan-K orteweg, R.C., Staveley-Smith, L., Donley, J., Koribalski, B., & Henning, P.A. 2004, in IAU Symposium 216, Maps of the Cosmos, eds. M. Colless, and L. Staveley-Smith, ASP Conf. Ser. (ASP: San Francisco), in press, astro-ph/0311129

K raan-K orteweg, R  $\mathcal{L}$ , O choa, M ., W oudt, P ., & Andemach, H .2004, these proceedings M am on, G A .1994, Ap& SS 217, 237 (astro-ph/9312036)

M am on, G A . 1998, in X IV th IAP A strophysics M eeting, W ide Field Surveys in Cosmology, eds. S. Colombi, Y. Mellier, & B. Raban, (Gif-sur-Yvette: Editions Frontieres), 323 (astro-ph/9809376)

M cPherson, A M ., Craig, S.C., & Sutherland, W .2003, Large G round-based Telescopes, eds. J.M. Oschmann, L.M. Stepp, Proceedings of the SPIE, Volume 4837, 82

Nagayama, T., Woudt, P.A., Nagashima, C., Nakajimia, Y., et al. 2004, MNRAS (in press)

Nagayama, T., Woudt, P.A., Nagashima, C., Nakajimia, Y., et al. 2004, these proceedings

Schroder, A., Kraan-Korteweg, R.C., & Mamon, G. 1999, PASA 16, 42

Schroder, A., Mamon, G., Kraan-Korteweg, R.C., & Woudt, P.A. 2004, A.A. (in preparation)

Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ 500, 525

Tashiro, M., Kaneda, H., Makishima, K., Iyomoto, N., et al. 1998, ApJ 499, 713

Two Micron All Sky Survey team 2003, 2M ASS extended objects. Final release. volp.

Visvanathan, N., & Yamada, T. 1996, ApJS 107, 521

W arren, S. 2004, in IAU Symposium 216, Maps of the Cosmos, eds. M. Colless, and L. Staveley-Smith, ASP Conf. Ser. (ASP: San Francisco), in press

West, R.M., & Tarenghi, M. 1989, A&A 223, 61

Woudt, P.A., & Kraan-Korteweg, R.C., 2001, A&A 380, 441

XMM -Newton Survey Science Centre (SSC) 2003: The First XMM -Newton Serendipitous Source Catalogue, Version 1 (1XMM)