Principal Component Analysis of RR Lyrae light curves

S.M.Kanbur¹? and H.Mariani¹

 ^1D epartm ent of A stronom y, U niversity of M assachusetts Am herst, M A 01003, U SA

Received XX XXX 2003 / A ccepted XX XXX 2003

ABSTRACT

In this paper, we analyze the structure of RR ab star light curves using Principal Com ponent A nalysis. We nd this is a very e cient way to describe m any aspects of R R ab light curve structure: in m any cases, a P rincipal C om ponent t with 9 param eters can describe a RR ab light curve including bum ps whereas a 17 parameter Fourier t is needed. As a consequence we show show statistically why the amplitude is also a good sum mary of the structure of these RR Lyrae light curves. We also use our analysis to derive an empirical relation relating absolute magnitude to light curve structure. In comparing this formula to those derived from exactly the same dataset but using Fourier parameters, we d that the Principal Component Analysis approach has distinct advantages. These advantages are, rstly, that the errors on the coe cients multiplying the tted parameters in such formulae are much smaller, and secondly, that the correlation between the Principal C om ponents is signi cantly sm aller than the correlation between Fourier am plitudes. These two factors lead to reduced form al errors, in some cases estimated to be a factor of 2, on the eventual tted value of the absolute m agnitude. This technique will prove very useful in the analysis of data from existing and new large scale survey projects concerning variable stars.

Keywords: RR Lyraes { Stars: fundam entalparam eters

1 IN TRODUCTION

K anbur et al (2002), H endry et al (1999), Tanvir et al (2004) introduced the use of P rincipal C om ponent A nalysis (PCA) in studying C epheid light curves. They showed that a m a jor advantage of such an approach over the traditional Fourier m ethod is that it is m uch m ore e cient: an adequate Fourier description requires, at best, a fourth order t or 9 param eters, w hilst a PCA analysis requires only 3 or 4 param eters w ith as m uch as 81% of the variation in light curve structure being explained by the rst param eter. Later, Leonard et al (2003) used the PCA approach to create C epheid light curve tem plates to estim ate periods and m ean m agnitudes for H ST observed C epheids. The purpose of this paper is to apply the PCA technique to the study of RR Lyrae light curves.

The m athem atical form ulation and error characteristics of PCA are given in K 02 and will only be sum m arized here.

2 DATA

The data used in this study were kindly supplied by Kovacs (2002 private communication) and used in Kovacs and W alker (2001, hereafter KW). These data consist of 383 RRab stars with well observed V band light curves in 20 di erent globular clusters. KW perform ed a Fourier t to these data, which, in some cases, is of order 15.D etails concerning the data can be found in KW. The data we work with in this paper is this Fourier t to the magnitudes and we assume that the Fourier parameters published by KW are an accurate t to the actual light curves. W e start with the data in the form used in KW : a list of the mean magnitude, period and Fourier parameters for the V band light curve. The light curve can thus be reconstructed using an expression of the form

$$V = A_0 + A_k \sin(k!t + k);$$
(1)

where A_0 is the mean magnitude, ! = 2 = P, P the period, A_k ; the Fourier parameters given in KW. These light curves are then rephased so that maximum light occurs at phase 0 and then rewritten as

$$V = A_0 + (a_k \cos(k!t) + b_k \sin(k!t)):$$
(2)

The a_k ; b_k are the light curve characteristics entering into the PCA analysis (K 02). We then solve equation (4) of K 02, either after, or before removing an average term from the Fourier coe cients in equation (2). W ith PCA, the light curve is written as a sum of "elementary" light curves,

$$V (t) = PCA1L_1(t) + PCA2L_2(t) + PCA3L_3(t) + \dots$$
(3)

where V (t) is the magnitude at time t, PCA 1; PCA 2:: etc. are the PCA coe cients and the $L_i(t); i = 1; 2; 3:::$ are the elementary light curves at phase or time t. These elementary light curves are not a priori given, but are estimated from the dataset in question. Each star has associated with it a set of coe cients PCA 1; PCA 2; ::: and these can be plotted against period just as the Fourier parameters in equation (1) are plotted against period. We also note that the PCA results are achieved as a result of the analysis of the entire dataset of 383 stars whereas the Fourier method produces results for stars individually. This feature of PCA is particularly useful when perform ing an ensem ble analysis of large num bers of stars obtained from projects such as OG LE, MA – CHO and GAIA.

3 RESULTS

Solving equation (4) of K 02 yields the Principal C om ponent scores and the am ount of variation carried by each com ponent. W hat we mean by this is the following: if we carry out an Nth order PCA t, then PCA will assume that all the variation in the dataset is described by N components and sim ply scale the variation carried by each component accordingly. Table 1 shows this "am ount of variation" quantity with and without the average term rem oved. We see that in the case when we do not rem ove the average term the rst PC explains as much as 97% of the variation in the light curve structure. In the case when we do rem ove the average term from the Fourier coe cients, the rst PCA coe cient explains as much 81 percent of the variation in light curve structure. In either case, the rst four components explain m ore than 99:99% of the variation.

Figures 1 and 2 show some representative light curves from our RR ab dataset. In each panel of these two gures, the solid line is the Fourier decom position of order 15 (that is 31 parameters) used by KW, whilst the dashed line is a PCA generated light curve of order 14 (that is 15 parameters). Straightforward light curves such as the one given in the bottom and top left panels of gures 1 and 2 respectively are easily reproduced by our method. The top left panel of

gure 1 provides an example of an RR ab light curve with a dip and sharp rise at a phase around 0.8. This is well reproduced by PCA. It could be argued that PCA does not do as well as Fourier in m im icking this feature, for example, in the bottom right panel of gure 2. How ever, the di erence in the peak magnitudes at a phase of around 0.8 is of the order of 0.02m ags. It is also in portant to rem em ber that the PCA method is an ensemble method and analyzes all stars in a dataset simultaneously. With Fourier, it is possible to tailor a decomposition to one particular star. This di erence can be seen either as a positive or negative point about either technique. G iven this, we contend that PCA does rem arkably well in describing the full light curve m orphology of R R ab stars. On the other hand, the Fourier curve in the bottom left panel of qure 2 at this phase is not as smooth as the PCA curve.

In fact the PCA curves do not change much after about

8 PCA parameters. Even though table 1 in plies that the higher order PCA eigenvalues are small, we feel justimed in carrying out such a high order PCA to because its only after about 8 PCA components that the tted light curve assumes a stable shape. The left panel of gure 3 displays an eighth order PCA to (9 parameters, dashed line) and a fourth order FCA to (9 parameters, solid line). The Fourier curve still has some numerical wiggles whilst the PCA curve is smoother. In addition, the two curves disagree at maximum light. The right panel of gure 3 shows, for the same estar, the same order PCA curve as the left panel and an eighth order Fourier to (17 parameters). Now the two light curves agree very well. Note that in portraying the PCA and Fourier ts of reduced order in this gure, we simply truncated the original representations to the required level.

We suggest that gures 1-3 and table 1 provide strong evidence that PCA is an e cientway to describe RR ab light curve structure without comprom ising on what light curve features are captured by this description.

Figures 4-6 display plots of the st three PC scores plotted against log period for our sample. The errors associated with these PCA scores are discussed in section 4 of K 02 and given in equation 6 of that section. The orthogonal nature of these scores m ay well provide insight into the physical processes causing observable features in the light curve structure. A detailed study of these plots, in conjunction with theoretical models, is left for a future paper.

Figure 7 graphs V band am plitude against the rst PCA coe cient (after averaging). W e see a very tight correlation. Since table 1 im plies that PCA1 explains about 81% of the variation in light curve structure, gure 6 shows that the am plitude is a good descriptor of RRab light curve shape, at least for the data considered in this paper. A lthough the Fourier amplitudes are also correlated with amplitude, with PCA, we can quantify, very easily, the am ount of variation described by each PCA component. This has implications for both m odeling and observation. On the m odeling side, a com puter code that can reproduce the observed am plitude at the correct period, will also do a good job of reproducing the light curve structure. On the observational side, this provides insight into why we can use the am plitude, rather than a fullblown PCA or Fourier analysis, to study the general trends of light curve structure. This is why com paring theoretical and observational RR ab light curves on periodam plitude diagram s works reasonably well, though we caution that a careful analysis should consider the ner details of light curve structure.

Figures 6 and 7 display plots of the rst two PCA coe cients and Fourier am plitudes, respectively, for our data, plotted against each other. W hilst A_1 and A_2 are correlated with each other, PCA1 and PCA2 are not, by construction. A similar situation would occur had we plotted A_1 or A_2 against A_3 . This is another advantage of PCA analysis of variable star light curves: the di erent PCA com ponents are orthogonal to each other. A practical advantage of this feature is outlined in the next section.

4 LIGHT CURVE LUM INOSITY RELATIONS

A major goal of stellar pulsation studies is to nd form ulae linking global stellar parameters such as lum inosity or




Figure 1. Light curve reproduction using Fourier (solid lines) and PCA (dashed lines) m ethods

m etallicity to structural light curve properties. If we are interested in the V band magnitude, then we can write,

M_v = f (lightcurvestructure);

where, since we do not know the function f, we try to estimate it empirically. Two di erent approaches to quantifying light curve structure will, in general, yield di erent form ulations of the function f, but if there does exist a true underlying function f, then both m ethods should give similar answers for M_v, given the same input data. W ith a Fourier based m ethod, the function f is related to the Fourier am – plitudes and phases, A_k ; $_{k1}$, usually with a linear relation. W ith a PCA approach, we use the PCA scores plotted in grunes 2-4. Hence a PCA relation, though also linear, will be di erent. The nature of PCA in plies that the error structure in such form ulae will be simpler and we quantify this below. B oth form ulations should, of course, give similar numbers for the nal estimated value of the physical parameter in question, in this case, M_v.

 $K \, \mathbb{W} \,$ used the Fourier \mathbb{m} ethod and found relations of the form ,

$$M_v = const: 1:82 \log P \quad 0:805A_1;$$
 (4)

and,

$M_v = const: 1.876 \log P \quad 1.158A_1 + 0.821A_3:$ (5)

W e note that these relations were obtained through an iterative procedure whereby outliers were rem oved and the relations re- tted (K ovacs 2004). In this paper, we use the PCA m ethod, but also, we use the entire dataset C m entioned in KW, consisting of 383 stars, and t the relations just once.W e do not rem ove any outliers. This m ay be why we obtain slightly di erent versions of the t using Fourier parameters than that published in KW. For ease of com - parison, we include in table 2 results obtained using both PCA and Fourier parameters. This table gives the name for the relation, the independent variables considered and co-e cients together with their standard errors. The value of




Figure 2. Light curve reproduction using Fourier (solid lines) and PCA (dashed lines) m ethods

chi-squared in the table is de ned as

$${}^{k}_{X} {}^{N} (M_{v} M_{v})^{2} = (N p);$$
 (6)

where M_{ν} is the tted value of M_{ν} and N; p are the num ber of stars and parameters respectively in the t.An examination of this table strongly suggests that

1) Similar relations to equations (4) and (5) between M $_{\rm v}$ and the PCA coe cients exist.

2) We can use an F test (W eisberg 1980) to test for the signi cance of adding a second and then a third PCA parameter to the regression. The F statistic we use is

$$\frac{(R S S_{N H} R S S_{A H}) = (df_{N H} df_{A H})}{R S S_{A H} = df_{A H};}$$
(7)

where $R\,S\,S_{N\,\,H}$; $R\,S\,S_{A\,\,H}$ are the residual sum of squares under the null and alternate (NH and AH) hypothesis respectively. Sim ilarly, df_{N\,\,H} and df_{A\,\,H} are the degrees of freedom

under these two hypotheses. For this problem, the null hypothesis is that the model with the smaller number of param eters is su cient w hilst the alternative hypothesis is that them odelw ith the greater num ber of parameters is required. Under the assumption of normality of errors, equation (7) is distributed as an F $_{(df_{\rm N~H}~df_{\rm A~H}~);df_{\rm A~H}}$, (W eisberg, 1980, p. 88). Applying this F test implies rstly, that adding the rst parameter PCA1 is a signi cant addition to log P and secondly, that adding a second and third param eter, PCA2 and PCA3 are also highly signi cant with a p value less than 0.0004. In the case of Fourier parameters, adding the A₁ parameter to log P is highly signi cant and adding the A₃ parameter to this is also highly signi cant. However, a form ula involving (log P; A1; A2) has a p value of 0.0058 and a form ula involving all 3 Fourier am plitudes and log P is not a signi cant addition to a form ula involving (log P; A 1; A 3).

3) The standard deviation of the ts given in the last column is generally slightly higher for the PCA case, when considering sim ilar num bers of parameters. This is perhaps

PrincipalComponentAnalysis of RR Lyrae light curves 5




Figure 3. Light curve reproduction using Fourier (solid lines) and PCA (dashed lines) m ethods. The left panel is a fourth order (9 param eters) Fourier t and an eight order PCA (9 param eters) t. The right panel is an eight order (17 param eters) Fourier t and an eight order PCA (9 param eters) t. (

(8)





Figure 4.Plot of rst PrincipalComponent against log period.



4) The errors on the coe cients in the PCA ts are always signi cantly smaller. This is an important point when we evaluate the errors on the nal tted value of the absolute m agnitude.

5) If we write the absolute m agnitude as a function of parameters, $x_1\,;x_2\,;::;x_N$,




Figure 5.P lot of second PrincipalC om ponent against log period.

then the error on the absolute m agnitude is given by,

$${}^{2} (M_{V} + \text{const:}) = \sum_{\substack{k=1\\k=1}}^{k \times N} {}^{2} (\mathbf{x}_{k}) \left(\frac{\theta f}{\theta \mathbf{x}_{k}}\right)^{2} + \frac{1}{2} \left(\mathbf{x}_{i}; \mathbf{x}_{j}\right) \left(\frac{\theta f}{\theta \mathbf{x}_{i}}\right) \left(\frac{\theta f}{\theta \mathbf{x}_{j}}\right) = 0$$

$$(9)$$

As table 2 indicates, ² (x_k) is always smaller when the x_k are PCA coe cients rather than Fourier amplitudes. Figure 8 and 9 portray graphs of PCA1vsPCA2 and A₁ verses A₂ respectively. We note that _{i;j} (x_i) (x_j) = ² (x_i ; x_j). Table 3 presents sample correlation and covariance coe cients




Figure 6.Plot of third PrincipalComponent against log period.




Figure 7. Plot of V band amplitude against the $\mbox{ rst PCA }\infty\mbox{-}e$ cient.

between the period and PCA parameters and period and Fourier parameters. Table 3, and gures 6 and 7 demonstrate that the correlation coe cient amongst any pair of PCA coe cients is smaller than between any pair of Fourier coe cients. Hence the error on the tted value of M_v , 2 (M_v), has to be smaller when using a PCA based form ula. We can use table 3 and equation (9) to form ally calculate the error on $M_v + \text{const. Table 4 presents these results. The label in the top row of this table (P1, F1, etc.,) refers to the appropriate relation in table 2. We see clearly that the PCA form ulae do better than their Fourier counterparts with a similar num ber of parameters. When we consider the (log P; PC 1; PC 2) and (log P; A_1; A_3) variables, then the "error advantage" using a$




Figure 8. Plot of rstPrincipalComponent against second PrincipalComponent.




Figure 9. Plot of rst Fourier am plitude against second Fourier am plitude.

PCA based m ethod is a factor of two. This occurs not just because the PCA coe cients are orthogonal to each other, but also because the errors on the coe cients in a PCA based form ula are signi cantly smaller than in the Fourier case.

Figure 10 displays a plot of the predicted absolute magnitudes obtained using a two parameter $(\log P; A_1; A_3)$ Fourier t and the three parameter $(\log P; P C A 1; P C A 2; P C A 3)$ PCA t. The two approaches are displaced from each other because we do not consider the constants in this study. D is regarding this, it can be seen that




Figure 10. Plot of tted $M_{\rm v}$ + const values when using Fourier and PCA methods.




Figure 11. Plot of absolute magnitude di erence versus fractional change in light curve parameters for Fourier (open squares) and PCA (closed squares).

the slope of this plot is 1: hence the two m ethods produce sim ilar relative absolute m agnitudes.

5 CONCLUSION

W e have shown that the method of PCA can be used to study RR Lyrae light curves. It has distinct advantages over a Fourier approach because a) It is a more e cient way to characterize structure since fewer parameters are needed. A typical Fourier t requires 17 parameters whereas a PCA t may only need 9.
b) U sing the PCA approach, we see clearly why the

am plitude is a good descriptor of R R ab light curve shape. c) The di erent PCA components are orthogonal to each other whereas the Fourier am plitudes are highly correlated with each other. This leads to relations linking light curve structure to absolute m agnitude using PCA having coe cients with sm aller errors and leading to m ore accurate estim ates of absolute m agnitudes. This can reduce the form al error, in som e cases, by a factor of 2.

In the present formulation of our PCA approach, the input data is a Fourier analysis. If these input data, that is the Fourier decompositions, contain signi cant observational errors, the error bars on the resulting PrincipalCom ponents will be larger. N either the PCA or Fourier approach can compensate fully for noisy data. In this sense, the sensitivity of PCA to noisy data should be sim ilar to Fourier, though the fact that PCA is an ensemble approach in which we initially remove an average term does quard against individual points having too much undue in uence. As an exam ple, table 4 of KW gives 17 outliers (in term s of their Fourier parameters), which KW removed in their analysis relating absolute m agnitude to Fourier param eters. W e do not remove these outliers, yet, in terms of the nal tted magnitudes presented in gure 10, PCA and Fourier produce very sim ilar results. Further, even with the inclusion of these 17 stars, the PCA method still produces PCA coe cients with sm aller errors as given in tables 2 and 3.K anbur et al (2002) discuss in detail the nice error properties of the PCA method as applied to variable stars and give a recipe with which to calculate errors on PCA coe cients. Their

gure 2, albeit for Cepheids, displays error bars on these coe cients. W e see that even with noisy data, the progression of PCA parameters with period is preserved, though of course, the error bars on the PCA coe cients are larger.

N geow et al (2003) developed a simulated annealing m ethod which can reduce num erical wiggles in Fourier decom position of sparse data. N geow et al (2003) give speci c exam ples of how such an approach in proves Fourier techniues using OGLE LMC Cepheids. A similar result will hold true for RR Lyraes. Hence this annealing technique couple with a Principal C om ponent analysis should prove very useful when dealing with noisy RR Lyrae data and will be treated in detail in a subsequent paper.

OurPCA results are based on a sample of 383 stars in globular clusters. How transferable are our results and how can our results be used to obtain PC coe cients for a new RR Lyrae light curve which appears to be norm al (ie no signs of B lazhko e ects etc.)?

O ur results are transferable to the extent that the original 383 stars are a good representation of the entire population of R R ab stars, including variation in m etallicity and di erences between eld and cluster variables. G iven this caveat, we suggest two m ethods to reproduce the light curve of a new R R ab star. Firstly, it is straightforw and to include the new star in the PCA analysis with the existing dataset. T his is our recommended approach and preserves the "ensem ble analysis" property of our PCA m ethod. Our second m ethod will be the subject of future paper but brie y it is

8 Kanbur and Mariani

this.We the progression of the PCA coe cients with period, such as given in gures 4 and 5, with sim ple polynom ial functions.As an aside, we remark that gure 4 contains signi cant scatter, perhaps associated with m etallicity, so that it would be best to include m etallicity in such polynom ial

ts.For a new star, we then guess its period and read o , for that period, the value of the PCA coe cients. Equation (3) then allows us to generate the light curve. We iterate this until a speci ed error criterion is satis ed.We can then use existing form ulae relating absolte m agnitude to light curve structure as de ned by PCA. This PCA tem plate approach has been used, with considerable success, in analysing H ST C epheid data (Leonard et al 2003).

W e note from table 2 that the chisquare on the tted relations are sim ilar for PCA and Fourier. Does this mean that despite the sm aller form alerrors with PCA, both m ethods' ability to predict R R ab absolte m agnitudes is limited by the intrinsic properties of R R ab stars them selves? To some extent this is true. Jurcsik et al (2004), in analysing accurate data for 100 R R ab stars in M 3, show that for som e 16 stars, am ongst which there exist som e pairs whose absolute m ean m agnitudes di er by about 0.05 m ags (the accuracy of the photom etry is about 0.02m ags), the Fourier param eters and periods are very sim ilar. That is, an empirical method relating absolute m agnitude to period and Fourier param eters in one waveband could not distinguish between these stars. Since, as Jurcsik et al (2004) point out, their data contains a sm all range of both m ass and m etallicity, tem perature is the only other variable, it may be the case that multiwavelength inform ation is needed. It is worthwhile to investigate how PCA fares with this dataset. Here we give an outline that suggests that PCA can be more e cient at extracting information from the light curve.

For the sixteen stars which had di ering absolute magnitudes but very sim ilar Fourier parameters, we can perform the following procedure: for every pair, $j \in k$, we calculate

(a1(j) a1(k))=a1(k) + (a2(j) a2(k))=a2(k)
+ (a3(j) a3(k))=a3(k) = diff1;
 (pca1(j) pca1(k))=pca1(k)+
 (pca1(j) pca2(k))=pca2(k) = diff2;

and

(vm ean(j) vm ean(k)) = diff3;

where al(j);a2(j);a3(j) are the Fourier amplitudes and pcal(j);pca2(j) are the PCA coe cients and vm ean(j) are the m ean m agnitudes. In the above, we always take the absolute value of the di erences. W e need to take fractional changes because the Fourier amplitudes and PCA coe – cients have di erent ranges. W e now plot di 3 against di 1 and di 2. This is presented in gure 11, where the open squares are di 1 and the closed squares are di 2.W e see that with PCA (closed squares), the di erences between light curve structure param eters are greater than with Fourier (open squares). This could im ply that PCA can be m ore e cient though the limitations associated with using a single waveband are still present. A m ore rigorous, quantitative discussion of this, in a F isher inform ation sense, will be given in a future paper.

Table 1. Percentage of variation explained by PC components

	PC1	PC2	PC3	PC4	PC5	PC6
w ithout average	81.4	7.8		2.3	0.74	0.57
w ith average	96.9	1.9		0.25	0.07	0.006

In other future work we plan to investigate the applicability of this method to light curve structure-metallicity relations, RRc stars and a comparison of observed and theoretical light curves using PCA.

ACKNOW LEDGM ENTS

SM K thanks G eza K ovacs for stimulating discussions and for kindly supplying the R R ab dataset. SM K thanks D. Jono for help writing the P C A and least squares program and C. N geow for help with latex. HM thanks F C R AO for providing a sum mer internship in 2001 when part of this work was com pleted. W e also thank the referee for constructive com m ents.

REFERENCES

- Hendry, M. A., Tanvir, N. A., Kanbur, S. M., 1999, ASP Conf. Series, 167, p. 192
- K anbur, S., Iono, D., Tanvir, N. & Hendry, M., 2002, MNRAS, 329, 126
- Jurcsik, J., Benko, J., Bakos, G., Szeidl, B., Szabo, R., 2004, ApJL, 597, 49
- Kovacs, G., Walker, A.R., 2001, A&A, 371, 579
- K ovacs, G ., 2004, private com m unication
- Leonard, D., Kanbur, S., M., Ngeow, C., Tanvir, N., 2003, ApJ, 594, 247
- N geow, C., K anbur, S.M., N ikolaev, S., Tanvir, N., H endry M., 2003, 586, 959
- Tanvir, N.R., Hendry, M.A., Kanbur, S.M., 2004, in preparation
- W eisberg, S., 1980, Applied Linear Regression, John W iley & Sons, 1st Ed.

	log P		rst		second		third		chisquare	
PCA										
Р0	-1.134	0:059							0.00321	
Ρ1	-1.550	0:082	0.269	0:038					0.00283	
Р2	-1.609	0:082	0.290	0:038	0.291	0:082			0.00274	
Р3	-1.744	0:088	0.329	0:039			-0.539	0:107	0.0027	
P4	-1.829	0:088	0.359	0 : 039	0.336	0 : 079	-0.583	0:105	0.00253	
Fourier										
F1	-1.677	0:083	-0.472	0 : 054					0.00266	
F2	-1.700	0:082	-0.726	0:092			0.613	0:179	0.00258	
FЗ	-1.740	0:085	-0.758	0:116	0.536	0:193			0.00261	
F 4	-1.720	0:085	-0.790	0:117	0.215	0:243	0.490	0:227	0.00258	

Table 2. Light curve lum inosity relation using PCA and Fourier m ethods.

T able 3. Sam ple correlation and covariance coe cients between period, PCA and Fourier coe cients

	log P;PCA1	log P ; P C A 2	log P;PCA3	PCA1;PCA2	P C A 2;P C A 3	PCA1;PCA3
correlation	0.631	0.099	-0.299	< 10 ⁶	< 10 ⁶	< 10 ⁶
covariance	0.0038	0.0002	-0.0006	< 10 ⁶	< 10 ⁶	< 10 ⁶
	log P ;A 1	log P ; A ₂	log P ;A 3	A ₁ ;A ₂	A ₂ ;A ₃	A ₁ ;A ₃
correlation	-0.655	-0.529	-0.562	0.926	0.931	0.902
covariance	-0.0028	-0.0012	-0.0011	0.0030	0.0013	0.0024

Table 4.Form al error on M $_{\rm V}$ + const: for PCA and Fourier relations

P1	P 2	Р3	P 4	F 1	F 2	F 3	F 4
0.0139	0.0142	0.0216	0.0240	0.0156	0.0313	0.0394	0.0311