New light on Dark Cosm os

Enrique Gaztanaga^{1;2}, Marc Manera¹, Tuom as Multam aki³ ¹ Institut d'Estudis Espacials de Catalunya, IEEC/CSIC, F. de Ciencies, Torre C5 Par 2a, UAB, Bellaterra (08193 BARCELONA)

² V isiting: IN AOE, Tonanzintla, P.O. Box 51, 7200 Puebla, M exico

³ NORDITA, Blegdam svej 17, DK-2100, Copenhagen, Denmark

19 February 2022

ABSTRACT

Recent studies by a num ber of independent collaborations, have correlated the CM B tem peratures measured by the WMAP satellite with di erent galaxy surveys that trace the matter distribution with light from the whole range of the electrom agnetic spectrum : radio, far-infrared, optical and X -ray surveys. The new data system atically nds positive correlations, indicating a rapid slow down in the growth of structure in the universe. Individual cross-correlation m easurem ents are of low signi cance, but we show that com bining data at di erent redshifts introduces in portant new constraints. Contrary to what happens at low redshifts, for a xed $_{\rm m}$, the higher the dark energy , the lower the ISW cross-correlation amplitude. At 68% con dence level, contend, the data nds new independent evidence of dark energy: = 0:42 1:22 . It also con m s, to higher signi cance, the presence of a large dark m atter component: m =0:34, exceeding the density of baryonic matter, but far from the critical value. 0:18 Combining these new constraints with the prior of a at universe, or the prior of an accelerating universe provides strong new evidence for a dark cosm os. Combination with supernova data yields $= 0:71 \quad 0:13, m = 0:29 \quad 0:04.$ If we also assume a at universe, we nd = 0.70 0.05 and w = 1.02 0.17 for a constant dark energy equation of state.

IN TRODUCTION

In the last few years a new cosm ological scenario with a signi cant sm ooth D ark E nergy (D E) com ponent has em erged. The Cosm ic Concordance Model (CCM, from now on) is a spatially at universe with baryons (b 4%). cold dark m atter ($_{CDM}$ 23%) and a signicant DE component (73%). The model is well supported by the supernova type Ia observations (SN IA) (Riess et al. 1998; Perlmutter et al. 1999), observations on large scale structure (LSS) (Tegm ark et al. 2004; Percival et al. 2001) and the cosm ic m icrow ave background experim ents (CMB), in particularby the recent W M A P experim ent (Bennett et al. 2003). The energy density of the universe seems dom inated by the unknown DE component, presenting a form idable observational and theoretical challenge. The three key observational probes m easure com plem entary aspects of the cosm ological parameter space. The SNIA indicate that the universe is accelerating but present data is degenerate for alternative cosm ological scenarios. The LSS observations constrain m but leave the D E question unanswered. Constraints from primary anisotropies in the CMB indicate that we live in a at universe but require a prior on the value of the local Hubble rate H₀ (Blanchard 2003). A ssum ing that the universe is well described by a CDM model, combining all these three observations gives us the cosm ological CCM m odel.

The Integrated SachsW olfe e ect, ISW , (Sachs & W olfe 1967) is a direct probe for the (linear) rate of structure formation in the universe. Secondary anisotropies in the CMB appear because of the net gravitational redshifts a ecting CMB photons that travel through an evolving gravitational potential . These secondary tem perature an isotropies are therefore correlated with local, evolving, structures on large scales. The correlation is negative when structures grow, as increasing potential leaves a cold spot in the CMB sky, and positive otherwise. In a at universe without DE (E insteindeSitter, or EdS, model) this cross-correlation is expected to be zero because the gravitational potential rem ains constant, despite the linear growth of the matter uctuations.

The rate of structure form ation in the universe can also be measured by galaxy peculiar velocities or galaxy redshift distortions, on very large scales through the so-call param eter determ ination (Peacock et al. 2001; Pope et al. 2004). The ISW e ect provides an independent and com plem entary probe of the same e ect. Independent, because it uses tem perature anisotropies instead of the velocity eld, and com plem entary, because of the di erent assum ptions and system atics that relate m easurem ents with theory. D espite recent advances in the size of galaxy redshift surveys such as SD SS and 2dFGRS, the spectrum of matter uctuations $P(k) / \langle k \rangle^2$ is quite di cult to measure directly over very large scales (Tegm ark et al. 2004; Percival et al. 2001;

2 Gaztanaga, Manera & Multamaki

G aztanaga & Baugh 1998). Part of the problem is that m atter correlations fall quickly to zero on scales larger than 30 M pc/h (k < 0.1 h/M pc). In contrast, uctuations in the gravitational potential go as (k) / (k)= k^2 and therefore extend over larger distances, which m akes the signal m ore detectable (see also comments to Fig. 4). The ISW cross-correlation traces the gravitational potential, , and thus provides a new window to study the largest structures, extending over several degrees in the sky or tens of M pc/h at the survey depth.

2 GROW TH OF DENSITY PERTURBATIONS

G ravitational evolution of matter uctuations, = = 1, is dependent on the cosm obgical model via the evolution of the scale factor $a = a(t) \cdot C$ om pared to a static background, a rapidly expanding background will show down the collapse of an over dense region. In the linear regime, a sm all initial perturbation $_0$ grows according to the growth factor D (t):

$$(t) = D (t)_{0}$$
 (1)

which, under quite generic assumptions, eg (Gaztaraga & Lobo 2001; Multamaki, Gaztaraga & Manera 2003; Lue & Starkman 2004), follows a simple harmonic equation:

$$\frac{d^2 D}{d^2} + 2 + \frac{H}{H^2} \frac{dD}{d} + 3c_1 D = 0; \qquad (2)$$

where = $\ln(a)$ is the conform altim e and H = H() <u>a</u>=a is the background Hubble rate (<u>a</u> and H- are proper time derivatives). For a at cosm obgical model with a generic dark energy equation of state

$$p = w (z) \tag{3}$$

we then have:

$$H^{2} = \frac{a}{a}^{2} = H_{0}^{2} \qquad m (1 + z)^{3} + e^{3} \frac{R_{z}}{0} \frac{dz^{0}}{1 + z^{0}} (1 + w (z^{0}))$$
(4)

where and m are the dark energy and dark matter densities today in units of the critical density c 3 H²=(8 G). And q_i is given by:

$$c_{1} = \frac{1}{2} \frac{H_{0}^{2} m (1 + z)^{3}}{H^{2}(z)}$$
(5)

In this paper we study two cases. A generic (not necessarily at) CDM m odelwhere DE density is constant over the evolution of the universe (w = 1); and a at CDM m odelwith a constant equation of state parameter. For those m odels we have

$$c_1 = (1=2)_m = (_m + a^3)$$
 (6)

O ne m ay choose to compare the results to the EdS m odel: = 0; $_{\rm m}$ = 1, in which case the solution to Eq. (2) is D / a. This means that grows linearly with the scale factor, / a, while the corresponding gravitational potential uctuation, =a, remains constant as the universe expands. For non EdS m odels would change during the expansion of the universe which would turn into a galaxy-CM B tem perature cross-correlation signal.

2.1 The ISW e ect

ISW temperature anisotropies are given by (Sachs & W olfe 1967):

$$4 T_{T}^{ISW} (\hat{\mathbf{n}}) \frac{T(\hat{\mathbf{n}}) T_{0}}{T_{0}} = 2 dz \frac{d}{dz} (\hat{\mathbf{n}};z)$$
(7)

where is the Newtonian gravitational potential at redshift z.0 neway to detect the ISW e ect is to cross-correlate tem – perature uctuations with galaxy density uctuations projected in the sky (C rittenden & Turok 1996).0 n large linear scales and sm all angular separations, the cross-correlation $w_{T_{\rm G}}^{\rm ISW}$ () = < 4 $_{\rm T}^{\rm ISW}$ (\hat{n}_1) _G (\hat{n}_2) > is (Fosalba & G aztamaga 2004):

$$w_{TG}^{ISW}() = \frac{1}{2} \frac{Z}{c} \frac{dk}{k} P(k)g(k)$$

$$g(k) = dz W_{ISW}(z) W_{G}(z) \frac{H(z)}{c} J_{0}(k r_{A})$$

$$W_{ISW}(z) = 3_{m} (H_{0}=c)^{2} \frac{dP(z)=a}{dz}$$
(8)

$$W_{G}(z) = b(z)_{G}(z)D(z);$$

where J_0 is the zero order B essel function, $_{\rm G}$ is the survey galaxy selection function along the line of sight z and $r_{\rm A}$ = $r_{\rm A}$ (z) the com oving transverse distance. The power spectrum is P (k) = A $k^{n_{\rm S}}$ T 2 (k), where $n_{\rm S}$ ' 1^1 and T (k) is the CDM transfer function, which we evaluate using the tting form ulae of E inseintein & H u 1998.W em ake the com - m on assumption that galaxy and matter uctuations are related through the linear bias factor, $_{\rm G}$ (f;z) = b(z) m (f;z).

For the CDM case the ISW e ect is non-zero, and the kernel W $_{\rm ISW}$ can be well approxim ated by W $_{\rm ISW}$ (z) = 3 $_{\rm m}$ (H $_0$ =c)^2D (z) (f 1), where f is the relative growth factor, f ' $_{\rm m}$ (z) $^{6=11}$. W $_{\rm ISW}$ decreases as a function of increasing redshift and goes to zero both for $_{\rm m}$! 0 and for $_{\rm m}$! 1. At low redshifts, the ISW e ect is larger for larger values of $_{\rm m}$, but the redshift evolution depends on the curvature (ie how quickly the H and D evolve to the EdS case). This is illustrated in Fig.1 which shows how W $_{\rm ISW}$ depends on z for di erent values of and $_{\rm m}$. At high redshifts, the ISW amplitude.

In Figures 2 and 3 we also shown for a given at cosmology model the dependence of the W $_{\rm ISW}$ on redshift and on the equation of state parameter w. For a given redshift and $_{\rm m}$ there exists a maximum of W $_{\rm ISW}$ around w = 0.5. Thism aximum would translate into a maximum in the cross-correlation signal w $_{\rm TG}$. If data turns out to be greater than this maximum this would clearly disfavor models with constant equation of state.

 $^{^1}$ Throughout the paper we m ade the assumption of scale invariant primordial uctuations (n $_{\rm s}$ ' 1). For other possibilities see, eg, (Barriga etal 2001).



Figure 1.Redshift dependence of W $_{\rm ISW}$ (z) in Eq.[8] for di erent values of $_{\rm m}$ and .Bottom left, top right and top left panels shows a xed $_{\rm m}$ = 0:5, $_{\rm m}$ = 0:3 and $_{\rm m}$ = 0:1 respectively. In all cases: = 0:0 (dotted blue line), = 0:7 (continuos black line) and = 1:0 (dashed red line).Bottom right panel shows a xed = 0:7 and $_{\rm m}$ = 0:3 (continuos black line), $_{\rm m}$ = 0:5 (dotted blue line) and = 0:1 (dashed red line).



Figure 2. Redshift dependence of W_{ISW} in eq.[8] for at models with constant equation of state. Right panel shows a xed m = 0.2 and left panel m = 0.3. In both cases w = 2 (black continuous line), w = 1.5 (red dashed line), w = 1 (green dot-dashed line), w = 0.5 (blue doubledot-dashed line), and w = 0 (brow n dotted line)

2.2 Bias Self-calibration

Linear bias is used to study how well light traces the underlying statistics of linear matter uctuations. On these very large scales, uctuations are small and linear theory works very wellboth for biasing and gravity. We remove the e ects of biasing in our parameter estimation by comparing the observed galaxy-galaxy correlation w_{GG} , in the very same sam ples used for the cross-correlation, to the matter-



Figure 3.Dependence of W $_{\rm ISW}$ on the equation of state parameter w for at models at di erent redshifts and values of $_{\rm m}$. = 0.3 (back continuous lines), $_{\rm m}$ = 0.25 (red dashed lines), $_{\rm m}$ = 0.2 (green dotted lines). Redshifts are 0.1, 0.25, 0.5, 1.0, 1.5 from top to bottom.

m atter correlation $w_{m,m}$ predicted by each model (Fosalba, G aztanaga & C astander 2003). The e ects of bias are also redshift dependent, but given a galaxy selection function $_{G}(z)$, picked at z = z, we approximate the bias with a constant b = b(z) for that particular survey. We then have: $w_{TG} = b(z)w_{Tm}$ and $w_{GG} = b^{2}(z)w_{m,m}$, so that an e ective linear bias b can be estimated as the square root of the ratio of galaxy-galaxy and matter-matter correlation functions:

$$b = \frac{W_{GG}}{W_{mm}};$$
(9)

Such prescription has been shown to work well in a variety of galaxy models (eg see (Berlind, Naratanan & Weinberg 2001)). The values of $w_{m m}$ can be computed similar to (8) by

$$w_{mm}() = \frac{1}{\frac{2}{Z}} dkkP(k)g(k)$$

$$g(k) = dzW_{m}^{2}(z)\frac{H(z)}{c}J_{0}(kr_{A})$$
(10)

$$W_{m}(z) = _{G}(z)D(z);$$

where the only di erence between W $_{\rm m}$ and W $_{\rm G}$ is the bias factor b(z) in Eq.[9]. Note how the estimation of b in Eq.[9] depends on the normalization of the power spectrum in w $_{\rm m\,m}$. We choose to normalize each model by xing $_{\rm 8}$. To make our results independent of this normalization we will marginalize over $_{\rm 8}$ and h. Taking at priors and ranges $_{\rm 8}$ = 0.8 1:0 and h = 0:72 0:77. We compare the predictions with the observational data w $_{\rm TG}$ normalized to the CCM model bias, ie w $_{\rm TG}$ =b, where b is estimated from Eq. (9) using w $_{\rm m\,m}$ in the CCM model. Consequently, for other models, we will need to renormalize each of the theoretical predictions to the CCM model bias using a \relative bias":

 $w_{T_G}^{m \text{ od}}=b=b_r w_{tm}^{m \text{ od}}$, where $b_r^2 = w_{m m} = w_{mm}^{m \text{ od}}$ is the ratio of the concordance m odel prediction to the one in the corresponding m odel. We choose to estimate this relative bias at R = 8 M pc/h, but the actual num ber has little e ect in our nal conclusions.

3 OBSERVATIONAL DATA

Recent analysis by independent collaborations, have crosscorrelated the CMB an isotropies measured by WMAP with di erent galaxy surveys. The median galaxy redshifts expand over a decade (ie 0:1 < z < 1:0) and trace the matter distribution with light from the whole range of the electrom agnetic spectrum : radio, far-infrared, optical and X-ray surveys (see Table 1). The cross-correlation and error estim ation techniques used are also quite di erent but they yield com parable results over the scales of interest. C om pare for example the Montecarlo errors to jackknife errors in Fig.3 in Fosalba & Gaztamaga (2004). In our compilation of the di erent data sets, we average the results on xed angular scales around = 6. This corresponds to proper distances of ' 25M pc/h at z ' 0:1 and ' 100 M pc/h at z ' 1:0 in the CCM model and avoids possible contam ination from the sm all scale SZ and lensing e ects, eg see Fig. 3 in Fosalba, Gaztanaga & Castander (2003).

Radio galaxies from NVSS (Condon etal 1998) and hard X -ray background observed by HEAO -1 (Boldt 1987), have been cross-correlated with W MAP data (Boughn & Crittenden 2004a; Boughn & Crittenden 2004b), to nd a signal of 1:13 0:35 tim es the CCM model prediction at z 0:9. The di erent biases for X -rays, $b^2 = 1:12$, (Boughn & Crittenden 2003) and for radio galaxies, b = 1:3 1:7, (Boughn & Crittenden 2002) have been taken into account. A com patible signal has also been found with the NVSS data by the W MAP team (Nolta et al. 2004).

The cross correlation of W MAP with galaxies (17 < $b_J < 20$) in the APM Galaxy Survey (M addox et al. 1990) (covering about 20% of the South Galactic Cap, SGC) was found to be $w_{TG} = 0.35$ 0:13 K at scales = 4 10 with b' 1 (Fosalba & Gaztanaga 2004). The cross-correlation ofW MAP with the SDSS DR1 (Tegm ark et al. 2004) (covering about 10% of the North Galactic Cap, NGC) have been done for several subsamples (Fosalba, Gaztaraga & Castander 2003). The rst sample (z 0:3) contains 5 m illion objects classi ed as galaxies in SDSS (with r < 21and low associated error). For this sample, which has b' 1, $w_{TG} = 0.26$ 0:13 K at scales = 4 10. The high redshift sample (z 0:5) has $w_{TG} = 0:53 0:21 \text{ K and } b^2 \prime 6$. The SD SS data has also been cross-correlated with W M A P by the SDSS team (Scranton et al. 2003) using nearly 25 million galaxies in four redshift samples. Their results are sim ilar with those obtained earlier by Fosalba, Gaztanaga & Castander (2003) but no bias from galaxy-galaxy auto correlation function is given. The infrared 2M ASS Galaxy Survey (Jamet et al. 2000), with z 0:1, show a W M A P cross-correlation of 1:53 0:61 times the CCM prediction, with a bias of b = 1:18 (A fshordi, Loh & Strauss 2004).

We have selected independent measurements for which the bias CCM b (from w_{GG}) is known, so that we can applied

Z	w _{TG} =b		b	catalog, B and
0.1	0 : 70	0:32	1.1	2M ASS, infrared (2 m)
0.15	0:35	0:17	1.0	APM , optical (b _j)
0.3	0:26	0:14	1.0	SD SS, optical (r)
0.5	0:216	0:096	2.4	SD SS high-z, optical (r+ colors)
0.9	0:043	0:015	1–2	NVSS+HEAO,Radio & X-rays

Table 1. Observed cross correlation w $_{T\,G}$ =b (averaged for ' 4 10 .) of W MAP anisotropies with di erent catalogs.Errors in w $_{T\,G}$ =b includes 20% uncertainty in b.Errors in the median redshift z are about 10% .

the bias \self-calibration " proposed in section x2.2. The data is summarized in Table 1 and displayed in Fig.1. In the results below we also include a 10% uncertainty in the median redshift.W e chose the values of NVSS+HEAO-1 quoted by (Boughn & Crittenden 2004b) as representative of both the Nolta et al. (2004) and Boughn & Crittenden (2004a) analysis. For the SDSS, we chose the values in Fosalba & Gaztanaga (2004) where the CCM bias b is estimated using ${\tt Eq.(9)}$. Note how the selected samples are complementary. The samples which have large sky overlap (eg 2M ASS and NVSS+HEAO-1) have negligible redshift overlap.W hen the redshift overlap is signi cant (ie in 2MASS-APM or SDSS-NVSS could be up to 20%) the sky overlap is sm all (less than 10%). Consequently, the di erent sam ples in Table 1 have less than 1% volume in common. This is negligible, given that individual sam pling errors (which are proportional to volum e) are of the order of 30% .

The most signi cant detection in Table 1 seems to be the one quoted by Boughn & Crittenden (2004b) for the NVSS+HEAO-1 samples. Given the system atic uncertainties involved in the bias and selection function of both of these samples, we have checked that our results do not changed much (less than 20% in the area of the contours in Fig.3) when we double the quoted errorbar. D oubling this errorbar corresponds to an additional 50% system atic uncertainty in the value b or to a 40% uncertainty in the median redshift of the samples.

The observational data not included in Table 1 is in good agreem ent with the values in the table, but is excluded to avoid redundancy. The agreem ent of the redundant data provides further con rm ation and indicates that errors are dom inated by sam pling variance rather than by the m ethodology or the system atics.

4 RESULTS

Fig.4 compares the w_{IG} observations with predictions for a xed value of $_{\rm m}$ = 0.3 and three di erent values of . W e can see how the shape of the prediction depends on the am ount of dark energy. E ven though W _{ISW} at z = 0 depends only weakly on , the evolution with redshift dependsmore strongly on .For a xed $_{\rm m}$, models with larger values of

evolve m ore rapidly with redshift to the EdS case, where the ISW e ect vanishes. Thus, contrary to what happens at z = 0, the lower the value of (for a xed _m) the larger the ISW amplitude at high redshifts (see also Fig.1).



mean redshift (\overline{z})

F igure 4. Symbols with error bars correspond to the di erent m easurem ents $w_{TG} = b$ in Table 1.As an illustration of the shape, the continuous, short-dashed and long-dashed lines show the concordance ($_m = 0:3; = 0:7$), opened ($_m = 0:3; = 0:0$) and closed ($_m = 0:3; = 1:1$) m odel predictions (at = 6). The dotted line corresponds to the galaxy-galaxy prediction (and also the dust contam ination m odel). All lines have arbitrary norm alization.

To test m odel prodictions with the data, we use a standard ²-test, ² = $(O_i - T_i)^2 = {2 \atop i}$, where O_i and i correspond to the di erent m easurements and errors and T_i correspond to the m odel. The label i runs for i = 1 to i = 5 m arking the di erent data points (column 1 in Table 1) as we move in redshift. In order to take into account the error in the median redshift we take:

$${}^{2}_{i} = {}^{2}_{w} + {}^{2}\frac{d(w_{tg}=b)}{dz} {}^{2}_{z}$$
 (11)

where z and w are the errors in the w_{TG} =b and z respectively (see Table 1). We use the relative ² values, z^{2} to do no one dones buck in parameter of z

 $_{\rm m\ in}^2$, to de ne con dence levels in parameter estimation. Top panel of Fig.5 shows the resulting con dence contours. Taking $T_i = 0$ we evaluate the signi cance of the combined ISW detection. We nd that this null hypothesis is rejected with a very high probability: P ' 99:997% (from P $_{=4}$ (2 > 26) ' 3 10 5). We next compute the expected ISW e ect and com pare it with the observational data within the CDM family of models, where m, and h 2 are free parameters (we x the baryonic content $_{\rm b}$ ' 0:05 and the primordial spectral index $n_{\rm s}$ ' 1). We choose to norm alize each model by xing 8. To make our results independent of this norm alization we will marginalize $_8$ over the range $_8 = 0.8$ 1.0 (at prior used). As we com pare w_{GT} norm alized to the CCM model bias, we need to compute the relative bias for other LCDM models (see section x2.2). We choose to estim ate this at R = 8M pc=h, but the actual number has little e ect in the conclusions. We have also marginalized over h in a at prior range h = 0:72 0:77). Our results are not very sensitive to the

 2 WeuseH $_{0}$ 100 h km/s/Mpc.



Figure 5.0 ne, two and three sigm a con dence contours in the ($_m$;) plane (m arginalized over h) for the CDM m odel.Top: constraints from only ISW .Bottom : constraints from SN Ia (blue) and ISW (green) along with the combined contours (purple).

ranges used for $_{8}$ and h:increasing these ranges by a factor of two change our contours in less than 20 % .

The best tusing only ISW data corresponds to m'' 0:82 0:40, in good agreement with 0.26 0:08, other cosm ological probes m entioned above. Bottom panel of Fig. refcom binedconts we show the con dence contours for a CDM model along with the constraints from recent SN IA (Barris et al. 2004) observations. From the gure it is clear how the ISW e ects gives new com plem entary inform ation about the cosm ological param eters. The EdS m odel is ruled out to high signi cance. The con dence contours are alm ost perpendicular to the SN IA contours, allowing to constrain the parameter space of the model well with just these two observations. Combination of ISW with supernova $= 0.71 \quad 0.13 \text{ and } m = 0.29 \quad 0.04.$ data yields

4.1 Uncertainties in the selection function

W e explore here how robust are our results to the uncertainties in the galaxy selection function.W e take a generic parametric form of the type:

$$(z)dz = \frac{1}{(\frac{m+1}{2})} \frac{z^{m}}{z_{0}^{m+1}} e^{-\frac{z}{z_{0}}} dz$$
(12)

so that it is normalized to unity. Parameters and m control the shape of the function and are treat as x parameters; z_0 is being changed accordingly to the median redshift z we want for the selection function. When computing our results we use = 1.5 and m = 2, in which case $z = 1.41z_0$.



F igure 6. Two di erent selection functions with the same median redshift z = 1:41. Both have the generic form given by the equation [12]. The black continuous line corresponds to = 1:5 and m = 2 while the red dashed line is for = 2:5 m = 4

In order to clarify the role of the selection function shape we recalculate our results with a much more peaked selection function. This second selection function have = 2.5 and m = 4 and it is plotted together with the ducial one in Figure 6.B oth cases have the same median redshift z = 1:41. Top panel of Fig.7 shows the contours in the ($_m$;) plane for the more peaked selection function (with = 2.5 and m = 4). The contours are similar to the ducialm odel (ie com pare to Fig.5) but favoring slightly lower values for and $_m$.

Besides the uncertainty on the shape of the selection function there is also uncertainty in the median redshift. We have checked what happens if this uncertainty is not taken into account. We just set the redshift errors $_z = 0$ in Eq.[11]. Contours for the ($_m$;) plane are plot in the bottom panel of gure 7, which are also to compare with gure 5. There is hardly any di erence because the theoretical values of wrg change very little within the median redshift error range.

4.2 Equation of state

The ISW e ect can also be used to constrain the dark energy equation of state parameter. In this case, as suggested by the CCM, we assumed a at universe. We focus on a constant w parameter and maintain the same at priors for h and $_8$ (0:72 < h < 0:77 0.8 < $_8$ < 1:0). Top panel of F igure 8 show the one,two and three sigm a contours for the ($_m$;w) plane using only the ISW data. Join contours with the SN Ia data are shown in the bottom panel of F igure 8. Both datasets are also com plan entary for the w determ ination. The SN Ia data is from (Barris et al. 2004).

Making a join ISW + SN Ia analysis with the at prior reduces notably the allowed space for the parameters to $w = 1.02 \quad 0.17$ and $= 0.70 \quad 0.05$. The contours are comparable with other analysis in literature (Sandvik et al 2004) which combines SN Ia with W M AP and SD SS data.



Figure 7.0 ne, two and three sigm a con dence contours in the ($_m$;) plane for the CDM m odel. Top panel: contours using a more peaked selection function (= 2.5, m = 4) but with the same m ean redshift as the ducial case (ie com pare to Fig.5). Bottom panel: contours when errors in the median redshift of the selection functions are neglected.

The results we found are still in full agreem ent to the CCM with ' 0:7 and w = 1.

4.3 Possible Contam inants

The constraining power of the new ISW data comes from the simultaneous tting of data at dierent redshifts, that is from the shape information in Fig.4.Because of the uncertainties in the relative normalization due to a relative bias, any given point alone does not constrain well the cosm ological parameters. But the combination of the data gives us a new powerful tool for cosm ological parameter estimation.

The shape of the curve as a function of redshift also provides an important test for system atics. CMB and galaxy m aps are both m asked and corrected from galactic absorption/extinction, but any residual contam ination could produce a cross-correlation signal. Em ission and absorption by our own galaxy produce patchy hot spots in the CMB m aps and negative density uctuations in the galaxy distribution (because of extinction). In principle, this should therefore result in a negative cross-correlation, but overcorrecting for the e ects of galactic absorption could also result in a positive signal. This possibility have been tested for each of the sam ples, by comparing the cross-correlation to W MAP m aps at di erent frequencies. M ost analysis use the W M A P Kp0 m ask, which excludes about 30% of sky on the basis of galactic or extra-galactic (eg radio sources) contam ination. In all cases the contam ination seems smaller than the errors



F igure 8.0 ne, two and three sigm a con dence contours in the (;w) plane (m arginalized over h and $_8$.Top:constraints only from ISW from ISW .Bottom:constraints from SN Ia (blue) and ISW (green) along with the combination (purple).

(eg see Fig. 2 in Fosalba & Gaztanaga (2004)). Moreover, one does not expect this e ect to have any redshift dependence, contrary to the measurem ents in Fig. 4.

Cold dust in distant galaxies, will also produce patchy hot spots in the CMB maps and positive density uctuations in the galaxy distribution (could also be negative because of internal extinction). The resulting cross-correlation should trace the galaxy-galaxy auto correlation function, w_{GG} , and should therefore have a very di erent redshift dependence to the ISW e ect. The dotted line in Fig. 4 shows the predicted shape dependence for $w_{G,G}$ contam ination with arbitrary norm alization. The shape is clearly incom patible with the actual cross-correlation m easurem ents. It is also worth noting how w_{GG} goes quickly to zero at z ' 0.2, while the ISW cross-correlation remains positive. This is due to the fact that at these corresponding large scales, > 40 M pc/h, m atter-m atter correlations w_{mm} e ectively decays to zero, while w_{TG} , which traces the gravitational potential, has a less rapid decay with distance.

5 CONCLUSION

The cross-correlation of CMB anisotropies with very different galaxy surveys provides consistent detections. Their combination follows the CCM predictions with a probability of only ' 3 10 ⁵ for being a false detection. This provides new and independent evidence for dark energy and dark matter, ruling out the EdS m odel to a high signi cance (for any value of H₀). Combination with SN IA data results in

strong constraints to $= 0.71 \ 0.13$ and $_{\rm m} = 0.29 \ 0.04$. This in good agreement with the at universe $_{\rm m}$ + ' 1 found independently by CMB data (Bennett et al. 2003; Teqm ark et al. 2004). If we assume a at universe, we nd

= 0.70 0.05 and w = 1.02 0.17 for a constant dark energy equation of state. The data shows, for the rst time, statistical evidence of a recent slow down in the growth of structure formation on linear scales, just as expected in a at accelerated universe. The new ISW constraints rely in a totally di erent physicale ect that previous cosm ological constraints, providing new light on a dark cosm os.

N ote added in proof: A fler this paper was originally subm itted to astro-ph (astro-ph/0407022) a related analysis using our data com pilation have been published by C orasanniti, G iannantonio and M elchiorri (astro-ph/0504115).

ACKNOW LEDGEMENTS

A cknow ledgem ents: W e acknow ledged support from the Spanish M inisterio de Ciencia i Tecnologia, project AYA 2002-00850 with EC-FEDER funding, and from the Catalan D epartam ent d'Universitats, Recerca i Societat de la Inform aci.

REFERENCES

- A fshordi, N., Loh, Y., Strauss, M.A., 2004, Phys. Rev. D 69, 083524
- A fshordi,N ., 2004 astro-ph/0401166.
- Barriga, J., Gaztanaga, E., Santos, M. G., Sarkar, S., 2001, MNRAS 324, 977
- Barris, B.J. et al., 2004, ApJ 602, 571.
- Bennett C L. et al., 2003, ApJ Suppl., 148,1
- Berlind, A., Naratanan, V., Weinberg, D., 2001, ApJ 549, 688
- B lanchard, A ., D ouspis, M ., R ow an-R obinson, M ., Sarkar, S ., 2003, A & A , 412, 35.
- Boldt, E., 1987, Phys. Rep. 146, 215.
- Boughn, S., Crittenden, R. 2004, Nature 427, 45
- Boughn, S., Crittenden, R. 2004, astro-ph/0404470
- Boughn, S., Crittenden, R. 2003, astro-ph/0305001

Boughn, S., Crittenden, R. 2002, Phys. Rev. Lett. 88, 021302

- Condon J.J.et al., 1998, AJ 115, 1693
- Crittenden, R.G. Turok, N., 1996, Phys. Rev. Lett. 76, 575.
- D.J.Einseinstein & W.Hu, 1998, Astrophys. J. 496, 605
- Fosalba, P.Gaztanaga, E., 2004, MNRAS 350, 37
- Fosalba, P.Gaztanaga, E., Castander, F.J., 2003, ApJ 597, L89
- Gaztanaga, E. & Baugh, C. M., 1998, MNRAS, 294, 229
- Gaztanaga, E., & Lobo, J.A., 2001, ApJ 548, 47
- Jarret, T.H., et al., 2000, AJ 119, 2498
- Lue, A., Scoccim arro, R., Starkm an, G., 2004, Phys. Rev. D, 69, 044005
- M addox, S. J., E fstathiou, G., Sutherland, W. J., Loveday, J., 1990, M N R A S 242, 43P
- M ultam aki, T ., G aztamaga, R ., M anera, M ., 2003, M N R A S 334, 761
- Nolta, M.R. et al. 2004, ApJ 608, 10
- Peacock, J.A. et al., 2001, N ature, 410, 169
- PercivalW J.et al, 2001, MNRAS, 327, 1297
- Perlm utter, S. et al., 1999, ApJ 517, 565
- Pope, A.C. et al., 2004, ApJ, 607, 655
- Riess A.G. et al., 1998, AJ 116, 1009

8 Gaztanaga, Manera & Multamaki

Sachs,R.K.,Wolfe,A.M., 1967,ApJ 469,437
Sandvik H B., Tegm ark M.,W ang X., andZaldarriaga M., Phys.
 Rev.D 69,063005
Scranton,R.et al., 2003 astro-ph/0307335.
Tegm ark, M.et al., 2004, ApJ, 606, 70