Revisiting the M agni cation of Type Ia Supernovae with SDSS

Brice Menard and NealDalal

Institute for Advanced Study, E instein D rive, P rinceton N J 08540 U SA

19 February 2022

ABSTRACT

We cross-correlate the sample of Type Ia supernovae from Riess et al. (2004) with the SDSS DR2 photom etric galaxy catalogue. In contrast to recent work, we nd no detectable correlation between supernova magnitude and galaxy overdensity on scales ranging between 1 10° . Our results are in accord with theoretical expectations for gravitational lensing of supernovae by large-scale structure. Future supernova surveys like SNAP will be capable of detecting unambiguously the predicted lensing signal.

Keywords:

IN TRODUCTION

W eak gravitational lensing by large-scale structure can both distort and m agnify the im ages of background sources. The form ere ect, cosm ic shear, has been detected (see R efregier 2003 for a review) and holds great prom ise for the study of matter uctuations on large scales. The latter e ect, cosm ic m agni cation, has yet to be detected convincingly but has been theoretically investigated in detail (Jain et al. 2003; M enard et al. 2003; Takada & Ham ana 2003). In particular, a num ber of authors have studied m agni cation of Type Ia supernova, which have been employed as standard candles to measure the expansion history of the universe (e.g. Tonry et al. 2003; Knop et al. 2003). The RMS magni ca-1% level on tion of supernovae is expected to be at the arom inute scales, with the largest m agni cations occurring near the centers of massive halos.

On scales larger than arcm inutes, only modest shear and magnication at the sub-percent level are expected. The theoretically expected magnitude of shear uctuations has been con med observationally in cosm ic shear surveys. A coordingly, it would be extremely surprising if observations contradicted the expected level of cosm ic m agni cation, since statistical shear and m agni cation are related in a simple way. In principle, standard candles like Type Ia supernovae should o er a means of measuring the magni cation caused by large scale structure, for example by cross-correlating supernova brightness with foreground galaxy overdensity. In practice, all but the very highest redshift supernovae (which su er the most lensing) are not expected to provide useful m easurem ents of cosmic magni cation, since even Type Ia SNe are not perfect standard candles. The observed dispersion in supernova

m agnitudes is roughly m 20%, whereas uctuations of only . 1% are expected from lensing (M enard et al. 2003; Takada & Ham ana 2003). Even ifm agni cation and galaxy overdensity were perfectly correlated, we would expect to see only a weak correlation between supernova brightness and galaxy overdensity, $r = \frac{1}{h} \frac{1}{2} = h m^2 i$. 0.05. Therefore a very large sam ple of SN Ia would be needed to observe such an e ect.

Recently, W illiams & Song (2004) have reported a highly signicant (> 99% con dence) detection of correlation between the magnitude of 55 supernovae and foreground galaxy overdensity on 10° scales, corresponding to magni cations at the 10% level. W ang (2004) has claim ed further evidence for gravitational magni cation of the R iess et al. supernovae. If con m ed, these results would confound much of the conventional wisdom on magnication e ects and would call into question previous cosm ic shearm easurem ents which detected much lower shear variance on these scales. Additionally, if such a correlation exists, it is possible to correct for it and therefore signi cantly reduce the dispersion in supernova lum inosities: the magnitude variance can be reduced by a factor $1 r^2$ (D alal et al. 2003). This would be especially important for modern supernova surveys, which can calibrate supernova m agnitudes nearly to 0.1 m ag.

In this paper, we revisit this issue by cross-correlating the most recent sample of cosm ological Type Ia SN e com – piled by Riess et al. (2004), with low redshift galaxies in the Sloan D igital Sky Survey (SD SS) photom etric catalogue. The plan of the paper is as follows. In section 2, we brie y review theoretical predictions for cosm ic m agni cation on sm all scales. Next, we describe the supernova sample and galaxy catalogue employed in our analysis. Results are presented in x4, and our conclusions are discussed in x5.

2 Menard & Dalal

2 COSM IC MAGNIFICATION

The large scale distribution of matter in the Universe can give rise to measurable gravitational lensing e ects (shear and magnication) on background objects. The magnication factor depends on whether matter along the line of sight is preferentially over-or underdense compared to the mean.

The local properties of the gravitational lens m apping are characterised by the convergence , which is proportional to the surface m ass density projected along the line-of-sight, and the shear , which is related to the gravitational tidal eld of the lensing m ass distribution. The SD SS population of galaxies with m $_{\rm r}$ < 21 has a m edian redshift near 0.3 (e.g. D odelson et al. 2002), so that angular scales of arcm inutes correspond to length scales of M pc. On these scales, the convergence and shear uctuations are small, j j j j 1. To leading order the m agni cation is approxim ately

$$= 1 + 2 + 0 \left({2 ; j ; j} \right);$$
(1)


see M enard et al. (2003) for higher order corrections. Observable e ects are due to departures from the m ean value of the m agni cation. It is therefore convenient to use the overm agni cation = 1. Then, the cross-correlation between and the foreground m atter overdensity traced by galaxies at an angular separation reads:

h gali() 2 h
$$_{DM}$$
 i()
 $\frac{3}{2} \circ \frac{H_{0}}{c}^{2Z} \frac{dw}{a(w)} \frac{D_{d} D_{ds}}{D_{s}} W_{gal}(w)$
 $\frac{1}{2} \frac{sds}{2} P \frac{s}{f_{v}(w)}; w J_{0}(s): (2)$

In this expression, w is the comoving distance along the line-of-sight, D $_{\rm d;s;ds}$ are angular diam eter distances to lens, source, and from lens to source respectively, W gal is the normalised distribution of galaxies along the line-of-sight, a is the scale factor and P is the matter power spec-= 0:19 and the tting function trum, calculated using of Peacock & Dodds (1996), and we assume that the linear bias $b_{gal} = 1$ for the galaxies and scales of interest (Verde et al. 2002). In principle, the bias of the detected galaxies should vary with redshift, but because the galaxy distribution is dom inated by a narrow redshift interval, the use of a constant bias seems reasonable. If instead of the two-point correlation, we wanted the cross-correlation with galaxy overdensity smoothed on scale we would replace J_0 (s) by $2J_1$ (s)=s .

Figure 1 shows the amplitude of this cross-correlation betw een and gal (Eq. 2). The solid line shows h gali com puted for di erent angular separations and the dashed line shows the integrated correlation, i.e. the correlation between supernova magnitude and the number of galaxies within radius . For estimating these quantities we have used a galaxy redshift distribution peaking at z = 0.3 and a WMAP cosmology with m = 0.27, = 0:73, n = 1,and $_8 = 0.9$ (Spergel et al. 2003). Black lines have been computed for supernova redshifts of $z_s = 0.6$ (the median redshift of our sample) and gray lines for $z_s = 2.$ A s can be seen in this gure, magnications of . 1% can be generated by structures on scales of a few arom inutes or larger.

As an aside, it is perhaps worth noting that high redshift sources are expected to be more strongly perturbed



F igure 1.C ross-correlation between gravitational magnication and galaxy overdensity as a function of angular scale. A linear bias of unity has been used in this calculation.

by lensing uctuations, both because the lensing e ciency increases with redshift, and because more distant sources are subject to lensing from structures along a greater path length. For example, sources at z = 2 are typically magnied by 2% by structures on scales of f. This has ram i cations for pencil-beam surveys attempting to constrain dark energy with supernovae at high redshift, z 1:5. For example, R iess et al. (2004) report the discovery of 16 SN Ia in the GOODS eld, over an area of 0.1 sq. degrees. Lensing uctuations are signi cantly correlated across this narrow

eld. If these high redshift supernovae played an important role in the constraints derived by R iess et al., then their derived dark energy parameters could be signi cantly biased, for example by 5% in w. Fortunately, the 16 high redshift GOODS supernovae were negligible within the larger sam – ple of 177 SN e reported by R iess et al., im plying that their dark energy constraints were not appreciably biased. Future wide area surveys, like the SNAP survey which expects to cover 20 sq. degrees, should also be undisturbed by correlated structure across their elds.

3 THE DATA

W e attempt to detect a correlation between supernovam agnitude and galaxy overdensity using following data sets:

For supernovae, we use the \gold" sample of 156 SN Ia com piled by R iess et al. (2004).For each supernova, the redshift, (dust-corrected) distance m odulus, and extinction correction have been provided by these authors. The SN redshifts range from z = 0.014 to 1.75 with an average value of 0:6.

For foreground galaxies, we use the second data release (DR2) of the Sloan D igital Sky Survey (SDSS). As shown by Scranton et al. (2002) this dataset provides an e cient star/galaxy separation for objects with r < 21 and a homogeneous density by requiring a seeing value greater than $1.5^{\circ0}$.

O fthe 156 SN e listed by R iess et al., 55 fallw ithin the SD SS D R 2 footprint. G iven the magnitude cut mentioned above, the average galaxy density is roughly 1 arcm in 2 . The list of SN Ia that we use is given in table 1.

W illiams & Song used magnitude cuts in each eld in order to restrict their galaxy counts to low redshifts objects and maxim ize the signal. In our case, the bulk of SD SS photom etric galaxies are at redshifts z . 0:3, while the median SN redshift in our sample is $z_{m\,ed} = 0.6$. A coordingly, the num ber of galaxies that will overlap the supernova redshift distribution of our sample will be largely negligible.

4 ANALYSIS

For each of the 55 SN Ia we rst count the num ber of neighbours within an angular radius . The correlation between this number of neighbours and the absolute SN Ia m agnitude is shown in Fig. 2. The left panel shows, as a function of supernova m agnitude, the distribution of galaxies located within 10° from the SN Ia position, i.e. the angular scale where W illiams & Song (2004) found the strongest correlation. The horizontal error bars correspond to the m easurement error in the SN Ia m agnitudes quoted by R iess et al. (2004), and the vertical errors correspond to the Poisson noise of the galaxy counts. There is no apparent correlation in this scatter plot; this will be quantied below. The right panel shows the same correlation but using stars instead of galaxies and allows us to check for possible system atics.

For each angular bin, we have computed the crosscorrelation coe cient

$$r = \frac{p (N N) (M M)}{h(N N)^{2} ih(M M)^{2} i};$$
(3)

where N is the number of neighbours (galaxies or stars) within a given angular separation from the supernova and M is the SN absolute m agnitude. Neglecting the m easurem ent uncertainties in each object, and accounting only for the scatter in the population, the error in the cross-correlation coe cient reads

$$r = \frac{1}{P} \frac{r^2}{N_{SN}}; \qquad (4)$$

where N_{SN} is the number of SN Ia. For cells of 10^{0} radius, we nd r = 0.03 0.13 and r = 0.04 0.26 for galaxies and stars respectively. The result obtained by W illiam s & Song (2004) is therefore not reproduced in our analysis: no significant correlation is detected.

In Fig. 3 we plot the cross-correlation \cos cient r as a function of . In this gure, we have counted galaxies within an annulus centred on each , so that the plotted errors are uncorrelated. For both galaxies and stars, no signi cant correlation has been detected on any scale.

5 D ISC U SSIO N

U sing the \gold" sample of type Ia supernovae given by R iess et al. (2004) and the galaxy distribution m easured by SD SS D R 2, we have investigated correlations between the brightness of the supernovae and the density of foreground galaxies, on scales of $1-10^{\circ}$. O ur analysis does not reproduce




Figure 3.C ross-correlation coe cient r between SN m agnitude and galaxy counts (solid) or star counts (dashed), as a function of radius. The overplotted curves show the theoretical prediction for source redshifts of 0.6 and 2.


the recent results of W illiam s & Song (2004), who detected a cross-correlation at high signi cance (> 99% con dence) on 10^{0} scales. Our result is in line with theoretical predictions: the . 1% rm s m agni cation caused by large-scale structure on arcm inute scales is negligible com pared to the observed 20% dispersion in supernova m agnitudes.

The number of SN e used by W illiams & Song closely m atches the number used in our analysis. The main difference between the two analyses is that W illiams & Song derive galaxy counts from the APM catalogue, whereas we have used counts from the SD SS photom etric catalogue. W hile the APM catalogue is based on scans of photographic plates taken with the UK Schmidt telescope, the SD SS uses m odern CCD s with high-quality in aging reaching down to m uch deeperm agnitudes, providing a higher galaxy density. Therefore we expect the SD SS to be m ore suitable for investigating weak correlations.

Thus there is no evidence for a signi cant correlation between supernova brightness and foreground galaxy overdensity. Such a correlation would be expected for a sample of weakly lensed supernovae. In this light, the recently claim ed detection of signi cant weak lensing in the R iess et al. (2004) sample by W ang (2004) is som ew hat surprising. Further exam ination of the data presented by W ang indicates that the e ect appears to be dom inated by a few outliers in the magnitude distribution, which W ang ascribes to magni cation factors 2, i.e. strong lensing. This would imply a superisingly high strong lensing rate (5%) am ong high z supernovae, inconsistent with the strong lensing rate (10^{-3}) of distant quasars (B row ne et al. 2003).

Lastly, we note that the theoretically expected SNgalaxy correlation (solid curves in gure 3) would be detectable with larger samples of supernovae with sm aller ux errors. A lthough the signal is not currently detectable using the R iess et al. (2004) supernovae and SD SS galaxies,

4 Menard & Dalal



F igure 2. Scatter plot of residual magnitude vs. num ber of galaxies (left) or stars (right) within 10^0 .

future datasets (e.g. SNAP) should unambiguously exhibit the lensing signal on small scales.

ACKNOW LEDGMENTS

BM is supported by a fellow ship of the F.G ould foundation. ND is supported by NASA through Hubble Fellow ship grant # HST HF-01148.01-A, awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for R esearch in A stronom y, Inc., for NASA, under contract NAS 5-26555.

REFERENCES

- B landford, R.D., Saust, A.B., B rainerd, T.G. and Villum sen, J.V., 1991, MNRAS, 251, 600
- Browne, I.W .A., et al. 2003, MNRAS, 341, 13
- Dalal N., Holz D.E., Chen X., and Friem an J.A. 2003, ApJ, 585, L11
- D odelson, S., et al. 2002, ApJ, 572, 140
- Jain, B. and Scranton, R. and Sheth, R.K. 2003, MNRAS, 345, 62
- Knop, R.A., et al. 2003, ApJ, 598, 102
- M enard, B., Ham ana, T., Bartelm ann, M., & Yoshida, N. 2003, A & A, 403, 817
- Peacock, J.A. & Dodds, S.J. 1996, MNRAS, 280, L19
- Refregier, A. 2003, ARA & A, 41, 645
- Riess, A. G., Strolger, L., Tonry, J., Casertano, S., Ferguson, H. C., M obasher, B., Challis, P., Filippenko, A. V., Jha, S., Li, W., Chomock, R., Kirshner, R. P., Leibundgut, B., Dickinson, M., Livio, M., Giavalisco, M., Steidel, C. C., Benitez, N., & Tsvetanov, Z. 2004, accepted to ApJ, astro-ph/0402512
- Scranton, R., et al. 2002, ApJ, 579, 48
- Spergel, D. N., Verde, L., Peiris, H. V., K om atsu, E., N olta, M. R., Bennett, C. L., H alpern, M., H inshaw, G., Jarosik, N., K ogut, A., Lim on, M., M eyer, S. S., Page, L., Tucker,

G.S., W eiland, J.L., W ollack, E., & W right, E.L. 2003, ApJS, 148, 175

- Takada, M . and H am ana, T . 2003, M N R A S, 346, 949
- Tonry, J.L., et al. 2003, ApJ, 594, 1
- Verde, L., et al. 2002, MNRAS, 335, 432
- W ang, Y .2004, astro-ph/0406635
- W illiam s, L.L.R. and Song, J.2004, accepted to MNRAS, astro-ph/0403680

Magni cation of Type Ia Supernovae 5

SN	RA	DEC	redshift	0		host A $_{\rm v}$
1999dk	01 31.5	+ 14 17	0.0141	34.43	0.26	0.20
1993ae	01 29.8	-01 59	0.0180	34.29	0.23	0.00
1994M	12 31.2	+ 00 36	0.0244	35.09	0.20	0.23
1994Q	16 49.9	+ 40 26	0.0290	35.70	0.19	0.33
1998cs	16 30.7	+ 41 13	0.0327	36.08	0.19	-0.03
1994T	13 19.0	-02 09	0.0360	36.01	0.20	0.09
1997I	04 59.6	-03 09	0.172	39.79	0.18	
1999fw	23 31.9	+ 00 10	0.278	41.00	0.41	0.26
1997bj	10 42.4	+ 00 02	0.334	40.92	0.30	0.34
1996K	08 24.7	-00 21	0.380	42.02	0.22	0.02
1995ba	08 19.1	+ 07 43	0.3880	42.07	0.19	
1997am	10 57.5	-03 14	0.416	42.10	0.19	0.00
1997bh	13 44.6	-00 20	0.420	41.76	0.23	0.60
1997Q	10 56.9	-03 59	0.430	41.99	0.18	
1998ba	13 43.6	+ 02 20	0.430	42.36	0.25	
1997aw	10 23.5	+ 04 07	0.440	42.57	0.40	0.80
1997ai	10 49.0	+ 00 32	0.450	42.10	0.23	
1999	02 33.9	+ 00 33	0.455	42.29	0.28	0.19
1997P	10 55.9	-03 57	0.472	42.46	0.19	
2002dc	12 36.8	+ 62 13	0.475	42.14	0.19	0.23
1995ay	03 01.1	+ 00 21	0.480	42.37	0.20	
1996ci	13 45.9	+ 02 27	0.495	42.25	0.19	i
1997cj	12 37.1	+ 62 26	0.500	42.74	0.20	0.15
1999U	09 26.7	-05 38	0.500	42.75	0.19	0.04
1997as	08 24.2	-00 48	0.508	41.64	0.35	0.85
1997bb	12 29.0	+ 00 09	0.518	42.83	0.30	0.11
2001.iy	10 52.4	+ 57 17	0.570	42.88	0.31	-0.04
1996I	12 00.7	-00 16	0.570	42.81	0.25	0.14
1997af	08 23.9	+ 04 09	0.579	42.86	0.19	
1995ax	02 26.4	+ 00 49	0.615	42.85	0.23	i
1996H	12 28.9	-00 05	0.620	43.11	0.30	0.09
1998M	11 33.7	+ 04 05	0.630	42.62	0.24	0.75
2003be	12 36.4	+ 62 07	0.64	43.07	0.21	0.23
1997R	10 57.3	-03 55	0.657	43.27	0.20	
2003bd	12 37.4	+ 62 13	0.67	43.19	0.28	0.27
2001.ix	10 52.3	+ 57 07	0.710	43.05	0.32	0.53
1998bi	13 47.7	+ 02 21	0.740	43.35	0.30	
1997ez	08 21.6	+ 03 25	0.778	43.81	0.35	i
2001hx	08 49.4	+ 44 02	0.798	43.88	0.31	0.31
2001hy	08 49.8	+ 44 15	0.811	43.97	0.35	0.03
1999fj	02 28.4	+ 00 39	0.815	43.76	0.33	0.23
2001 j	02 28.1	+ 00 27	0.815	44.09	0.28	0.23
1996cl	10 57.0	-03 38	0.828	43.96	0.46	1
1997ap	13 47.2	+ 02 24	0.830	43.85	0.19	i
2003eq	12 37.8	+ 62 14	0.839	43.86	0.22	0,22
2001 jh	02 29.0	+ 00 21	0.884	44.23	0.19	-0.01
2003eb	12 37.3	+ 62 14	0.899	43.64	0.25	0.26
2003lv	12 37.5	+ 62 11	0.94	43.87	0.20	0.15
2002dd	12 36.9	+ 62 13	0.95	44.06	0.26	0.24
2002aa 2003es	12 36.9	+ 62 13	0.954	44.28	0.31	0.07
2003es 2002ki	12 30.5	+ 62 21	1.140	44.84	0.30	0.09
1999fv	23 30.6	+ 02 21 + 00 17	1.140	44.04	0.34	0.24
2003az	12 37.3	+ 62 19	1.265	44.19	0.20	0.24
2003az 2003dy	12 37.3	+ 62 19 + 62 11	1.340	45.20 45.05	0.20	0.25
Loosay	10 01.0	+ 62 11	1.010	45.53	0.35	0.00

Table 1. Supernovae used in our analysis. Here, $_0$ and $_$ are the distance modulus and its error. All quantities are taken from the larger \gold" sample of R iess et al. (2004).