# CONNECTING COSMOLOGY AND COLLIDERS

MARK TRODDEN Department of Physics Syracuse University Syracuse, NY 13244-1130, USA. trodden@physics.syr.edu

The broad connections between cosm ology and collider physics, particularly precision m easurem ents at the high-energy frontier, are discussed. These proceedings sum m arize a colloquium delivered to a general audience of experim ental and theoretical particle and collider physicists at the International Conference on Linear Colliders (LCW S2004) in Paris.

#### 1 Introduction

The vast majority of talks at a conference like LCW S2004 are, naturally, focused on the physics of the very small, operating in regimes in which quantum eld theory is thought to be a completely adequate description. Gravity, and particularly its application to the universe, are generally not part of the discussion. However, in this talk I would like to argue the point that particle physics and cosm ology, as disciplines independent of one another, no longer exist; that our most fundam ental questions are now the same and that we are approaching them in complem entary ways.

I will begin by inventorying the energy budget of the universe, and pointing out the places where our understanding is seriously ham pered by issues that are mly rooted in particle physics. I will then go on to describe in broad terms the current status of our approaches to these issues. In some cases, most notably dark matter and baryogenesis, a linear colliderm ay rule out or provide evidence for existing proposals. On the other hand, if this is not the case, then precision measurements of physics at the TeV scale may very well point the way to a new understanding of these fundamental cosm ological conundrum s.

Beyond these topics, I will brie y speculate on possible connections between collider experiments and one of the most esoteric cosm obgical concepts - dark energy.

G iven space constraints, I will necessarily be more brief than in the actual colloquium, and will om it my very short discussions of cosm ic rays and new -old in ation and some of the more peripheral comments. Further, my referencing will be very sparse, restricted to a few experimental results and some review articles from which the reader can nd more complete references. I apologize in advance to any colleagues who may feel slighted by this necessary decision.

#### 2 The New Cosm ological Paradigm

The data-driven revolution in cosm ology cannot have escaped the notice of particle physicists. During the last decade a host of new precision measurements of the universe have provided a clear and surprising accounting of the energy budget of the universe. There now exists compelling evidence, from multiple techniques, that the universe is composed of 5% baryonic matter, 25% dark matter and a whopping 70% dark energy, with negative pressure, su ciently negative to cause the expansion of the universe to accelerate.

The best known evidence for this comes from two sources. The rst is from Type Ia supernovae studies<sup>1;2</sup>. These data are much better t by a universe dom inated by a cosm obgical constant than by a at matter-dom inated model. This result alone allows a substantial range of possible values of  $_{\rm M}$  and

. However, if we independently constrain  $_{\rm M}$  0:3, we obtain 0:7, corresponding to a vacuum energy density 10  $^8$  erg=cm  $^3$  (10  $^3$  eV) $^4$ .

The second is from studies of the small anisotropies in the Cosm ic M icrow ave Background Radiation (CMB), culm inating in the W MAP satellite<sup>3</sup>. One very important piece of data that the CMB uctuations give us is the value of total. For a at universe (k = 0, total = 1) we expect a peak in the power spectrum at 1' 220. Such a peak is seen in the W MAP data, yielding 0:98 total 1:08 (95% c.l.) { strong evidence for a at universe.

## 3 The Baryon A sym m etry of the Universe

O ne would think that the baryonic component of the universe was well understood; after all, we are made of baryons. However, from the point of view of cosm ology, there is one fundam ental issue to be understood.

D irect observation shows that the universe around us contains no appreciable primordial antimatter. In addition, the stunning success of big bang nucleosynthesis rests on the requirement that, de ning  $n_{b(b)}$  to be the number density of (anti)-baryons and s to be the entropy density,

2:6 10<sup>10</sup> < 
$$\frac{n_b n_b}{s}$$
 < 6:2 10<sup>10</sup> : (1)

This number has been independently determined to be  $= 6:1 \ 10 \ {}^{10} + 0:3 \ 10 \ {}^{10} \ 0:2 \ 10 \ {}^{10}$  from precise measurements of the relative heights of the rst two microwave background (CMB) acoustic peaks by the WMAP satellite. Thus the natural question arises; as the universe cooled from early times, at which one would expect equalam ounts of matter and antimatter, to today, what processes, both particle physics and cosm ological, were responsible for the generation of this very speci c baryon asymmetry? (For a review and references see  $^{4;5}$ .)

If we're going to use a particle physics model to generate the baryon asym – metry of the universe (BAU), what properties must the theory possess? This question was rst addressed by Sakharov in 1967, resulting in the following criteria

V iolation of the baryon number (B) symmetry.

V iolation of the discrete sym m etries C (charge conjugation) and C P (the composition of parity and C )

A departure from therm al equilibrium.

There are m any ways to achieve these. One particularly simple example is given by G rand Uni ed theories (GUTs). However, while GUT baryogenesis is attractive, it is not likely that the physics involved will be directly testable in the foreseeable future.

In recent years, perhaps the most widely studied scenario for generating the baryon number of the universe has been electrow eak baryogenesis and I will focus on this here. In the standard electrow eak theory baryon number is an exact global symmetry. How ever, baryon number is violated at the quantum level through nonperturbative processes. These e ects are closely related to the nontrivial vacuum structure of the electrow eak theory.

At zero tem perature, baryon num ber violating events are exponentially suppressed. However, at tem peratures above or comparable to the critical tem perature  $T = T_c$  of the electrow eak phase transition, B-violating vacuum transitions may occur frequently due to therm allactivation.

Ferm ions in the electrow eak theory are chirally coupled to the gauge elds. In terms of the discrete sym m etries of the theory, these chiral couplings result in the electrow eak theory being m axim ally C -violating. How ever, the issue of CP -violation is m ore com plex.

CP is known not to be an exact sym metry of the weak interactions, and is observed experimentally in the neutralK aon system through K<sub>0</sub>, K<sub>0</sub> mixing. However, the relevant e ects are parametrized by a dimensionless constant which is no larger than 10<sup>20</sup>. This appears to be much too small to account for the observed BAU and so it is usual to turn to extensions of the minimal theory. In particular the minimal supersymmetric standard model (MSSM).

The question of the order of the electroweak phase transition is central to electroweak baryogenesis. Since the equilibrium description of particle phenomena is extremely accurate at electroweak temperatures, baryogenesis cannot occur at such low scales without the aid of phase transitions.

For a continuous transition, the associated departure from equilibrium is insu cient to lead to relevant baryon number production. For a rst order transition quantum tunneling occurs around  $T = T_c$  and nucleation of bubbles of the true vacuum in the sea of false begins. At a particular temperature below  $T_c$ , bubbles just large enough to grow nucleate. These are term ed critical bubbles, and they expand, eventually lling all of space and completing the transition. As the bubble walls pass each point in space, the order param eter changes rapidly, as do the other elds and this leads to a signi cant departure from therm alequilibrium. Thus, if the phase transition is strongly enough rst order it is possible to satisfy the third Sakharov criterion in this way.

There is a further criterion to be satis ed. As the wall passes a point in space, the Higgs elds evolve rapidly and the Higgs VEV changes from h i = 0 in the unbroken phase to h  $i = v(T_c)$ , the value of the order parameter at the symmetry breaking global minimum of the nite temperature e ective potential, in the broken phase. Now, CP violation and the departure from equilibrium occur while the Higgs eld is changing. A flerwards, the point is in the true vacuum, baryogenesis has ended, and baryon number violation is exponentially supressed. Since baryogenesis is now over, it is in perative that baryon number violation be negligible at this tem perature in the broken phase, otherw ise any baryonic excess generated will be equilibrated to zero. Such an e ect is known as washout of the asymmetry and the criterion for this not to happen m ay be written as

$$\frac{v(T_c)}{T_c}$$
 1: (2)

It is necessary that this criterion be satis ed for any electroweak baryogenesis scenario to be successful.

In the m inimal standard model, in which experiments now constrain the H iggs mass to be  $m_H > 114.4 \text{ GeV}$ , it is clear from numerical simulations that (2) is not satistical. This is therefore a second reason to turn to extensions of the m inimalm odel.

O ne important example of a theory beyond the standard model in which these requirements can be met is the MSSM. In the MSSM there are two Higgs elds, 1 and 2. At one bop, a CP-violating interaction between these elds is induced through supersymmetry breaking. A lternatively, there also exists extra CP-violation through CKM-like elects in the chargino mixing matrix. Thus, there seems to be su cient CP violation for baryogenesis to succeed.

Now, the two Higgs elds combine to give one lightest scalar Higgs h. In addition, there are also light stops t (the superpartners of the top quark) in the theory. These light scalar particles can lead to a strongly rst order phase transition if the scalars have m asses in the correct region of parameter space. A detailed two loop calculation  $^{6}$  and lattice results indicate that the allowed

region is given by

$$m_{\rm h}$$
 120G eV (3)

$$m_t m_t;$$
 (4)

for tan  $h_2$  i=h\_1i > 5. In the next few years, experiments at the Tevatron and the LHC should probe this range of H iggs masses and we should know if the M SSM is at least a good candidate for electroweak baryogenesis.

W hat would it take to have con dence that electroweak baryogenesis within a particular SUSY model actually occurred? First, there are some general predictions: If the Higgs is found, the next test will come from the search for the lightest stop at the Tevatron collider. Important supporting evidence will come from CP-violating e ects which may be observable in B physics. For these, the preferred parameter space leads to values of the branching ratio BR (b ! s ) di erent from the Standard M odel case. A lthough the exact value of this branching ratio depends strongly on the value of the and  $A_t$  parameters, the typical di erence with respect to the Standard M odel prediction is of the order of the present experimental sensitivity and hence in principle testable at the BaBar, Belle and BTeV experiments.

However, what is really necessary is to establish a believable model. For this we require precision measurements of the spectrum, masses, couplings and branching ratios to compare with theoretical requirements for a su cient BAU. Such a convincing case would require both the LHC and ultimately the LC if this is truly how nature works.

# 4 Dark Matter

Theorists have developed m any di erent m odels for dark m atter, som e of which are accessible to terrestrial experim ents and som e of which are not. There is not space to review all of these here. R ather, I will focus on a speci c exam ple that is of interest to collider physicists (for a review and references see<sup>7</sup>).

A prime class of dark matter candidates are W eakly Interacting M assive Particles (W MPs). Such a particle would be a new stable particle . The evolution of the number density of these particles in an expanding universe is

$$\underline{\mathbf{n}} = 3Hn \quad h \, \mathrm{vi}(\mathbf{n}^2 \quad n_{eq}^2); \tag{5}$$

where a dot denotes a time derivative, H is the Hubble constant, is the annihilation cross-section and  $n_{\rm eq}$  is the equilibrium value of n .

In the early universe, at high tem perature, the last term in this equation dominates and one nds the equilibrium number density of particles. If



Figure 1: The co-m oving num ber density of a dark m atter particle.

this were always the case then today we would nd negligible numbers of them and their energy density would certainly be too little to account for the dark matter. However, as the universe expands it reaches a temperature, known as the freeze-out temperature, at which the evolution equation become dominated by the rst term on the right-hand side - the damping due to the the Hubble expansion. A fler this point, annihilations cease and the distribution of

particles at that time is merely diluted by the expansion at all later times, leading to an abundance that is much higher than the equilibrium one at those temperatures. This is illustrated in gure  $1^8$ .

In fact, to a rst approximation, the dark matter abundance remaining today is given by

$$_{\rm DM}$$
 0:1  $\frac{_{\rm weak}}{_{\rm cak}}$ ; (6)

where  $w_{eak}$  is the typical weak interaction cross-section. From this one can clearly see why it is that W IM Ps get their nam  $e - w_{eak}$  interacting particles yield the correct order of m agnitude to explain the dark m atter.

W hat I have just described is a generic picture of what happens to a W IM P.O byiously, a speci c candidate undergoes very speci c interactions and a detailed calculation is required to yield the correct relic abundance. The most popular candidate of this type arises in supersymmetric extensions



Figure 2: A portion of the m SUGRA parameter space with  $A_0 = 0$ , tan = 10, and > 0.

of the standard model. Supersymmetry, of course, is attractive for entirely independent particle physics reasons. However, a natural prediction of SUSY with low-energy SUSY breaking and R-parity is the existence of the lightest superpartner of the standard model particles. This Lightest Supersymmetric Particle (LSP) is typically neutral, weakly interacting, with a weak scale mass, and hence can be a compelling dark matter candidate.

W eak scale SUSY has a large number of parameters. A detailed analysis requires us to focus on particular models. It is common to use a model – m inim al supergravity (m SUGRA) – described by just 5 parameters, the most important of which are the universal scalar mass m<sub>0</sub> and the universal gaugino m ass M<sub>1=2</sub>, both de ned at the scale M<sub>GUT</sub> ' 2  $10^{16}$ GeV.

W hat m ight the LSP be in this fram ework? As can be seen from gure 2  $^9$  the LSP is typically the the lightest neutralino or the right-handed stau  $\gamma_R$ . If it is a neutralino, it is almost purely B ino over a large region of parameter space, with a reasonable H iggsino component for m $_0$  1TeV.

It is, of course, very important to go beyond m SUGRA to understand all the possible ways for an LSP to be the dark m atter. However, m SUGRA does provide a crucial and m anageable set of com m on m odels.

If SUSY is discovered at colliders, one would like to determ ine the relic

density of such a particle to an accuracy of a few percent, in order to compare with the known dark matter abundance. This requires a precise determination of the masses and couplings in the theory, a goal that, although challenging, may well be possible with the LHC and a linear collider.

### 5 Dark Energy

As I have mentioned, it is hard to see how one might make measurements directly relevant to the dark energy problem in colliders. Nevertheless, in the interest of not giving up hope, and because we appear to be extremely ignorant about this problem, I would like to mention at least one connection between the cosm ological constant, a candidate for the dark energy, and collider physics.

In classical general relativity the cosm obgical constant is a completely free parameter. However, if we integrate over the quantum uctuations of all modes of a quantum eld in the vacuum, we obtain a natural expectation for its scale. Unfortunately this integral diverges, yielding an in nite answer for the vacuum energy. Since we do not trust our understanding of physics at extrem ely high energies, we could introduce a cuto energy, above which ignore any potential contributions, expecting that a more com plete theory will justify this. If the cuto is at the P lanck scale, we obtain an estim ate for the energy density in this component

$$_{ac}$$
 M  $_{P}^{4}$  (10<sup>18</sup> G eV)<sup>4</sup> : (7)

Unfortunately, a cosm ological constant of the right order of m agnitude to explain cosm ic acceleration must satisfy

$$v_{ac}$$
 (10 <sup>3</sup>eV)<sup>4</sup>; (8)

which is 120 orders of m agnitude sm aller than the above naive expectation.

A second puzzle, the coincidence problem arises because our best-t universe contains vacuum and matter densities of the same order of magnitude. Since the ratio of these quantities changes rapidly as the universe expands. there is only a brief epoch of the universe's history during which we could observe the transition from dom ination by one type of component to another.

To date, I think it is fair to say that there are no approaches to the cosm ological constant problem that are both well-developed and compelling (for reviews see  $^{10;11;12}$ ). In addition, given the absurdly small mass scales involved, it is generally thought unlikely that collider physics will have any impact on this problem. While I think this is probably true, I would like to emphasize a particular connection between collider experiments and this problem.

As I have mentioned, a prime motivation for the next generation of accelerators is the possibility that supersymmetry might be discovered. At the risk of insulting some of my colleagues, when one is constantly dealing with supersymmetric theories in the context of collider signatures, it is easy to forget that supersymmetry is much more than a symmetry implying a certain spectrum and speci c relationships between couplings and masses. Supersymmetry is, of course, a space-time symmetry, relating internal symmetry transformations with those of the Poincare group. There is a direct connection between this fact and the vacuum energy.

The power of supersymmetry is that for each fermionic degree of freedom there is a matching bosonic degree of freedom, and vice-versa, so that their contributions to quadratic divergences cancel, allowing a resolution of the hierarchy problem. A similar elect occurs when calculating the vacuum energy: while bosonic elds contribute a positive vacuum energy, for fermions the contribution is negative. Hence, if degrees of freedom exactly match, the net vacuum energy sums to zero.

W e do not, however, live in a supersymmetric state (for example, there is no selectron with the same mass and charge as an electron, or we would have noticed it long ago). Therefore, if supersymmetry exists, it must be broken at some scale M  $_{\rm SUSY}$ . In a theory with broken supersymmetry, the vacuum energy is not expected to vanish, but to be of order

$$_{\rm vac}$$
 M  $_{\rm SU\,SY}^4$  (10<sup>3</sup> G eV)<sup>4</sup>; (9)

where I have assumed that supersymmetry is relevant to the hierarchy problem and hence that the superpartners are close to experimental bounds. However, this is still 60 orders of magnitude away from the observed value.

It is a crucial aspect of the dark energy problem to discover why it is that we do not observe a cosm ological constant anything like this order of m agnitude. If we nd SUSY at colliders and understand how it is broken, this m ay provide much needed insight into how this occurs and perhaps provide new inform ation about the vacuum energy problem.

## 6 Conclusions

In this colloquium I have tried to argue that particle physics and cosm ology, as disciplines independent of one another, no longer exist; that our most fundam ental questions are the same and that we are approaching them in com plem entary ways. I have emphasized the deep connections between results obtained in existing colliders and expected from future ones and the puzzles facing cosm ology regarding the energy budget of the universe. From the fam iliar baryonic matter, through the elusive dark matter and perhaps all the way to the mysterious dark energy, collider experiments are crucial if we are to construct a coherent story of cosm ic history. In conjunction with observational cosm ology such experiments hold the key to unlock the deepest secrets of the universe.

### A cknow ledgm ents

I would like to thank all the organizers of LCW S2004, and in particular D avid M iller and H enri V ideau, for a wonderful conference and for their hospitality in Paris. I also owe thanks to m y co-m embers of the ALCPG W orking G roup on C osm ologicalC onnections and in particularm y co-editors M arco B attaglia, Jonathan Feng, N orm an G raf and M ichael Peskin for m any useful conversations. This work was supported in part by the NSF under grant PHY-0094122. M T is a C ottrell Scholar of R esearch C orporation.

- 1. A.G. Riess et al. [Supernova Search Team Collaboration], Astron.J. 116,1009 (1998) [arX iv astro-ph/9805201].
- 2. S. Perlm utter et al. [Supernova Cosm ology Project Collaboration], Astrophys. J. 517, 565 (1999) [arX iv astro-ph/9812133].
- 3. G. Hinshaw et al, Astrophys. J. Suppl. 148, 135 (2003) [arXiv:astroph/0302217].
- 4. M. Trodden, Rev. M od. Phys. 71, 1463 (1999) [arX iv hep-ph/9803479].
- 5. A. Riotto and M. Trodden, Ann. Rev. Nucl. Part. Sci. 49, 35 (1999) [arX iv:hep-ph/9901362].
- 6. M. Carena, M. Quiros, M. Seco and C. E. M. Wagner, Nucl. Phys. B 650, 24 (2003) [arX iv hep-ph/0208043].
- 7. J.L.Feng, eConfC 0307282, L11 (2003) [arX iv hep-ph/0405215].
- 8. G. Jungman, M. Kam ionkowski and K. Griest, \Supersymmetric dark matter," Phys. Rept. 267, 195 (1996) [hep-ph/9506380].
- 9. J. L. Feng, K. T. M atchev and F. W ikzek, \Neutralino dark m atter in focus point supersymmetry," Phys. Lett. B 482, 388 (2000) [hepph/0004043].
- 10. S.M. Carroll, Living Rev. Rel. 4, 1 (2001) [arX ivastro-ph/0004075].
- 11. P.J. Peebles and B.Ratra, arX iv astro-ph/0207347.
- 12. V. Sahni and A. A. Starobinsky, Int. J. M od. Phys. D 9, 373 (2000) [arX iv astro-ph/9904398].