# SN Ib 1990I: Clumping and Dust in the Ejecta?

A.Elm ham di<sup>1;2</sup>, I.J.D anziger<sup>3</sup>, E.C appellaro<sup>4</sup>, M.D ella Valle<sup>5</sup>, C.G oui es<sup>6</sup>, Mark M.P hillips<sup>7</sup>, M.Turatto<sup>8</sup>

<sup>1</sup> SISSA / ISAS , via Beinut 4 - 34014 Trieste - Italy

<sup>2</sup> European Southern Observatory, Karl-Schwarzschild-Str 2, 85748 Garching b. Munchen, Germany

<sup>3</sup> INAF, Osservatorio Astronom ico di Trieste, V ia G B Tiepolo 11 - I-34131 Trieste - Italy

<sup>4</sup> INAF, Osservatorio Astronom ico di Capodim onte, via Moiariello 16, I-80131 Napoli, Italy

<sup>5</sup> INAF, O servatorio A stro sico di A roetri, largo E. Ferm i 5, 50125 Firenze, Italy

<sup>6</sup> CEA /DSM /DAPN IA /Service d'Astrophysique, Saclay, Paris, France

 $^7$  Las C am panas O bservatory, C amegie O bservatories, C asilla 601, La serena, C hile

<sup>8</sup> IN AF, O sservatorio A stronom ico di Padova, vicolo dell'O sservatorio 5, 35122 Padova, Italy

Abstract. Photom etry and spectra of the type Ib SN 1990I are presented and analysed, covering about 400 days of evolution. The good quality of the data allow one to set reliably the supernova age (i.e. both date of maxim um and explosion time estimates are given). The presence of optical helium lines is shown. SN 1990I seems to show higher velocities compared to a sample of type Ib events. The nebular emission lines display a high degree of asymmetry and the presence of ne structures, suggestive of non-spherical clumping in the ejecta of SN 1990I. U sing the [O I] 6300,64 A  $\,$  ux, we estimate a lower limit on the oxygen mass to fall in the range (0:7 1:35) M  $\,$  . The oxygen mass requires a lling factor as small as  $10^2$  on day 254, indicating a highly clum py distribution of the oxygen m aterial. The amplitude and evolution of the [Ca II]/[D I] ux ratio is similar to that observed in a sam ple of SN e Ib/c events and thus suggests similar progenitor masses. A blueshift of the order 600 km s  $^1$  is reported in the [0 I] 6300,64 A after day 254. The [Ca II] 7307.5 A em ission pro le appears blueshifted as well at late epochs.We recover the quasi-bolom etric \BVRI" light curve of SN 1990I.The constructed bolom etric light curve shows a change of slope at late phases, with an e-folding time of 60 2 d in the [50:200] d time interval, considerably faster than the one of  $^{56}$ C o decay (i.e. 111.3 d), suggesting the -rays escape with lower deposition, owing to the low mass nature of the ejecta. A fier day 200, an e-folding time '47 2.8 d is measured. W hile the light curves of SN e 1990I and 1993J are similar in the [30:100] d time range, they tend to behave di erently after day 200. A simpli ed -ray deposition model is applied after adding a contribution of about 35% to the computed pseudo-bolom etric light curves to account for near-IR lum inosities to estimate the ejecta and  $^{56}$ Ni m asses (M ( $^{56}$ N i) = 0:11 M and M<sub>ej</sub> = 3:7 M ). The de cit in lum inosity is estimated to be about 50% around day 308. The observed spectral blueshift combined with the dram atic and sudden drop in the pseudo-bolom etric light curve and (B V) colour is interpreted to be a consequence of dust condensation in the ejecta of SN 1990I around day 250.

Key words. Supernovae: type Ib/c, light curve, spectra; Nucleosynthesis: <sup>56</sup>N im ass, oxygen m ass; D ust form ation

## 1. Introduction

O ne of the interesting topics in supernovae (SN e) research for the last few years has been the study of hydrogen de-

cient events, namely type 1b (helium -rich) and type Ic (helium -poor) events (W heeler & Harkness. 1990). The particular interest in these two subclasses com es from the fact that they are the least understood among supernovae varieties in their observational properties, progenitors and therefore the physics of the explosion (N om oto et al. 1990; B aron. 1992). The possible connection of the strongly energetic ones with G am m a-R ay Bursts (i.e. the em ergence of H ypernovae; W oosley et al. 1999; N om oto et al. 2000) m akes them exciting events. In addition, the appearance of the transition events such SN e 1993J and 1987K suggests a physical connection between type Ib/c and type II objects; thus one needs to understand the di erences as well as the sim ilarities am ong them.

Type Ib and Ic SN e are classi ed on the basis of their spectra. These events are hydrogen de cient at maximum light. They lack also the deep Si II absorption near 6150 A characterizing type Ia events. SN e Ic are spectroscopically sim ilar to SN e Ib at late epochs where the spectra are dom inated by oxygen and calcium em ission lines, whereas

Send o print requests to: A . E lm ham di, elm ham di@ sissa.it

at early phases SN e Ic do not show the H e I lines (W heeler Table 1. Spectroscopic observations of SN 1990I & Harkness.1990).

Di erent models, evolution and explosion, have been suggested to explain the main observational properties of SNe Ib/c. The most accepted scenario is related to the mechanism of the hydrogen rich SNe II, namely core collapse in massive stars. There are many arguments reinforcing the idea of core collapse events as the favoured in the progenitors of type Ib/c SNe:

They are associated with population I stars, and are observed to occur in spiral galaxies ongoing star form ation occurs (association with H II regions).

The presence of strong oxygen em ission lines.

The discovery of transition events (e.g. SN e 1987K and 1993J; Filippenko 1988, Swartz et al. 1993), whose spectral properties changed from type II to type 1b/c as y since the maximum light time: 29/04/1990 they aged.

The early detection of such events in the radio (e.g. SN 1983N and SN 1984L), suggests interaction with a dense circum stellar material, hence strong mass loss.

Here we present and analyse the photom etry and spectralevolution of type Ib SN 1990I. The paper is organized as follows.We describe rst the spectral behaviour of SN 1990I during both the photospheric and nebular phases. We use the [O I] 6300,64 A lum inosities to estimate the ejected mass of oxygen (Sect. 2.1.2). Then we study the light curves and colour evolution of SN 1990I. The photom etry is com pared to a selected sam ple of type lb/c, Ilb events (Sect. 2.2). We nally describe a simple model of ray deposition from radioactive decay used to the bolom etric evolution of SN 1990I, providing  $^{56}\mathrm{N}\,\mathrm{i}$  and ejecta massestimates (Sect. 3.2).

#### 2.0 bservations

SN 1990I was discovered by Pizarro et al. (IAUC 5003) in NGC 4650A. Initially it was classi ed as type Ia on the basis of an early spectrum taken by Sivan and Marseille on April 30 (IAUC 5003). Based on three early spectra (M ay 18, June 7 and June 8) taken with the Cerro Tololo 1.5-m telescope, SN 1990I has been re-classi ed as a type Ib by Phillips (IAUC 5032). The SN is located in the polar ring of its host ring galaxy NGC 4650A.

W e present spectra and photom etry covering about 400 d of evolution. The data are collected both from ESO -KP (Key Project) database and from CTIO observations.

# 2.1. Spectroscopic evolution

The 'pumalof the spectroscopic observations of SN 1990I are reported in Table 1. Figure 1 illustrates the spectroscopic evolution, covering both the photospheric and nebularphases. The well sam pled data of SN 1990I, especially light curves, allow us to estim ate the maximum light time with some con dence (see Sect. 2.2 for discussion regarding the SN age). For each spectrum the corresponding days

Date (dd/mm/yy)	Phase <sup>y</sup>	Range (A)	Instrum ent	
29/04/90	m ax	5370-9100	ESO-2.2m	
09/05/90	+ 12d	3610-9510	ESO-NTT	
18/05/90	+ 21d	2950–7620	CTIO-1.5m	
07/06/90	+ 43d	2950–7600	CTIO-1.5m	
08/06/90	+ 44d	3310-9840	CTIO-1.5m	
19/06/90	+ 53d	3320-9210	CTIO-1.5m	
02/07/90	+ 67d	2960–7620	CTIO-1.5m	
24/07/90	+ 88d	2960–7620	CTIO-4.0m	
26/07/90	+ 90d	6230-9460	C T IO -4 .0m	
21/12/90	+ 237d	3580-6940	ESO-3.6m	
11/01/91	+ 258d	3590-7010	ESO-3.6m	
20/02/91	+ 298d	3700-9590	ESO-3.6m	
21/04/91	+ 358d	3530–9390	ESO-3.6m	

have been connected for the recession velocity (cz = 2902km s  $^{1}$ ) of the parent galaxy NGC 4650A.

The early spectrum (around maximum) displays a blue continuum with broad P-Cygnipro les, indicating both high velocities and tem perature. The He I 5876 A is unambiguously recognized with a velocity of the order 13400 km s  $^{1}$ , as measured from the absorption m in im um, clearly high com pared to other SN e Ib at sim ilar phase ( 10000 km s<sup>1</sup>; Branch et al. 2002, hereafter B02). For the 12 d spectrum, V 12500 km s<sup>1</sup> is measured, and this is used as reference to infer the possible location of the various He I lines, namely 4471 A, 5015 A, 6678 A and 7065 A.Figure 2 dem onstrates the evidence of the prom inent optical He I lines in the ejecta of SN 1990I. The relative strengths of the He I lines are such that strong departures from LTE are apparent as suggested by W heeler et al. (1987) for other type 1b SN e. Lucy (1991) has shown that strong departures from LTE result from excitation by non-therm al electrons originating from radioactive decay of <sup>56</sup>C o dispersed throughout the envelope. It must be kept in m ind, nevertheless, that since the Fe II lines, especially 4924 A, 5018 A, a ect mainly the spectra shortward 5200 A, the absorption features are subject to line blending in that portion of the spectra. In addition the He I4471 A line may be a ected by the Mg II 4481 A.

The photospheric velocity, as inferred from Fe II 5169 A line, appears to be greater with respect to values derived from the SNe Ib sample in the B02 work. From the 12 d spectrum we measured V<sub>phot</sub> 9000 km s<sup>1</sup>.At com parable phase B 02 provided V<sub>phot</sub>  $6500 \text{ km s}^{-1}$ . If we use the derived velocity as an indicator of the epoch, and hence try to t he B 02 sample data (their Figures 22 and 23), one estim ates the epoch to be around m axim um . This seems unlikely, taking into account both the nature of the spectrum and the shape of the light curves (Sect. 2.2). Therefore the high velocity nature of SN 1990I ejecta seem s unavoidable.

The Call near-IR triplet, displaying broad P-Cygni since maximum light are indeed reported . The spectra pro les, is the strongest emission feature at red wave-



Fig.1. Spectral evolution of SN 1990I during the photospheric phase (A), and during the nebular phase (B). Days since maximum are shown. All the spectra have been corrected for the recession velocity of the host galaxy and scaled arbitrary for clarity. The strongest features are also reported.

lengths at this phase. The Ca II H & K is also present. A lthough features in the blue part of the spectrum are hardly discernible because of the noise, we remark in the 21 d spectrum a weak absorption feature centered at 3890 A. This feature has been a subject of discussion for early spectra of SN Ib 1999dn, and has been attributed to N i II 4067 A in order to t the 3930 A absorption (D eng et al. 2000; their Sep. 14 spectrum). This best t at 3930 A with N i II 4067 A , for the case of SN 1999dn, leads the authors to believe it is early evidence of  $^{56}$ N im ixing.

In Figure 2 two absorption features near 6050 A and 6250 A are marked (long-dashed lines). Both these features are present in the three early spectra, while the 6250 A feature is no more seen in the 43 d spectrum .D eng et al. (2000), through spectrum synthesis of SN 1999dn around maxim um, have argued for evidence of both C II and H , attributing the feature around 6300 A to the blend of the two lines, and uniquely due to C II later on.B enettiet al. (2002), on the other hand, have included N e I lines to t early spectra of SN 1991D, providing in provement in tting both absorption features, sim ilar to the ones marked by long dashed lines in Fig. 2.

W hether hydrogen is present or not in type lb/c events is an important issue, since it is directly related to the nature of the progenitor stars and their evolution (F ilippenko 1997).

The possible identication of the feature around 6250 A as H may point to the presence of a thin hydrogen

layer above the helium layer with high velocity, which disappears or is overwhelm ed by C II 6580 A as H optical depth decreases as a consequence of the envelope expansion (D eng et al. 2000). In fact, if correctly identi ed, we measure from the 12 d spectrum an H velocity of 15000 km s<sup>-1</sup>, higher compared to that of He I 5876 A ( 12500 km s<sup>-1</sup>). Note that since SN 1990I is very similar to SN 1991D with respect to the presence and the shape of the two absorption features one m ight expect N e I lines to provide good t (Benetti et al. 2002), while the presence of H still remains a possible interpretation. D etailed synthetic spectra are needed to test both possibilities.

The absorption near 7500 A is attributed to O I 7774 A line in type Ib SNe, nevertheless in some cases the introduction of the blend with Mg II 7890 A is needed for tting using synthetic spectra (e.g. SN 1999dn, Deng et al. 2000).

As the supernova evolves in time the features display a shift towards the red part of the spectrum . This shift is caused by the retreat of the photosphere into the slower m oving layers.

The last three spectra in Figure 1 (left panel) present evidence of  $[D \ I]$  6300,64 A and  $[C \ a \ II]$  7291,7324 A em issions, suggesting the onset of the nebular phase. The right panel, on the other hand, displays the real nebular phase spectra. As illustrated, the 237 d spectrum, displays a strong  $[D \ I]$  6300,64 A em ission line, which is quite broad.

F ig.2. Possible location of H  $\in$  I lines in the photospheric phase of SN 1990I (short-dashed lines). The absorption features near 6050 A and 6250 A are marked (longdashed lines; Section 2.1). Phases since maximum light are reported.

The other prom inent features are N a ID 5893 A doublet and the broad M g I] 4571 A line. A dditional nebular lines are di cult to discern. Regrettably, the 237 d spectrum does not extend redward of 7000 A, whereas the 298 d spectrum does. In fact, although the spectra are noisy, the [D I], [C a II], N a I and M g I] em ission lines are still stronger compared to the continuum, and as time goes on they tend to shrink with respect to the continuum. This continuum may come from stars very close to the SN. In fact the presence of narrow H em ission in at least

ve photospheric and either H  $\,$ em ission or absorption in three of the nebular spectra suggests the presence of an H II region inadequately subtracted. The narrow H feature (EW ' 3.8 A on day 21) is located at the rest wavelength of the host galaxy and is seen to not broaden with time, which re ects its association with H II region rather than with the SN .

Interestingly, at this phase, the [C a II] and [O I] uxes seem to be comparable. This is of interest since it is related to the nature of the progenitor star. Fransson & Chevalier (1987, 1989) have shown that the forbidden emission line ratio [C a II]/[O I] is weakly dependent on density and temperature of the emitting zone, and is expected to show an alm ost constant evolution at late epochs. Furtherm ore the ratio seem s to be very sensitive to the core m ass and hence to the main-sequence one. The ratio increases with decreasing main-sequence m ass. In Figure 3 we show the evolution of the computed ratio in SN 1990I together with SN e 1985F (Ib), 1998bw (Ic), 1993J (IIb), 1996N (Ib) and 1987A (II). The plot is indicative of the sim ilar main-

F ig.3. The time evolution of the ratio F ([CaII])=F ([DI]) of SN 1990I together with that of a sample of SN e. The di erence between SN 1987A and the remaining SN e is clear. For convenience we refer to phases since explosion date (see Sect. 2.2.).

sequence-m ass-nature of SN 19901 com pared to the other lb/c events.

It is note worthy, how ever, that in type 1b/c SN e there is no hydrogen rich C a II em itting zone as is the case of type II events (de K oolet al. 1998). We note as well that the values reported for SN 1990Im ay be even lower if we correct for the depletion behaviour blueward of the [D I] em ission (see Section 2.1.1).

# 2.1.1. Nebular em ission line pro les

In Figure 4 we show, in velocity space, the late evolution of the most prom inent nebular lines, namely M g I] 4571 A, [O I] 6300,64 A, [Ca II] 7307.5 A and O I 7774 A. The [D II] 7322 A, on the other hand, may contribute in the red side of the [Ca II] 7307.5 A.

At 90 d the at top of the [D I], [C a II] and O I em ission lines is indicative of shell structure. An approximate position of the blue edge of plateau in [D I] line is at 2600 km s<sup>1</sup>. The [D I] line on day 237 is still peaked at zero velocity, whereas on day 258 it appears blueshifted by 600 km s<sup>1</sup>. Unfortunately, the spectra in this transition phase, from +237 d to +258 d, do not show [C a II] and O I pro les in order to check whether there is any dependence of the shift on wavelengths. The M g I] 4571 A , at that phase, is noisy which makes any conclusion about a possible blueshift.

This behaviour was also seen in SN 1993J, and was interpreted as an indication of large scale asymmetries and clum py distribution of the ejecta (Spyrom ilio.1994). The [Ca II] 7307.5 A line shows a blue shift the magnitude of







Fig.4. The time evolution, in velocity space, of the most prom inent nebular lines: Mg I] 4571 A, [O I] 6300,64 A, [Ca II] 7307.5 A and O I 7774 A. The [O II] 7322 A may contribute in the red side of the [Ca II] 7307.5 A. Shown as well are the coe cient by which the spectra have been multiplied for clarity. Rest wavelengths are marked by vertical long-dashed lines.

which seem s to decrease from day 90 to day 358. Since this interval embraces the epoch of dust form ation (Section 2.2) and spectral observations are so sparse it is not possible to disentangle the two possible competing e ects. This scarcity of data also inhibits a more detailed discussion of the blueshift seen in the 0 I 7774 A line at day 90. The strength of the 0 I 7774 A feature, on the other hand, arising as it does from high levels in an envelope that should be relatively cool, could indicate non-therm al electron excitation analogous to what has been proposed for the H e I lines at an earlier phase.

The \early-tim e" blueshifts a ecting several lines have been reported for SN 1996N by Sollerm an et al. (1998) and extensively discussed there. Sim ilar e ects have also been reported for SN e 1987A, 1999em and 1988A. Various hypothesis have been reviewed without m conclusions being drawn although large scale asymmetries are favoured. In spite of their reported lack of temporal evolution of the blueshifts in SN 1996N, evolution was evident in SN 1987A and seems evident in our sparse observations of  $[D\ I]$  6300,64 A prior to dust formation. We consider a decrease in line optical depth, possibly combined with different excitation conditions for di erent lines, as the most promising explanation.

On the other hand, we note the presence of ne structures superimposed clearly on the proles of O I 7774 A (90 d spectrum), [Ca II] 7307.5 A (90 d and 298 d) and Ca II 8662 A. In Figure 5 two structures on the top of [Ca II] 7307.5 A and Ca II 8662 A are marked. The bum ps have comparable velocities (-800 and 500 km s<sup>-1</sup>) in both lines. Sim ilar bum ps have been reported in the 1993J spectrum around day 154 (since maximum) on the top of [O I] 6300,64 A and O I 5577 A lines, and have been attributed to asymmetry, whereas the lack of similarity in line proles of permitted and forbidden transitions of the same element is a result of spatially diment excitation conditions (Spyrom ilio. 1994). In the longer wavelength lines fringing in the CCD  $\,$  could complicate and confuse a clearer interpretation.

The situation of SN 1990Im ight be more com plicated in view of further information from the photom etric evolution (Section 2.2), where the coexistence of both large scale asymmetries and dust condensation at later epochs (after day 237) is more probable. In fact a second blueing e ect was reported in SN 1987A spectra, after 500 d, where the emission line pro les of Mg I] 4571 A , [O I] 6300,64 A and [C I] 9824,9850 A became asymmetric with peak emission blueshifted by about 500 600 km s<sup>1</sup>, and was attributed to extinction by dust within the m etal-rich ejecta (Danziger et al. IAUC 4746, Lucy et al. 1991). An analysis of SN 1999em data has indicated a sim ilar conclusion about dust form ation around day 500 (E lm ham diet al. 2003).

The [D I] 6300 A pro lein the last two spectra displays a depletion of emission in the [ 2500:0] km s <sup>1</sup> region. In Figure 6 we present, at three di erent phases of evolution, the [D I] 6300 A pro le of SN 1990I compared to SN e 1993J and 1998bw .W e shift arbitrarily the peaks to m atch each other for the clarity of the comparison. There are some points of interest on this gure. First, SN 1990I shows a broader pro le with respect to SN 1993J, and is similar to that of SN 1998bw . Second, the depletion of the blue wing is easily visible in the bottom box, pointing to di erent excitation conditions compared to that in the other two objects. A gain a clum ping of the radioactive m aterial in the oxygen shell with velocities belonging to [ 2500:0] km s <sup>1</sup> zone is a possible explanation.

Finally, the presence of the depletion structure in the 298 d and 358 d  $[D \ I]$  pro les prevents us from checking whether the blueshift seen in the 258 d  $[D \ I]$  emission increases or not with time.

## 2.1.2.0 xygen mass

E stim ating the oxygen m ass is important as it can be indicative of the core m ass and hence related to the nature of the progenitor star. The  $[0 \ I] 6300,64 \text{ A}$  doublet line em ission is known to be the prim ary coolant in the ejecta of 1b/c events, at least at late epochs, in view of the absence of hydrogen in this class of objects as opposed to type II events (U om oto 1986; Fransson 1987).

In principle, the mass of ejected oxygen can be recovered using the  $[0 \ I]$  6300,64 A ux. U om oto (1986) has shown that the oxygen mass, in M , is given by:

$$M_{O_X} = 10^8 \quad D^2 \quad F([O I]) \quad \exp(2:28=T_4)$$
 (1)

where D is the distance to the supernova (in M pc), F is the [D I] integrated ux (in ergs s<sup>1</sup> cm<sup>2</sup>) and T<sub>4</sub> is the tem – perature of the oxygen-em itting gas (in 10<sup>4</sup> K). Equation 1 holds for the high density lim it (N<sub>e</sub> 10<sup>6</sup> cm<sup>3</sup>) where the density is above the critical density for the [D I] line (7 10<sup>5</sup> cm<sup>3</sup>). This condition, on the other hand, is clearly achieved in the ejecta of type Ib/c SN e. In fact a rough estim ate of the mass density can be computed at



Fig.5. The strongest lines, in velocity space, of the 90 d spectrum. The features are clearly asymmetric and at topped, with the presence of ne structures. Two bumps with similar velocities in the [Ca II] 7307.5 A and Ca II IR-triplet lines are marked.

day 237.W e have attempted to t the [0 I] 6300,64A feature with 2 G aussians, xing the wavelength separation of the doublet and with both having the same velocity width. It is not possible to obtain a particularly good overall t since one is left with either a residual in the blue wing of [] I] 6300 A for narrow er lines, or little evidence of the [0 I] 6364 A feature for broader lines. Thus a one-com ponent Gaussian t m ay not represent the real situation where clum ping is present. On the other hand by assuming that there is a contribution from an Fe II 6239 A em ission, and allowing its velocity width to be a free parameter we obtain the tshown in Fig. 7. The FW HM for the [O I] lines 60 A while for the Fe II 6239 A line we obtain 47 A . is We also show residuals which might represent the e ects of clumping in the [0 I] em itting region. Conveniently, in this the velocity width for the Fe II feature is less than that for [0 I] as m ight be expected. W e adopt the velocity widths for [0 I] determ ined in this way but note that sim ilar values would be obtained if we accepted the best t of the [0 I] doublet alone around the peak em ission of both components of the doublet.

U sing then the derived F W H M of the [0 I] line at 237 d (2860 km s<sup>1</sup>), and adopting the ejecta m ass of M<sub>ej</sub> = 3:7 M (Sect. 3.2) we derive a value of N<sub>e</sub> 1:16 10<sup>7</sup> cm<sup>3</sup>. We have assumed abundance fractions of He, O and heavier elements in the following proportions: 0.1, 0.22 and 0.68, respectively. We also assume that half of the He and O and all of the other elements are singly ionized. Since we assume matter uniform ly distributed this should be a lower limit for regions where clumping occurs.

1.2



A more accurate estimate of the density is based on the relative strengths of the [0 I] doublet components (Leibundgut et al. 1991; Spyrom ilio & Pinto. 1991). The authors of both papers have com puted the variation of the line doublet ratio as function of the density at a given time and found it insensitive to the adopted tem perature, especially at late epochs. From the 237 d [O I] line pro le twe m easure a relative intensity I (6300)=I (6364) ' 1:69, sim ilar to that of SN 1987A at sim ilar phase ( 1.45 around day 250). The uncertainty in the tem perature leads to the Ne following range 2:3  $10^{9}$ 4 10<sup>9</sup> according to Fig. 6 of Leibundgut et al. (1991). The large di erence (a factor of about 100) between this density estim ate and the one from the simple approach using the volum e calculated from the [O I] velocity and M ej indicates, indeed, that the oxygen material lls the volume at day 237 in a clumpy way rather than hom ogeneously. This high density would favour the presence of Fe II features m entioned previously and identi ed in Figure 1.

Equation 1 implies time-dependence because of the variation of F ([O I]) and the temperature. Schlegel & K irshner (1989), when estimating the oxygen masses ejected by SN e Ib 1984L and 1985F using this method, have adopted a constant temperature  $T_4 = 0.4$  at the neb-

Fig.7. Fit to the 237 d [0 I] 6300,64 A line pro le of SN 1990I. Shown in thick line is the sum of the three Gaussians for [0 I] 6300 A , [0 I] 6364 A and Fe II 6239 A (dashed lines; see text). The residuals are displayed in the lower panel.

ular phase. The assumption of a constant temperature at di erent late phases provides di erent oxygen masses as one may expect, since earlier nebular epochs are hotter compared to latter ones. W orth noting is the exponential dependence of the oxygen mass on temperature.

A method for estimating temperature at a given time is based on the  $[D \ I] 5577 \ A$  to  $[D \ I] 6300,64 \ A$  ux ratio. This comes from the assumption that the O I lines are formed mainly by collisional excitation, and the ratio is given by the following expression (Houck & Fransson 1996):

$$\frac{F (6300)}{F (5577)} = 0.03_{6300} [1 + 1.44T_3^{0.034} (\frac{10^8}{N_e})]$$
(2)  
exp (25.83=T\_3)

where  $T_3$  is the tem perature of the oxygen-em itting gas (in  $10^3$  K),  $_{6300}$  is the [O I] 6300 A Sobolev escape probability ('1) and N<sub>e</sub> is the electron density.

Because of line blending, mainly with Fe II lines, the [D I] 5577 A ux measurements are less precise, especially for the last three spectra. We use the 237 d spectrum because it is less noisy and does not show the depletion feature seen later in the blue side of the [D I] 6300,64 A emission pro le.

A lower lim it on the ratio (left hand term in Eq.2) can be estimated using an upper lim it on the  $[0 \ I] 5577 \text{ A}$  ux by integrating over the same velocity interval of the  $[0 \ I]$ 6300,64 A pro le.





Table 2. The estimated ejected oxygen mass.

SN	SN	М <sub>ох</sub>	R eferences
nam e	type	(M)	
1987A	II	1.5-2	1,2
1984L	lb	0 <b>:</b> 3°	3
1985F	Ib	03 <sup>°</sup>	3
1987M	Ic	0.4?	4
1990I	Ib	0.7-1.35 <sup>?</sup>	th is work
1993J	IIb	0:5	5
1996N	lb	0.1-0.3?	6

Refs:

1-Fransson et al.1993; 2-Chugai1994; 3-Schlegel& Kirshner1989; 4-Filippenko et al.1990; 5-Houck & Fransson 1996; 6-Sollerm an et al. 1998.

? Estim ated adopting the same method (i.e. using Eq. 1)

W hen using the density range estim ated on the basis of the [0 I] 6300,64 A doublet ratio we derive tem peratures of the order of 3200 3500 K.We obtain then 0:7 1:35 M as oxygen m ass. Here we note that since we are adopting an upper lim it on the [O I] 5577 A ux, the recovered oxygen m ass is a lower lim it. The very weak [0 I] 5577 A feature in the last three spectra is an indication of low tem peratures in the oxygen-em itting zone. Indeed, although the determ ination of oxygen m ass following this method su ers from several uncertainties, it seems that SN 1990I has reached lower tem peratures earlier and has a higher oxygen m ass com pared to typical Ib/c SNe. In Table 2 we report the amount of ejected oxygen m ass for a selected sam ple of type 1b/c SN e. The table includes as welldata of SN II 1987A and SN IIb 1993J.One should note how ever that adopting a di erent m ethod, Tutukov & Chugai (1992) have argued for an amount of 0:8 M as total ejected oxygen mass<sup>1</sup> (neutral and ionized) for type 10 SN 1984L, and concluded that a typical value of 1 M should be ejected as well by SNe Ib 1983N and 1985F.

The som ewhat higher oxygen m ass and lower tem perature with respect to other SN e Ib/c m ay be related to other observational peculiarities in SN 1990I. First the higher expansion velocity (i.e. faster cooling), and second the possible dust condensation at a time as early as 250 d (be deduced photom etrically later).

# 2.2. Photom etric evolution

The photom etry data of SN 1990I are reported in Table 3 together with the di erent instruments with which the observations were obtained. The SN magnitudes have been measured with the ROMAFOT package in M IDAS (Buonanno et al. 1983). A fler measuring the point spread function from the eld stars, the programme allows the simultaneous tting of the SN and the background, which

has been approximated by a tilted plane. In Figure 8, the B, V, R and I light curves are displayed. The well sam pled light curves, following maximum, coupled with the reported early spectral properties (i.e. broad P-C yqni pro les superim posed on a blue continuum ) provide the possibility of estimating the time of maximum light and roughly the explosion date. In fact based on the shape of the light curves during the rst 30 days of evolution and bearing in m ind that the typical rise time of type Ib/c SN e is about 15 20 days to reach m axim um light, we estim ate both the date of maximum light and the explosion time to be, respectively, April 29 (1990; JD 48010 5d) and April 11 (1990; JD 47992 5d). These dates are taken as references through this work. How ever, because the point ofmaximum was not de ned (see Fig. 8) we estimate that maximum light could have occured at most 5 days earlier.

SN 1990I su ers high galactic extinction,  $A_V^{gal} = 0.374$ m ag, according to m aps of the galactic dust distribution by Schlegel, Finkbeiner & D avis.(1998). This corresponds to a colour excess about E (B V) = 0.12, where the standard reddening laws of C ardelli et al. (1989) have been used. In addition we note the presence of a narrow N aI D interstellar absorption feature in the 12 d spectrum at the redshift of the parent galaxy, with an equivalent width (EW) 0.26 A. This can be used to estimate the reddening in the host galaxy. U sing correlations relating EW (N a I D) to the reddening within the parent galaxy from the literature we obtain E (B V) 0.06 (B arbon et al. 1990) and E (B V) 0.034 (B enetti et al., in preparation). In what follows, we adopt  $A_V = 0.4$  m ag as an estimate of the total extinction.

A fier the fast fall from maximum light ( $_{\rm V}$  5.6 m ag (100 d) <sup>1</sup>), SN 1990I settles onto a linear decline with a rate in the V band  $_{\rm V}$  1.92 m ag (100 d) <sup>1</sup> up to about 230 days since explosion (dotted line in Fig. 8). The <sup>56</sup>C o to <sup>56</sup>Fe decay slope, 0.98 m ag (100 d) <sup>1</sup>, corresponding to full -ray trapping is also shown. Except for rare cases where the late slope approaches the full trapping rate (e.g. SN lb 1984L; Schlegel & K irshner 1989), the steeper decline rate of about 1.9 m ag (100 d) <sup>1</sup> is common for type lb/c SN e at late epochs (e.g. SN lb 1983N, SN lc 1983V, SN lc 1994I and SN IIb 1993J, C locchiatti & W heeler 1997; SN lb 1996N, Sollerm an et al. 1998). This rapid decline, on the other hand, is indicative of signi cant -ray escape resulting from a low mass ejecta in this class of object.

Interestingly, how ever, SN 1990I shows an even steeper decline starting at about day 230 with a V-band decline rate  $_{\rm V}$  2.7 m ag (100 d) <sup>1</sup>. In addition, while the R and I light curves con m this behaviour, the B light curve tends to atten. The least-squares t to the light curves at the three di erent phases of evolution are reported in Table 4. At early phase, the B light curve declines faster com pared to the other bands, whereas for the two follow – ing phases it exhibits a lower decline rates. The V band, on the other hand, is the faster fading band during the follow ing two phases.W e com pute as well the third to second phase decline slope ratio for each band. Results are show n

 $<sup>^{1}</sup>$  0.2 M neutral and 0.6 M as ionized-oxygen m ass.

D ate	JD	B()	V()	R()	I( )	Instrum ent
(UT)	2400000+					
27/04/90	48008.5		15.42 .03			ESO -2.2m
29/04/90	48010.5	16.68 .05	15.58 .03	ĺ	ĺ	ESO-2.2m
30/04/90	48011.5	16.75 .05	15.65 .03	ĺ		ESO-2.2m
01/05/90	48012.5		15.73 .03	ĺ		ESO-2.2m
02/05/90	48013.5	17.01 .05	15.80 .03	15.28 .03		CTIO-0.91m
10/05/90	48021.5	17.35 .05	16.18 .03	15.60 .03		ESO-NTT
14/05/90	48025.5	17.65 .05	16.44 .03	15.87 .03	15.38 .03	CTIO-0.91m
18/05/90	48029.5		16.50 .03			ESO-NTT
28/05/90	48029.5	17.85 .05	16.78 .03			ESO-NTT
08/06/90	48050.5	18.18 .05	17.06 .03	16.55 .03	15.96 .03	CTIO-0.91m
12/06/90	48054.5	18.22 .05	17.16 .03	16.64 .03	16.05 .03	CTIO-0.91m
16/06/90	48058.5	18.26 .05	17.23 .03	16.71 .03	16.12 .03	CTIO-0.91m
19/06/90	48061.5	18.27 .05	17.26 .03	16.75 .03	16.13 .03	ESO-1.54m
20/06/90	48062.5	18.24 .05	17.27 .03	16.72 .03		ESO-1.54m
21/06/90	48063.5	18.34 .05	17.30 .03	16.83 .03	16.25 .03	CTIO-4.0m
25/06/90	48067.5	18.40 .05	17.37 .03	16.92 .03	16.32 .03	CTIO-4.0m
29/06/90	48071.5	18.44 .05	17.43 .03	16.97 .03	16.37 .03	CTIO-4.0m
04/07/90	48076.5	18.53 .05	17.58 .03	17.09 .03	16.44 .03	CTIO-0.91m
12/07/90	48084.5	18.63 .05	17.71 .03	17.20 .03	16.56 .03	CTIO-0.91m
26/07/90	48098.5	18.78 .05	17.94 .03	17.42 .03		ESO-3.6m
01/08/90	48104.5	18.98 .05	18.15 .03	17.55 .03	16.92 .03	CTIO-0.91m
12/08/90	48115.5	19.11 .05	18.30 .03	17.73 .03	17.08 .03	CTIO-0.91m
22/08/90	48125.5	19.32 .05	18.49 .03	17.86 .03	17.26.03	CTIO-0.91m
23/11/90	48218.5		20.47 .1			CTIO-0.91m
11/12/90	48236.5	22.26 .3	21.46 .1	19.98 .03	19.82 .03	CTIO-4.0m
14/12/90	48239.5		21.11 .1	20.04 .1		ESO-1.54m
15/12/90	48240.5		21.32 .1		19.72 .03	CTIO-0.91m
21/12/90	48246.5		21.34 .1			ESO-3.6m
28/12/90	48253.5	22.42 .3	21.88 .1	20.60 .1	20.11 .1	CTIO-0.91m
15/01/91	48272.5			21.02 .1		CTIO-0.91m
16/01/91	48273.5		22.42 .2	21.15 .1	20.99 .1	CTIO-0.91m
12/02/91	48300.5	23.28 .35	23.01 .25	21.70 .1	21.40 .1	CTIO-4.0m
20/02/91	48308.5	22.90 .35	22.90 .25	21.85 .1	21.56 .1	ESO-3.6m
09/04/91	48356.5			23.37 25		C T IO -4.0m

Table 3. SN 1990IP hotom etric Observations

in Table 4. The change in slope behaviour at late phases is seen also in other events, nam ely SN 1993J, SN 1998bw and SN 1996N with di erent occurrence tim es. SN 1996N, between 184 d and 312 d, seem s to have sim ilarV, R and I decline rates as SN 1993J (Sollerm an et al. 1998). The last photom etric point of SN 1996N (around day 342) is clearly deviating from the linear trend of the previous data, indicating the occurrence of a phase with steeper decline after day 312. In Table 4 we present the SN 1996N third to second decline rate ratios using the available photom etric data from Sollerm an et al. (1998). Interesting points can be read out from Table 4. A comparison of ratio values dem onstrates the peculiar behaviour of SN 1990I with an increasing ratio towards the red bands, indicating som e kind of dependence of the de ned ratio on wavelength. A lthough there is a paucity of SN 1996N data, it is sim ilar to SN 1990I in having a \greater than one" ratio, whereas SNe 1993J and 1998bw tend to atten. Furtherm ore, SN 1993J has an opposite behaviour with respect to SN 1990I, in the sense that its B light curve declines m ore rapidly in

the third phase while the V, R and I light curves atten. The opposite holds for SN 1990I.

In Figure 9 the absolute V light curve of SN 1990I is com pared with other type Ib/c core collapse events. For SN 1990I a heliocentric velocity corrected for the Local G roup infall onto the V irgo C luster is 2751 km s  $^1$  (as reported in the LEDA database) translates into 32:97. Table 5 sum m arizes the m ain param eters adopted for the selected sam ple. The gure reveals a strong sim ilarity between SN 1990I and 1993J, from maximum to about day 230, with SN 1990I about 0.2 m ag brighter. The maximum brightness is, on the other hand, quit comparable to those of SN e 1987M and 1994I and interm ediate between the bright SNe 1999bw and 1991D and the faint SN 1983I.M oreover SN e 1990I, 1993J, 1991D and 1998bw decline from maximum with a similar rate to reach the exponential decay and indicate sim ilar peak-to-tail contrast, whereas SN e 1983I, 1987M and 1994I display narrower peak widths and greater peak-to-tail contrast. The sample we are studying, although small, shows a corre-



Fig.8. B, V, R, and I light curves of SN 1990I. The light curves have been shifted by the reported am ounts. A lso shown are the ts to the V-light curve tail (dotted lines) and as well the  ${}^{56}$ C o to  ${}^{56}$ Fe decay slope (long-dashed line).

Table 4. Decline rates of SN	1990I. Data for events with two	
slopes at late tim es are also rep	orted (see text for discussion).	

P hase range	В	V	R	I
(days)	m ag (100 d) $^{ m 1}$	m ag (100 d) $^{ m 1}$	m ag (100 d) $^{ m 1}$	m ag (100 d) $^1$
[16:40]	6.2	5.6	4.78	
<b>[</b> 50 <b>:</b> 150]	1.53	1.92	1.77	1.7
[200:340]	1.2	2.7	2.6	2.6
The third to second phase decline slopes ratio :				
[200:340]/ [50:150] (90I)	0.79	1.4	1.467	1.53
[150:350] / [45:150] (93J)	1.47	0.82	0.62	0.75
<sub>[310:490]</sub> / <sub>[40:330]</sub> (98bw)	0.86	0.56	0.78	0.745
<sub>[t 312]</sub> / <sub>[t 312]</sub> (96N)		1.6	1.4	

lation between the peak width and the peak-to-tail contrast. SN 1984L seems to be the extrem e case of the sam – ple, with the smaller peak-to-tail contrast and a possibly broader peak. The decline rate from maximum and the peak-to-tail contrast are computed and shown in Table 5 for events with well sam pled early data. Interestingly, the events with steeper decline rates (i.e. narrow widths) and greater peak-to-tail contrast are classi ed as type Ic SN e, while the remaining events are type Ib. This is not surprising since the width and the way the light curve declines from maximum during this early phase of evolution are related to the progenitor star properties, namely the mass,



Fig.9.C on parison of the V absolute light curve of SN 1990I with those of other Ib/c events. The <sup>56</sup>C o to <sup>56</sup>Fe decay slope is shown (short-dashed line). The used parameters are reported in Table 5.

radius, energy and the degree of m ixing. These parameters, especially the ejected m ass, controls the time the trapped decay energy takes to di use through the envelope (Ensman & W oosley 1988). In fact, for smaller m ass the maximum is reached earlier because of larger radioactive heating and higher expansion velocity. The smaller the ratio  $M_{ej}$ =E " and the closer the <sup>56</sup>N i to the surface the faster the light curve declines (Shigeyam a et al. 1990). O ne scenario for SN e Ic shows, indeed, that they originate from slightly smaller m asses at the time of explosion compared to type Ib objects (Y am aoka & N om oto.1991). If how ever SN e Ic have been stripped of helium the main sequence progenitors could have larger m asses than those of type Ib SN e.

As discussed before, and except for SN 1984L, all the SN e of the sample decay, on the exponential tail, faster than  $^{56}$ Co (short-dashed line in Figure 9). If we assume the V light curves trace approximately the bolom etric ones, they all rejects maller ejecta masses allowing greater

ray escape. A fler 230 d, SN 1990I deviates clearly from its previous slope and the deviation from the SN 1993J

tail increases with time, approaching even the SN 1996N light curve. In fact, while around day 225 the SN 1990I light curve coincides with SN 1993J tail a deviation about 1.3 m ag is measured around day 310. W e note the rapid change seen in the B V colour evolution at the same age. V intrinsic colour of SN 1990I Figure 10 reports the B together with that of other type Ib/c events for com parison. (B V) $_0$  of SN IIb 1993J is also shown (dashed line). The comparison indicates that SN 1990I follows a similar evolutionary trend as the other SNe, displaying a rapid reddening during the st 40 days, re ecting the cooling due to the envelope expansion . A fler a short duration peak it turns to the blue with a slope in the [40:120] days time range of about 0.43 m ag (100 d)  $^{1}$ , sim ilar to SN e 1984L, 1987M and 1991D, although it is about 0.2 m ag bluer. At this epoch, SN 1993J exhibits a faster decline ( 0.92 m ag (100 d) <sup>1</sup>). The colour peak, around day 40, corresponds to the beginning of the exponential radioactive tail in the V-absolute light curves (Fig. 9). Then in Fig. 10, the SN e settle on an alm ost  $\rho$  has of long duration, wellsam pled for SN e 1993J and 1990I. Unfortunately, late



Fig.10. The evolution of the B V intrinsic colour of SN 1990I, compared to those of other Ib/c events. A dopted reddening for each event is reported in Table 5.

observations of SN e 1987M and 1991D are not available, however late phase data of SN e 1984L and 1985F m ay indicate the presence of a plateau phase.

Again, the SN 1990I intrinsic (B V) colour displays a dram atic fall after day 245. It decreases by 0.8 m ag in about 75 days. The time of this rapid and steep decline re ects the occurrence of the previously seen deviation in the V-absolute light curve. These facts point to the presence of a phenom enon able to cause a de cit in the optical light curves and also to cause the line pro les to be blueshifted in a short time (t=t 0.1; Section 2.1). This rapid transform ation of pro les during the nebular epoch has also been reported in SN e 1987A and 1999em, and when combined with a sudden increase in the decay rate of the optical light curves, it has been interpreted as an indication of dust form ation. We believe that this the case for SN 1990I also. In fact, bearing in m ind that dust condensation is expected to correlate with the cooling of the ejecta, we interpret the form ation of dust earlier in SN 1990I com pared to SN e 1987A ( 530 d) and SN 1999em ( 500 d) to be due to lower ejecta m ass and rapid cooling for this event. It might also result from the higher m etallicity content in the envelope of SN 1990I.SN 1987A shows a slight wiggle around day 550 in the optical colour evolution (B V) (Suntze & Bouchet 1991). That was interpreted as a blueing associated with dust form ation, but was not so dram atic as dem onstrated by SN 1990I. Such a blueing e ect at the time of dust form ation has been discussed quantitatively in the case of SN 1987A by Lucy et al. (1991). The authors suggest that the blueing,

reported in both (B V) and (U B) colours at the onset of dust condensation, resulted from an increase of dust albedo at shorter wavelengths. In fact one knows from the work of D raine (1985) that for both graphite and astronom ical silicates the albedo does increase with decreasing wavelength in the optical range. A nother explanation, in the case of SN 1987A, has been provided by Phillips & W illiams (1991), ascribing the increasing ux in the blue to the increasing contribution from [Fe II] lines, however even including all available Fe lines, D anziger et al. (1991) did not succeed in getting such a strong e ect. The blueing at the time of dust form ation could be a result of both dust albedo e ect and em ission lines (both Fe and others). In any case the \dram atic" blueing seen in SN 1990I depends quantitatively on the accuracy or reliability of the V) m easurem ent at day 240.M odelling are required **(**B to quantitatively check this e et.

## 3. Physical param eters estim ate

In this section we construct the bolom etric light curve of SN 1990I, and adopt a very simple model for radioactive powering. A reasonable t to the data, especially during the initial part of the radioactive tail after the steep decline from the maximum brightness, provides quantitative insights on the basic parameters responsible for the bolom etric light curve shape, namely the ejecta and  $^{56}N$  i m asses.

## 3.1. Bolom etric light curve

In Figure 11 we present the \bolom etric" light curve of SN 1990I together with that of SN 1993J (Richmond et al. 1994, 1996) for com parison. The lum inosities were com puted by integrating the uxes using the available photometric data bands. The conversion of the magnitudes to uxes was carried out assuming the calibrations of Bessell (1979). The adopted parameters, i.e. distance and reddening, are reported in Table 5. The derived lum inosities, how ever, should be corrected for the near-IR (JHK) uxes. To account for this contribution, in the case of SN 1998bw, a fraction of 42% at day 370 was estimated to be added to the  ${\rm L}_{\rm BVRI}$  (Sollerm an et al. 2002), whereas Patat et al. (2001) have derived a fraction of about 35% on day 65.4.A fraction of about 50%, on the other hand, has been estimated around day 440 for SN IIP 1999em (Elm ham di et al. 2003). Though not precise, in view of the possible increasing in portance of the near-IR fraction with time (e.g. SN 1998bw; Sollerm an et al. 2002), here we simply adopt 35% as a constant percentage contribution of the near-IR uxes. The bolom etric light curves of SN 1990I and 1993J shown in Figure 11 have been indeed scaled up to account for this constant contribution.

The constructed bolom etric light curve of SN 1990I can be divided into three phases. F irst phase (the initial 40 days): although the rising part is not covered, the slow drop from the maximum compared to SN 1993J is con med.We note the sim ilarity starting around day 30

## Table 5.M ain parameters of the SNe sam ple.

SN nam e	SN type	H ost galaxy	D istance (M pc)	A <sub>v</sub> (mag)	$v^{z}$ (m ag (100 d) <sup>1</sup> )	M ax-to-tail contrast (m ag)	R eferen <i>c</i> es
1990I	Ib	NGC 4650	39 <b>.</b> 3°	0.4	5.7	1.05	this work
1983I	Ic	NGC 4051	13 <b>.</b> 17 <sup>?</sup>	0.043	8.4	2.2	1,2
1984L	Ib	NGC 0991	20 <b>.</b> 44 <sup>?</sup>	0.0			3,4,5
1987M	Ic	NGC 2715	22 <b>.</b> 7 <sup>?</sup>	1.3			6
1991D	Ib	PGC 84044	$178.52^{?}$	0.186			7
1993J	IIb	M 81	3.64	0.93	7	1.23	8
1994I	Ic	NGC 5194	8.32	1.4	11	2.42	9
1996N	Ib	NGC 1398	22	0.0			10
1998bw	Ic-H ypemova	ESO 184-G82	35 <b>.</b> 16 <sup>?</sup>	0.2	6	1.64	11,12,13,14

R efs:

1-T svetkov 1985; 2-T svetkov 1988; 3-M eikle et al. 1984; 4-B uta 1984; 5-T svetkov 1987;

6-Filippenko et al. 1990; 7-Benetti et al. 2002; 8-Lew is et al. 1994; 9-Richm ond et al. 1996;

10-Sollerm an et al. 1998; 11-G alam a et al. 1998; 12-M cK enzie & Schaefer 1999; 13-P atat et

al. 2001; 14-Sollerm an et al. 2000.

? Estimated using \LEDA " database and adopting H  $_0$  = 70 km s  $^{1}$ M pc  $^{1}$ .

z Early decline rate from maximum .

with SN 1993J, not only in the decline rate but as well in the computed lum inosity. A round day 20 SN 1990Iwas

0.15 dex fainter. S econd phase (the [50:200] days tim e interval): the light curve settles on the exponential decay tail, with a match with SN 1993J at least until day 100, whereas around day 130 SN 1990I seems to be 0.06 dex brighter. At this epoch, an e-folding time of 60 2 days is measured, considerably faster than the one of  $^{56}$ C o decay (i.e. 111.3 d), suggesting the -rays escape with decreased deposition, ow ing to the low mass nature of the ejecta. Third phase (after day 200): at this phase both SN e (1990I & 1993J) tend to behave di erently. W hile SN 1993J atten (e-folding time ' 85 2 d), SN 1990I shows the dram atic fall seen in the VR I broadband photom etry (e-folding time ' 47 2.8 d).

The observed late time attening in SN 1993J (and also in SN 1998bw) can be explained as due to the interaction of the ejecta with the circum stellar material (Sollerm an et al. 2002), although for SN 1998bw Chugai (2000) has found good agreem ent with observations adopting complete  $^{56}$ N im ixing and a higher matter density in the central part of the envelope. On the other hand, the sudden drop in lum inosity after day 200 combined with spectral features in the case of SN 1990I point to dust condensation.

# 32. Simple -ray deposition model

The principal goal of the simple light curve modeling is to t the bolom etric behaviour starting at the transition phase (i.e. the early part of the radioactive decay tail) up to about 100 d. In the time range of interest, the contributions from radioactive elements other than  ${}^{56}N$  i and  ${}^{56}C$  o are not taken into account, for the simple reason that the m ore long-lived elem ents, like  $^{57}$ Co, shown to be important at late phases (W oosley et al. 1989) are not su ciently abundant to a ect the light curves at day 100 (D anziger et al. 1991).

The model describes the lum inosity evolution in an homologously expanding spherical ejecta with a point source ray deposition from <sup>56</sup>N i! <sup>56</sup>C o! <sup>56</sup>Fe. It is based on the simple approach adopted by Swartz & W heeler (1991) and C locchiatti & W heeler (1997) when studying, respectively, the late time light curve of SN 1984L and type lb/c SN e representative light curves. In fact, assuming free expansion of the ejecta (v (r;t) = r=t), a power law density distribution ( (r;t) / r <sup>n</sup> (t)) and a purely absorptive opacity ( ) being constant throughout the ejecta, the equations for the mass, kinetic energy and ray optical depth can be solved, giving rise to an expression for the total ray optical depth relating the main physical parameters (C locchiatti & W heeler 1997):

$$= C \qquad M_{ej}^{2} = E \quad t^{2} \tag{3}$$

where t refers to time since explosion and C is a constant depending on the density function. For a power index n = 7 we have C' 0.053, adopted in our calculations. A typical value for the opacity,  $= 0.033 \text{ cm}^2 \text{ g}^{-1}$ , is also assumed (W oosley et al. 1989).

W ithin this simpli ed picture, the emergent -ray lum inosity, at a given time t, reads:

$$L(t) = L_0(t) [1 \exp(((t))]$$
 (4)

where  $L_0\left(t\right)$  is the total -ray lum inosity from radioactive decay.

In order to estimate  $L_0$  at a given time, a compilation of the radioactive decay energy based on the properties of the  ${}^{56}N$  i!  ${}^{56}Co$ !  ${}^{56}Fe$  decay is needed. For this purpose,



F ig.11. The computed \BVRI" bolom etric light curve of SN 1990I together with that of SN 1993J for comparison. SN e 1990I and 1993J lum inosities are shifted upward to account for a \JHK " contribution of the order of 35%. The best twith a simplified -ray deposition model from <sup>56</sup>N i! <sup>56</sup>Co! <sup>56</sup>Fe is shown. It corresponds to M<sub>ej</sub> 3.7 M and M (N i) 0.11 M. The window displays the evolution of the fraction of the -ray lum inosity deposited in the envelope corresponding to the best t.

we adopt the main param eters presented by Je ery (1999) and N adyozhin (1994). In this scheme the total released energy via -photons per  ${}^{56}$ N i decay is calculated to be 1.73 M ev, while the mean photon plus positron kinetic energy per  ${}^{56}$ C o decay is computed to be 3.74 M ev. The positron fraction in energy is only about 3.5% of the total energy per  ${}^{56}$ C o decay, while the neutrinos are assumed to escape the ejecta without any energy contribution (Je ery 1999).

F inally, we compute the total rate of radioactive energy production at a given time and combine it with equations 3 and 4, thus providing a general description of the simple radioactive decay energy deposition model in a spherical geometry. The model, although simple, might be a good indicator for  $^{56}$ N i and ejecta mass estimates, especially at the transition phase (i.e. in the [40:100] days time range).

Figure 11 displays the best t of SN 1990I bolom etric light curve (dotted line). A dopting, in fact, an energy E ( $10^{51}$ ergs) = 1, our best model t indicates the follow – ing values: M ( $^{56}$ N i) = 0:11 M and M<sub>ej</sub> = 3:7 M . The window in the gure displays the evolution of the fraction of the -ray lum inosity deposited into the envelope corresponding to the best t. O f course the model t does not account for the late drop in lum inosity, how ever it reproduces well the initial part of the radioactive tail. SN e 1990I and 1993J, in this context, seem to have similar

 $^{56}\text{N}$  iejected m ass, whereas in view of the slight di erence in brightness in favour of SN 1990I in the [90 : 130] d time range, SN 1993J has a slightly smaller ejected m ass ( 3.4 M from our best t to SN 1993J data). The best t parameters for SN 1993J are in good agreement with those from m ore complicated light curve calculations (M ( $^{56}\text{N}$  i) = 0:1 0:14 M and M  $_{\rm ej}$  = 1:9 3:5 M ; Young, B aron & Branch.1995)

We note, however, that owing to the observed higher velocity nature of SN 1990I compared to the Ib SN e sam – ple of B ranch et al (2002), a higher kinetic energy might be anticipated though it is not, a priorim andatory. An adopted higher energy of about E ( $10^{51}$  ergs) = 1.2 can be compensated with a mass M<sub>ej</sub> ' 3:95 M to keep equal the -ray optical depth, in accordance with equation 3 (with an energy dependence given by: M ( $^{56}$ N i) ' 0:11 E  $^{0:23}$  M and M<sub>ej</sub> ' 3:7 E  $^{0:36}$  M ). Therefore, bearing in mind as well the possible uncertainties in the adopted parameters (i.e. distance and reddening), an ejected mass in the range [3 4:5] M is possible.

# 4.D iscussion and conclusions

W e have presented and analysed photom etric and spectroscopic data of SN 1990I, covering about 400 days of evolution. The observations were carried out at ESO and CTID. The classi cation of the SN as a type 1b event discovered nearmaximum light is con rm ed. of some strong em ission lines. In fact, contrary to SN 1993J where some bum ps have been identied on the [D]

The optical H e I lines are identied in the photospheric spectra of SN 1990I, displaying higher velocities com pared to those derived, at similar epochs, from Branch et al. (2002) type Ib sample. The high velocity nature of SN 1990I is exem pli ed rst from photospheric velocities, inferred from Fe II lines and He I lines, and second to a lesser extent from com paring [0 I] 6300,64 A em ission line pro les with those of SN e 1993J and 1998bw. The phase of the spectral observations is uncertain because the st photom etric point m ay not de ne the point of m axim um . But it cannot be pre-m axim um . This means that the spectra could only be around maximum and later in phase but not earlier. Thus the abnorm ally high expansion velocity of SN 1990I seem s beyond doubt if the phases for the other objects in the published sample are well determ ined. We provided then an estim ate of the maximum time and explosion date that are taken as references throughout the paper.

During the rst 30 days the spectra at 6000 6500 A range show the presence of two absorption features near 6050 A and 6250 A, while around 43 days since maximum light no presence of the 6250 A trough is seen. The possible identication of this trough with H may point to the presence of a thin hydrogen layer. The presence of hydrogen, expelled at high velocities, in SN 1990I and in the ejecta of type Ib events in general is of course a crucial issue in understanding the progenitor properties. D exper investigation of these line identications requires detailed synthetic spectra and preferably a large sam ple study.

Entering the nebular phase, SN 1990I shows strong em ission features of M g I], N a I, [D I] and [C a II], typical for type Ib/c SN e at late epochs. We note the presence of a depletion in the [ 2500 : 0] km s<sup>-1</sup> region of the [D I] 6300,64 A em ission pro le in the 298 d and 358 d spectra, which m ay point to large scale clum ps in the oxygen ejecta at late phases. On the other hand the possible presence of Fe II 6239 A em ission line feature (which seem s com m on in m any type Ib/c SN e at this phae) com plicates the interpretation of the depletion feature.

The similarity in the main-sequence-mass between SN 1990I and a sample of SN e Ib/c is shown through the evolution of [Ca II] to [O I] ux ratio. It must be kept in mind, when comparing to type II SN e, that in Ib/c events there is no hydrogen rich [Ca II] emitting zone as is the case for type II objects and thus any rm conclusion should be taken with considerable caution.

We estimate a lower limit on the ejected oxygen mass to fall in the range  $(0:7\ 1:35)$  M , although this estimate is subject to large uncertainties (i.e. density, temperature and uxes). Clumping of the oxygen materials is indicated on the basis of two independent estimates of the density of oxygen.

The large asymmetry seen, especially, in the oxygen and calcium forbidden lines is a result of radioactive material being distributed asymmetrically. Furthermore we identify possible ne structures superimposed on the top of some strong emission lines. In fact, contrary to SN 1993J where some bumps have been identi ed on the [0 I] 6300,64 A and O I 5577 A pro les with similar velocities, two ne structures, on day 90, are found to repeat in the Ca II 8662 A and [Ca II] 7307.5 A in SN 1990I.We interpret this to be an indication of di erent excitation conditions, in space and time, between SN 1990I and SN 1993J.

Interestingly, SN 1990I spectra in addition to asymmetric proles, display a blueshift change during the time period from 237 d to 258 d (t=t 0:1), seen clearly in the [O I] 6300,64 A prolew ith a velocity of 600 km s<sup>1</sup>. Because of the presence of the depletion and poor S/N it is di cult to con m the blueshift in the last two spectra. On the other hand, [C a II] 7307.5 A appears blueshifted at the last two epochs.

A photom etric com parison of SN 1990Iw ith other type Ib/c and IIb events, also dem onstrates som e peculiarities. D uring the early tailphase SN 1990I shows a decline rate sim ilar to other SN e Ib/c, and faster than that of  $^{56}$ C o decay, indicating a low mass ejecta that allows the -rays to escape w ith low er deposition. Furtherm ore, light curves of SN 1990I show a marked change of slope. In fact, around day 230 the \V, R and I" light curves drop suddenly, while the B light curve tends to atten. A comparison w ith events that show changes in slope is made introducing the third to second slope ratio as a m easure of the how the SN changes its slope at late epochs. SN 1990I behaves in a completely di erent way compared to SN 1993J.

As a consequence the rapid and sudden drop is also seen in the colour evolution of SN 1990I, at an epoch sim – ilar to that of the already reported blueshift in the nebular lines. We interpret this rapid change, photom etrically and spectroscopically, to be due to dust condensation in the ejecta of SN 1990I. The dust condensation at a time as early as 250 d (since explosion) could be due to the rapid cooling and low ejecta in SN 1990I com pared to SN 1987A and SN 1999em which show evidence for dust form ation around 530 d and 500 d, respectively.

We constructed the \BVRI" quasi-bolom etric light curve, and compared it to that of SN 1993J, revealing a similarity in the [30:100] time interval. After 200 d, the two SN e evolve di erently: while SN 1993J tends to atten, SN 1990I falls dram atically. Then, in order to estim ate physical parameters of the SN, we applied a simple -ray deposition model. The model allows us to estimate the ejecta and <sup>56</sup>N i masses (M (<sup>56</sup>N i) = 0:11 M and M<sub>ej</sub> = 3:7 M ). Still to note how ever that more accurate light curve modelling should include asymmetry and clumping e ects.

D raw ing conclusions about the progenitor nature of SN 1990I (type Ib/c in general), based upon our present understanding of stellar evolution and the properties of the SN-explosion, is still speculative. The great unknown is whether this class of object has progenitor stars of sim - ilar (ie 15 20 M ) or higher (ie > 20 M ) m asses com pared to type II SN e.

The oxygen mass derived for SN 1990I is of the order

1 M , which is predicted to form in ZAMS stars of 15 20 M according to the standard theory of evolution in massive stars. We note here that SN 1987A was found to erect 1:5 2 M of oxygen. This would mean that the SN 1990I progenitor was slightly lower in mass with respect to SN 1987A. Further support for the lower mass case com es from the estimated helium core (  $5^2$  M for SN 1990I and 6 M for SN 1987A). W hat remains, how ever, is that SN e Ib/c eject a m ass of <sup>56</sup>N i in the range  $0:\!15~\text{M}$  , a higher amount compared to that from 0:1 typicaltype II SNe (i.e. 0:07 M for SN 1987A).

A lthough how the <sup>56</sup>N im ass scales with the progenitor m ass remains unclear in core collapse SN e, a possible solution for this (56N i yield M (progenitor)<sup>00</sup> paradox can be related to the binary nature of type Ib/c progenitors (Tutukov & Chugai. 1992) whereby the iron yield is very sensitive to rotation induced in the close binary system. In deed, very recently H irschi et al. (2004) have tested the e ects of rotation on evolution of m assive stars. The direct consequence of rotation on pre-SN m odels is o increase the core sizes especially for stars in the 15 25 M m ass interval. The <sup>56</sup>N i m ass ejected in the explosion is found to increase by a factor of 1.5.

It appears then that low m ass progenitors within the context of close binary system evolution (m ass-loss due to m ass transfer ) are m ore \probable" com pared to single m assive stars (i.e. W olff ayet stars losing m ass through w inds).C learly, large and w ellsam pled type lb/c SN e data (spectra and photom etry) together with m ore precise estim ates of M  $_{\rm ej}$ , M ( $^{56}$ N i), M (oxygen) are needed in order to build an accurate physical understanding of SN lb/c explosions.

The sudden fall in lum inosity at day 230 results from the fact that the light curve in Fig. 11 is not a real bolom etric light curve. A fler day 230 the dust absorbs at shorter wavelengths and re-em its at much longer IR wavelengths, consistent with the dust having an am bient tem perature which is probably lower than 1500 K and decreasing with tim e. This contribution of IR radiation from the dust has not been included. W hat remains to be understood, how ever, is why dust has form ed in SN 1990I but has not been recorded in other type Ib/c. Possibly inadequate sam pling of light curves and spectra of type Ib/c at nebular phases is at least partially responsible.

This paper dem onstrates the need to obtain good tem – poral coverage in both photom etry and spectroscopy with su cient S/N to avoid comprom ising diagnostic capatibilities.

A cknow ledgements. This work is based on data obtained at CTIO and ESO-La Silla.We would like to thank all observational astronomers, in particular N. Suntze, M. Hamuy, M. Navarrete, H. Bond, A. Crotts and L. wells. A. Elm ham di thanks B. Leibundgut for discussion and constructive com – ments on the original manuscript. A. Elm ham di is grateful to the ESO (European Southern Observatory) support that allowed his stay at Garching, where this work was concluded. W e are grateful to the referee (S.E.W oosley) for helpfulcom - m ents.

## References

- Barbon R. et al. 1990, A & A S, 237, 79
- Baron E. 1992, MNRAS, 255, 267
- Benetti S., Branch D., Turatto M. et al. 2002, MNRAS, 336, 91
- BesselM .S.1979, PASP, 91, 589
- Branch D., Benetti S., Kasen D. et al. 2002, ApJ, 566, 1005.
- Buonanno R., Buscem a G., Corsi C. E, Ferraro I. & Iannicola G. 1983, A & A, 126, 278
- Buta R. 1984, IAU Circ., No.3981
- Cardelli J.A., Clayton G.C. & Mathis J.S. 1989, ApJ, 345, 245
- ChugaiN N.1994, ApJ, 428, 17
- ChugaiN.N.2000, AstL, 26, 797
- ClocchiattiA.& W heeler J.C. 1997, ApJ, 491, 375
- Danziger I.J., Bouchet P., Goui es C., Lucy L.B. 1989, IAUC 4746
- Danziger IJ., Lucy L.B., Bouchet P., Goui es C. 1991, In \Supernovae", p. 69, ed. W oosley S.E., Springer-Vergal, New York.
- de KoolM ., LiH . & M cC ray R . 1998, ApJ, 503, 857
- D eng J.S., Q iu Y.L., H u J.Y., H atano K. & Branch D. 2000, A pJ, 540, 452
- Elm ham di A., Danziger I. J., Chugai N. N. et al. 2003, MNRAS, 338, 939
- Ensman L.M. & Woosley S.E. 1998, ApJ, 333, 754
- Filippenko A.V. 1988, AJ, 96, 1941
- Filippenko A.V., Porter A.C. & Sargent W . 1990, AJ, 100, 1575
- Filippenko A.V. 1997, ARA & A, 35, 309
- Fransson C., 1987, in ESO Workshop on \the SN 1987A", Garching, European Southern Observatory, p. 467
- Fransson C., Chevalier R.A. 1987, ApJ, 322, 15
- Fransson C., Chevalier R.A. 1989, ApJ, 343, 323
- Fransson C. et al. 1993, in \Supernova and Supernova rem nants", M CCRAY and W ANG, X ian, China, Colloquium 145 p.211
- Galam a T.J. et al. 1998, Nature, 395, 670
- HirschiR, Meynet G. & A. Maeder. 2004, accepted for publication in A&A (astro-ph/0406552)
- Houck J.C. & Fransson C., 1996, ApJ, 456, 811
- Je ery D.J.1999, astro-ph/9907015.
- Leibundgut B. et al, 1991, ApJ, 372, 531
- Lew is J.R. et al. 1994, MNRAS, 266, 27
- Lucy L.B. 1991, ApJ, 383, 308
- Lucy L.B., Danziger, I.J., Goui es, G., Bouchet, P. 1991, in \Supernovae", ed. S.E.Woosely (New York: Springer), p.82
- M cK enzie E.H.& Schaefer B.E. 1999, PASP, 111, 964
- Meikle W .P.et al. 1984, IAU Circ., No.4003
- Nadyozhin D.K. 1994, ApJS, 92, 527
- Nomoto K., Filippenko A.V.& Shigeyam a T.1990, A & A, 240
- N om oto K " M aeda K " N akam ura T . et al. 2000, hgrb sym p, 622
- Patat F. et al. 2001, AJ, 555, 900
- Phillips M.M. 1990, IAUC .5032.

 $<sup>^2</sup>$  The ejected m ass 3.7 M  $\,$  and an assum ed rem nant of 1.3–1.6 M  $\,$ 

- Phillips M . M . & W illiam s R . E ., 1991, In \Supernovae", p.
- 36,ed.W oosley S.E., Springer-Vergal, New York.
- Pizarro O., Miranda J., Pasquini L. et al. 1990, IAUC .5003.
- Richm ond M .W .et al. 1994, AJ, 107, 1022
- Richm ond M .W .et al. 1996, A J, 112, 732  $\,$
- SchlegelE.M.& KirshnerR.P.1989,AJ,98,577
- SchlegelD.J., FinkbeinerD.P. & Davis M. 1998, 500, 525
- Shigeyam a T ., N om oto K ., T su jin oto T ., H ashim oto M .1990, A pJ, 361, 23
- Sollerm an J., Leibundgut B. & Spyrom ilio J. 1998, A & A, 337, 207
- Sollerm an J. et al. 2000, ApJ, 537, 127
- Sollerm an J. et al. 2002, A & A , 386, 944
- Spyrom ilio J. & Pinto P.A. 1991, In Proc. ESO=EPIC W orkshop on \SN 1987A and other Supernovae", p.423, ed.Danziger I.J., ESO, M unich.
- Spyrom ilio J. 1994, M N R A S, 266, 61
- Suntze N.B.& Bouchet P.1991, In \Supernovae", p.12, ed. W oosley S.E., Springer-Vergal, New York.
- Swartz D.A.& W heeler J.C. 1991, ApJ, 379, 13
- Swartz D.A. et al. 1993, Nature, 365, 232
- T svetkov D . Y . 1985, A stron Zhur., 62, 365
- T svetkov D.Y. 1987, PAZh, 13, 894
- T svetkov D.Y. 1988, Perem. Zvezdy, 22, 653
- Tutukov A.V.& ChugaiN.N.1992, SvAL, 18, 242
- U om oto A.1986, ApJ, 310, 35
- W heeler J.C. et al. 1987, ApJ, 313, 69
- W heeler J.C. & Harkness R.P. 1990, Rep. Prog. Phys., 53, 1467
- W oosley S.E., Hartmann D., Pinto P.A. 1989, ApJ, 346, 395
- W oosley S.E., Eastman R.G. & Schmidt B.P. 1999, ApJ, 5167, 88
- Yamaoka Y. & Nomoto K. 1991, In Proc. ESO=EPIC Workshop on \SN 1987A and other Supernovae", p. 193, ed.Danziger I.J., ESO, Munich.
- Young.T.R., Baron E.& Branch D. 1995, AJ, 449, 51.