A Radio and M id-Infrared Survey of Northern Bright-R in m ed C buds

LK.Morgan, MA.Thompson, JS.Urguhart, Glenn J.White & J.Miao

Centre for A strophysics & Planetary Science, School of Physical Sciences, University of Kent, Canterbury, Kent CT27NR,UK e-m ail: 1km8@kent.ac.uk

Abstract. We have carried out an archival radio, optical and infrared wavelength in aging survey of 44 Bright-Rimmed Clouds (BRCs) using the NRAO/VLA Sky Survey (NVSS) archive, in ages from the Digitised Sky Survey (DSS) and the M idcourse Space eX perim ent (MSX). The data characterise the physical properties of the Ionised Boundary Layer (IBL) of the BRCs. We have classi ed the radio detections as: that associated with the ionised cloud rims; that associated with possible embedded Young Stellar Objects (YSOs); and that unlikely to be associated with the clouds at all. The stars responsible for ionising each cloud are identi ed and a com parison of the expected ionising ux to that measured at the cloud rims is presented. A total of 25 clouds display 20 cm radio continuum em ission that is associated with their bright optical rims. The ionising photon ux illum inating these clouds, the ionised gas pressure and the electron density of the IBL are determ ined. We derive internal m olecular pressures for 9 clouds using m olecular line data from the literature and com pare these pressures to the IBL pressures to determ ine the pressure balance of the clouds. We nd three clouds in which the pressure exerted by their IBLs is much greater than that measured in the internal molecular material. A comparison of external pressures around the remaining clouds to a global mean internal pressure shows that the maprity of clouds can be expected to be in pressure equilibrium with their IBLs and hence are likely to be currently shocked by photoionisation shocks. We identify one source which shows 20 cm emission consistent with that of an embedded high-m ass Y SO and con m its association with a known infrared stellar cluster. This embedded cluster is shown to contain early-type B stars, in plying that at least som e BRCs are intim ately involved in interm ediate to high m ass star form ation.

cbuds - ISM :molecules

1. Introduction

It has bong been suspected that the illum ination of dense, quiescent clum ps within m olecular clouds by nearby OB stars could be responsible for their triggered collapse and subsequent star formation. BRCs associated with old ($\& 10^6$ yr) HII regions are potential examples of Radiatively Driven Implosion (RDI) in which the UV ux of the associated OB star ionises the external layers of the cloud and causes the BRC to collapse. The ionisation front moves slowly into the globule, creating a dense outer shell of ionised gas (the Ionised Boundary Layer or IBL), which streams radially away from the cloud surface. The high density in the shell causes a high recombination rate which shields the globule from quickly evaporating (Repurth, 1983). The increased pressure in the ionised gas causes an expansion of the BL into the intercloud medium and an ionisation front preceded by a shock in the neutral gas propagates into the

Send o print requests to: L.K. Morgan

Key Words.stars:formation - ISM : HII regions - ISM : cbud (White et al., 1997; Le och & Lazare, 1994). The UV radiation from the OB stars often sweeps the molecular material of the cloud into a cometary morphology with a dense core boated at the head' of the com etary globule.

> The ratio of the internal pressure of the molecular cloud (mostly due to turbulent motions, but with a small therm al contribution) to the external pressure of the BL is predicted by RD Im odels to be key to the evolution of the cloud. If the BL has a pressure greater than, or equal to, the turbulent and thermal pressure of the cloud then photo-ionisation shocks and a D-critical ionisation front propagate into the cloud interior, compressing and heating the molecular gas (Bertoldi, 1989; Le och & Lazare, 1994). However, if the interior pressure of the cloud is greater than the pressure created by the ionised gas within the BL then the ionisation front stalls at the surface of the cloud until the e ects of mass evaporation from the cloud (due to the boiling o ' of particles by the ionising radiation) and increasing recombination within the BL raises the ionised gas pressure to equilibrium with the interior pressure (Le och & Lazare, 1994). As this equilibrium is reached, the evolution of the

cbud follows the same path as the initially underpressured cbud case. An analysis of the pressure balance present in BRCs thus allows a separation of shocked and unshocked cbuds. Our aim is the identi cation of shocked cbuds, as these cbuds are in portant subjects for future study and m odelling to enable a greater understanding of the processes involved in triggered star form ation.

The general conditions prevalent within the IBLs of BRCs are not well known as most studies have, to date, accurately determined the conditions within only a few individual clouds (e.g.Thompson et al. 2004; White et al. 1999; Megeath & Wilson 1997; Le och et al. 1997; Le och & Lazare 1995). Recently, Thompson, Urquhart, & White (2004) measured the IBL pressures and electron densities for 18 clouds from the Sugitani and Ogura southern hem isphere BRC catalogue (Sugitaniand Ogura, 1994). As a companion paper to Thom pson, Urquhart, & White (2004) we have utilised archival data to extend their analysis of the Sugitani and Ogura southern catalogue to incorporate the northern hem isphere Sugitani, Fukui & Ogura (SFO) catalogue (Sugitaniet al, 1991) which identied 44 BRCs associated with RAS point sources which are therefore candidates for RD I.

From observations of the free-free emission associated with BRCs identied in the SFO catalogue we are able to determ ine the ionised gas pressure, the ionising photon ux in pinging upon the cloud surface and the electron density of the BL. This can be achieved using data obtained from the NRAO VLA 20 cm Sky Survey (NVSS) (Condon et al., 1998). M idcourse Space eXperim ent (M SX) in ages at 8.3 m allow us to trace the Photon D om inated Regions (PDRs) associated with the BRCs, con ming the identication of individual sources of 20 cm em ission as BLs. The identi cation of the prim ary ionising stars associated with each BRC through a literature search allows us to compare the 20 cm em ission m easured toward each BRC with that predicted by the incident ionising ux. Through further quanti cation of these properties of BLs at the edge of BRCs we hope to clarify the role of the RD I scenario in global star form ation. To this end we have collated information using a thorough literature search and the SIM BAD database of astronom icalcatabgues (http://simbad.u-strasbg.fr) aswellas analysing optical, radio and infrared in ages to identify the stars responsible for ionising the clouds.

2. Survey Procedure

Wе obtained 20 am radio in ages from the radio 1998) NVSS catabque (Condon et al., dow n baded using the postage stam p server (http://www.cv.nrao.edu/nvss/postage.shtml). The NVSS was a 20 cm sky survey complete north of = -40 carried out using the VLA in its D con guration. The angular resolution of the NVSS is 45^{00} (Condon et al., 1998) and the limiting 1 noise of the survey is 0.5 m Jy. Our sample comprises 44 15° 15° in ages centred upon the coordinates of each $\mathbb{R}\mathsf{A}\mathsf{S}$ source associated with the 44 BRCs contained in the SFO catalogue.

Ionising radiation excites Polycyclic A rom atic Hydrocarbon (PAHs) which re-emit the absorbed energy at infrared wavelengths (Leger & Puget, 1984). MSX 8.3 m in ages incorporate the PAH spectral features at 7.7 m and 8.6 m as well very small dust grain continuum emission. In ages of the cloud sample were acquired in order to trace any associated PAH emission and con m that the 20 cm emission at the cloud rims is free-free continuum emission from the BL. The MSX surveyed the entire galactic plane within the range joj 5 in four mid-infrared spectral bands between 6 and 25 m at a spatial resolution of 18.3⁰⁰ (Price et al., 2001). The A band of the MSX corresponds to a wavelength of 8.3 m and is the most sensitive of the MSX bands. The A band of the MSX ranges from 6.8 - 10.8 m and includes the broad silicate band at 9.7 m and the PAH emission features at 7.7 m and 8.6 m, which are of particular interest to this study as they trace the interface between molecular material and ionisation features associated with 20 cm emission. In ages were downbaded from the NASA/PAC Infrared Science Archive (http://irsa.ipac.caltech.edu) and were analysed using GAR, part of the Starlink Software Collection. The MSX 8.3 m images were smoothed to a resolution of $45^{\circ\circ}$, to match that of the NVSS images and improve the signal to noise ratio. The rm .s. noise of each individual in age was determined from o -source sky m easurements and ranged from 1.5 – 39.1 $\,$ 10 5 W m 2 sr^1 .

3.Results and Analysis

3.1. Source Identi cation and Classi cation

Using NVSS data we have studied the BRCs from the survey of Sugitaniet al. (1991). Identi cation of bright rims was initially achieved by overlaying contours of the 20 cm em ission onto the R-band in ages of the clouds obtained from the Digitised Sky Survey (DSS) and searching for 20 cm em ission that is positionally coincident with the bright optical rims of the clouds. We detected no 20 cm emission (to a level of three times the rm .s. noise) associated with the rims of the clouds SFO 3, 8, 9, 19, 20, 22, 23, 24, 26, 33, 34 and 39. The images of SFO 16, 17 & 18 were disregarded due to their relatively low quality, which is likely to be due to the sidebbes of nearby confusing sources com bined with the low surface brightness of the 20 cm rim emission from these cbuds. For the cbuds SFO 2 & 44 there are no M SX data to assist with the identi cation, however, as the emission is clearly extended the inclusion of these sources was felt to be justi ed. A total of 26 radio sources were determ ined to be positionally associated with the BRCs from the SFO catabgue and the coordinates of the peak emission, peak uxes and integrated ux densities of these sources are presented in Table 1.DSS in ages overlaid with NVSS and MSX 8.3 m emission contours are presented in Fig. 2, centred on

the RAS source position given in Sugitaniet al. (1991) with a eld of view of 2^0 unless otherwise stated. In ages of the clouds SFO 2 and 44 are presented separately in Fig. 3 as they consist of NVSS emission contours only.

The radii of the optical bright rins (taken from Sugitaniet al. 1991) of the clouds toward which 20 cm emission has been detected range from $13 - 133^{00}$, corresponding to a physical range of 0.04 - 0.83 pc. From the source counts of Condon et al. (1998) we determ ine that there is a probability of P (< r) 0.2 of nding a background radio source within 133^{00} of the IRAS point. This in plies that up to 5 sources m ay potentially be due to confused background radio galaxies. In order to rule out any possible confusion the NVSS archival data were compared with opticalD igitized Sky Survey (DSS) in ages and m id-infrared M SX 8.3 m in ages.

PAH emission is a tracer of UV-dominated PDRs (Leger & Puget, 1984) as the PAHs are transiently heated by the absorption of UV photons, therefore M SX in ages of the northern SFO catalogue can help to identify PDRs that are associated with the detected radio emission. Comparing emission in the optical, radio and infrared regimes helps to eliminate chance associations and can identify true emission from the bright rim. The radio sources detected in the NVSS survey images were classified according to the scheme of Thom pson, Urguhart, & White (2004):

- 1. Bright-rim em ission clouds with 20 cm and 8.3 m em ission positionally coincident with their bright optical rims.
- 2. Broken-rim med clouds, in which the 20 cm and M SX 8.3 m emission is positionally coincident with the rim of the cloud (as Type 1) but the rim has a reverse curvature with respect to the norm all orientation, i.e. the rim is curved towards the molecular cloud, rather than the ionising star (e.g. the well known broken-rim med globule CG4 in the Gum Nebula (Reipurth, 1983). No clouds of this type were identied in this survey, the classi-cation has been retained to maintain consistency with Thom pson, Urquhart, & White (2004).)
- 3. Embedded objects with compact and coincident 20 cm and mid-infrared emission that is set back from the rim toward the centre of the cloud.
- 4. 20 cm em ission that is uncorrelated with either the bright optical rim or M SX 8.3 m em ission.

3.2. Confusing Sources

We have tentatively labelled the sources SFO 6 & 27 as Type 1 sources though there is some confusion as to their identication. These objects shall now be examined in more detailand justication for their nominal identications given.

321.SFO 6

SFO 6 has strong (9) 20 cm em ission associated with the optical rim which appears to follow the optical bright rim

(Fig 2). There is a small region of 8.3 m emission centred at the peak of the optical rim which, due to the confused nature of the emission, is not de nitive enough to infer any clear association with the cloud rim but does not contradict an identication of the source as Type 1. However, the 20 om emission near the optical rim is confused by a stronger unassociated source that has no optical or infrared counterpart and is thus likely to be an extragalactic background source.W e suggest that this source may be NEK 135.2+00.3, identied in the Clark Lake 30.9 MHz galactic plane survey (Kassin, 1988). The source is unresolved in the survey and only marginally resolved in the NVSS in ages.

322.SFO 27

SFO 27 has strong (6) 20 cm emission associated with the optical rim (Fig.2). There is widespread M SX 8.3 m emission associated with the IRAS source and extending along the optical rim which supports an identi cation of the source as Type 1. However, the 20 cm emission near the optical rim is confused by a stronger unassociated source that has no obvious optical or infrared counterpart.

We have continued the analyses of SFO 6 and SFO 27 as Type 1 objects and have masked the emission from the background sources as much as it is possible to do so.

3.3. Ionised R in s Associated W ith SFO Objects

For those objects that have been identi ed as Type 1 sources and are likely true bright-rim candidates we have evaluated the ionising photon ux impinging upon the clouds, the electron densities and the pressures in the IBLs. These quantities were determined by using the general equations from Le och et al. (1997). Rearranging their Eq.(6), the ionising photon ux arriving at the cloud rim may be written in units of cm 2 s 1 as:

$$= 1.24 \quad 10^{10} \text{ S} \quad \text{T}_{0}^{0.35 \quad 0.1 \quad 2} \tag{1}$$

where S is the integrated radio ux in m Jy, T_e is the electron temperature of the ionised gas in K, is the frequency of the free-free emission in GHz and is the angular diameter over which the emission is integrated in arcseconds.

The 20 cm ux m easured at the ionised rim m ay possibly be overestin ated due to nebula emission from the H II regions in which the BRCs are embedded. This emission forms a background which m ay add to our actual rim brightness at 20 cm. However, due to the poor (u,v) coverage of the NVSS snapshot observations any structure larger than 5^0 is litered out (Condon et al., 1998), therefore if any large scale nebula emission is present we m ay expect it to be at a low level. This is supported by comparison of the DSS in ages, in which nebula emission shows up as large, di use, red regions, with NVSS data in which we nd no correlating 20 cm emission.

The electron density $(n_{\rm e})$ of the BL surrounding the cloud may also be derived from the integrated radio $\,$ ux

Table 1	. P roperties	ofRadio Em	ission	associated	with	the SFO	0 bjects.
---------	---------------	------------	--------	------------	------	---------	-----------

SFO object	peak em ission	peak em ission	Peak Flux	Integrated F lux	Source C lassi cation
	(2000)	(2000)	(m Jy/beam)	(m Jy)	
1	23 59 33.7	+ 67 23 54	69.2	728.4	1
2	00 03 50.3	+ 68 32 09	4.4	13.2	1
4	00 58 60.0	+ 60 53 16	5.3	26.6	1
5	02 29 01.8	+ 61 33 17	11.5	21.4	1
6	02 34 45.0	+ 60 48 18	4.6	42.9	1^{T}
7	02 34 47.8	+ 61 49 14	11.7	86.3	1
10	02 48 07.3	+ 60 25 35	30.7	519.6	1
11	02 51 33.0	+ 60 03 43	4.0	10.1	1
12	02 55 00.9	+ 60 35 44	12.0	127.0	1
13	03 00 52.5	+ 60 40 19	3.7	6.3	1
14	03 01 24 2	+ 60 29 12	8.1	41.9	3
15	05 23 26.5	+ 33 11 54	3.3	3.0	1
16	05 20 00.9	-05 50 22	1.1	0.6	4
17	05 31 27.1	+12 04 54	1.9	7.7	4
18	05 44 28.7	+ 09 08 40	8.1	9.1	4
21	05 39 38.3	-03 36 25	1.3	1.3	4
25	06 41 06.3	+10 14 30	3.0	15.5	1
27	07 04 03.8	-11 23 19	3.6	39.0	1^{T}
28	07 04 41.4	-10 22 15	1.3	0.6	4
29	07 04 54.5	-12 09 42	2,2	0.6	4
30	18 18 46,1	-13 44 39	57.1	125.6	1
31	20 50 48.8	+ 44 21 29	12.8	345.5	1
32	21 32 34.9	+ 57 24 08	2.9	5.0	1
35	21 36 09.3	+ 58 31 53	1.8	12.1	1
36	21 36 10.6	+ 57 26 34	3.8	11.5	1
37	21 40 29.0	+ 56 36 13	3.5	11.0	1
38	21 40 44.3	+ 58 15 01	6.0	26.6	1
40	21 46 09.1	+ 57 09 59	4.9	14.6	1
41	21 46 29.0	+ 57 18 10	2.7	28.5	1
42	21 46 38.6	+ 57 11 39	4.9	12	1
43	22 47 50 3	+ 58 02 51	35.3	204.5	1
44	22 28 59.1	+ 64 12 11	37.0	95.8	1

^T Tentative classi cation, see Section 3.2 for details

S by substituting for the ionising photon ux in Eq.(6) of Le och et al. (1997). The electron density in cm 3 is given by:

$$n_{e} = 122.41 \frac{S T_{e}^{0.35 \ 0.1 \ 2}}{R}$$
(2)

where those quantities common to both Eq.(1) and (2) are in the same units, R is the radius of the cloud in pc and is the elective thickness of the BL as a fraction of the cloud radius (typically 0.2, Bertokli 1989).

As the ionised ow is sonic at the cbud surface (Le och & Lazare, 1994) we must take into account the ram pressure of the ow, as well as the bulk pressure of the ionised layer. Assuming a totally ionised gas the total pressure of the ionised layer with respect to the internal neutral gas is:

$$P_{T} = P_{i} + {}_{i}c_{i}^{2} = 2 {}_{i}c_{i}^{2}$$
(3)

where c_i is the isothermal sound speed in the ionised interclump gas (assumed to be 11.4 km s¹ Bertoldi (1989)) and _i is the density within the BL.

Values for , n_{e} and $P_{\rm T}$ were calculated using Eqs.(1)–(3), assuming a boundary layer thickness fraction of =0.2 (Bertoldi, 1989) and an electrice electron temperature of $T_{e}=10^{4}$ K.As the relatively by angular resolution of the NVSS means that a number of BRCs are either unresolved or marginally resolved at 20 cm the cloud radii were taken from the original BRC survey paper of Sugitaniet al. (1991)). Values for , n_{e} and $P_{\rm T}$ are presented in Table 2.

In our derivation of the ionising uxes of these clouds we have in plicitly assumed that the bright rins are resolved, this is not always the case (e.g. SFO 15). A comparison of the ratio between predicted and measured ionising uxes in the resolved (e.g. SFO 38) and non-resolved case does not reveal any consistent e ect associated with this beam dilution'. Our analysis of the cloud SFO 5 reveals an ionising ux of 7.5 10^8 cm⁻² s⁻¹, whereas the (resolved) observations of Le och et al. (1997) nd 4.8 x 10^9 cm⁻² s⁻¹ using the same analysis. The di erence in results appears insigni cant considering the di erent frequencies of observation and total area of integrated em ission.Le och et al. (2002) nd a factor of 6 between predicted and measured ionising ux in their resolved observations of the Tri d nebula.However, the (also resolved) observations of SFO 5 presented in Le och et al. (1997) nd a sm all (0.8) di erence between their predicted and measured ionising uxes. W hile the non-resolution of the bright rims in som e cases is certainly undesirable and an important consideration in future observations it does not appear to be a signi cant factor in the ux com parisons presented here.

3.4. Identi cation of Ionising Stars

We searched the SIMBAD astronom ical database in order to identify possible ionising stars of each BRC.AllO or B type stars boated within each HII region were considered as possible candidate ionising stars. For each star the predicted ionising uses incident on the rim of the relevant SFO object were determ ined using the tables of Panagia (1973) and the stellar spectral type. If there was disagreement in the literature as to the speci cation of a star, or it was simply not quoted, it has been assumed that the star in question is an evolved main sequence star (Type V). A list of stars with high predicted ionising uxes (i.e. nearby OB stars) was drawn up. Stars that contributed less than 50% of the ux of the most dom inant star were discounted. The stars that have been identied as the main ionising stars of each SFO object are listed in Table 3 along with their positions, the HII region with which they are associated and their spectral type.

3.5. Flux Comparison

A comparison of the 20 cm ux measured from the NVSS in ages with the predicted Lym an continuum ux allows us to determ ine the likelihood that the optical bright rims are due to ionisation by the nearby stars identi ed in Section 3.4. From the tables of Panagia (1973) we have identied the uxes of Lyman continuum photons associated with each star and thus calculated the subsequent ux we expect to impinge on the optical bright rims. We have assumed that there are no bases due to absorption by intervening material between the star and cloud, also, the star-cloud distance used in our calculations is that seen in projection in the plane of the sky and together these assumptions lead to the predicted ionising ux being a strict upper limit. It should be noted that a m isclassi cation of an ionising star may lead to large di erences in the ionising ux that we predict, a m isclassi cation of half a spectral class m ay lead to an increase or decrease of a factor of two in the predicted Lym an photon ux. There are sometimes disagreements in the literature on the spectral classi cation of a particular star, two examples of this are the stars HD 5394 and BD + 60 502 that we believe to be ionising the surfaces of SFO 4 and SFO 5 respectively. The form er has been identied as BOIV by Morgan et al. (1955) but as B3IV by Racine (1968) and the latter as 05 by Conti& Alschuler (1971) and B8 by Boulon & Fehrenbach (1958). In the cases of disagreement we have taken the type specified as the most reliable by the SIM BAD database.

The predicted ionising uses are presented along with the measured ionising uses in Table 2. It can be seen in the majority of cases that $_{\rm P}$ > as we expect, however, the dimension between $_{\rm P}$ and in some cases is much larger than we might expect. The disparity between $_{\rm P}$ and

in these cases is most likely due to the assumption that there is negligible absorption between the ionising star and the bright rim. Le och et al. (2002) nd an attenuation of a factor of 6 over an assumed distance of 1 pc. The assum ed distances in our sample range from 1 to 37 pc and we thus expect attenuation to have a signi cant e ect upon the ratio of predicted to measured ionising uxes. It is worth noting that the projected star-cloud distance is an underestim ate. The combination of these two facts illustrates that theoretical predictions of based upon the spectral type and projected distance of the ionising star offen overestim ate the true ionising photon ux illum inating these clouds. Another e ect which may a ect our results is that the 20 cm em ission from the bright rims was measured using an interferom eter which acts as a high pass liter and may thus liter out ux from large scale structures. In two cases (SFO 27 and SFO 31) the ux that has been predicted is signi cantly less than the ux that is observed. In the case of SFO 27 the 20 cm ux is confused with an unassociated source and so the measured 20 cm ux and hence the calculated value of are highly uncertain. In the case of SFO 31 the direction of the suspected ionising star is not supported by the morphology of the cloud and no other possible ionising stars have been identied in the region.

3.6. Upper Lin its to the Ionising Flux for Non-D etections

There are a total of 12 cbuds in our survey where no 20 cm em ission was detected to a level of 3 . We have checked the upper lim its of any possible 20 cm em ission from these clouds for consistency with the ux predicted from our candidate ionising stars. The predicted ionising ux was determined as in Section 3.3 using the stars and spectral types as laid out in Table 3. The 3 upper limits, subsequent maximum possible observed (i.e. 3) uxes and predicted ionising uxes are presented in Table 4. It can be seen that P > max in all but one case and that $P > m_{ax}$ by a factor of 10 in a large number (40%) of cases. These predictions are obviously inconsistent with the NVSS observations. In these cases the reasons for the discrepancies between theoretical prediction and measurement may mean that the predicted ionising starm ay be m isclassi ed orm isidenti ed, that extinction between the suspected ionising star and bright rim is signi cant, or that the projected distances between star and cbud are incorrect.

Table 2. Values for the measured ionising	ux, predicted ionising	ux, the m easured	electron density	and ionised gas
pressure for Type 1 radio sources detected	in the survey.			

	Measured ionising ux	Predicted ionising ux	E lectron D ensity	Ionised gas pressure
SFO Object	$(10^8 \text{ cm}^2 \text{ s}^1)$	$_{\rm P}$ (10 ⁸ cm 2 s 1)	n_e (cm 3)	$P_1 = k_B$ (10 ⁵ cm ³ K)
1	18.5	43.7	872	274.3
2	2.7	2.1	271	85.3
4	2.7	43.1	628	197.5
5	7.5	61.3	364	114.4
6	3.1	32.1	388	122.0
7	3.6	78.7	164	51.6
10	9.9	10.5	516	162.3
11	2.3	10.2	232	73.0
12	5.2	44.5	350	110.1
13	3.2	12.4	203	63.9
15	3.0	18.8	228	71.7
25	2.2	27.1	211	66.4
27	5.6	1.7	427	134.3
30	22.1	4065.9	454	142.8
31	8.7	3.5	483	151.9
32	2.2	11.4	396	124.6
35	1.2	6.9	204	64.2
36	2.4	59.6	175	55.0
37	2.0	10.2	357	112.3
38	2.8	13.3	248	78.0
40	2.5	8.1	348	109.5
41	1.5	8.0	248	78.0
42	1.3	7.3	250	78.6
43	4.8	145.8	277	87.1
44	2.1	4.6	269	84.7

3.7. Candidate Ultra-Com pact HII Regions

We have identied one 20 cm source in this survey as a Type 3 source, SFO 14. The 20 cm NVSS emission from SFO 14 follows a potential ionised cloud rin, although there is a localised peak coincident with the IRAS position which is clearly identiable in the MSX in age.

SFO 14 has the characteristics of a compact or ultracompact H II region: it is infrared-lum hous ($L_{\rm IR}$ $10^3 - 10^4$ L) and its IRAS colours match the criteria suggested by W ood & Churchwell (1989) for ultra-compact H II regions (nam ely that log F₆₀/log F₁₂ 1.3 and log F₂₅/log F₁₂ 0.57). To further investigate the possibility of this object being an embedded compact or ultra-compact H II region within the BRC we have used the Far InfraRed (FIR) and radio lum inosity of the source to estim ate the spectral class of any YSO that may be present. The FIR lum inosity of the YSO itself, reprocessed by the circum stellar dust in the region and re-em itted in the FIR.

In addition to HIRES IRAS uxes taken from in ages obtained from the NASA/IPAC Infrared Science Archive (http://isa.jpac.callech.edu) we have used the 450 m and 850 m uxes presented in Morgan et al. (2004, in prep.) as well as the 2 mm ux presented in Sugitaniet al. (2000) to estimate the FIR lum inosities of SFO 14. The lum inosity was estimated by integrating under a grey-body t using the IRAS 60 m ux together with the 450 m, 850 m and 2 mm ux measurements, the distance (1.9 kPc) was taken from Sugitaniet al. (1991). Further details about the SCUBA observations and grey-body modelling may be found in M organ et al. (2004, in prep.). The assumption has been made that all of the lum inosity arises from a single embedded YSO. This assumption may be rather crude but enables a rst-order estimate. W ood & Churchwell (1989) showed that, for a realistic initialm ass function, the spectral type of the most massive member in a cluster is only 1.5-2 spectral classes lower than that derived for the single embedded star case.

In a compact or ultra-compact H II region, the 20 cm ux is predominantly due to thermal free-free emission. If this emission is assumed to be optically thin and we assume that the region is in photoionisation equilibrium then we can relate the integrated radio ux of the region to the total number of ionising photons from the embedded YSO via Eq.(7) of Carpenter, et al. (1990).

 $N_{i} = 7:7 \quad 10^{43} \text{ S} \quad D^{2} \quad 0:1$ (4)

where N_i is the total number of photons per second ionising the cloud material, S is the integrated radio ux density in mJy, D is the distance to the source in kpc and is the frequency of the observation in GHz. Note that the 5 GHz term of Eq.(7) of Carpenter, et al. (1990) has been rem oved and the equation coe cient adjusted accordingly. Using the

SFO object	H II region	Ionising Star			Spectral	P rojected D istance
			(2000)	(2000)	Туре	to Bright Rim (Pc)
1	(S171/NGC 7822)	BD + 66 1675 ^a	00 02 10.3	+ 67 24 32	0 7V	3.7
2	(S171/NGC 7822)	BD +661675	00 02 10.3	+ 67 24 32	0 7V	16.9
3	(S171/NGC 7822)	BD +661675	00 02 10.3	+ 67 24 32	0 7V	5.3
4	(S185)	HD 5394 ^b	00 56 42.3	+ 60 43 00	BOIV	1.1
5	(S190/IC 1805)	BD +60 502 ^c	02 32 42.5	+ 61 27 22	0 5V	14.9
		$BD + 60 504^{\circ}$	02 32 49.4	+ 61 22 42	0 4V	16.1
		BD +60 507°	02 33 20.6	+ 61 31 18	0 5V	17.1
6	(S190/IC 1805)	BD + 60 504	02 32 49.4	+ 61 22 42	0 4V	20.5
7	(S190/IC 1805)	BD + 60 502	02 32 42.5	+ 61 27 22	0 5V	14.6
		BD + 60 504	02 32 49.4	+ 61 22 42	0 4V	16.6
		BD + 60 507	02 33 20.6	+ 61 31 18	0 5V	11.4
8	(S190/IC 1805)	BD + 60 502	02 32 42.5	+ 61 27 22	0 5V	12.4
		BD + 60 504	02 32 49.4	+ 61 22 42	0 4V	11.3
		BD + 60 507	02 33 20.6	+ 61 31 18	0 5V	11.1
9	(S190/IC 1805)	BD + 60 502	02 32 42.5	+ 61 27 22	0 5V	15.0
		BD + 60 504	02 32 49.4	+ 61 22 42	0 4V	14.4
		BD + 60 507	02 33 20.6	+ 61 31 18	0 5V	13.1
10	(S199/IC 1848)	HD 17505 ^d	02 51 08.0	+ 60 25 04	0 6V	12.3
11	(S199/IC 1848)	HD 17505	02 51 08.0	+ 60 25 04	0 6V	11.9
12	(S199/IC 1848)	BD +60586 ^d	02 54 10.7	+ 60 39 03	O 8III	3.9
13	(S199/IC 1848)	BD +59578 ^d	02 59 23.2	+ 60 33 59	0 7V	7.0
14	(S199/IC 1848)	BD + 59 578	02 59 23 2	+ 60 33 59	0 7V	8.6
15	(S236/IC 410)	HD 242908 ^e	05 22 29.3	+ 33 30 50	0 5V	22.1
		HD 242926 ^e	05 22 40.1	+ 33 19 10	0 6V	12.0
16	(S276)	0 rf	05 32 04.1	+ 00 21 55	09 . 5I	37.3
		¹ 0 ri ^f	05 35 15.5	-05 23 19	0 7V	26.7
		0 rĺ	05 35 26.8	-05 54 31	08.5II	26.9
17	(S264/ 0 ri)	0 ri ^f	05 35 08.2	+ 09 56 04	0 5V	16,3
18	(S264/ 0 mi)	0 ri	05 35 08,2	+ 09 56 04	0 5V	17.0
19	(S277/IC 434)	HD 37468 ^g	05 38 44.8	-02 36 01	09.5V	13,3
20	(S277/IC 434)	HD 37468	05 38 44.8	-02 36 01	09 . 5V	83
21	(S277/IC 434)	HD 37468	05 38 44.8	-02 36 01	09 . 5V	7.1
22	(S281)	¹ O ri ^h	05 35 15.5	-05 23 19	0 7V	6.5
		0 rh	05 35 26.8	-05 54 31	08.511	8.1

results of this equation we can use the tables of Panagia (1973) to nd the spectral type of any central YSO. The results of the above analyses are presented in Table 5. There is reasonably good agreem ent between the spectral type predicted through the di erent m ethods suggesting that there is a B0.5 - B1.5 YSO embedded within the molecular material of the clouds.

A literature search was carried out in order to identify the nature of the candidate ultra-com pact H II region. SFO 14 is a well studied region within the W 3/W 4/W 5 m olecular cloud com plex and is associated with the infrared stellar cluster AFGL 4029 (Deharveng et al., 1997). Snellet al. (1988) report the presence of a strong m olecular out ow and Carpenter, Heyer, & Snell (2000) nd that the region contains an embedded cluster containing 240 10 stars. The region has been found to contain a lum nous red YSO invisible in the optical, associated with a cluster of m assive red stars; an optical and IR re ection nebula; a high velocity ionised stellar wind; an optical jet and a bright H_2 em ission knot (Deharveng et al., 1997). The spectral type of the m ost massive star in this cluster inferred from the NVSS 20 cm $\,$ ux is consistent with that derived by D ehaveng et al. (1997) from infrared observations.

The BRC identied as a potential high mass star-forming site has been con im ed as containing an em bedded cluster. The spectral class of the star within this cluster has been determ ined as being early B type and thus relatively young with a main sequence lifetime of 10⁶ years. A molecular out ow is present also (Snellet al, 1988) and so star formation is still occurring in this region. The region of em ission is larger than the NVSS 45⁰⁰ beam (corresponding to 0.4 pc at the assumed distance) at the 3 level and so is not an ultra-compact HII region, the main sequence lifetime for the most massive predicted YSO is 10^6 years and so we can assume an age of the embedded cluster that is younger than this. The radii of the 3 contours of the embedded cluster is 1.1 pc. Given a propagation speed of 0.903 pc M yr^1 (Le och & Lazare , 1994) we nd that the radio region would have taken 12x106 years (sim ilar to

Table 3. (cont.)	Ionising stars of the SFO	objects, their rela	evant H II regions, i	positions and ionising	uxes.
------------------	---------------------------	---------------------	-----------------------	------------------------	-------

SFO object	H II region	Ionising Star			Spectral	P rojected D istance
			(2000)	(2000)	Т уре	to Bright Rim (Pc)
23	(S249)	BD +22 1303 ⁱ	06 22 58 2	+ 22 51 46	0 9V	83
24	(S275/NGC 2244)	HD 46223 ^j	06 32 09.3	+ 04 49 25	0 5V	19.3
25	(S273/NGC 2264)	HD 47839 ^k	06 40 58.6	+ 09 53 45	0 7V	4.7
26	(S296/CM a0 B I)	HD 54662^1	07 09 20 2	-10 20 48	O 7III	29.7
27	(S296/CM a0 B I)	HD 53456 ¹	07 04 38.3	-11 31 27	BOV	3.9
28	(S296/CM a0 B I)	HD 53367 ¹	07 04 25.5	-10 27 16	BOIV	2.1
29	(S296/CM a0 B I)	HD 53975	07 06 36.0	-12 23 38	07.5V	9.5
30	(S49)	BD -13 4927 ^e	18 18 40.1	-13 45 18	0 5V	1.0
31	(S117)	HD 199579 ^m	20 56 34.7	+ 44 55 29	0 6V	20.5
32	(S131/IC 1396)	HD 206267 ⁿ	21 38 57 . 6	+ 57 29 21	0 6V	11.3
33	(S131/IC 1396)	HD 206267	21 38 57 . 6	+ 57 29 21	0 6V	10.5
34	(S131/IC 1396)	HD 206267	21 38 57 . 6	+ 57 29 21	0 6V	12.1
35	(S131/IC 1396)	HD 206267	21 38 57 . 6	+ 57 29 21	0 6V	14.5
36	(S131/IC 1396)	HD 206267	21 38 57 . 6	+ 57 29 21	0 6V	4.9
37	(S131/IC 1396)	HD 206267	21 38 57 . 6	+ 57 29 21	0 6V	11.9
38	(S131/IC 1396)	HD 206267	21 38 57 . 6	+ 57 29 21	0 6V	10.4
39	(S131/IC 1396)	HD 206267	21 38 57 . 6	+ 57 29 21	0 6V	12.6
40	(S131/IC 1396)	HD 206267	21 38 57 . 6	+ 57 29 21	0 6V	13.4
41	(S131/IC 1396)	HD 206267	21 38 57 . 6	+ 57 29 21	0 6V	13.5
42	(S131/IC 1396)	HD 206267	21 38 57 . 6	+ 57 29 21	0 6V	14.1
43	(S142/NGC 7380)	HD 215835	22 46 54 2	+ 58 05 04	0 5V	5.4
44	(S145)	HD 213023°	22 26 52.4	+ 63 43 05	0 9V	8.5
		BD 62 2078 ^p	22 25 33.6	+ 63 25 03	07 . 5V	13.8

Stars found to be at sim ilar distances to, form ally declared to be associated with or described as probable m embers' of relevant H II regions in the following references: ^a C ram pton & F isher (1974); ^bSavage, et al. (1977); ^c Ishida (1970); ^d Hughes, et al. (1978); ^e H um phreys (1978); ^f Reynolds & Ogden (1979); ^g K uiper (1975); ^h O 'D ell & D oi (2003); ¹H ardie, et al. (1960); ^j O gura & Ishida (1981); ^k G ies et al. (1997); ¹C laria (1974); ^m Bally & Scoville (1980); ⁿ M atthews (1979); ^o C havarria-K. et al. (1994); ^p Sugitani et al. (2000)

Table 4. Values for the 3 levels, maximum possible measured ionising ux and predicted ionising ux, for radio sources not detected in the survey.

	3	ux upper lim it	Max.observed ionising ux	Predicted ionising ux
SFO Object		(m Jy)	$_{\rm max}$ (10 ⁸ cm 2 s 1)	$_{\rm P}$ (10 ⁸ cm 2 s 1)
3		2.04	2.02	21.15
8		1.52	1.78	118.31
9		2.10	2.02	78,22
19		0.75	0.51	0.57
20		1.40	1.47	1.47
22		1.78	1.93	22.83
23		1.91	2.07	2,52
24		2.69	2.17	11.52
26		1.37	1,22	1.06
33		1.55	1.13	13,29
34		1.40	1.48	9.98
39		1.72	1.85	9.18

the main sequence lifetime of a BO .5 star) to have formed as a consequence of the embedded clusters' in uence. Using conservative estimates the cluster is su ciently old that it may have been in uenced by the propagation of photoionisation shocks into the cloud. The distance of the core from the optical bright rim is 0.56 pc, assuming a shock velocity of 1 km s¹ (Thompson et al., 2004; W hite et al., 1999) we not a shock crossing time of 5×10^5 years. This is not conclusive evidence that the star exciting the bright rim of this cloud has a ected the star-forming processes within

the cloud, how ever, neither can the possible in uence of this starbe ruled out.

Accurate ages are needed both for the star embedded within the cluster and for the star(s) that are ionising the region, which will allow a more certain link between the development of the embedded cluster and the ionising star(s) to be drawn. Theoretical work is underway to analyse and re-create the large scale properties of the clusters, this will lead to conclusive support for or against the RD I scenario. D exper observations at m id-infrared wavelengths are needed to identify any embedded clusters which are still unknown and m olecular observations will help to provide better constraints on the internal pressures of a larger sample of BRCs.

4.D iscussion

25 cbuds from this survey were classied as Type 1 radio sources, i.e. with 20 cm emission clearly associated with their photoionised cloud rims. The pressures from their BLs were estimated in Section 3.3 and, togetherwith data on the m olecular interior of the clouds these allow us to determ ine the pressure balance between the IBL and the internal molecular material of the clouds. In the RDI scenario a photoionisation driven shock propagates into the molecular cbud and if the internal pressure of the cbud is greater than or equal to the pressure in the developing IBL, the shock stalls at the surface of the cloud (Le och & Lazare, 1994). Evaporation of the cloud ensures that eventually (provided that the source of ionisation is maintained) the surface area of the cloud will decrease, leading to a relative increase in the BL pressure and hence allowing the propagation of a D-critical ionisation front.

If the clouds that we observe are currently overpressured with respect to their BL then we do not expect there to be (or have been) photoionisation shocks propagating into the cloud and it is therefore unlikely that any existing star form ation in these clouds could have been caused via the RD I scenario. If the clouds are underpressured with respect to the BL then we may expect photoionisation shocks to be propagating into the cloud and the possibility exists that RD I is the cause of star form ation in the region. Thus the pressure balance between the BL and m olecular m aterial of the clouds acts as a simple diagnostic of the clouds whose star form ation may have been triggered by their UV illum ination.

The data that we require on the molecular interior of these clouds may be derived through molecular line observations. There are a small number of such observations that have previously been published and these form the basis for our comparison of internal molecular pressures with those determined for BLs in this work.

4.1. Molecular Pressures

The pressure of the molecular gas within the clouds in our sample is a result both of turbulent velocities within the cloud interiors and the therm all pressure. As the clouds are composed mostly of cold gas there is a negligible therm all contribution to the internal pressure or any observed linewidths relating to the molecular material of these clouds. Given molecular line observations we may derive the internal molecular pressures of our cloud sample from the turbulent velocity dispersion (²) and the molecular gas density $_{\rm m}$: We use the relation between molecular pressure $P_{\rm m}$, the square of the turbulent velocity dispersion 2 and the density of the

molecular gas m; i.e. P_m ' 2 m. The turbulent velocity dispersion m ay be written in terms of the observed linew idth as $^2 = \langle \rangle ^2/(8\ln 2)$.

Observations of line em ission from the northern SFO objects were collected via a SIM BAD search speci c to the SFO catalogue. We have used the linew idth taken from the $C^{18}O$ (J = 1 ! 0) observations of Jansen et al. (1994) and their value for H₂ density to determ ine an internalm olecular pressure for SFO 4, the values for the internalm olecular pressure and density of SFO 5 are derived from the results of the $C^{18}O$ (J = 1 ! 0) observations of Le och et al. (1997). Values for the internal pressure and density of SFO 11 have been taken from Thom pson et al. (2004).

De Vries, et al. (2002) conducted a millimetre and submillimetre molecular line survey of BRCs selected from the SFO catabque. In their survey they observed $C^{18}O$, HCO^+ and N_2H^+ (in addition to other species) emission from the cbuds. The core densities of the cbuds SFO 16, 18 and 25 were determined from the N_2H^+ observations of De Vries, et al. (2002) as it is likely to be optically thin in our cloud sample and hence a good tracer of high colum n density. W e have assum ed spherical geom etry and used the radii from Sugitaniet al. (1991). SFO 13 was not detected in N_2H^+ by De Vries, et al. (2002) and its density was thus determ ined from their HCO⁺ observations. Internal molecular pressures for these four clouds were then determined from the linewidths of the $C^{18}O(J = 1 ! 0)$ observations presented in the same work. $C^{18}O(J = 1 ! 0)$ linewidths and core densities from Cernicharo et al. (1992) and Duvert, et al. (1990) were used to calculate internal pressures for SFO 20 and 37 respectively in preference to the observations of D e V ries, et al. (2002) due to their superior angular resolution.

The IBL pressures calculated from the NVSS data range from 51.6 to 274.3 10^5 cm 3 K/k_B with a mean of 107.7 10^5 cm 3 K/k_B while the molecular line data that has been collected show a range of internalmolecular pressures of 0.3 $-32.3 \ 10^5$ cm 3 K/k_B with a mean of 12.3 10^5 cm 3 K/k_B (Table 6).

Factors contributing to the uncertainties in our values for internal molecular pressure include the accuracy involved in m easuring linew idths and the uncertainty for radii for these clouds taken from Sugitaniet al. (1991) (including the assumption of spherical symm etry made in determining volum e densities from column densities). We estimate that uncertainties in the internal molecular pressures presented total no more than a factor of 5.

The masses derived from the column densities given in D e Vries, et al. (2002) concurwellw ithin these uncertainties with values for cloud masses derived from submillimetre observations presented in Morgan et al. (2004, in prep.). Of the six clouds for which we have analysed the pressure balance three (SFO 4, SFO 13 and SFO 37) exhibit large enough di erences in the pressures determined due to their internal turbulent motions and due to the IBL that it is highly likely that they are in a state of photoionisation induced pressure in balance (see Fig. 1), even accounting for the large enous inherent within this study. These clouds are

Table 5. Infrared and radio-derived spectral types for the Type 3 radio sources detected in the survey.

SFO Object	IRASPSC ID	IR Lum inosity	SpectralType	F lux D ensity	Ionising Photon Flux	SpectralType
		(L_{IR}/L)	(IR)	S (m Jy)	$Log(N_i)(s^1)$	(Radio)
14	02575+ 6017	8321	B0.5	427.7	47.1	B15

therefore excellent candidates for testing the theories of RD I and comparison to BRCs which are known to be in pressure balance (or in balance) with their BLs.

In addition to the clouds that we have identi ed here as being in pressure in balance with their IBL's, Cernicharo et al. (1992) present observational evidence that SFO 20 (OR FH2) is undergoing RD I. They nd a bipolar out ow emanating from the source and while their evidence is not de nitive they conclude that the UV eld present due to the illum inating star -O ri has strongly modi ed the evolution of the primitive globule and strongly increased its star form ation potential.



F ig.1. G raph of internal molecular pressure versus IBL pressure. Error bars in IBL pressure represent an approximate 75% uncertainty in the assumption that $T_e = 10^4$. The solid line indicates pressure balance between the internal and external pressures.

The mean internal pressure of the clouds for which we have molecular line data is $12.3 \ 10^5 \ cm^{-3} \ K/k_B$ (consistent with the results of Thompson, Urquhart, & W hite (2004) who nd an internal molecular pressure of $15 \ 10^5 \ cm^{-3} \ K/k_B$), the remainder of our clouds for which we have 20 cm data but no molecular data have been found to have ionised gas pressures higher than this without exception. All of the clouds within our sample are therefore candidates for the possible propagation of photoionisation induced shocks into their interiors, although as this determ ination is based upon a globalm ean of internal pressure no de nitive statements can be made. However, with pressures of only 2 $10^7 \ cm^{-3} \ K/k_B$ the BRCs SFO 1 and SFO 10 are highly likely to

be underpressured with respect to their BLs. Together with SFO 7 and SFO 36, which are the two clouds most likely to be overpressured with respect to their BLs with BL pressures only a factor of 4 times the globalm ean, these clouds represent good candidates for future molecular observations to accurately determ ine their internal pressures.

5.Sum m ary and Conclusions

From archival studies of radio, optical and infrared observations we have analysed 44 BRCs for signs of triggered star formation.We detected 20 cm emission to a level of 3 or greater in 32 of the sample and using a comparison of various wavelength in ages have sorted the clouds into the four distinct types identied by Thompson, Urquhart, & White (2004). i.e. Type 1 BRCs, Type 2 broken-rim med clouds, Type 3 em bedded radio sources and Type 4 unassociated radio sources. We have compared the ionising uses from the candidate ionising star(s) and that determ ined from the 20 am em ission of the rim itself for our Type 1 sources. The num ber of Type 1 sources identied within our sample is 25 and the conditions prevailing within their BLs have been analysed, including the incident ionising uxes, electron density and ionised gas pressures. This analysis has made a signi cant contribution to (alm ost doubled) the num ber of BRCs that have known BL conditions. Through comparison of BL pressures to interior molecular gas pressures we have identied three clouds which are highly likely to be in a state of pressure in balance and thus are ideal candidates for future observations and theoretical studies in RD I. Observations of higher resolution and greater sensitivity are required in order to identify possible further em bedded sources within our sam ple as well as furtherm olecular observations to determ ine internal pressures for a greater sample.

A comparison of the results that we have derived for SFO 5 with those of Le och et al. (1997) reveals a di erence in derived IBL pressure of a factor of . 3, this is believed to be due to the di erences in observational data (frequency, integration area, etc.), we do not feel that these di erences unduly a ect our conclusions as the other errors inherent in our derivation are as large or larger than that arising from this beam dilution.

We draw the following conclusions:

1. A comparison of individual pressures determ ined for BLs from the NVSS data and internal molecular pressure of the BRCs leads us to believe that the majority of our sample is in approximate pressure equilibrium. It is therefore likely that photoionisation induced shocks are propagating into the interior of the clouds. Of the clouds for which we have detected 20 cm radio em ission and for which molecular line data exist the three clouds SFO 4, SFO 13 and SFO 37 have the highest likelihood of being underpressured with respect to their IB Ls.

- 2. In many cases the predictions of ionising ux from our candidate ionising stars are inconsistent with the measurements of 20 cm ux that we have measured. D iscrepancies may be due to the assumption that there is negligible absorption being invalid, the misclassi cation of the spectral type of the ionising star and the possible existence of as-yet undiscovered additional OB stars within the region or an inaccuracy in the distance measurements to the objects.
- 3. The Type 3 radio source in our survey has been identied with a known young embedded cluster containing massive stars.With the 20 cm NVSS data we have con rm ed that this cluster contains early B type stars. Further observations and modelling are required to investigate whether this cluster may have been induced to form by the presence of nearby, external massive stars.

A cknow ledgm ents The authors would like to thank an anonym ous referee for their suggestions which considerably in proved this paper. LKM and JSU are both supported by a PPARC doctoral studentship. This research would not have been possible without the SMBAD astronom ical database service operated at CCDS, Strasbourg, France and the NASA Astrophysics Data System Bibliographic Services. The Digitized Sky Survey was produced at the Space Telescope Science Institute under U.S. Government grant NAG W-2166. The images of these surveys are based on photographic data obtained using the Oschin Schmidt Telescope on Palom ar Mountain and the UK Schmidt Telescope. The plates were processed into the present compressed digital form with the permission of these institutions. This research made use of data products from the Midcourse Space Experim ent. Processing of the data was funded by the Ballistic M issile D efense O rganization with additional support from NASA 0 ce of Space Science. This research has also made use of the NASA / PAC Infrared Science Archive, which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Adm inistration.

References

- Bally, J. & Scoville, N.Z. 1980, ApJ, 239, 121
- Bertoldi, F., 1989, ApJ, 346, 735
- Bica, E., Dutra, C.M., & Barbuy, B. 2003, A&A, 397, 177
- Boulon, J., & Ferhenbach, C. 1958, Journal des Observateurs, 42, 149
- Carpenter, J.M., Snell, R.L., & Schberb, F.P. 1990, ApJ, 362, 147
- Carpenter, J.M., Heyer, M.H., & Snell, R.L. 2000, ApJS, 130, 381
- Cernicharo, J., Bachiller, R., Duvert, G., Gonzalez-Alfonso, E., & Gom ez-Gonzalez, J. 1992, A& A, 261, 589
- Chavamia-K., Lama, E. 1994, A&A, 283, 963
- Claria, J. J. 1974, , A&A, 37,229
- Condon, J.J., Cotton, W D., Greiæn, E.W., Yin, Q.F., Perley, R.A., Taylor, G.B., & Broderick, J.J., 1998, AJ, 115, 1693
- Conti, P.S., & Alschuler, W.R., 1971, ApJ, 170, 325
- Crampton, D. & Fisher, W. A. 1974, Publications of the Dominion Astrophysical Observatory Victoria, 14, 283
- De Vries, C.H., Narayanan, G., & Snell, R.L. 2002, ApJ, 577, 798
- Deharveng, L., Zavagno, A., Cruz-Gonzalez, I., Salas, L., Caplan, J., & Carrasco, L. 1997, A&A, 317, 459
- Duvert, G., Cemicharo, J., Bachiller, R., & Gom ez-Gonzalez, J. 1990, A& A, 233, 190
- Gies, D.R. et al. 1997, ApJ, 475, L49
- Gonzalez-Alfonso, E., Cemicharo, J., & Radford, S. J. E. 1995, A& A, 293, 493
- Hardie, R. H., Seyfert, C. K., & Gulledge, I. S. 1960, ApJ, 132, 361
- Hughes, V.A., Vallee, J.P., & Viner, M.R. 1978, BAAS, 10, 661
- Humphreys, R.M. 1978, ApJS, 38, 309
- Ishida, K. 1970, PASJ, 22, 277
- Jansen, D, J, van Dishoeck, E, F, & Black, J, H, 1994, A&A, 282, 605
- Kassim, N.E. 1988, ApJS, 68, 715
- Kraemer, K. E. et al. 1999, Bulletin of the American Astronom ical Society, 31, 1446
- Kuiper, T.B.H. 1975, A&A, 42, 323
- Le och, B. & Lazare, B. 1994, A&A, 289, 559
- Le och, B. & Lazare, B. 1995, A&A, 301, 522
- Le och, B., Lazare, B., Castets, A., 1997, A&A, 324, 249
- Le och, B., Cemicharo, J., Rodr guez, L. F., Mivile-Deschênes, M. A., Cesarsky, D., & Heras, A. 2002, ApJ, 581, 335
- Leger, A. & Puget, J. L. 1984, A&A, 137, L5
- Matthews, H. I. 1979, A&A, 75, 345
- Megeath, S.T. & Wilson, T.L. 1997, AJ, 114, 1106
- Morgan, W, W., Code, A, D., & Whitford, A, E. 1955, ApJS, 2, 41
- Morgan, LK., Thompson, MA., Urquhart, J.S., White, G.J & Miao, J. In preparation
- O'Dell, C.R.& Doi, T. 2003, AJ, 125, 277
- Ogura, K. & Ishida, K. 1981, PASJ, 33, 149
- Panagia, N. 1973, AJ, 78, 929





F ig.2. Im ages from the survey of clouds with radio detections. Each cloud is represented by a pair of im ages from the D igitised Sky Survey. The left im ages has NVSS 20 cm contours overlaid, while the right im age has MSX 8.3 m em ission contours overlaid. The MSX im ages used have been sm oothed to the same resolution as the NVSS im ages, 45^{00} . The arrow s show the direction of the prim ary ionising sources with the base of the arrow placed at the coordinates of the IRAS source catalogued in Sugitaniet al. (1991) NVSS contours start at 3 and increase in increm ents of 20% of the peak value. MSX contours start at 3 unless otherwise speci ed and increase in increm ents of 20% of the peak





Fig.2. (cont.) SFO 6 M SX contours start at 6, SFO 7 M SX contours start at 6 and SFO 10 M SX contours start at 9 for clarity.









Fig.2. (cont.) SFO 15 M SX contours start at 12 for clarity.





Fig.2. (cont.) SFO 27 M SX contours start at 18 and SFO 31 M SX contours start at 21 for clarity.





Fig.2. (cont.) SFO 32 M SX contours start at 9 and SFO 35 M SX contours start at 9 for clarity









Fig.2. (cont.) SFO 42 M SX contours start at 15 for clarity



Fig.3.SFO 2 and 44 NVSS contours overlaid on DSS im ages

- Price, S. D., Egan, M. P., Carey, S. J., Mizuno, D. R., & Kuchar, T. A. 2001, AJ, 121, 2819
- Racine, R. 1968, AJ, 73, 588
- Repurth, B. 1983, A&A, 117, 183
- Reynolds, R.J. & Ogden, P.M. 1979, ApJ, 229, 942
- Savage, B. D., D rake, J. F., Budich, W., & Bohlin, R.C. 1977, ApJ, 216, 291
- Sknustkie, M .F., M eyer, M .R., Canasco, L, Cnuz-Gonzalez, L, Salas, L., Johnson, P., Spillar, E., 1990, BAAS, 25, 1374
- Snell, R.L., Huang, Y.-L., Dickman, R.L., & Claussen, M.J. 1988, ApJ, 325, 853
- Sugitani, K., Fukui, Y., & Ogura, K. 1991, ApJS, 77, 59
- Sugitani, K., & Ogura, K. 1994, ApJS, 92, 163
- Sugitani, K., M atsuo, H., Nakano, M., Tamura, M., & Ogura, K. 2000, AJ, 119, 323
- Thom pson, M. A., White, G. J., Morgan, L. K., Miao, J., Fridlund, C. V. M., & Huldtgren-White, M. 2004, A&A, 414, 1017
- Thom pson, M. A., Urquhart, J. S., & White, G. J. 2004, A&A, 415, 627
- White, G.J., Le och, B., Fridlund, C.V.M., Aspin, C.A., Dahmen, G., Minchin, N.R., & Huldtgren, M. 1997, A&A, 323, 931
- W hite, G.J. et al. 1999, A&A, 342, 233
- Wood, D.O.S. & Churchwell, E. 1989, ApJ, 340, 265

Table 6. Ionised boundary layer pressures and internal pressures and densities of clouds observed in molecular line wavelengths. A dash indicates non-detection in the NVSS.

SFO Object	IBL pressure		Internal P ressure	Internal M olecular D ensity
	(10 5 cm 3 K /k $_{ m B}$)		(10 5 cm 3 K /k $_{ m B}$)	(10^3 cm^{-3})
4ª	197.5	>>	8.2	50.0
5 ^b	114.4		32.3	11.0
11 ^c	82.0		25.0	2.5
13^{d}	63.9	>>	16.3	2.0
16^{d}	-		0.3	2.0
18^{d}	-		7.8	16.6
20 ^e	-		6.9	20.0
25 ^d	66.4		12.5	7.4
37 ^f	112.3	>>	1.5	3.0

^aJansen et al. (1994);^bLe och et al. (1997); ^cThom pson et al. (2004); ^dD e V ries, et al. (2002); ^eC emicharo et al. (1992); ^fD uvert, et al. (1990)