

# STAR FORMATION AT REDSHIFT ONE: PRELIMINARY RESULTS FROM AN H SURVEY

Michelle Doherty<sup>1</sup>, Andrew Bunker<sup>1;2</sup>, Robert Sharp<sup>1;3</sup>, Gavin Dalton<sup>4</sup>, Ian Parry<sup>1</sup>, Ian Lewis<sup>4</sup>, Emily Macdonald<sup>4</sup>, Christian Wolf<sup>4</sup>

<sup>1</sup>Institute of Astronomy, Madingley Road, Cambridge, CB3 0HA, UK (e-mail:md@ast.cam.ac.uk)

<sup>2</sup>School of Physics, University of Exeter, Stocker Road, Exeter, UK

<sup>3</sup>Anglo-Australian Observatory, Sydney, Australia

<sup>4</sup>Astrophysics, NAPL, Keble Road, Oxford, OX1 3RH, UK

We present the first successful demonstration of multi-object near-infrared spectroscopy on high redshift galaxies. Our objective is to address the true star formation history of the universe at  $z \sim 1$ , a crucial epoch which some have suggested is the peak of star formation activity. By using H $\alpha$  – the same robust star formation indicator used at low- $z$  – redshifted into the J- and H-bands, we can trace star formation without the systematic uncertainties of different calibrators, or the extreme dust extinction in the rest-UV, which have plagued previous efforts. We are using the instrument CIRPASS (the Cambridge Infra-Red PA noramic Survey Spectrograph), in multi-object mode, which has been successfully demonstrated on the Anglo-Australian Telescope (AAT) and the William Herschel Telescope (WHT). CIRPASS has 150 fibres deployable over  $40^\circ$  on the AAT and  $15^\circ$  on the WHT. Here we present preliminary results from one of our fields observed with the WHT: H $\alpha$  detections of  $z \sim 1$  galaxies in the Hubble Deep Field North.

## 1 Introduction

### 1.1 Background

An important issue in studies of galaxy formation and evolution is determining the peak epoch of star formation in the universe. There is clear evidence that the star formation rate (SFR) was higher in the recent past – it is known to rise steeply to  $z \sim 1$  and perhaps to peak around redshifts  $1 \sim 2$ . (e.g. Lilly et al. 1996, Tresse et al. 2002). However there are large discrepancies between SFR estimates obtained from different methods: different indicators have uncertain relative calibration and are also differently affected by dust extinction. There is a factor of  $\sim 10$  uncertainty in the star formation history of the universe at  $z \sim 1$  (e.g. Hopkins et al. 2001), a discrepancy which needs to be addressed by using a uniform tracer of star formation across all redshift bins. The H $\alpha$  emission line is a reliable tracer of the instantaneous SFR as the H luminosity of a galaxy is directly proportional to the ionising flux from its massive stars. It is also relatively immune to metallicity effects and less susceptible to extinction than the rest-UV. However, tracing H $\alpha$  to early epochs forces a move to the NIR at  $z > 0.6$ . Previous H surveys in the NIR have so far been restricted to small samples, as they have relied on longslit spectroscopy which is inefficient for surveys, in terms of telescope time (e.g. Glazebrook et al. 1999, Tresse et al. 2002, Erb et al. 2003). A large sample (several hundred) of galaxies is needed to address this matter properly.

## 1.2 An H Survey with the CIRPASS multi-object spectrograph

We are using the fibre-fed CIRPASS spectrograph (Parry et al. 2000), to undertake an H survey of galaxies at  $z = 0.7 - 1.5$  (Doherty et al. 2004), in order to address the true star formation history of the universe at this epoch. We aim to target several hundred galaxies. We have carried out a successful pilot study for this project, performing multi-object spectroscopy of galaxies in the vicinity of the HDF-N. The instrument operates in the near-infrared, between 0.9 and 1.67  $\mu\text{m}$  and was used in multi-object mode, with 150 fibres of 1.1<sup>0</sup> diameter on the sky (comparable to the expected seeing convolved with typical galaxy profiles), deployable over 15<sup>0</sup> field of view. The plate was centroided to an accuracy better than 0.5<sup>0</sup> using six guidebundles centred on bright stars. A Hawaii 2K detector was used, and a grating of 831 lines/mm, producing a dispersion of 0.95Å/pixel. The full width at half maximum (FWHM) of each fibre extends over 2.7 pixels in both the spatial and spectral domain. The wavelength coverage was 1726Å, covering most of the J-band, and the grating was tilted to set a central wavelength of  $\lambda_c = 1.25 \mu\text{m}$ . The resolving power was  $R = \lambda / \Delta\lambda_{\text{FWHM}} = 5000$ . At this resolution the background is very dark between the OH sky lines in the J- and H-bands, allowing for high sensitivity to emission lines falling between the skylines (which contaminate only about 10% of the array). We used half of the fibres on target objects and half on sky, to allow for good sky subtraction. Thus with the capability of observing 75 objects simultaneously, CIRPASS offers a huge multiplex advantage, ideal for carrying out such a survey.

## 2 Initial Results: Bright H galaxies in the HDF-N

We detected H in a sub-sample of four targets in the HDF-N at  $> 5\sigma$  in just 3 hours of observing. Examples are shown in Figure 1. In an average fibre, between skylines, for a spectrally unresolved line (velocity FWHM  $< 60 \text{ km s}^{-1}$ ) we achieve a sensitivity of  $7.2 \times 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$  at  $5\sigma$  (in 3 hours, equivalent to a star forming rate of  $5 h_{70}^2 M_{\odot} \text{ yr}^{-1}$ ). The emission lines we detect are typically broader than this, with velocity full width at half maximum of  $100 - 250 \text{ km s}^{-1}$ , equivalent to  $\Delta v_{1D} = 40 - 100 \text{ km s}^{-1}$ . Some of this velocity width may be due to galactic rotation in the extended galaxies: the most spatially extended galaxies exhibit the broadest lines. H traces the instantaneous star formation rate as the H luminosity of a galaxy is directly proportional to the ionising flux from massive stars, which drops off only about 20 million years after star formation ceases. The UV luminosity, on the other hand, evolves in time with changes in stellar population, continuing to rise even after star formation has ended. We compared SFRs derived using the two different methods, for each of our  $> 5\sigma$  detections, using conversions given in Kennicutt (1998). We performed photometry on the HST/ACS images from GOODS v1.0 (Giallisco et al., 2003), with 1<sup>0</sup> apertures, for consistency with our infrared fibre size and used the B band magnitudes (4500Å) to calculate rest frame UV (2400Å) flux densities and corresponding star formation rates. We find that SFRs calculated from the UV luminosity densities are typically a factor of at least two lower than those derived from H. This is probably due to the differential effect of dust extinction in the redshift one galaxies between  $\lambda_{\text{rest}} = 2400\text{Å}$  and 6563Å. This result is consistent with values in Glazebrook et al. (1999), Tresse et al. (2002), and Yan et al. (1999), who all find SFR(H)/SFR(UV) ratios of around 2-3.

## 3 Summary

We have performed multi-object, near-infrared spectroscopy of  $z \sim 1$  galaxies in the Hubble Deep Field North, using CIRPASS-MOS. These observations form an important pilot study for an on-going survey to trace star formation rates at redshift one. The brightest H galaxies (about 10% of our sample) were detected at at least  $5\sigma$  in only 3 hours exposure time, demonstrating




Figure 1: Example spectra for two objects where H<sub>α</sub> was detected at more than 5 $\sigma$ . The night sky spectrum is overlaid in dotted lines for convenient identification of sky residuals which did not fully subtract out. The expected position of H<sub>α</sub> at the optical spectroscopic redshift (Cohen et al. 2000) is marked with a dashed line.

the success of this technique. We have shown that we can detect H $\alpha$  at sufficient signal to noise to obtain a good estimate of star formation rates in these galaxies. By pre-selecting galaxies with H $\alpha$  redshifted between the OH sky lines, we can detect star formation rates of  $5 h_{70}^2 M_{\odot} \text{ yr}^{-1}$  (5 in 3-hours). The star formation rates obtained are higher than those estimated from UV continuum by a factor of about two, due to dust obscuration in the UV. This is consistent with previous work in this area. The results of this pilot study are discussed further in Doherty et al (2004, submitted to MNRAS). The full survey will be presented in Doherty et al. (2004b, in prep.) We have another 60 sources in this field, at known redshifts, therefore by stacking the spectra we will be able to obtain a global star formation rate for  $z \sim 1$  galaxies in the vicinity of the HDF-N. We aim to build up a sample of several hundred galaxies in total, over several different fields, which will allow us to determine the H $\alpha$  luminosity function at  $z \sim 1$  and hence address the true star formation rate at this important epoch.

#### Acknowledgements

CIRPASS was built by the instrumentation group of the Institute of Astronomy in Cambridge, UK. We thank the Raymond and Beverly Sackler Foundation and PPARC for funding this project. We thank the WHT staff, in particular Danny Lennon, Kevin Dee, Rene Ruten, Juerg Rey and Carlos Martin, for their help in enabling CIRPASS to be used as a visitor instrument. MD is grateful for support from the Fellowship Fund Branch of AFUW Qld Inc., the Isaac Newton Studentship, the Cambridge Commonwealth Trust and the University of Sydney. ECM acknowledges the C.K. Marr Educational Trust.

#### References

1. J.G. Cohen et al, ApJ 538, 29 (2000).
2. M. Doherty et al, MNRAS, submitted, (2004).
3. M. Doherty et al, in preparation, (2004b).
4. D.K. Erb et al, ApJ 591, 101 (2003).
5. M. Giavalisco and GOODS team, AAS meeting, 202, 2003.
6. K. Glazebrook et al, MNRAS 306, 843 (1999).
7. A.J. Hopkins, AJ 122, 288 (2001).
8. R.C. Kennicutt, ARA&A 36, 189 (1998).
9. S.J. Lilly et al, ApJ 460, L1 (1996).
10. I.R. Parry et al, in Proc. SPIE Vol. 4008, p. 1193-1202, ed. Masanori Iye; A.F. Moorwood (2000)
11. L. Tresse et al, MNRAS 337, 369 (2002).
12. L. Yan et al, ApJ 519, L47 (1999).