# The young pulsar PSR B0540-69.3 and its synchrotron nebula in the optical and X-rays?

N.I. Sera movich<sup>1,2</sup>, Yu.A. Shibanov<sup>1</sup>, P.Lundqvist<sup>2</sup>, and J. Sollerm an<sup>2</sup>

<sup>1</sup> Io e Physical Technical Institute, Politekhnicheskaya 26, St. Petersburg, 194021, Russia

<sup>2</sup> Stockholm Observatory, A lbaN ova Science Center, Department of Astronomy, SE-106 91 Stockholm, Sweden

Received 22 M arch 2004; accepted 22 June 2004

Abstract. The young PSR B0540-69.3 in the LMC is the only pulsar (except the C rab pulsar) for which a near-UV spectrum has been obtained. However, the absolute ux and spectral index of the HST/FOS spectrum are signi cantly higher than suggested by previous broad-band time-resolved groundbased UBVRI photom etry. To investigate this di erence, observations with ESO /VLT/FORS1 and analysis of HST/WFPC2 archival data were done.W e show that the HST and VLT spectral data for the pulsar have > 50% nebular contam ination and that this is the reason for the above mentioned di erence. The broadband HST spectrum for the range 3300{8000 A =  $1.07^{+0.20}_{0.19}$ . This is dierent from the is clearly nontherm aland has a negative spectral index, F / with alm ost at spectrum of the C rab pulsar, and also steeper than for the previously published broadband photom etry of PSR B0540-69.3. We have also studied the spatial variations of the brightness and spectral index of the Pulsar W ind Nebula (PW N) around the pulsar, and nd no signi cant spectral index variation over the PW N. The HST data show a clear asym m etry of the surface brightness distribution along the major axis of the torus-like structure of the PW N with respect to the pulsar position, also seen in Chandra/HRC X -ray in ages. This is di erent from the Crab PW N and likely linked to the asymmetry of the surrounding SN ejecta. The HST/W FPC2 archival data have an epoch separation by 4 years, and this allows us to estim ate the proper motion of the pulsar. We nd a motion of 4.9 2.3 mas yr<sup>1</sup> (corresponding to a transverse velocity of 1190 560 km s<sup>1</sup>) along the southern iet of the PW N. If this is con rm ed at a higher signi cance level by future observations, this makes PSR B 0540-69.3 the third pulsar with a proper motion aligned with the jet axis of its PWN, which poses constraints on pulsar kick models. To establish the multiwavelength spectrum of the pulsar and its PWN, we have included recent Chandra X -ray data, and discuss the soft pulsar X -ray spectrum based on spectral ts including absorption by interstellar gas in the Milky Way, LMC as well as the supernova ejecta. We have compared the multiwavelength spectra of PSR B0540-69.3 and the C rab pulsar, and nd that both PSR B0540-69.3 and the C rab pulsar have a weaker ux in the optical than suggested by a low -energy power-law extension of the X -ray spectrum. This optical depression is more severe for PSR B0540-69.3 than for the Crab pulsar. The same trend is seen for the PW Ne of the two pulsars, and continues for low energies also out in the radio band. We discuss possible interpretations of this behavior.

Key words.pulsars: individual: PSR B0540-69.3 { ISM : supernova rem nants { supernovae: general { A strom etry

#### 1. Introduction

PSR B0540-69.3 in the Large Magellanic C bud (LMC) was discovered as a pulsed (P = 50.2 ms) Xray source by Seward et al. (1984). Pulsations have since also been detected in the optical and at radio wavelengths (M iddleditch & Pennypacker 1985; M anchester et al. 1993a). The pulse pro le in the optical (B oyd et al. 1995) is broad and double-peaked, with a sep-

aration of 0.2 in phase between the twom axim a, consistent with what is also seen in X-rays (Seward et al. 1984; de P has et al. 2003). The pro le is also broad in the radio (the duty cycle is > 80%), and there is a hint of a double structure (M anchester et al. 1993a).

Parameters of PSR B0540-69.3 are compiled in Table 1. The pulsar spins rapidly, is young (spin down age 1660 yr), and sits in a compact synchrotron nebula (see Fig. 1), which we will henceforth refer to as its pulsar wind nebula (PW N). The similarities with the C rab pulsar and its nebula are such that PSR B0540-69.3 with its supernova remnant, SNR 0540-69.3, are sometimes referred to as the  $\$  rab twin". Even the structures of the

Send o print requests to:N.Sera movich; e-mail: natalia@astro.su.se

<sup>&</sup>lt;sup>?</sup> Based on observations performed at the European Southern Observatory, Paranal, Chile (ESO Program 67D-0519).

Table 1. Parameters of PSR B0540-69	.3 (M anchester et al. 1993a,	unless speci ed otherwise).
-------------------------------------	-------------------------------	-----------------------------

	0 bærved								D	erived			
Р	P		a	l <b>,</b> b <sup>b</sup>	DM	с	d	В	е	E	_	d	f
m s	10 15	5	m as yr $^{1}$		am <sup>3</sup> pc		yr	G		erg s <sup>1</sup>		kpc	
50.2	479 <b>:</b> 04	0:10	5	279:7 31:5	146	4	1660	4 <b>:</b> 96	10 <sup>12</sup>	1 <b>:</b> 495	10 <sup>38</sup>	51	1:3

<sup>a</sup> Indirect proper motion estimates (Manchester et al. 1993b) <sup>d</sup> Spin-down age P = 2P-

<sup>b</sup> G alactic coordinates (K aaret et al. 2001)

° D ispersion m easure

<sup>e</sup> Magnetic eld for a 10 km radius NS 3:2  $10^{19} (P P)^{1=2}$ 

 $^{
m f}$  D istance to the LM C (Panagia 2004)

PW Ne appear to be sim ilar. Both have a torus and jets (G otthelf & W ang 2000), although the proper motion for PSR B0540-69.3, suggested by M anchester et al. (1993b) based on a displacement between the pulsar optical position and the center of the PW N as seen in radio, seem s not to be along the spin axis as it is in the C rab case.

There are, however, di erences on a larger scale. While the PWN of PSR B0540-69.3 is surrounded by an X-ray and radio emitting outer shell of 30<sup>00</sup>, or 7.3 pc (Manchester et al. 1993b; radius Gotthelf & W ang 2000), an outer shell around the C rab is still not con med (although high-velocity gas has been revealed in the UV, Sollerm an et al. 2000). A nother di erence is that SNR 0540-693 is oxygen-rich (e.g., Kirshner et al. 1989; Sera movich et al. 2004), whereas the Crab Nebula has nearly norm al solar abundances of m etals (B lair et al. 1992 and references therein). It is therefore believed that the progenitor to PSR B0540-69.3 was a much more massive star than the Crab progenitor (Kirshner et al. 1989).

PSR B0540-69.3 is one of few pulsars for which a near-UV or optical spectrum has been reported. Hill et al. (1997) obtained a time-integrated near-UV spectrum with HST/FOS and Middleditch & Pennypacker (1985) used tim e-resolved photom etry to establish a broadband ground-based UBVRI spectrum in the optical. These two spectra show, however, a signi cant di erence in absolute ux in the spectral range where they overlap. To investigate this mism atch we have added two recent sets of data, one is the ESO /VLT/FORS spectroscopy of SNR 0540-69.3 analyzed by Sera movich et al. (2004), and the other is HST /W FPC2 in aging (Caraveo et al. 2000; Morse 2003) retrieved from the HST archive. A bonus of our study is that we also obtain the optical spectrum of the PW N around PSR B0540-69.3. This was rst studied quantitatively by Chanan et al. (1984), albeit at a low spatial resolution which did not allow them to resolve the pulsar from the PWN.

To connect the optical pulsar emission to the emission at other wavelengths, we have also included recent results from radio and X-rays. Previous attempts to establish the multiwavelength spectrum of PSR B 0540-69.3 have assumed a rather high hydrogen column density for the X-ray absorption, N<sub>H</sub> 4:6  $10^{21}$  cm<sup>2</sup> (K aaret et al. 2001). W ith this value for N<sub>H</sub> it is possible to t the soft

X -ray spectrum with a single power-law. This suggests a non-therm al nature of the emission, likely to be form ed in the magnetosphere of the rotating neutron star (NS). There are, how ever, reasons to reinvestigate this since the spectral ts have not considered the fact that a large fraction of the absorbing gas has LMC abundances rather than M ilky W ay abundances. It could even be that the supernova ejecta can contribute to the absorption of the X -ray emission. Taking these considerations into account, we show that the situation is more complicated than assuming a single power-law for the optical/X -ray spectrum. W e have also done the same exercise for the PW N.

The outline of the paper is as follows: in Sect. 2 we describe the optical spectroscopic and photom etric observations of PSR B0540-69.3 and its PW N, as well as a reinvestigation of the X-ray data of K aaret et al. (2001). In Sect. 3 we discuss these results and put them in a multiwavelength context. We also discuss results we nd for the proper motion of the pulsar, and their possible interpretation.

#### 2.0 bservations, data analysis and results

# 2.1.VLT observations and data reduction

Spectroscopic observations of PSR B 0540-69.3 were performed on 2002 January 9 with the FO cal Reducer/low dispersion Spectrograph (FORS1) on the 8.2m UT3 (MELIPAL) of the ESO/VLT, using a slit width of  $1^{00}$  and

Table 2.Log of VLT observations of PSR B 0540-69.3 on 2002 January 9.

No.	Time	Exposure	A im ass	Seeing <sup>a</sup>
	UΤ	S		arcsec
1	01:59:32.0	1320	1.43	1.25
2	02:23:25.0	1320	1.42	1.30
3	02:52:38.0	1320	1.41	1.18
4	03:16:31.0	1320	1.41	1.12
5	03:42:32.0	1320	1.41	1,24
6	04:06:25.0	1320	1.43	1.09
7	04:32:42.0	1320	1.45	0.88
LTT 3864	08:51:03.0	20	1.03	0.67

<sup>a</sup> Fullwidth at halfmaximum of the stellar pro le.

the grism GRIS\_ $600B^1$ . This grism has a dispersion of 50 A=mm, or 1.18 A=pixel, and a wavelength range of 3605{6060 A. The optical path also includes a Linear A tm ospheric D ispersion Corrector that compensates for the e ects of atm ospheric dispersion (Avila et al. 1997). The pixel scale of the detector is  $0^{\circ}$ 2 per pixel. We obtained 7 exposures of 1320 seach (see Table 2), in total 154 m inutes of exposure time. The position angle, PA = 88, was the same in all these exposures. The slit crosses the pulsar and its PW N as shown in Fig. 1. The mean seeing was  $1^{\circ}$ 15.



Fig.1.A  $15^{00}$   $15^{00}$  in age of the eld around PSR B0540-69.3 obtained in the F547M band with the HST/W FPC2 (M orse 2003). The pulsar was exposed on the PC chip and its position is marked by an arrow. The slit position of the VLT observations is marked by thin parallel lines. The slit width is  $1^{00}$ . The di use emission surrounding the pulsar is the pulsar wind nebula (PW N). Note its elongation in the NE-SW direction.

The spectroscopic in ages were bias-subtracted and at- elded using standard procedures within the NOAO IRAF Longslit package. We used the averaged sigm a clipping algorithm avsigclip with the scale parameter set equal to none to combine the images. Wavelength calibration of the combined images was done using arc frames obtained with a helium argon lamp. The spectra of the objects were then extracted from the 2D image using the apalland background tasks. Flux calibration of the spectra was accomplished by comparison with the spectrophotometric standard star LTT 3864 (Hamuy et al. 1994) observed on the same night (see Table 2). A tm ospheric extinction corrections were perform ed using a spectroscopic



F ig.2. Spatial proles of PSR B0540-69.3 and its PWN along the slit in Fig.1 (in counts, VLT data) at the continuum wavelengths 4600.0 A, 5248.7 A and 5450.0 A, from top to bottom, respectively. In each panel the pulsar is assumed to sit at the main peak of the prole. This lines mark the six-pixel wide extraction window chosen for the spectral analysis of the pulsar (where 1 pixel corresponds to  $0^{\circ}$ 2). The regions outside the this lines, but within the solid lines have also a width of six pixels, and mark the regions used for the nebular subtraction in the spectral analysis. The spatial prole of a background star is shown in the inset in the bottom panel for comparison.

extinction table provided by  $ESO^2$ . A signi cant ux (response) degradation was found at wavelengths below 3860 A where the standard star spectrum displays many

<sup>&</sup>lt;sup>1</sup> http://www.eso.org/instruments/fors1/grisms.html

<sup>&</sup>lt;sup>2</sup> http://www.eso.org/observing/dfo/quality/UVES/ les/ /atm oexan.t ts



Fig. 3. Spatial pro les as in Fig. 2, but using the HST/W FPC2/F547M in age shown in Fig. 1. The pivot wavelength is 5483 A.W e show pro les obtained along the VLT slit with the spatial resolution of 0<sup>o</sup>,046 provided by the PC chip pixel-size (top), rebinned to the VLT CCD-pixelscale of 0<sup>o</sup>,2 (m iddle), and sm oothed using a G aussian with the FW HM of 1<sup>o</sup>,2, com parable to the seeing value of the VLT observations, and then rebinned to the VLT CCD-pixel scale (bottom). The VLT six pixel extraction window is marked by vertical lines.

deep spectral features, as well as in the very red end at > 6030 A.W e therefore excluded these wavelength regions from our analysis.

#### 2.2. Spectroscopy of the pulsar

As seen from Fig.1 the pulsar is in the center of a bright compact nebula, which contam inates the ux from the pulsar. This contam ination is particularly strong in nebular lines from the supernova remnant, but is also signi cant at continuum wavelengths where the spectrum of the extended object is expected to be dom inated by synchrotron em ission from the PW N.

In Fig. 2 we have plotted spatial proles for the em ission from the pulsar+ nebula along the slit at several wavelengths where the contribution of nebular lines is negligible. D espite som em inor variation of the shape of the pro-

le with wavelength, the pro le has a strong peak around the pulsar position. However, with a seeing of 1<sup>00</sup> the pulsar and nebular em issions are strongly blended. This is partially con med by a comparison of the pro lewith the PSF of a background star shown in the inset of the bottom panel of Fig. 2. To illustrate this further we have com pared the VLT pro lesw ith the pro les obtained from the HST/W FPC2/F547M image, shown in Fig. 1. From the F 547M in age we rst extracted the data covered by the VLT slit, and then we averaged the emission across the slit for each position along the slit. The peak position of the ux along the slit varies with the wavelength of the spectroscopic image by < 1 VLT pixel. To take this into account in our test we extended the slit width by one pixel. In the st test case we kept the high spatial resolution in the HST in age  $(0^{\circ}, 046, Fig. 3 \text{ top})$ , while in the second case we averaged over a coarser pixel scale  $(0^{0}.2)$  to simulate the VLT pixel size (Fig. 3 m iddle). In a third experiment we sm oothed the initial image using a Gaussian with a FW HM of  $1^{\circ}_{:2}$  to model the VLT seeing conditions and we then rebinned it to the VLT CCD-pixel scale (Fig. 3 bottom).

The pivot wavelength of the F547M lter is 5483A, which is close to 5450 A, chosen for the pro le shown at the bottom of Fig. 2. As seen from Fig. 3, the pulsar is clearly resolved from the extended PWN at the PC chip spatial resolution as a narrow central peak on a broad  $7^{00}$  in size) asymmetric pedestal formed by the PW N. ( It is still resolved at the VLT CCD -pixel scale of  $0.2^{\circ}$ , while it is not resolved after the 1.2 sm oothing. i.e., close to the seeing conditions of the VLT observations. A though the pulsar should contribute signi cantly to the ux within the spatial strip of six VLT pixels centered on the main peak of the whole pro le, it is obvious that the nebula will contam inate severely the spectral VLT observations.

W ith this in m ind, we extracted a 1D spectrum averaged over 6 pixels, equal to  $1^{\circ}_{\cdot \cdot 2}$ , along the slit centered at the pulsar position, as shown by thin vertical lines in Figs. 2 and 3. To subtract the nebular contribution we extracted 1D spectra averaged over six adjacent pixels to the east and an equal number to the west of the central strip, as indicated by thick lines in Fig. 2. An averaged spectrum was constructed from the two adjacent spectra and subtracted from the spectrum for the central region. The resulting spectrum contains no signi cant em ission from

1,6

1,4

1,2

nebular lines, except for [0 III] 4959, 5007, which has a high spatial variability within the nebula. The resulting spectrum, with the [0 III]-feature rem oved, is presented in Fig. 4. As can be seen from this gure, our spectroscopic data agree well with the result by Hillet al. (1997) at the lower boundary of their 1 uncertainty range, but give about 2 4 times higher ux than the photom etric data of M iddleditch et al. (1987). This can partly be understood from our method of correcting for the PW N em ission.We averaged this by using the em ission  $1^{\circ}_{\cdot}2^{\circ}_{\cdot}4$  away from the center of the spatial pro le.0 ne problem is that Fig.2 shows that the center of the pro le in the VLT data may not coincide exactly with the pulsar position. A second, and m ore serious problem, is that the seeing spreads out much of the weak pulsar em ission from the central region whereas seeing makes the PW N emission peak toward the center regardless of whether there is a pulsar or not. This willm ost likely lead to erroneous background subtraction so that the VLT spectrum shown in Fig. 4 is contam inated with signi cant PW N em ission. The agreem ent with Hill et al. (1997) indicates that also their analysis overestim ated the pulsar em ission.

# 2.3.HST observations

The pulsar eld has been in aged with the HST/W FPC2 several times in various bands mainly to study SNR 0540-69.3. For additional analysis of the optical emission from PSR B0540-69.3 and its PW N we retrieved some of these im ages from the HST archive.

The data using the wide and medium band lters  $F336W^3$ , F547M, and F791W, obtained on 1999 O ctober 17 with 600 s, 800 s, and 400 s total exposure times, respectively (M orse 2003), are particularly useful for the continuum emission analysis since these lters do not cover any bright emission lines from the LM C or the supernova rem nant. W e also retrieved data sets for the narrow band F 658N and wide band F 555W lters, both of which were obtained on 1995 O ctober 19 with 4000 s, and 600 s exposures, respectively (C araveo et al. 2000).

The pulsar and its PW N are clearly detected on the PC chip in all these images. This is illustrated in Figs. 1 and 3 which present the data for the F547M band. The F658N liter includes the 6576 6604 A range, which m eans that contam ination from high-velocity H em ission from the supernova remnant at central wavelength

6578 A (Sera movich et al. 2004) can enter into the lter passband, as well as [N II] em ission from the LM C. This can also be seen in Fig. 3b of C araveo et al. (2000) where it is shown that a lam ent passes across the pulsar along the NW direction. A lthough the pulsar stands out rather clearly on the im age, the uneven background introduces som e uncertainty to the estim ated pulsar ux. The background contam ination is more severe in the F555W band as it captures the bright [O III] 4959, 5007 A lines



which are much more di cult to spatially disentangle from the pulsar. The uneven background is most clearly seen in the in age obtained with the F 502N narrow band liter centered at these lines and overlapping with the F 555W band. Therefore, the pulsar and PW N continuum ux m easurements in the F 555W band can only be considered as upper limits.

#### 2.4. Photom etry of the pulsar

For the photom etry of the pulsar in the F336W, F547M, F555W, F658N, and F791W bands we used the Daophot and Photcal RAF packages. To measure the pulsar ux we used circular apertures whose radii were determ ined from maxim izing the signal-to-noise ratio S/N of the detected pulsar, and hence minim izing the magnitude error m

<sup>&</sup>lt;sup>3</sup> http://www.stsci.edu/instruments/wfpc2/

<sup>/</sup>W fpc2\_hand\_current/ch3\_optical lters2.htm 1# 474439

in each lter. For a point-like source, m and S/N are Table 3. Broad-band uxes from PSR B0540-69.3. de ned in the standard way as

$$m = \frac{2.5}{\ln 10} \frac{S}{N}^{-1}; \qquad (1)$$

$$\frac{S}{N} = \frac{N_{c}}{G} + A_{N}^{2} (1 + \frac{A}{A_{sky}})^{1=2} N_{c}; \qquad (2)$$

where N<sub>c</sub> is the ux in counts for a given aperture,  $\frac{2}{N}$ is the standard deviation in counts per pixel, A is the num ber of pixels in the aperture, G is the gain, and A sky is the num ber of pixels in the annulus used for background m easurem ents. The optim al aperture radii were found to be 2.5, 3.0, 4.0, 2.5, and 3.5 pixels, providing S/N of 8, 10,25,7.5, and 8 in the F336W , F547M , F555W , F658N, and F791W bands, respectively. A perture corrections for the magnitudes were derived with the mkap le package using several relatively bright nearby eld stars in the PC im ages.

Pipeline-provided zeropoints PHOTFLAM ( ux densities in wavelength units) and pivot wavelengths PHOTPLAM (in A) taken from the ts header of each in age were used for the ux calibration (see handbook for W FPC2<sup>4</sup>). Fluxes in units of Jy at the pivot frequencies =  $10^8 \text{c/PHOTPLAM}$  (in Hz) were derived from the F<sub>0</sub> (= 0) aperture corrected source counts using the expression

$$F = 10^{21} \frac{N_c}{c t_{exp}} P H O T F LAM P H O T P LAM^2$$
(3)

where c is the speed of light in cgs units, and  $t_{exp}$  is the exposure time in seconds. W e also calculated dereddened uxes by multiplying F with the extinction correction factors  $k_{ext} = 10^{0.4k_i}$ , where  $k_i$  is the interstellar extinction m agnitude for the \i-th<sup>00</sup> passband determ ined using the extinction color excess toward SNR 0540-69.3. In order to compare with Middleditch et al. (1987) we have used the same values for E (B V) and R, i.e., 0.20 and 3.1, respectively, for both data sets, but a m ore recent extinction curve (Cardelli et al. 1989). W e will discuss the extinction in greater detail in Sect. 2.8.

The results of the pulsar photom etry are presented in Table 3 and shown in Fig. 4. The measured uxes are about a factor of 2 4 low er than the spectroscopic results in Sect. 2.2, except for the F555W lter which includes contamination from [O III]. The uxes in this lter are presented as upper lim its (see above). On the other hand, the HST photom etry is compatible with the results obtained by M iddleditch et al. (1987), but the uncertainties are several tim es sm aller. Considered separately, our spectral and photom etric m easurem ents of the pulsar ux are in good agreem ent with previous results. How ever, they do not erase the signi cant discrepancy between these results (see Fig. 4), which is much larger than the uncertainty of our photom etric m easurem ents. The only plausible explanation to the discrepancy is that both the VLT and HST

Band	P ivot	M easured	D ereddened		
	frequency	ux	ux		
	log [Hz]	log F [Jy]	log F [Jy]		
F336W	14.950 (13)	0.243 (51)	0.648 (51)		
F 547M	14.737 (8)	0.625 (42)	0.873 (42)		
F 555W	14.741 (21)	0.69	0.94		
F 658N	14.658(1)	0.793 (54)	0.995 (54)		
F791W	14.581 (15)	0.874 (50)	1.027 (50)		

<sup>a</sup> Hereafter num bers in the parentheses are uncertainties of the respective values. For exam ple, 14.950 (13) m eans 14:950 0:013.

Table 4. Broad-band uxes from a 10 pixel aperture around PSR B0540-69.3.

Band	P ivot		M eas	sured	D ereddened		
	frequ	frequence		ıx	ux		
	log	[H z]	log F	[Jy]	log F	[Jy]	
F336W	14.95	14.950 (13)		2 (60)	0.967 (60)		
F 547M	14.73	14.737 (8)		1.013 (20)		2 (20)	
F 555W	14.74	14.741 (21)		1.141		.392	
F791W	14.58	31 (15)	1,271	L(18)	1.424	4(18)	

Table 5.Parameters of the power law spectral ts (F =) of the pulsar data shown in Fig. 4.

0 bservations		Log F <sub>o</sub> ,
		$_0 = 5:47  10^{14} \text{ Hz}$
		[Jy]
VLT spectrum	1.88 0.01	1.469(2)
(this work)		
HST/FOS spectrum <sup>a</sup>	1.6 0.4	1,228 (115)
(Hillet al. 1997)		
HST photom etry	$1.07^{+0:20}_{0:19}$	0.879 (25)
(this work)		
10 pixelm easurem ents	$1.24^{+0.19}_{0.18}$	1,241 (22)
(this work)		
T in e-resolved photom etry	$0.26^{+0.45}_{0.43}$	0.880 (54)
(Middleditch et al. 1987)		

<sup>a</sup> Log F o from Hillet al. (1997).

spectroscopy are strongly contam inated by the PW N, as already suggested in Sect. 2.2. As an additional test we m easured the pulsar ux in the F336W, F547M, F555W, and F791W bands, using a circular aperture with a radius of 10 PC-pixels (corresponding to a total diam eter of 0.92) centered on the pulsar, without subtraction of the background. These conditions should approximately reproduce the param eters of the spectral m easurem ents within a circular aperture of alm ost the same diam eter  $(0^{\circ}.86)$  m ade with the HST /FOS by Hillet al. (1997). The measured uxes are presented in Table 4 and shown by triangles in Fig. 4. They are much closer to the HST and VLT spectral uxes. The 20% {70% discrepancy between these HST/W FPC2 and FOS uxes must be considered as sm all com pared with the large uncertainties of the FOS ux. Using a W FPC2 im age Hill et al. (1997) estim ated

www.stsciedu/instruments/wfpc2/

<sup>/</sup>W fpc2\_hand\_current/ch8\_calibration9.htm 1# 464103

the PW N contribution to be 30% in their FOS aperture. Note that this has not been accounted for in the spectrum in Fig. 4. How ever, their estim ate assumed a uniform nebular background and was performed on an image with a PSF a ected by spherical abberation.We have redone the exercise with the F574M image from 1999, and estimate that within a 10 pixel radius, the PW N contributes at least 50%. This can also be seen by com paring our accurate pulsarphotom etry with our 10 pixel test. The contam ination of the HST/FOS spectrum would be even larger if the centering was not perfect. A though form al uncertainties of the VLT spectral ux appear to be smaller, the contam ination from the nebula in the optical range is even stronger than in the UV, exceeding 250% at the redmost wavelengths of the VLT spectral range. This is natural, since the brightness of the PW N increases more steeply with the wavelength than does the pulsar brightness (see below).



Fig. 5.  $10^{00}$   $10^{00}$  region of the eld around PSR B0540-69.3 as observed in the F547M band with the HST/W FPC2 (M orse 2003). Background stars and the pulsarprojected on the PW N have been subtracted o (cf. Fig. 1). We have marked six selected areas in the plane of the assumed torus as well as two areas in the jets. The ellipse shows the aperture used to measure the ux from the whole PW N. The pulsar position is marked by a white cross.

To sum m arize this part, we conclude that the current spectral data on the pulsar em ission cannot be considered as reliable. Hopefully, narrow -slit spectroscopy with the HST/STIS could help to solve the contam ination problem .T im e resolved photom etry with a high signal-to-noise is also a powerful tool to obtain the pulsed em ission in a contam inated area, as can be seen even for the C rab pulsar (Sollerm an 2003). On the other hand, our broadband HST spectrum of the pulsar, where the background from the nebula has been accurately subtracted o , can be considered as a fair estim ate of the pulsar spectral energy distribution. As seen from Fig. 4, it is signi cantly steeper than found by Middleditch et al. (1987). If we de ne the spectrum as  $F = F_0 (=_0)$ , then our power-law index is  $= 1.07^{+0.20}_{0.19}$  while that of Middleditch et al. is

= 0:33 0:45 (see Table 5) using updated dereddening corrections. (The method how to calculate these powerlaw indices is described in Sect. 2.5.) The atter spectrum of M iddleditch et al. is partially due to a spectral jump upwards for the U band, whereas our ux toward the overlapping F336W band goes down sm oothly, as would be expected from the extrapolation of the longer wavelength data.Thisuptum in Middleditch et al. (1987) could be due to a system atic error in their U band ux. W e note that also the C rab-pulsar broadband spectrum by M iddleditch et al. (1987) has a signi cant excess in the U band, which has not been con med by more recent spectral observations extending even further into the UV (Sollerman et al. 2000). However, possible variability of the pulsar em ission in the UV range cannot be excluded as another cause of the di erence between our results for PSR B0540-69.3 and the result by M iddleditch et al. (1987). W e also note that while we are studying the time-integrated ux, M iddleditch et al. studied the tim e-resolved ux, albeit with a poor signal-to-noise and spatial resolution.

## 2.5. Photom etry of the PW N

U sing the HST images discussed in Sect. 2.3 we have also made aperture photom etry of the continuum emission from the whole PW N, as well as di erent parts of it selected on the basis of the morphology of the nebula. For this analysis we have discarded the F555W and F658N

Iters due to the risk of contam ination by the [O III] and H emission from the SNR. To make an accurate analysis we rst subtracted o all stars in the immediate vicinity of the nebula as well as stars which overlap with the PWN.We also subtracted o the pulsar. For the subtraction we used the allstar task within the Daophot package (Stetson 1987) and for a comparison also the credit task within the Crutil package. There was no signi cant difference in the subtraction results using the two m ethods. The photom etry was perform ed with the Isophote package (Jedrzejew ski 1987).

The PW N of PSR B0540-69.3 has a rem arkable structure that can be seen clearly in Figs. 1 and 5. To sam ple di erent regions, we have put circle apertures at eight different positions, one in each jet (with an aperture radius of 12 pixels, i.e.  $0^{\circ}$ , 55), and three (with an aperture radius of 7 pixels, i.e.  $0^{\circ}$ , 32) on each side of the pulsar in the plane of the presum ed torus. We also constructed an elliptical aperture with a 74 pixel (i.e.  $3^{\circ}$ , 4) sem im a jor axis, ellipticity 0.3, and positional angle 45 degrees, that encapsulates the entire PW N (cf. Fig. 5), except for the weak northem



Fig.6. Broad-band uxes from the full PW N obtained with the HST (thick errorbars) using the elliptical aperture shown in Fig.5. The thick line and led area provide the best power-law t to the data and its 1 uncertainties, respectively (see Table 6). Previous data by Chanan et al. (1984) in the I and B bands (thin errorbars and dashed line), as well as the slope of the pulsar spectrum (cf. Fig. 4), norm alized arbitrary (dot-dashed line), are shown for comparison. Numbers near the lines denote spectral indices. Note that Chanan et al. (1984) did not correct for contam inating ux from background stars and nebular line em ission.

jet. The apertures are marked and identi ed in Fig. 5.A ll regions show emission in all lters, except for the northerm jet (N orth Jet") and A rea 6" which are not detected in the F 336W band. For these two regions we provide 3 upper limits, based on the standard sky deviations per pixel within the respective areas. The results are presented in Table 6.

As expected, the measured broadband spectra from the whole PW N and its di erent parts are well described by power-laws with negative spectral index, which con-

m the nontherm alorigin of the continuum nebular em ission. To derive the spectral indices we used the following method.For each data point we made Gaussian ts in log space to the ux and the lter function. We then simulated 10,000 sets of data using a M onte C arlo code which uses a fast portable random generator<sup>5</sup>, and for each data set we made a linear t to obtain a power-law. The power laws were then ordered in increasing value of the power law index, and the median value was chosen to represent the best t power law. In order not to be dependent on the seed value for the random number series, we ran the code 500 times with dierent seed values, and then took the average value for the power-law index to be the nal estim ate of the index. The led hexagons in Figs. 6 and 7 show 1 errors estim ated from the constraint that 68% of the constructed power laws must lie within a 1 area.

The advantage of using a M onte C arbo code rather than simple weighted means to estimate power law indices is that we can allow and test for non-G aussian distributions in log space. This is obviously not the case with a steep spectrum, non-G aussian liter functions, as well as upper limits. O ur tests, how ever, show that the liters are narrow enough to get good ts from G aussian ts to the points with estimated uxes. From our M onte C arb code approach it is also easy to estimate the error of the derived power-law index. The same approach was also used to t the pulsar spectrum in Sect. 2.4. The results are presented in Tables 5, 6 and shown for the emission from the whole PW N in Fig. 6 and from its di errent regions in Fig. 7.

According to Chanan et al. (1984) the uxes (in Jy) from the whole PW N are  $\log F_B$ 2:19 and  $\log F_{I}$ 2:63. Ourvalues are much lower ( 60% and 40%, respectively, see Fig. 6). The main reason is that we have subtracted o the pulsar and stars overlapping with the PW N, whereas these objects are not resolved from the PWN in the B and I im ages of Chanan et al. (1984) which were obtained at 1°2-1°4 seeing. Hence, their B and I uxes are contam inated by non-PW N em ission and this changes significantly the derived spectral slope of the PW N: Chanan et al. (1984) obtain = 1:06 (after dereddening with  $= 1.48^{+0.09}_{0.08}$  from V = 0.20 whereas we get Ε(B the HST data.

C om pared with the pulsar the whole PW N is more than an order of magnitude brighter, and its spectrum is signi cantly softer (cf. Tables 3 and 6 and thick solid and thin dot-dashed lines in Fig. 6). This shows that the NS spindow n power is transformed to optical emission more e ciently in the PW N than in the pulsar magnetosphere (see below).

A more detailed study shows that the spectrum may vary over the PW N. This is seen from Fig. 7 where we have plotted the results of the HST photom etry of di erent parts of the PW N and the respective power-law spectral ts (thick lines) with their 1 uncertainties (uniform ly

led regions).Stripe- led regions show extensions of the ts in cases when only upper limits in one of the three bands were obtained. The spectral hardness of som e parts of the nebula is comparable or even higher to that of the pulsar, as it is for A reas 1, 5, 6", and both jets". The spectra of the N  $\pm$  part of the torus-like structure appear to become harder tow and the outer boundary of the nebula. On the other hand, A rea 2", which is the brightest among the three selected areas S-W of the pulsar, has a steeper spectrum than its closest neighbors.

A nother feature of the spatial ux and spectral variations of the PW N is demonstrated by Figs. 8 and 9 where we have plotted the distributions of the frequencyintegrated optical uxes F, derived from the above spectral ts, and the spectral indices versus the angular distance from the pulsar along the m a jor axis of the toruslike structure of the PW N. There is a signi cant decrease of the surface brightness going from the brightest area, area 2, tow and the N-E edge of the torus (Fig. 8). The brightness di erence exceeds the 6-sigm a level of the un-

<sup>&</sup>lt;sup>5</sup> http://www.ntnu.no/ jpem/t3e-asm/vranm.ar.html

9

Table 6.B road-band uxes and power-law t ( $F = F_0$  (= 0) ) parameters of the emission from di erent regions of the PW N marked in Fig.5.Their o sets from the pulsar and areas are given in the 2nd column.Upper and lower values for each entry of the uxes are the measured and dereddened uxes with E (B V) = 0.20 ( $A_V = 0.62$ ), respectively.

	Source			Band			Power-law t
R egion	0 set	A rea arcsec²	F336W Log uz	F547M x from the are	F791W Ea Jy		log F $_{0}$ , $_{0} = 5.47  10^{14} \text{ H z}$ Jy
FullPW N	0°°000 N 0°°000 W	10.92	1.688 (6) 2.093 (6)	2.177 (7) 2.426 (7)	2.482 (8) 2.640 (8)	{ 1.48 <sup>+ (9)</sup>	{ 2.414 (12)
A rea 1	0°°455 S 0°°409 W	0.33	0.350 (116) 0.755 (116)	0 <b>.</b> 647 (94) 0 <b>.</b> 896 (94)	0.991 (48) 1.144 (48)	{ 1.09 <sup>+ (33)</sup> (30)	{ 0.949 (50)
A rea 2	1 <sup>∞</sup> 001 S 0 <sup>∞</sup> 910 W	0.33	0.224 (146) 0.629 (146)	0.766(35) 1.015(35)	1.054 (30) 1.207 (48)	{ 1.58 <sup>+ (43)</sup> (42)	{ 0 <b>.</b> 980 (45)
A rea 3	1°456 S 1°365 W	0.33	0.176 (147) 0.581 (147)	0 <b>.</b> 628 (60) 0 <b>.</b> 877 (60)	0.892 (45) 1.045 (45)	{ 1.28 <sup>+ (43)</sup> (41)	{ 0 <b>.</b> 856 (49)
A rea 4	0°°409 N 0°°364 E	0.33	0.114 (236) 0.519 (236)	0 .582 (128) 0 .831 (128)	0.868 (102) 1.021 (102)	{ 1.49 <sup>+ (67)</sup> (59)	{ 0 <b>.</b> 806 (58)
A rea 5	1°046 N 0°865 E	0.33	0.252 (140) 0.657 (140)	0.485 (106) 0.734 (128)	0.791 (64) 0.944 (64)	{ 0.92 <sup>+ (37)</sup> (33)	{ 0 <b>.</b> 786 (58)
A rea 6	1⁰775 N 1⁰365 E	0.33	0.032 <sup>a</sup> 0.437	0.239 (98) 0.488 (98)	0 .377 (102) 0 .530 (102)	{ 0.27 <sup>+ (91)</sup> (90)	{ 0.488 (98)
North Jet	1°001 N 2°366 W	0.96	0.265ª 0.670	0.642 (50) 0.891 (50)	0.875 (68) 1.028 (68)	{ 0.87 <sup>+ (56)</sup> (54)	{ 0.891(51)
South Jet	1°274 S 1°547 W	0.96	0.483 (128) 0.888 (128)	0.796 (43) 1.045 (43)	1.105 (33) 1.258 (33)	{ 1.03 <sup>+ (35)</sup> (30)	{ 1.077 (38)

<sup>a</sup> 3 upper lim it.

certainty level of the dimmest area, area 6, and shows an asymmetry of the ux distribution with respect to the pulsarposition which is similar to what is also seen in X-rays with  $Chandra/HRC^{6}$ . In the Crab PW N, the brightness di erence between the near and far sides of the torus, for a given viewing angle, is usually explained by Doppler boosting and relativistic aberration of the synchrotron radiation from relativistic particles owing at subrelativistic velocities from the pulsar in the torus plane, assuming an axisymmetric distribution of the pulsar wind around the pulsar rotation axis (e.g., Kom issarov & Lubarsky 2004). However, the considerable asymmetry of the brightness distribution between the two sides (N-E versus S-W ) of the torus of PSR B0540-69.3, as seen in projection, makes axial symmetry less obvious in a general picture. This is further strengthened by a sim ilar asym m etry in the torusplane, albeit less pronounced, seen in recent X -ray in ages of the C rab PW N (e.g., M oriet al 2004). The asym m etry

can be produced either by breakdown of axial sym m etry in the pulsar wind (e.g., due to plasm a instabilities) or by inhom ogeneity of the PW N environm ent, i.e., an asym m etry of the SN ejecta. The latter is indeed indicated by the asym m etric distribution of optical lam ents projected on the PW N of PSR B0540-69.3 (M orse 2003), as well as the general redshift of the gas em itting optical lines (K irshner et al. 1989; Sera m ovich et al. 2004).

The spatial distributions of F and in Figs. 8 and 9 appear to have sim ilar shapes and m ay suggest a harder spectrum from the dimmer outer areas of the PW N. The data are, how ever, rather uncertain and a constant spectral index of 1 seem s to be compatible with all errorbars in Fig. 9. To check that more thoroughly we analyzed the {F distribution presented in Fig. 10 which also includes

both jet areas. A linear regression t to versus log (F) using the m ethod described in this Sect. above yields

<sup>&</sup>lt;sup>6</sup> chandra harvard edu/photo/2004/snr0540



F ig.7.B road-band optical spectra for all the selected areas of the PW N shown in Fig.5.The led hexagons show 1 uncertainty regions around the best-t power laws indicated by thick lines. The uxes of the \jets" are normalized to a 7-pixel aperture area for convenience in order to compare their brightnesses with the brightness of other parts of the PW N. The spectral slopes of the whole PW N (thin lines) and the pulsar (dot-dashed lines) are shown for comparison. The dot-dashed and thin lines have been shifted arbitrarily in the vertical direction.

This is shown by a solid line in Fig.10, and indicates that brighter structures of the PW N have a steeper spectral index. The gray area shows the 1 uncertainty of the t. The range of the line slopes within this area is  $1:35^{+1:02}_{0:98}$ . A zero slope, and thus no correlation, is allowed only if we increase the uncertainty area to 1.4. This value can be considered as an overall signi cance of a possible  $-\log(F)$  correlation.We also checked the spectral index- ux correlation using the spectral t parameter F<sub>0</sub> as well as the measured ux value F in the F547M band (see Table 6) instead of the derived F and got only a slightly tighter correlation.

Therefore, deeper observations of the PWN are needed to probe a correlation between the brightness and the spectral index which is indeed only marginally indicated by the current optical data. A study of the index- ux distribution in X-rays would also be useful for the PWN of PSR B0540-69.3, as has recently been done for the Crab PWN (M oriet al. 2004). The study of M ori et al. shows that for brightnesses above 0.7 counts s<sup>1</sup> arcsec<sup>2</sup> there is a hint that (cf. their Fig. 3) the spectral index of the torus region increases with the surface brightness, as marginally indicated also in our case for PSR B0540-69.3. However, including

0.4 counts  $s^1$  arcsec  $^2$ , the brightnesses down to spectral index { surface brightness distribution appears at. The indices of the Crab jet are generally smaller com pared to those of peripheral PW N regions although their surface brightnesses are comparable (Mori et al. 2004). In the case of PSR B0540-69.3 the correlation may be enhanced by a larger brightness asymmetry of the PWN.We note that such a correlation, as well as а at index versus ux distribution, contradicts simple expectations from synchrotron cooling of relativistic particles which suggest a softening of the underlying electron spectrum toward the PWN boundary. In this picture, the fainter outer regions of the nebula would em it softer spectra. However, this simple picture does not work even for the much better studied C rab, where the PWN torus does not change its size signi cantly from radio to hard X -rays, whereas the respective cooling times dier by many orders of a magnitude implying much larger extents in the radio and optical than in X-rays. The same appears to be true for PSR B0540-69.3 (Caraveo et al. 2000). This is not yet explained, neither by the classical isotropic pulsar-wind model of a PWN by Kennel & Coroniti (1984), nor by modern MHD versions of it (Bogovalov & Khangoulyan 2002;



Fig.8.0 ptical uxes F in the (3:8 8:9)  $10^{4}$  Hz frequency range from di erent parts of the PW N marked according to notations in Fig.5 versus the distance from the pulsar along the major axis of the torus-like structure of the PW N.N egative and positive labels on the horizon-tal axis correspond to the N-E and S-W directions from the pulsar, respectively. The bold errorbar at the center shows the ux from the whole PW N rescaled to the 7 pixel aperture area used for measurements in \A reas 1 { 6".



F ig.9. Sam e as in F ig.8 but for spectral indices. Sym bols denote the sam e area num bers as in F ig.8.

K om issarov & Lubarsky 2004; DelZanna et al.2004) despite the fact that these also include anisotropy of the wind along the pulsar rotation axis, and can qualitatively explain the observed tonus-jet structure invoking a com – plicated axisymmetric picture of the wind term ination shock in the internal region of the PWN.

# 2.6. Propermotion and astrometry of PSR B0540-69.3.

The position of PSR B 0540-69.3 is de ned on the HST PC chip frames with an accuracy of better than 0.17 PC pixels which corresponds to  $0^{\circ}$ 0077. This is only a factor of 1.5 larger than the yearly proper motion value reported by M anchester et al. (1993b), see Table 1. This allows a direct estimate of the proper motion of the pulsar using accurate superposition of the F 555W and F 547M in ages taken at epochs separated by 4 years (see Sect. 2.3). We used the



Fig.10. Spectral index versus optical ux,  $\log(F)$ , in the (3.8 8.9)  $10^{14}$  Hz range for the di erent areas of the PW N marked according to Fig.5. The thick solid line shows the best t linearm odel given by Eq. (4). The point marked by bold errorbars shows the parameters for the full PW N rescaled to a 7-pixel aperture (cf. Fig. 8). The gray area shows the 1 uncertainty area for possible linear ts to the data points using the method described in Sect. 2.5. A zero slope correlation is excluded at the 1.4 level.

positions of 9 reference stars to construct the coordinate transform ation between the two in ages with the IRAF routines geom ap and gregister. The ms errors of the transform ation twere 0.078 and 4:6 10<sup>4</sup> of the PC pixel size in RA and DEC, respectively, with residual sbeing 0.156 pixels in RA and 8:1 10<sup>4</sup> pixels in DEC. U sing in centroid for m easuring the coordinates of the pulsar we nd a shift of 0:431 0.203 pixels between its positions for a time di erence of 3.995 years, where the error accounts for the centroid and transform ation uncertainties. This corresponds to a proper motion = 4:9 2:3 m as y<sup>1</sup> in the South-East direction at a position angle of 108:7 32:9 (along the southerm jet). The signi cance of this result is low and can be considered only as an attempt to make a

rst direct m easurem ent of the proper m otion. B ased on the displacem ent between the pulsar optical position and the center of the PW N, as seen in radio, M anchester et al. (1993b) argued for a sim ilar value of the proper m otion but in the South-W est direction (in the plane of the torus). W e note that the proper m otion of the C rab pulsar is aligned with the sym m etry axis of the inner C rab nebula, as de ned by the direction of the X -ray jet discovered by RO SAT (C araveo et al. 1999), and that a sim ilar situation applies to the Vela pulsar (D e Luca et al. 2000; C araveo et al. 2001; D odson et al. 2003). If our estim ates are close to reality, we have the intriguing situation that

Table 7. Coordinates of PSR B0540-69.3.

R ange	R A .(J2000)	Dec. (J2000)
	$05^{h} 40^{m} +$	-69 19 <sup>0</sup> +
X —ray <sup>a</sup>	11 <sup>s</sup> 0 0 <sup>s</sup> 7	57 <b>°</b> 4 2°0
0 ptical <sup>b</sup>	10 <sup>s</sup> .99 0 <sup>s</sup> .18	55°°1 0°°5
R adio <sup>c</sup>	11 <sup>s</sup> 1	57 <b>°</b> 5
X —ray <sup>d</sup>	11°221 0°13	2 54 <sup>00</sup> 98 0 <sup>00</sup> 7
0 ptical <sup>e</sup>	11 <sup>s</sup> :173 0 <sup>s</sup> :12	1 54 <sup>00</sup> +41 0 <sup>00</sup> +7

<sup>a</sup> Seward et al. (1984);<sup>b</sup> Caraveo et al. (1992);

<sup>c</sup> M anchester et al. (1993a); <sup>d</sup> K aaret et al. (2001);

<sup>e</sup> this work

all these three young pulsars appear to m ove along the jet axis. A di erence is, how ever, that while the C rab and Vela pulsars both have transverse velocities of 130 km s<sup>1</sup>, our results for PSR B0540-69.3 indicate a higher transverse velocity 1190 560 km s<sup>1</sup>, assum ing a distance to the LM C of 51 kpc (P anagia 2004). A third epoch of HST im aging to con rm the large value and direction for the transverse velocity is clearly needed to establish this result at a higher signi cance level. B ased on our proper m otion estim ates a level of > 3 can be achieved starting from the beginning of 2005.

We also determ ined the coordinates of the pulsar in the F547M image using seven GSC-II stars visible within the PC chip frame and the IRAF routines comap, octran, and cosetwics. The form alims errors of the astrometric t are  $0^{\circ}423$  and  $0^{\circ}472$  in the RA and DEC, respectively. Combined with the nominal GSC-II catalog accuracy of  $0^{\circ}5$ , this gives an accuracy of the position of the pulsar of  $0^{\circ}655$  and  $0^{\circ}688$  in RA and DEC, respectively. In Table 7 we compare our astrometry with previous results. We note the good agreement between our measurement and the latest Chandra result (K aaret et al. 2001).

#### 2.7.X-ray spectrum

#### 2.7.1. Interstellar absorption

The new optical results allow us to update the multiwavelength picture of the emission of PSR B0540-693. We will discuss this in greater detail in Sect. 3. Before establishing the overall spectrum, one rst needs to accurately correct for absorption and scattering by the interstellar gas and dust. In a recent analysis of time-resolved Chandra data, K aaret et al. (2001) argue that the pulsed emission of PSR B0540-69.3 can be approximated by a power-law  $F_E = F_{E_0} (E=E_0) \stackrel{E}{=} with \stackrel{E}{=} = 0.83 \quad 0.13$ within the photon energy range  $E = 0.6 \quad 10.0 \text{ keV}$  range, provided that the foreground photoelectric absorption is caused by gas with Milky W ay (MW) abundances and a column density of N<sub>HI</sub> = 4.6  $10^{21}$  cm<sup>-2</sup>. A similar num ber (N<sub>HI</sub> =  $4.0^{+0.6}_{0.4} \quad 10^{21}$  cm<sup>-2</sup>) was estimated by Finley et al. (1993) analyzing ROSAT X-ray data.

However, the use of MW abundances is obviously a simpli cation for PSR B0540-69.3. As a matter of fact, only a fraction of the photoelectric absorption is likely to

occur in the M ilky W ay. The recent Parkes 21 cm multibeam survey of the LMC (Staveley-Sm ith et al. 2003) shows that the MW contribution to the column density in the direction to PSR B0540-69.3 is just  $N_{HT}$  $0.6 \quad 10^{21}$  cm<sup>2</sup>. This survey also shows that the maximum value of  $N_{HI}$  in the LMC is 5:6  $10^{21}$  cm<sup>2</sup>, and that this occurs close to the position of PSR B0540-69.3. This is consistent with the hydrogen column density found by thing the wings of the Ly absorption pro le for the neighboring LMC star Sk 69 265,  $N_{HI} = (5 0;5)$  $10^{21}$  cm<sup>2</sup> (G ordon et al. 2003). It is quite likely that PSR B0540-69.3 could have a similarly high column density, especially since its dispersion measure (see Table 1) 50% higher than for any other pulsar in the LM C is (Craw ford et al. 2001). A rough estim ate of the LM C part of N<sub>HT</sub> for PSR B0540-69.3 can be obtained from scaling of the estimated column density for SNR 1987A and its neighboring star, \Star 2". M ichael et al. (2002) used LM C abundances to derive N  $_{H I}$  (LM C) =  $2 5^{+0.4}_{0.3}$  $10^{21}$ cm  $^2$  for SNR 1987A, and Scuderiet al. (1996) obtained  $N_{HI}$  (LM C) =  $3.4_{0.9}^{+1.0}$  10<sup>21</sup> cm<sup>2</sup> for Star2 allowing for a foreground MW contribution of  $0.6 \ 10^{21}$  cm<sup>2</sup>.We adopt  $N_{HI}(LMC) = 2.9^{+1:0}_{0:9}$  10<sup>21</sup> cm<sup>2</sup> for SNR 1987A and Star 2. Assuming that SNR 1987A and PSR B0540-69.3 su er sim ilar am ounts of LM C absorption in proportion to the LM C 21 cm emission (Staveley-Sm ith et al. 2003) along their respective lines of sight, we obtain  $N_{H_{T}}^{0.540}$  $5 \cdot 5^{+1:9}_{1:7}$   $10^{21}$  cm<sup>2</sup>, where H  $1^{0540}$  =  $N_{H_{T}}^{87A}$  H I<sup>0540</sup>=H I<sup>87A</sup> 32:00 Jy/beam and  $HI^{87A} = 16:75$  Jy/beam are the line ux densities toward the pulsar and SNR 1987A, respectively, according to the Parkes survey data  $base^7$ . The assumptions used to obtain this result for N  $_{\rm H\,I}^{0540}$  are of course uncertain, but the result points in the sam e direction as the estimates from the 21 cm emission, Sk 69265 and the pulsar dispersion measure discussed above, i.e., the column density for PSR B0540-69.3 is high. A ssum ing that the 21 cm em ission at the position of PSR B 0540-69.3 marks an upper lim it to its N  $_{\rm H\,I}$  (LM C ), we can lim it the range to  $N_{HI}^{0540} = (4.6 \ 1.0) \ 10^{21} \text{ cm}^2$ . This is sim ilar to what was used by K aaret et al. (2001), but with the important di erence that the X-ray absorption is not mainly Galactic, but arises in the LMC.

To illustrate the e ect of LM C abundances we show in Fig. 11 the ratio of photoelectric cross sections (per hydrogen atom) in the LM C and MW for the energy range 0.1 (10 keV.W e will refer to this ratio as f (E). The drop in f (E) at E > 0.28 keV (the K-shelledge of carbon) just re ects the low erm etal content in the LM C compared to the Galaxy. For the MW we have used the abundances in M orrison & M cC am m on (1983, henceforth MM 83), and for LM C we have adopted the abundances of H e, C, N, O, M g, Si and Fe from K orn et al. (2002). We have also included the elements N e, N a, A l, S, A r and C a for which we have assumed that the LM C abundances in M M 83, respectively. The exact numbers for these elements are not

www.atnf.csiro.au/research/multibeam/release/

in portant for our analysis since the absorption is dom inated by C, N, O and Fe in the energy range we are most interested in.W e assume that the interstellar gas is neutral, and we disregard dust. Photoionization cross sections were taken from the code used in Lundqvist & Fransson (1996) with further updates for Na, Mg, Al, Ar, & Causing the TOP base archive (C unto & M endoza 1993), as well. as for He (Sam son et al 1994; Pont & Shakeshaft 1995). W e have tested this code against the results of M M 83 for solar abundances, and the cross sections agree to the sam e level of accuracy as the recent cross sections com piled by W ilms et al. (2000). The comparison against MM 83 is relevant as K aaret et al. (2001) did their analysis using XSPEC Ver. 10.0 which uses data fully compatible with M M 83. Figure 11 shows that LM C abundances strongly suppress the photoelectric absorption, and that C and O are particularly in portant constituents at the energies for which K aaret et al. (2001) claim photoelectric absorption is most important for the observed pulsar spectrum, i.e., at < 1 keV.



F ig.11. Ratio of photoelectric absorption per hydrogen atom in the LMC and M ilky W ay for the energy range 0.1{ 10 keV. A rrow sm ark at which energy each element starts to contribute to the photoelectric absorption. The absorbing gas is assumed to be neutral. The gradual changes of f (E) across absorption edges are due to moderate zoning of the photon energy in the code used for the cross section calculations. See text for further details.

We have used the function f(E) in combination with the results of M M 83 to see how LM C abundances may a ect conclusions about the derived spectrum of PSR B0540-69.3. If we assume that the pulsar emits a pulsed power-law spectrum with slope  $_{\rm E}$  = 0.83 in the range 0.6 10:0 keV, as argued for by K aaret et al. (2001) and marked in Fig.12 as a straight dotted line, the attenuated spectrum should look like the solid line in Fig.12 after passage through a column density of neutral hydrogen in the M ilky W ay with a value of N<sub>HI</sub>(MW) = 4:6 10<sup>21</sup> cm<sup>2</sup>. If we disregard possible elects of an accurate treat-

ment for the response matrix of Chandra, we can deabsorb this spectrum with a more likely composition for the photoelectrically absorbing gas. For this we have chosen  $N_{HT}$  (MW) = 0.6 10<sup>21</sup> cm<sup>2</sup> and  $N_{HT}$  (LMC) = 5.0 10<sup>21</sup>  $\,\mathrm{cm}^{-2}$  . The deabsorbed spectrum is marked by the upper dashed line in Fig. 12. At the lower energy lim it of the t by K aaret et al. (2001), i.e., at 0.6 keV, the deabsorbed spectrum undershoots by a factor of 2:9 com pared to the power-law, but on the other hand overshoots by orders of m agnitude at energies below the K-shell edge of carbon. The latter can be xed by just low ering  $N_{HI}$  (LMC) to  $10^{21}$  cm  $^2$  , i.e., still consistent with the likely < 4:3 range argued for earlier in this section. The spectrum would in that case undershoot by a factor of > 3:8 at 0.6 keV com pared to the power-law. The assumption of a power-law spectrum at energies below a few keV, where the photoelectric absorption sets in, is of course uncertain. Our results could indicate that the intrinsic spectrum is not a power-law, but actually falls below the power-law at 0.6 keV. However, before jumping to such a conclusion we need to check another possible source of X-ray absorption, namely the supernova ejecta.



F ig.12. Soft X-ray spectrum of the pulsed em ission from PSR B0540-69.3. The dotted line shows the intrinsic power-law spectrum argued for by K aaret et al. (2001), and the solid line shows the attenuated spectrum after passing through a column of gas with MW abundances, N<sub>HI</sub>(MW) = 4.6  $10^{21}$  cm<sup>2</sup>. The lower dashed line shows deabsorption using a more realistic value for the MW contribution, N<sub>HI</sub>(MW) = 0.6  $10^{21}$  cm<sup>2</sup>, and the upper dashed line shows further deabsorption with the value N<sub>HI</sub>(LMC) = 5.0  $10^{21}$  cm<sup>2</sup> for LMC, which is close to the maximum value found for the LMC as a whole according to Staveley-Sm ith et al. (2003). The vertical bar at 0.6 keV marks the energy above which K aaret et al. (2001) tted their power-law to the deabsorbed spectrum.



F ig.13. Same as F ig. 12, but with a lower value for the LMC deabsorption,  $N_{\rm H\,I}$  (LMC) = 4:1  $10^{21}$  cm<sup>2</sup>. D eabsorption due to supernova ejecta has also been included, assuming the same structure as in SN 1987A. The structure of SN 1987A on day 118 (B linnikov et al. 2000) was expanded hom ologously until 500 years (m odel \A", long-dashed) or 750 years (m odel \B", short-dashed) after the explosion. N ote that the age-parameter in these m odels do not necessarily relate to the sam e age for SN R 0540-69.3 as it depends on the explosion energy, the envelope m ass and the density structure. See text for further details.

# 2.7.2. Supernova ejecta absorption

Observations of SNR 0540-69.3 show that the remnant is oxygen-rich (Kirshner et al. 1989; Sera movich et al. 2004), which m eans that the progenitor star probably was massive. Kirshner et al. (1989) suggest it to have had a mass similar to that of the SN 1987A progenitor, i.e., around 20 M , and the results of Sera m ovich et al. (2004) show that a likely range is 15 22 M . To check the im portance of supernova ejecta for possible X -ray absorption, the well-studied SN 1987A therefore serves as a good tem plate. W e have used the m ixed m odel 14E1 displayed in Figs. 2 and 6 in B linnikov et al. (2000), which has 2 M of pure oxygen spread out over the innerm ost 5 M of supernova ejecta. We assume that the post-explosion structure expands hom ologously. At an age of 1000 years after explosion, the hydrogen column density through the ejecta, from the center, is only  $N_{HI}$  (SN) 1:1  $10^{19}$  cm  $^2$ , ie, far less than even the MW contribution in the direction toward LM C. However, the very high m etal content in the supernova ejecta, and its concentration toward the center of the ejecta, boosts the X-ray absorption at energies above 300 eV. Above the K-shell ionization edge of oxygen, i.e. at 600 eV, the cross section per hydrogen atom is 40 times higher than for the MW absorption in M M 83. Because N  $_{\rm H~I}$  (SN ) scales with the time since explosion as  $/ t^2$ , a less evolved supernova remnant can make a signi cant contribution to the X-ray absorption of em ission from the pulsar and its  ${\tt PW}$  N .

The importance of the supernova ejecta is illustrated in Fig.13.Here we have used the same value for N  $_{\rm H \, I}\,(\!M \,\,W\,$  ) as in Fig. 12, but lowered the LMC contribution to  $N_{HI}(LMC) = 4:1 \quad 10^{21} \text{ cm}^2$ . We have also included deabsorption due to supernova ejecta, assuming a hom ologously expanding SN 1987A at an age of 500 years (model A", long-dashed) and 750 years (model B", short-dashed). The increased oxygen column density com pared to that in Fig. 12 m akes it possible to retrieve the initial power-law spectrum above 0.6 keV for a supernova similar to SN 1987A at an age of 500 years, but it is also clear that a rem nant with an age closer to the spin-down age of PSR B0540-69.3 will not contribute signi cantly to the X-ray absorption. A lthough the ejecta of SNR 0540-69.3 could expand more slowly than those of SN 1987A, could contain more oxygen (and/or having it more concentrated to the center), could have a clum py and asymmetric structure (as is indicated for SN 1987A, W ang et al. 2002), or could have a low er age than the pulsar spin-down age, it seems that we have to stretch the parameters to claim that the X-ray absorption along the line of sight to the pulsar is largely a ected by the supernova ejecta. A possible way to test this is to check the spatial variations of the X-ray absorption over the larger PW N. The low m etal content of LM C, and the low MW foreground absorption, make such tests sensitive to any supernova ejecta contribution. X-ray spectral ts to data obtained with Chandra of the Crab pulsar have recently highlighted the importance of the abundance factor (W illingale et al. 2001; W eisskopf et al. 2003). U sing various abundances it was found that the line of sight to the Crab is signi cantly underabundant in oxygen.

To sum marize this section, it is evident that the X-ray spectral analysis of PSR B 0540-69.3 needs a revision. C ontrary to previous assumptions, the m etal abundance of the X-ray absorbing gas must clearly be sub-solar, unless the supernova ejecta contribute signi cantly. The latter, how ever, appears to be less likely. W hile a full analysis of the X-ray spectrum, i.e., a detailed reduction of the C handra data including various abundance com binations in X SPEC, is beyond the scope of this paper, we have argued that the power-law spectrum, which seems to be appropriate to use at energies above 1 keV, m ay experience a depression below 1 keV (cf. Sect. 3). This can be tested by how the X-ray spectrum connects to that in the optical. W e will discuss that in Sect 3.

# 2.8. Extinction toward PSR B0540-69.3

As for the column density toward PSR B0540-69.3, the foreground optical extinction in the MW is well established. From the study of Staveley-Sm ith et al. (2003), the color excess in the direction toward PSR B0540-69.3 is E (B V)<sub>MW</sub> = 0.06, and this agrees with the value found for SN 1987A (Scuderi et al. 1996) and that used by G ordon et al. (2003). The value used for the parameter R in these works is the standard value 3.1.

The extinction parameters E (B V) and R(V) for LM C along the line of sight to SNR 0540-69.3 and the pulsar have not yet been investigated in detail in the same way as for SN 1987A (Scuderiet al. 1996). However, in Sera movich et al. (2004) we studied SNR 0540-69.3 and H II regions close to it. The reduced spectra were analyzed using the total value E (B  $V)_{TOT} = 0:20$  and R (V) = 3:1, i.e., the same num bers we used in Sect. 2, and we found that the H /H /H line ratios are in good agreementwith Case B recombination theory (Baker & Menzel 1938; Hummer & Storey 1987). A llowing for higher extinction would boost H and H relative to H causing a disagreem ent with the Case B theory, which for these lines norm ally explains the observations of supernova rem nants well (e.g., Fesen & Hurford 1996).

Looking at the projected pulsar neighborhood, G ordon et al. (2003) obtain the average value R (V)<sub>LM C</sub> = 2:76 0:09 for the LM C 2 supershell, and it seems reasonable that this could be used also for PSR B0540-69.3. Out of the eight stars form ing this average, six of them can accom modate the standard value of 3.1 within 1. The spread in E (B V)<sub>LMC</sub> ranges between 0.12{0.24 (including 1 errors), so the values we have used for PSR B0540-69.3 and its PWN in Sect. 2.4 and 2.5 seem reasonable also from this comparison. However, to check the e ect of a di erent extinction curve on E (B V), and still being com patible with Case B line ratios for SNR 0540-69.3, we have compared  $k_{\text{ext}}$  (cf. Sect. 2.4) for the extinction used in Sect.2.4 (we call that case C1") to a case (called C2") with E (B  $V)_{LMC} = 0:14$  (R (V) = 2:76, G ordon et al. 2003) and E (B V)<sub>MW</sub> = 0:06 (R (V) = 3:1, Cardelliet al. 1989). We form ed the ratio  $g() = k_{ext}() = k_{ext}(H)$ for both cases, and found that  $jg()_{C1}$   $g()_{C2}$   $j=g()_{C1}$ does not exceed 2% within the interval 2620 {8480 A . O nly at the very blue end of the FOS spectrum of Hill et al. (1997), i.e., at 2500 A, does the ratio approach 5%. This justi as the use of E (B  $V_{LMC} = 0.14$  (assuming  $V_{MW} = 0.06$ ) regardless of whether we choose to E(B use extinction combinations like C1 or C2. The absolute

ux level of the dereddened spectrum of course depends on the exact value of R (V) being used. Further direct extinction studies of the pulsar and its neighborhood in the UV and optical bands are needed to pin down the detailed extinction corrections. It seems, however, that the steep spectral slopes we obtain for the pulsar and its PW N in the optical in Sect. 2.4 and 2.5 cannot be corrected by som e extrem e reddening corrections as this is neither justi ed by our observations of the supernova rem nant nor by the study by G ordon et al. for supposedly neighboring objects.

# 3.D iscussion

# 3.1. Multiwavelength spectrum of PSR B0540-69.3

To connect the optical pulsar em ission to the em ission at other wavelengths, we have com piled results for the radio (M anchester et al. 1993a), X -ray (K aaret et al. 2001; de P laa et al. 2003) and -ray (Thom son et al. 1994) spectral regions. The unabsorbed spectrum is displayed in F ig. 14. In comparison with previous compilations (e.g., H irayam a et al. 2002; de P laa et al. 2003) the accuracy is signi cantly improved due to the new high spatial resolution data obtained in the optical with HST and in X-rays with C handra. This reveals new features in the spectrum of the pulsar.

The optical and X-ray parts of the spectrum can be tted with power-laws (for speci cassum ptions about the intervening extinction and photoelectric absorption discussed in Sect. 2.7 and 2.8), which would suggest a nontherm al nature of the emission in both domains, likely to be form ed in the magnetosphere of the rotating neutron star. How ever, the connection between the emission in X-rays and in the optical seems far from trivial, especially if the slope for the optical spectrum derived from the archival HST/W FPC 2/F336W data is correct (spectralindices and other em ission param eters for PSR B0540-69.3, as well as for the C rab and Vela pulsars, in the optical and X-rays are presented in Table 8). The data for PSR B0540-69.3 suggest at least two spectral breaks between the optical and X-ray spectral bands. For a com parison, Fig. 14 also shows the multiwavelength spectrum of the total pulsed em ission from the C rab pulsar (K uiper et al. 2001; Sollerm an 2003). For the C rab, it seems that a sm ooth turn-over can be possible (dotted line) between the X -ray band and the optical. This is in contrast to spectra of the middle-aged pulsars Vela and PSR B0656+14, whose optical uxes are generally compatible with the low-frequency extrapolation of a power-law spectral tail for E > 1 2 keV (K optsevich et al. 2001; Shibanov et al.2003).

In Sect. 2.7 we noted that PSR B 0540-69.3 could have a non-powerlaw spectrum below 1 2 keV, and inspired by this we tried to use the shape of the C rab optical/X – ray tum-overto t the spectrum of PSR B 0540-69.3. This, how ever, fails for the W FPC 2/F 336W band where the ux falls below such a t. As we pointed out in Sect. 2.8, the depression in U is unlikely to be caused by insu – cient dereddening; to reach the C rab pulsar spectral slope

02, cf. Table 8) one has to apply E (B V) 0:55, ( which is at least twice as high as the most likely value (cf. Sect. 2.8). Does this mean that PSR B0540-69.3 experiences a spectral dip in the F336W band? W hile future deep and well-calibrated data in U and UV should reveal this, we note that such an explanation is not farfetched. As a matter of fact, the broad-band optical spectra of middle-aged pulsars (Vela, PSR B0656+14, and Geminga, cf. Shibanov et al. 2003) do show a dip in the U and B bands, which could indicate a multicomponent continuum spectrum, or the presence of unresolved em ission/absorption features, possibly related to electron/ion cyclotron lines originating in the magnetospheres of the neutron stars (Mignani et al. 1998; Jacchia et al. 1999). If the depression in U is of m ore general character, the multiw avelength spectrum of PSR B 0540-69.3 suggests a double break \knee" in the spectral region between the optical



F ig.14.M ultiwavelength unabsorbed spectrum of PSR B0540-69.3. The data were obtained with the di erent telescopes and instruments marked in the plot. The optical data are from this paper. Phase-averaged (upper polygon) and pulsed (lower polygon) X -ray emission spectra with their uncertainties are shown for the 0.6 { 10 keV range (K aaret et al. 2001). The pulsed emission spectra obtained with ROSAT and RXTE are from de Plaa et al. (2003), and the radio and -ray data are from M anchester et al. (1993a) and Thom son et al. (1994), respectively. All upper limits are de ned at 3 level. The unabsorbed spectrum of the total pulsed emission of the C rab pulsar is shown for com parison, where the high energy and optical data are from K upper et al. (2001) and Sollerm an (2003), respectively. The dotted line for the C rab pulsar indicates a possible connection between the optical/X -ray regions through the EUV. The same line (shifted to overlap with the soft X -ray spectrum) is overlaid on the optical/X -ray region of PSR B0540-69.3.N ote how the blue band uxes of PSR B0540-69.3 undershoot by a factor of 3 com pared to this overlay while it ts well in the X-ray band.

and soft X -ray bands.0 bærvations in the UV and reanalysis of the C handra X -ray data with accurate corrections for extinction and photoelectric absorption, as discussed in Sect. 2.7, will help us constrain the position of the breaks and to understand whether they are located just bluew ard of the U band and below 0.6 keV, or occur in the EUV range.

Taking into account the di erence in distance to the C rab pulsar ( 2 kpc) and PSR B 0540-69.3 ( 51 kpc), as well as the spectral energy distributions for both pulsars (see F ig. 14), we note that the overall intrinsic ux from PSR B 0540-69.3 is alm ost as high as that from the C rab

pulsar in the radio (while its slope is possibly steeper in this range), but that it is 1:4 and 3 timeshigher in the optical and X-ray ranges, respectively. This is also show n in Table 8.At the same time, the spindow n lum inosity, E-, of PSR B0540-69.3 is 3 times lower than for the C rab. Therefore, the e ciency of producing nonthermal optical and X-ray photons in the magnetosphere of the rotating neutron star from its spindow n power, = L=E-, is a factor

4 and 10 higher for PSR B 0540-69.3 in the optical and X-rays, respectively (Table 8). For a comparison we show also in Table 8 the parameters for the 10 times older Vela pulsar, which is much less lum inous and a less e -

Table 8.C om parison of the optical and X-ray spectral indices ( $^{\circ}$ , $^{X}$ ), lum inosities ( $L^{\circ}$ , $L^{X}$ ), e ciencies ( $^{\circ}$ , $^{X}$ ),
and a weighted ratio ( $L^{\circ} = E^{X} = L^{X} = C^{\circ}$ ) of the two young pulsars, C rab and P SR B 0540-69.3, and the older Vela
pulsar. Here $E^{0}$ and $E^{X}$ are the energy intervals in the optical <sup>a</sup> and X-rays <sup>b</sup> , respectively, used for the frequency
integration to obtain $L^0$ and $L^X$ . Information on the period P, dynamical age , spindown lum inosity E-, and distance
d for each pulsar is included.

PSR	Р		E-	d	0	L <sup>0;a</sup>	0	Х	L <sup>x</sup> ;b	Х	L <sup>O</sup> EX
	m s	kyr	$10^{37}$ erg s $^1$	kpc		$10^{33}$ erg s $^1$	10 5		$10^{36}$ erg s $^1$	10 <sup>3</sup>	
C rab	33.49	1.24	46	2	-0.21°	7 <b>.</b> 9°	1.7	0 <b>.</b> 87 <sup>d</sup>	1.0 <sup>d</sup>	2.17	34.9
0540	50.2	1.66	15	51	1.07 <sup>e</sup>	10 <b>.</b> 9 <sup>e</sup>	7.3	0 <b>.</b> 83 <sup>d</sup>	3.13 <sup>d</sup>	21	15.5
Vela	89	11	0.69	0.29	$0.25^{f}$	7 <b>:</b> 0 <sup>f</sup> 10 <sup>5</sup>	1:0 10 <sup>3</sup>	0 <b>.</b> 5 <sup>g</sup>	2:3 <sup>g</sup> 10 <sup>5</sup>	3 <b>:</b> 4 10 <sup>3</sup>	13.5
<sup>a</sup> For the optical range 1.57{3.68 eV.						<sup>e</sup> Th	is paper.				

<sup>a</sup> For the optical range 1.57{3.68 eV.

 $^{\rm b}$  For the X -ray range 0.6{10 keV .

 $^{\rm c}$  U sing data from Sollerm an (2003) for 1.57{3.68 eV .

<sup>d</sup> K aaret et al. (2001);  $L^{X}$  rescaled to the 0.6{10 keV range.

cient optical and X -ray em itter, but still capable of powering a weak and less extended PW N around it. The Vela pulsar has also a peculiar optical spectrum in com parison with the C rab pulsar with a possible excess in the near-IR and a dip in the U band (Shibanov et al. 2003). Based on the available data it is not yet clear whether these spectral peculiarities in the pulsar optical emission do indicate a spectral evolution with pulsar age (PSR B0540-69.3 has a spin-down age which is 400 years higher than the C rab pulsar) or whether they are connected to the pulsar optical e ciency, or just re ects speci c param eters (e.g., viewing angle and magnetic eld geometry) of each pulsar. Detecting PSR B0540-69.3 in the near-IR would allow us to understand to which extent its spectrum is sim ilar to that of the Vela pulsar and other m iddle-aged pulsars detected in the optical range. We also note that the ratio of optical to X -ray lum inosity, weighted by the observed spectral ranges (cf. the last column in Table 8), is for PSR B0540-693 about half the value for the Crab pulsar, but com parable with that of the Vela pulsar.

## 3.2. Multiwavelength spectrum of the PW N

A compilation of our optical data for the PWN around PSR B0540-69.3 together with radio and X-ray data is shown as an unabsorbed multiwavelength spectrum in Fig 15. The spectrum suggests the same double knee connection between the optical and X -ray spectral parts as for the pulsar (cf. above). How ever, the assumed knee-breaks appear to be less pronounced than for the pulsar because the order of m agnitude higher ratio for the intrinsic optical to X-ray ux for the PW N than for the pulsar. This is also re ected in the PWN lum inosities presented in Table 9, where we have also collected information on the spectral indices and lum inosities of the C rab and Vela PW Ns. The sm oothness of the knee in the PW N spectrum as compared to the pulsar spectrum may be explained by propagation e ects of the relativistic particles generated in the pulsar m agnetosphere and m oving through the PW N. If the particle energy distribution function contains features which are re ected in the pulsar spectrum, these features should  $^{\rm f}$  U sing data from Shibanov et al. (2003).

<sup>g</sup> Pavlov et al. (2001a); L<sup>X</sup> rescaled to the 0.6{10 keV range and d = 0.29 kpc.

become less pronounced due to such propagation e ects. In Fig. 15, we also note a smaller di erence between the radio and optical uxes for the PWN, compared to for the pulsar, which may be also be caused by such propagation e ects.

Unfortunately, there is no data of sim ilar quality on the spatially averaged optical spectrum of the C rab PW N, but spectra of speci c signatures of the C rab PW N (knots, wisps etc.) are signi cantly steeper than the pulsar spectrum (Sollerm an 2003). On a larger scale, the continuum optical em ission of the whole C rab N ebula, at  $10^{00}$  spatial resolution (Veron-Cetty & Woltjer 1993), shows a sim ilar spectral behavior. In particular, a bright area roughly overlapping in position and m orphology with the C rab X ray PWN was resolved. Its spatially averaged spectrum is steeper than the spectrum of the pulsar. W e see the same behavior for the whole PW N of PSR B0540-69.3 and for som e parts of it (cf. Figs. 6 and 7), whereas the much fainter Vela PW N has not yet been detected in the optical (M ignaniet al. 2003). How ever, som e hints of speci c structures in the Vela case, clearly detected in Xrays (Helfand et al. 2001; Pavlov et al. 2001), may have been detected also in the near-IR JH bands (Shibanov et al. 2003). There is no reliable spectral inform ation on them and their identication still have to be con med by deeper observations both in the optical and  $\mathbb{R}$ .

The opticale ciency of the PW N of PSR B0540-693 30 higher than for the pulsar. This is is a factor of markedly dierent from the situation in X-rays where the e ciency of the PWN is only 4 times higher than for the pulsar (K aaret et al. 2001). If we assume that the bright optical areas of Veron-Cetty & Woltjer (1993) are associated with the the optical emission of the whole C rab PW N, we obtain a similar situation also for this pulsar 50 higher optical e ciency of the PW N com with its pared to that of the pulsar, while the e ciency ratio is only a factor of 2{3 in X-rays. This may give a hint on the particle energy distribution responsible for the synchrotron em ission of the PW N.

To estimate the ratio of the e ciencies in the optical and X -rays we norm alized the lum inosity in each dom ain



F ig.15. Spatially averaged multiwavelength spectrum of the PSR B0540-69.3 PW N obtained with the di erent telescopes indicated in the plot. The optical data are from the present work, whereas the radio and X-ray data are from M anchester et al. (1993b) and K aaret et al. (2001), respectively. The led regions (for the HST and Chandra data) show 1 uncertainty regions around the best- t power laws indicated by thick lines. We have also included sim ilar data for the C rab PW N, where the data are from Velusamy et al. (1992), Veron-C etty & W oltjer (1993) and K aaret et al. (2001) for the radio, optical and X-ray bands, respectively. The dotted curved line is a cubic spline t for the multiwavelength spectrum of the C rab PW N, and the dashed-dotted line is a linear t for the optical/X-ray region of the C rab PW N. These ts have been shifted to the level of the X-ray em ission of the PSR B0540-69.3 PW N for a com parison. Note how the ts overshoot for PSR B0540-69.3 in both the optical and the radio.

Table 9. Same as in Table 8 but for the PW Ns of the same pulsars. Information on the PW N size and the ratios of the pulsar to PW N lum inosities in the optical and X-rays for each pulsar is included.

PW N	size	0	L <sup>0;a</sup>	0	Х	L <sup>x</sup> ;b	Х	L <sup>O</sup> EX	L <sup>0</sup> <sub>psr</sub> =L <sup>0</sup> <sub>pw n</sub>	L <sup>X</sup> <sub>psr</sub> =L <sup>X</sup> <sub>pw n</sub>
	рс		$10^{33}$ erg s $^1$	10 5		$10^{36}$ erg s $^1$	10 <sup>3</sup>			
C rab	1.5	0.92°	4240 <sup>c</sup>	920	1 <b>.</b> 14 <sup>b</sup>	21 <b>.</b> 8 <sup>d</sup>	47.5	867	0.0017	0.046
0540	0.6{0.9	1.5°	366 <sup>e</sup>	245	1.04 <sup>b</sup>	12 <sup>d</sup>	79.7	136	0.03	0.26
Vela	0.14	{	{	{	0.5 <sup>f</sup>	$6:8^{d}$ 10 $^{4}$	9 <b>:</b> 8 10 <sup>2</sup>	{	{	0.34

<sup>a</sup> For the optical range  $1.57\{3.68 \text{ eV}$ .

<sup>b</sup> For the X -ray range 0.6{10 keV.

<sup>e</sup> This paper.

<sup>f</sup> From Pavlov et al. (2001b);  $L^{X}$  rescaled to the 0.6{10 keV range and d = 0.29 kpc.

<sup>c</sup> Optical uxes are taken from Veron-Cety & Woltjer (1993).

 $^{\rm d}$  K aaret et al. (2001);  ${\tt L}^{\rm X}\,$  rescaled to the 0.6{10 keV range.

to the respective energy range for which the lum inosity was calculated. The ranges are 3370 { 7870 A (i.e., 1.58{ 3.68 eV) and 0.6 { 10 keV . As seen in Tables 8 and 9, the norm alized optical lum inosity ( $L^{\circ} = E^{\times} = L^{\times} = E^{\circ}$ , where the parameters are de ned in Table 8) is 16 and 140 times higher than that in X-rays for the pulsar and its PW N, respectively. The inferred values for the Crab are even higher ( 35 and 870, respectively). This is natural

since the rotational loss of the C rab pulsar is 3 times higher and its multiwavelength spectrum does not show a depression in the optical, as PSR B 0540-69.3 appears to have.

The PWN of PSR B0540-69.3 has similar sizes in the optical (this paper) and in X-rays (G otthelf & W ang 2000; K aaret et al. 2001) and extends up to  $4^{00}$  away from the pulsar, which corresponds to 1

pc at 51 kpc. However, the brighter emission is conned to 0.6 0.9 pc (cf. Fig. 5). This is a factor of 2 sm aller than the size of the C rab PW N which has also similar sizes in the optical and in X-rays (Hester et al.  $2002^8$ ). The sm aller size of the PSR B0540-69.3 PW N is in rough agreement with the expected (e.g., Kennel & Coroniti 1984) PW N size scaling with pulsar spindown luminosity, /  $E^{-0.5}$ . A similar proportionality has been found from the comparison of the C rab and Vela PW Ns (Helfand et al. 2001; Pavlov et al. 2001b).

The higher optical and X-ray e ciencies of PSR B0540-69.3, as compared with the Crab pulsar, are not re ected in a simple way in its PWN e ciencies. For instance, the X-ray e ciency of the PSR B0540-69.3 PW N is almost twice as high as that of the Crab PW N, while its optical e ciency is 4 tim es lower than in the C rab case. The reason for this is unclear and may be explained either by the propagation e ects discussed above, or by di erent pulsar environm ents, or a combination of these e ects. The pulsar contribution to the total pulsar+ nebula X-ray lum inosity is 4.5%, 21%, and 25% for the PSR B0540-69.3, Crab, and Vela pulsars, respectively. The contribution is smaller in the optical and ranges from 0.17% for the Crab to 3% for PSR B0540-69.3. The Vela PW N has not been detected in the optical range, but we note that even for this much older and fainter pulsar, the PW N dom inates the total X -ray lum inosity, and that its contribution is comparable to that of PSR B0540-69.3, although E- is an order of the m agnitude sm aller. This cannot be explained by a simple scaling with the spindown lum inosity, as in case of the PW N sizes, and requires additional studies.

Our results do not reveal any signi cant variation of the spectral index over the torus and 'et like structures of the PW N of PSR B0540-69.3, although a m arginal inverse correlation between the spectral hardness and brightness cannot be excluded at 1:4 level (cf. Fig. 10 and Sect. 2.5 form ore details). No signi cant variation of the spectral index was also found in recent X -ray studies of the Crab PW N along its torus and the cores of the PW N jets (M oriet al. 2004). This suggests that for both these PW Ns the energy spectra of the emitting electrons and positrons injected in these two di erent directions by the shocked pulsar wind are similar. This fact, as well as the similar sizes of the these parts of PW Ns in the optical and X-rays suggest that the particle spectra in the bright, central parts of PW Ns are not a ected by synchrotron cooling. In the Crab case the spectral softening and hence the synchrotron losses become signi cant only in the faint outerm ost regions of the PW N detected in X-rays (M oriet al. 2004; W eisskopf et al. 2000) and in the UV/optical ranges (Scargle 1969; Veron-Cetty & Woltjer1993; Hennessy et al. 1992). A sim ilar dependence may be seen in the Vela PW N (Pavlov et al. 2001b). This in poses in portant constraints on the pulsar w ind m odels.

D eeper observations in the optical and X -rays are needed to study the faint outer regions of the PW N of PSR B0540-69.3, and to test the reality and m agnitude of the inverse hardness/brightness correlation.W e note a signi cant surface brightness asymmetry with the respect to the pulsar position along the major axis of the torus structure of the PW N of PSR B0540-69.3 seen in both the optical and in X-rays. This asymmetry can hardly be explained within the fram ework of axisym metrical pulsar wind models by sim ply invoking D oppler boosting and relativistic aberration e ects, as has been done for the near and far sides of the C rab PW N torus. The asym m etry is m ore likely to be caused by plasm a instabilities in the internal parts of the pulsarwind ow, or by asym metry of the SN ejecta. The latter is also discussed from another point of view in Sect. 3.3.

#### 3.3. Proper motion and ejecta asymmetry

In Sect. 2.6 we found the indicative result that PSR B0540-69.3 has a high proper motion corresponding to 1190 560 km s<sup>1</sup>, and that the direction is consistent with the pulsarm oving along the southern jet axis. A pulsar velocity of 1000 km s<sup>1</sup> is high, but not exceptional. In a recent VLA study of pulsar proper motions, Brisken et al. (2003) show that there is a relatively large fraction of pulsars which have velocities in excess of 500 km s<sup>1</sup>. Peng et al. (2003) quantify this number to be 16%.

Even if our results for the proper motion of PSR B0540-69.3 have only a 2 signi cance, and a third epoch of HST imaging is needed to test whether or not PSR B0540-69.3 belongs to this high-velocity class of objects, we cannot avoid connecting the origin of the possibly large velocity of PSR B0540-69.3 to the origin of the signi cant redshift of the inner part of SNR 0540-69.3 which has been estimated to be several hundred km s<sup>1</sup> (K irshner et al. 1989; Sera m ovich et al. 2004). The m ost straightforward interpretation of this is that the 0540-69.3 system could be the result of a very asymmetric explosion.

Evidence of asymmetric supernova ejecta in corecollapse supernovae is abundant, and perhaps of speci c interest for the 0540-69.3 system with its asymmetric inner ejecta structure (e.g., K irshner et al. 1989), is that in general the asym m etry appears to increase with depth into the ejecta (see A kiyam a et al. 2003 and references therein). The prime example is SN 1987A where the powering by radioactive nucleids in the center occurs along bipolar jets that are likely to be aligned with the rotationalaxis of the presupernova (W ang et al. 2002). A ssum ing that also the explosion in core-collapse supernovae itself could be jetinduced, K hoklov et al. (1999) nd from 2D -m odeling that pulsar kick velocities of 1000 km s<sup>1</sup> can be achieved. The pulsar would move along the jet axis, consistent with our tentative nding for PSR B0540-69.3, but to reach a velocity as high as 1000 km s<sup>1</sup> a large di erence in momentum between the two jets is required. A spointed out

<sup>&</sup>lt;sup>8</sup> see also Chandra public in ages at chandra harvard edu/photo/2002/0052/0052\_xray\_opt.pg

by Lai (2000), it is not clear what could give rise to such a di erence.

Laiet al. (2001) discuss various models how to produce large pulsar kick velocities in the context of the aligned spin axis and pulsar propermotion in Crab and Vela (cf. Sect. 2.6), and their discussion may now also apply to PSR B0540-69.3. The conclusion is that spin-kick alignm ent requires fast, perhaps close to break-up rotation, at the pulsar birth. There is observational evidence pointing in the sam e direction at least for the C rab pulsar (A toyan 1999; Sollerm an et al. 2001). We note that am ong the various models reviewed by Lai et al. (2001), the hydrodynam ically driven kicksm ay face a problem in reproducing kick velocities of the order we infer for PSR B0540-69.3, and Fryer (2004) nds that even in his most asymmetric 3D-m odels of an exploding 15 M star, the neutrino em ission becomes asymmetrically emitted, thereby damping out the hydrodynam ical pulsar kick. Fryer suggests that one way to obtain fast pulsars is to rely on them being produced by low-mass progenitors (8 12 M). This is, how ever, not a likely explanation for the 0540-69.3 system, which most probably originates from a 20 M progenitor. The recent models of Scheck et al. (2004) appear to bem ore successful in producing pulsars with high kick velocities. It could also be that models including rotation will alter the results of the hydrodynam ically driven kick models (cf. Laiet al. 2001), perhaps providing a link to the results of K hok lov et al. (1999). Further observations of the 0540-69.3 system will show if it can add to C rab and Vela as a testbed for di erent kick scenarios.

A cknow ledgem ents. W e are gratefulto Jelle de P laa for sharing with us his reduced RO SAT and RXTE data of PSR B0540-69.3, to Lucien K uiper for high energy data of the C rab pulsar, and to the referee, Patricia C araveo, for useful com m ents allow ing us to im prove several im portant points in the text. Partial support for this work was provided by RFBR (grants 02-02-17668, 03-02-17423 and 03-07-90200). and support was also given by The Royal Swedish A cadem y of Sciences. The research of PL is further sponsored by the Swedish R essearch C ouncil. PL is a Research Fellow at the Royal Swedish A cadem y supported by a grant from the W allenberg Foundation. The work was initialized while N IS was supported by a stipend from The Swedish Institute.

#### References

- A kiyam a, S., W heeler, J.C., M eier, D.L., Lichtenstadt, I.2003, ApJ, 584, 954
- Atoyan, A.M. 1999, A&A, 346, L49
- Avila, G., Gero, R., & Beckers, J. M. 1997, in Optical Telescopes of Today and Tomonrow, SPIE 2871, ed. A. Ardeberg, 1135
- Baker, J.G., Menzel, D.H. 1938, ApJ, 88, 52
- Blair, W . P., Long, K . S., W ancura, O., et al. 1992, ApJ, 399, 611
- Blinnikov, S. I., Lundqvist, P., Bartunov, O. S., et al. 2000, ApJ, 532, 1132
- Bogovalov, S.V. & Khangoulyan, D.V. 2002, MNRAS, 305, 211

- Boyd, P.T., van Citters, G.W., Dolan, K.G., et al. 1995, ApJ, 448, 365
- Brisken, W . F ., Fruchter, A . S ., G oss, W . M ., et al. 2003, A J, 126, 3090
- Caraveo, P.A., Bignami, G.F., Mereghetti, S., & Mombelli, M. 1992, ApJ, 395, L103
- Caraveo, P.A., De Luca, A., Mignani, R.P., & Bignami, G. F.2001, ApJ, 561, 930
- Caraveo, P.A., Mignani, R.P., 1999 A&A, 344, 367
- Caraveo, P.A., Mignani, R.P., De Luca, A., et al. 2000, in A decade of HST science, eds. Mario Livio et al. (Baltimore) 105, [astro-ph/0009035]
- Cardelli, J.A., Clayton, G.C., M athis, J.S. 1989, ApJ, 345, 245
- Chanan, G.A., Helfand, D.J., & Reynolds, S.P. 1984, ApJ, 287, L23
- Chevalier, R.A., & Kirshner, R.P. 1979, ApJ, 223, 154
- Clark, D.H., Touhy, I.R., Long, K.S. 1982, ApJ, 255, 440
- Carw ford, F., Kaspi, V. M., M anchester, R. N., et al. 2001, ApJ, 553, 367
- Cunto, W ., Mendoza, C., Ochsenbein, F., & Zeippen, C.J. 1993, A & A, 275, L5
- DeLuca, A., Mignani, R.P., & Caraveo, P.A. 2000, A&A 354, 1011
- DelZanna,L.,Amato,E.,& Bucciantini,N.2004,A&A,421, 1063
- de Plaa, J., Kuiper, L., Herm sen, W. 2003, A&A, 400, 1013
- Dodson, R., Legge, D., Reynolds, J.E., & McCulloch, P.M. 2003, ApJ, 596, 1137
- Dopita, M . A ., & Tuohy, I. R ., 1984 ApJ, 282, 135
- Fesen, R.A., & Hurford, A.P. 1996, ApJS, 106, 563
- Finley, J.P., O gelm an, H., H asinger, G., et al. 1993, A pJ, 410, 323
- Fryer, C.L. 2004, ApJ, 601, L175
- Fukugita, M ., Shim asaku, K ., Ichikawa, T . 1995, PASP, 107, 945
- Gordon, K.D., Clayton, G.C., Misselt, K.A., et al. 2003, ApJ 594, 279
- Gotthelf, E.V. 2003, ApJ, 591, 361
- Gotthelf, E.V., & W ang, Q.D. 2000, ApJ, 532, L117
- H am uy, M ., Suntze , N .B ., H eathcote, S.R .1994, PA SP, 106, 566
- Hester, J., Mori, K., Burrows, D., et al. 2002, ApJ, 577, L49
- Hill, R.J., Dolan, J.F., Bless, R.C., et al. 1997, ApJ, 486, L99
- H irayam a, M ., N agase, F ., Endo, T ., et al. 2002, M N R A S, 333. 603
- Hummer, D.G., & Storey, P.J. 1987, MNRAS, 224, 801
- Hwang, U., Petre, R., Holt, S.S., & Szym kowiak, A.E. 2001, ApJ, 560, 742
- Jacchia, A., de Luca, F., Lazzaro, E., et al. 1999, A&A, 347, 494
- Jedrzejewski, R. I. 1987, MNRAS, 226, 747
- K aaret, P., M arshall, H.L., A kdcroft, T.L., et al. 2001, ApJ, 546, 1159
- Kennel, C.F., & Coroniti, F.V. 1984, ApJ, 283, 694
- Khokhlov, A.M., Ho ich, P.A., Oran, E.S., et al. 1999, ApJ, 524, L107
- Kirshner, R. P., Morse, J. A., W inkler, P. F., & Blair, W. P. 1989, ApJ, 342, 260
- Kom issarov, S.S., & Lubarsky, Y. 2004, MNRAS, 349, 779
- K optsevich, A , B , Pavlov, G . G , Zharikov, S . V , et al. 2001, A & A , 370, 1004

- Kom, A.J., Keller, S.C., Kaufer, A., et al. 2002, A&A 385, 143
- Kuiper, L., Herm sen, G., Cusum ano, G., et al. 2001, A&A 378, 918
- Lai, D. 2000, in Stellar Astrophysics, eds. L. S. Chang et al. (K luwer) 127, [astro-ph/9912522]
- Lai, D., Chemo, D.F., & Cordes, J.M. 2001, ApJ, 549, 1111
- Lundqvist, P., & Fransson, C. 1996, ApJ, 464, 924
- MacAlpine, G.M., Law rence, S.S., Sears, R.L., et al. 1996, ApJ, 463, 650
- Manchester, R.N., Mar, D.P., Lyne, A.G., et al. 1993a, ApJ, 403,L29
- 1993b, ApJ, 411, 756
- Maran, S.P., Sonneborn, G., Pun, C.S.J., et al. 2000, ApJ, 545, 390
- Mathewson, D.S., Dopita, M.A., Tuohy, I.R., & Ford, V.L. Wilms, J., Allen, A. & McCray, R. 2000, ApJ, 542, 914 1980, ApJ, 242, L73
- M ichael, E., M cC ray, R., Chevalier, R., et al. 2003, ApJ, 593, 809
- M ichael, E., Zhekov, S., M cC ray, R. 2002, ApJ, 574, 166
- M iddleditch, J., & Pennypacker, C.R. 1985, Nature, 313, 659
- Middleditch, J., Pennypacker, C.R., Burns, M.S. 1987, ApJ, 315,142
- Mignani, R.P., Caraveo, P.A., Bignami, G.F. 1998, A&A, 332, T.37
- Mori, K., Burrows, D. N., Hester, J. J., et al. 2004, ApJ, 609, 186
- Morrison, R., McCammon, D. 1983, ApJ, 270, 119
- Morse, J.A. 2003, Rev Mex AA (Conf. Series), 15, 243
- Oke, J.B., Gunn, J.E. 1983, ApJ, 266, 713
- Panagia, N. 2004, in the Proceedings of IAU Colloquium 192: Supernovae (10 years of SN 1993J), eds. J.M. M arcaide & K W .W eiler (Springer Verlag), in press [astro-ph/0309416]
- Pavlov, G.G., Kargaltsev, O.Y., Sandwal, D., et al. 2001a, ApJ, 552, 129
- Pavlov, G.G., Zavlin, V.E., Sandwal, D., et al. 2001b, ApJ, 554, 189
- Peng, Q., Zhang, L., Luo, X., & Chou, Ch. 2003, in Young neutron stars and their environm ent IAU Symp.218, eds. F.Camilo & B.M.Gaensler (ASP Conference Proceedings) 17, [astro-ph/0309090]
- Pont, M., & Shakeshaft, R. 1995, J. Phys. B, 28, L571
- Sam son, J.A.R., He, Z.X., Yin, L., & Haddad, G.N. 1994, J.Phys.B, 27, 887
- Scheck, L., Plewa, T., Kifonidis, K., Janka, H.-Th., & Muller, E.2004, in Proc.12th W orkshop on Nuclear A strophysics, Ringberg Castle, in press [astro-ph/0404355]
- Scuderi, S., Panagia, N., Gilmozzi, T., et al. 1996, ApJ, 465, 956.
- Sera movich, N. I., Lundqvist, P., Sollerman, J., & Shibanov, Yu.A.2004, A&A, to be submitted
- Seward, F.D., & Hamden, Jr.F.R. 1994, ApJ, 421, 581
- Seward, F.D., Hamden, Jr.F.R., & Helfand, D.J. 1984, ApJ, 287, L19
- Shibanov, Yu.A., Koptsevich, A.B., Lundqvist, P., Sollerm an. J.2003, A&A,406,645
- Sollerm an, J. 2003, A & A, 406, 639
- Sollerm an, J., Kozm a, C., & Lundqvist, P. 2001, A&A, 366, 197
- Sollerm an, J., Lundqvist, P., Lindler, D., et al. 2000, ApJ, 537, 861
- Staveley-Sm ith, L. 1994, PA SA u, 11, 83

- Staveley-Sm ith, L., K in, S., Calabretta, M. R., et al. 2003, MNRAS, 339, 87
- Stetson, P.B. 1987, PASP, 99, 191
- Thom son, D. J., A rzoum anian, Z., Bertsch, D. L. et al. 1994, ApJ, 436, 229
- van der Heyden, K. J., Paerels, F., Cottam, J., et al. 2001, A&A,365,L254
- Velusamy, T., Roshi, D., Venugopal, V.R., 1992, MNRAS, 255, 210
- Veron-Cetty, M. P., & Woltjer, P. 1993, ApJ, 270, 370
- W ang, L., W heeler, J.C., Hoe ich, P., et al. 2002, ApJ, 579, 671
- Manchester, R.N., Staveley-Smith, L., & Kesteven, M.J. Weisskopf, M.C., O'Dell, S.L., Paerels, F., et al. 2003, ApJ, accepted [astro/ph0310332]
  - W illingale, R., Aschenbach, B., Griths, R.G., et al. 2001, A&A,365,L212