

The nearest extreme starburst: bubbles, young star clusters, and outflow, in NGC 3256

S. L. Lipari¹, R. J. D'Az¹, J. C. Forte², R. Terlevich³, Y. Taniguchi⁴, M. P. Aguero¹, A. Alonso-Herrero⁵, E. Mediavilla⁶, S. Zepf⁷

- ¹ Cordoba Observatory and CONICET, Laprida 854, 5000 Cordoba, Argentina.
- ² La Plata Observatory and CONICET, La Plata, Argentina.
- ³ Institute of Astronomy, Madingley Road, Cambridge CB3 0HA.
- ⁴ Astronomical Institute, Tohoku University, Aoba, Sendai 980-8578, Japan.
- ⁵ Steward Observatory, University of Arizona, Tucson, AZ 85721, USA.
- ⁶ Instituto de Astrofísica de Canarias, 38205 La Laguna, Tenerife, Spain.
- ⁷ Department of Physics and Astronomy, Michigan State Univ., East Lansing, MI 48824, USA.

Received ; in original form

ABSTRACT

In this work we report, for the extreme starburst in the IR merger NGC 3256:

- (i) The detection of 4 galactic bubbles (using ESO NTT and HST WFC2 H images). These shells would be associated with previous SNe explosions.
- (ii) The first analysis of the spatial distribution of young star clusters (YSCs) candidates shows that more than 90 per cent of them are located in a complex starburst structure, including some of the bubble walls, three nuclei and three blue asymmetrical spiral arms.
- (iii) The kinematics study of the ionized gas in the core of the main optical nucleus, performed with HST STIS spectra. The shape of the rotation curve and the emission line profile could be explained by the presence in the core of YSCs without outflow. Any low luminosity AGN associated to this core would have a mass less than $10^7 M_{\odot}$. It is also probable that the compact X-ray and radio emission of ULX (7)N (detected in the main optical nucleus core) is associated to a few recent SN remnants.

Key words:

galaxies: individual (NGC 3256) { ISM : bubble { galaxies: star clusters { galaxies: kinematics { galaxies: interactions { supernova: general

1 INTRODUCTION

The study of extreme star formation and the associated galactic winds processes is an important issue in modern astrophysics. In particular, this issue plays an important role in the formation and evolution of galaxies, at high and low redshift (see for references Taniguchi & Shioya 2000; Lipari et al. 2004a,b,c). Motivated by this, we began a project of study of nearby star forming + GW galaxies and distant Ly emitters (for references see Lipari et al. 2004a,b,c). The first step for this project is to understand the extreme star formation process in nearby galaxies because we can obtain more detailed and unambiguous information. Thus, our group started a study of nearby IR mergers/QSOs, with galactic winds.

Luminous infrared (IR) galaxies ($L_{IR [8-1000 \text{ m}]} > 10^{11} L_{\odot}$, LIRGs) are dusty strong IR emitters $-L_{IR} / L_B \sim 5$ (300- where frequently a strong enhancement of star forma-

tion is taking place, and a very high per cent of LIRGs (70-95%) are mergers. Therefore, at low z , luminous IR mergers ($z < 0.1$) and IR QSOs ($z < 0.5$) are excellent laboratories for the study of extreme star formation processes. In these systems very high star formation and supernova (SN) rates are expected. Hubble Space Telescope (HST) observations confirmed that star cluster forming frequency is highest during violent burst of star formation, in IR mergers (for references see the reviews of Schweizer 2002; and Whitmore 2001). Recently, Lipari et al. (2003) reported several galactic shells and arcs, in nearby IR mergers and IR QSOs with galactic winds. SNe, shells/arcs and young star clusters are thus good tracers of extreme starburst events.

NGC 3256 is a nearby LIRG ($z = 0.0094$) which is a merger between (at least) two gas rich disk galaxies (Toomre 1977; Joseph & Wright 1985). The total IR luminosity of NGC 3256 is $L_{IR [8-1000 \text{ m}]} = 3.3 \times 10^{11} L_{\odot}$, consequently the bolometric luminosity makes this system the most luminous

nearby galaxy (for $V = 3000 \text{ km s}^{-1}$; Sargent, Sanders, & Phillips 1989). In particular, the strong 10 μm IR emission in NGC 3256 is "very extended", with most of it originating outside the central kpc, indicating clearly that the near to mid-IR emission is probably powered by a recent starburst and not by a compact object or an AGN (Graham et al. 1984; Joseph & Wright 1985). Throughout the paper, a Hubble constant of $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ will be assumed. For this merger, we adopted the distance of 37.5 Mpc ($V_{\text{Syst}} = 2817 \pm 15 \text{ km s}^{-1}$; Lipari et al. 2000), and thus, the angular scale is $1'' = 180 \text{ pc}$.

2 OBSERVATIONS

Optical HST Wide Field and Planetary Camera 2 (WFPC2) observations were analysed, which include broad-band images positioned on the Planetary Camera (PC) chip with scale of $0''.046 \text{ pixel}^{-1}$, using the filters F450W (4521 Å, 958 Å, B Cousin filter), F555W (5407 Å, 1223 Å, V), and F814W (7940 Å, 1758 Å, I). In addition, narrow band line ramp filter LRF (FR680N images (6631.4 Å, 76 Å) were obtained, positioned in the WFPC2 (WF2 CCD). They allowed to map H + [N II] line emission, in an effective field of the ramp filter of $10'' \times 10''$, with a scale of $0''.1 \text{ pixel}^{-1}$.

HST Near IR Camera and Multi-Object Spectrometry (NICMOS) archival data were studied, which include mainly in images with the filters F160W (1.60 μm , 0.4 m), F190N (1.90 μm , 0.01 m, Pa γ), F222M (2.20 μm , 0.14 m) and F237M (2.37 μm , 0.14 m); using the camera 2 with a scale of $0''.076 \text{ pixel}^{-1}$.

Near-UV and optical HST Advanced Camera for Surveys (ACS) archival images were analysed, obtained with the High Resolution Channel (HRC). They include images with the filters F220W (2228 Å, 485 Å), F330W (3354 Å, 588 Å), F555W (5346 Å, 1193 Å), and F814W (8333 Å, 2511 Å). The scale is $0''.027 \text{ pixel}^{-1}$.

HST STIS long slit spectroscopy of the main optical nucleus of NGC 3256 was obtained. The spectra were taken with the G750M grating, with the slit $0''.1 \times 52''.0$, giving a resolution of 50 km s^{-1} . The STIS CCD has a scale of $0''.05 \text{ pixel}^{-1}$. The slit was aligned at PA 91°.

The ESO NTT (SUSI data include H + [N II] narrow band images, using the ESO filters E629 (6571 Å, 115 Å, H α (redshifted) and E443 (7018 Å, 65 Å, H α continuum). These NTT images have seeing of $0''.7 \text{ FWHM}$; and the CCD scale is $0''.13 \text{ pixel}^{-1}$.

The IRAF and STSDAS² software packages were used to analyse the HST and ESO imaging and spectrophotometric data. The HST data were reduced at the Space Telescope Science Institute (STScI), using the standard process. All the bands were calibrated according to the procedures detailed in Lipari et al. (2004a). Specifically, K-band photometric calibration of the HST NICMOS was performed using published photometric solution for the F222M and F237M filters (HST NICMOS Data Handbook, version 4.0).

² IRAF and STSDAS are the reduction and analysis software facility developed by NOAO and STScI, respectively

3 RESULTS AND DISCUSSION

3.1 Galactic bubbles or shells and the morphology of the extreme star formation process

Using HST WFPC2 and ESO NTT data, Lipari et al. (2000) found that the extreme massive star formation process shows extended triple asymmetrical spiral arms ($r \approx 5 \text{ kpc}$), emanating from three different nuclei. We already proposed, that probably the interaction between dynamical effects of the merger process, the galactic wind (outflow), and the molecular ionized gas (in the inflow phase) could be the mechanism that generates the massive star formation process (at scales of 5–6 kpc) with this triple asymmetrical spiral arm structure.

Figures 1a and 1b show ESO NTT and HST WFPC2 (WF2H) images of this starburst structure. The main associations of giant H II regions were labelled according to our previous work. Figures 1a and 1b depict at least 4 interesting giant galactic shells or bubbles:

(i) Shell A: This is the more defined shell detected in NGC 3256, it is located to the south (west of the region 2 (R2: a possible second optical nucleus; Lipari et al. 2000)). This shell is clearly observed in the NTT H α image with a resolution of $0''.7$ (Fig. 1a), however more interesting details were detected in the HST WFPC2 and ACS images (Fig. 1b). Half of the circular shell was detected, and the other half is heavily obscured by a strong and wide dust lines system. The near IR NICMOS data confirm this last proposition, the bubble wall is more extended to the south at this bands, and shows a knotty structure. This obscured area is located in the south region of the main body of this merger (Lipari et al. 2000). The diameter of this galactic shell is $D_A = 1''.8 = 324 \text{ pc}$.

The HST ACS F220W and F330W UV images show remarkable details of the blue compact knots located at the border of this shell (which are brighter in the east side). This knots depict strong UV continuum, probably associated to the presence of a high number of massive O and B stars.

HST WFPC2 (H α , F450W, ACS (F330W, F555W, F814W) and NICMOS (F222M, F237M, Pa γ) images show exactly at the centre of this shell, a young star cluster (YSC) candidate plus a giant H II region. This configuration constrains the age of the star-forming complex to a few Myr, this kind of coincidences between H II region emissions and YSCs (for the most massive clusters of $10^6 - 10^7 M_\odot$) could only be detected during the first $7 - 10^6 \text{ yr}$ (Alonso-Herrero et al. 2002).

(ii) Bubble B: Two narrow arcs or shells were found in the west side of the region 2 (each one is marked with a line in Fig. 1b). Furthermore, HST ACS F220W and F330W UV images show that the core of this region (R2) is the brightest area of NGC 3256, in the near-UV. This fact suggests the presence of a very high number of massive O and B hot stars, which normally start the first phase of the galactic wind/bubble and also are the progenitors of core-collapse SNe.

The bubble structure detected around R2, was already noted by Lira et al. (2002). The diameter of the more external arc is about the same than Shell A: $D_B = 1''.9 = 342 \text{ pc}$.

(iii) Shell C: A less extended arc was detected to the west side of region 4. It is observed more clearly in HST WF2H

data (Fig. 1b). The diameter of this shell is $D_c = 0.9 = 162 \text{ pc}$.

(iv) Shell D: To the west, and adjacent to region 7, the NTT H image (Fig. 1a) shows another structure with bubble shape. However, more clear details of this structure are observed in the HST WFC2 and ACS images, showing several blue compact knots. These knots are located at the inner border of the shell. The diameter of this bubble is $D_D = 1.7 = 305 \text{ pc}$.

These "shells, bubbles, arcs" could be associated mainly to explosions of SNe and the final phase of the galactic wind, i.e., the blowout phase of galactic bubbles (Norman & Ikeuchi 1989). Multiple explosion of SNe, from massive progenitors, are the main galactic objects capable to generate this blowout phase (Norman & Ikeuchi 1989). Massive YSCs and associations of giant H II regions are the places, where multiple SN explosions are expected.

3.2 The young star clusters in NGC 3256 (HST WFC2, NICMOS and ACS images)

We have started a multiwavelength studies of the young stellar cluster candidates (YSC) in NGC 3256, using HST ACS/UV, WFC2/Optical, NICMOS/near-IR images. From these HST data we study the colours, ages, masses, subgroups, spatial distribution and metallicities of these YSC. Here we report new results from this study.

First, the colour distribution, the colour-magnitude diagram and the spatial distribution of YSC (of NGC 3256) were analysed. Using mainly the Catalogue of YSC published by Zepf et al. (1999: their Table 1). We note that from this catalogue, the positions of YSC in pixels (and also the corrected values of RA and DEC) were used; since the published data of RA and DEC have small errors, which were generated in the transformation from pixels to RA/DEC. An electronic corrected version of this Catalogue of YSC is available at zepf@pa.msu.edu and lipari@oac.uncor.edu.

From this study, the following results were found:

(i) Complete sample of YSC: The optical spatial distribution of all the YSC was first analysed. Fig. 2a depicts their interesting distribution. We found that the YSC are located in more than a 90 per cent in the starburst regions: they concentrate in the 3 arms, 3 nuclei, and in some of the bubble walls, tracing the extreme starburst.

For the southern obscured areas, especially for the obscured southern nucleus (R3), and for the obscured part of the arm III, we found a similar result. In particular, using our near-IR Catalogue of YSC and H II regions (with objects detected in the HST NICMOS H and Pa bands images; Alonso-Herrero et al. 2002), we also found that the YSC and H II regions are located mainly in the nuclei, the blue arms and, at some of the bubble walls.

(ii) Very Red YSC: The colour $(B-I)$ distribution of the YSC candidates shows bimodal shape with a main (strong: 920 objects) peak at $(B-I) = 0.75$ and a second (small: 40 objects) peak at $(B-I) = 2.6$. Thus, a first subgroup of YSC was selected including the very red objects found around this second peak. These YSC were selected using the condition $(B-I) > 2.3$, since at $(B-I) = 2.3$ starts the distribution of YSC associated with this second peak. Fig. 2b shows that these very red YSC are located mainly in the

obscured arm III and around the core of the main nucleus (at a radius $r = 0.8 = 144 \text{ pc}$).

Zhang, Fall, & Whitmore (2001) found a similar subgroup of red YSC, in the IR pre-merger "The Antennae". They suggested that these objects could be the youngest clusters, detected when they emerge from their dust cocoons. For NGC 3256, the location of the very red YSC suggest that these clusters candidate are mainly very dusty objects.

(iii) Blue and Very Blue YSC: The observed and theoretical YSC colour-magnitude diagrams ($B-I$ vs. m_B) (Zepf et al. 1999: their Figs. 3 and 4), clearly show that the stellar population models (of Bruzual & Charlot (2003) and Leitherer et al. (1999)) generate unique ages between 3 and 7 Myr, for YSC with colour in the range $0.55 < (B-I) < 0.5$. See for details, English & Freeman (2003).

Using this fact, we selected two subgroups of blue and very blue YSC: those with $(B-I) < 0.5$ and $(B-I) < 0.0$, respectively. Thus the first subgroup include the second one. Fig. 2c depicts the spatial distribution of these two subgroups. Both samples were found distributed mainly in the second optical nucleus (R2) and the closer asymmetrical arm II. This fact is more evident for the very blue YSC (Fig. 2c).

These results are in agreement with a "composite" multiple merger model proposed for NGC 3256 (Lipari et al. 2000). In this model the main merger could be between two gas rich massive spiral galaxies (Sc), and then with a satellite galaxy (of one of these Sc systems). In particular, we proposed that the kinematics and spectroscopic properties of the core of region 2 could be associated to a 3rd. nucleus, arisen in a satellite of the two original spiral galaxies that collided. In order to explore these type of scenarios we are working in detailed numerical simulations, which are in progress (Lipari et al. 2000, 2004, in preparation). The results of these kind of simulations suggest that the two massive galaxies are the main component of the merger event, and that the satellite generally merges later. Since the starbursts in each blue asymmetrical arms are probably related with the multiple merger event and associated with each nuclei, we already suggested that the starburst could be younger in R2 and the arm II (Lipari et al. 2000). The new results presented in this work support this suggestion: (i) the very blue YSC are located mainly in the region 2 and the arm II; and (ii) the brightest near-UV areas (in the HST ACS images) are also located in the region 2 and the arm II.

It should be noted that 5% of the YSC studied are objects with absolute magnitude less than -13 (luminosities greater than $10^{40} \text{ erg s}^{-1}$) which are strong candidates to be massive YSC with masses in the range $10^6 - 7 M_\odot$.

It is interesting to remark that a simple estimation of the kinematical energy of the Shells A and B, using a wall thickness of 20-40 pc, electron density $N_e = 1000 \text{ cm}^{-3}$ (Lipari et al. 2000), and filling factor 0.1, yields $3-6 \times 10^{52} \text{ erg}$. This is equivalent to 1/3 of the kinetic energy of the nuclear outflow in M82 (Bland & Tully 1988).

3.3 The core kinematics and the outflow of the main optical nucleus (HST STIS spectra)

HST STIS 0.1 - 5.0 long slit spectra show a rotational pattern in the central arcsecond (core) of NGC 3256 main

optical nucleus (Fig. 3a), with an amplitude of $V_r = 160 \text{ km s}^{-1}$. For this core, Ne et al. (2003) suggested that a compact X-ray and radio source ULX (7) North (Lira et al. 2002; our R1 region) would have a mass of at least $10^8 M_\odot$ and could be associated to a supermassive black hole. Nevertheless, another possibility must be considered, since there is an extranuclear compact X-ray and radio source, ULX (13) [which is located inside of the the H II region 5, called R5 in our Fig. 1a], with the same spectral properties than ULX (7)N but not associated to a nuclear mass concentration.

In addition, using HST STIS long slit spectra with the slit $0''.2 \times 52''_0$ (100 km s^{-1} of resolution), Ne et al. (2003) suggested the presence, of broadened H and [N II] emission lines, with linewidths of 450 km s^{-1} , as evidence for an AGN in the core. Figure 4 shows the HST STIS spectrum of the core, using the $0''.1 \times 52''_0$ long slit (giving a better spectral resolution: 50 km s^{-1}). From this figure is very clear that the broadening of the H and [N II] emission lines is due to the presence of a strong blue outflow component. This component was previously detected at large scale, and associated to a nuclear galactic wind, by Lipari et al. (2000). Moreover, this broad outflow component was observed also in the spectra located at several hundred pc from the nucleus. This fact rules out the possibility that this emission could be associated with the broad line region of an AGN. In the core and in several STIS spectra (with $0''.1$ slit) we measure for the main component of the emission lines H and [N II], FWHM in the range $90\{130 \text{ km s}^{-1}$, these are normal values for H II regions. Meanwhile, for the outflow component was measured FWHM in the range $200\{260 \text{ km s}^{-1}$, probably associated with shock processes.

The central rotation curve of region 1 (Figure 3a) is well fitted by a Plummer (Kuzmin law (Binney & Tremaine 1987)), that could correspond to a spherical (or disk-like) mass distribution with the following parameters: $V_{\text{sys}} = 2898 \text{ km s}^{-1}$, scale radius $a = 40 \text{ pc}$, $M_{\text{tot}} = 2.16 \times 10^8 M_\odot$. Moreover, any single component fitted model cannot have a scale radius smaller than 40 pc (the resolution is better than 18 pc). In fact, the central slope and maximum of the rotation curve could not be fitted by a point mass larger than $10^7 M_\odot$, even allowing for a beam smearing corresponding to a resolution $0''.15$, which is twice the slit width and the image quality of the STIS data used. This is well illustrated by the grey line in Figure 3a, which corresponds to a point central mass of $2 \times 10^7 M_\odot$, with a beam smearing of $0''.15$.

Considering that several young star cluster candidates of mass $10^6\text{--}10^7 M_\odot$ and luminosities of $10^{39}\text{--}10^{40} \text{ erg s}^{-1}$ are observed inside $r \approx 50 \text{ pc}$ and that the rotation curve corresponds to that of an extended mass distribution, it seems likely that any low luminosity AGN associated to ULX (7)N would have a mass less than $10^7 M_\odot$ and would not contribute significantly to the core luminosity. Indeed, the radial change of mass to F814W (I) luminosity ratio (Figure 3b), calculated in the way described in detail in Lipari et al. (2004a), does not allow the possibility of a hidden dark mass kinematically detected, down to the resolution of $0''.1$. There is no evidence of a strong central light or gravity point source, that would generate, respectively, a strong deep or peak in the M/L_I profile. Moreover, the central values of the mass to F814W (I) luminosity ratio

are similar to those found in other extended starbursts, e.g. our study of M/L_I in NGC 2623.

It seems more plausible that the compact X-ray and radio emission of ULX (7)N, as for ULX (13), is associated to a few recent SN remnants in the massive young star clusters of the nucleus (R1) core. As was pointed out by Ne et al. (2003) this would require several supernovae in the nucleus in the last 15 years, which seems a possibility to be considered.

3.4 The possible IR supernova in 1997 HST NICMOS images (in NGC 3256)

Using the HST NICMOS images, taken at F222M ($K_s, 2.2 \mu\text{m}$) and F237M ($2.4 \mu\text{m}$) bands we found a very bright source in the main body of NGC 3256. This object appears in the HST images obtained in 1997 Nov. 28.9 UT. We previously identified this object as a possible SN event (PSN 9711-001; Lipari et al. 2004d). Recently, we found that this object is associated to the emission from the coronagraphy hole of NIC-2 camera.

ACKNOWLEDGMENTS

Part of the HST observations were obtained from the archives at ESO Garching and STScI Baltimore. The authors thank J. A. Costa (Pulido, H. D. Ottori, and B. Garcia-Lorenzo for discussions and assistance. Finally, we wish to thank the referee, Dr. Richard de Grijs, for constructive and valuable comments, which helped to improve the content and presentation of the paper.

REFERENCES

- A Alonso-Herrero A., Rieke G., Rieke M., Scoville N., 2002, *AJ*, 124, 166
 Binney J., Tremaine S., 1987, *Galactic Dynamics*, Princeton Univ. Press, New Jersey
 Bland J., Tully R., 1988, *Nature*, 334, 43
 Bruzual G., Charlot S., 2003, *MNRAS*, 344, 1000
 Graham G., Wright G., Meikle W., Joseph R. D., Bode M., 1984, *Nature*, 310, 213
 English J., Freeman K., 2003, *AJ*, 125, 1124
 Joseph R. D., Wright G., 1985, *MNRAS*, 214, 87
 Leitherer C. et al., 1999, *ApJS*, 121, 3
 Lipari S. L., Colina L., Mucchetti F., 1994, *ApJ*, 427, 174
 Lipari S. L., Diaz R., Taniguchi Y., Terlevich R., D'ottori H., Carranza G., 2000, *AJ*, 120, 645
 Lipari S. L., Terlevich R., Diaz R., Taniguchi Y., Zheng W., Svetanov Z., Carranza G., D'ottori H., 2003, *MNRAS*, 340, 289
 Lipari S. L., Mediavilla E., Diaz R., Garcia-Lorenzo B., A. Costa-Pulido J., Aguero M. P., 2004a, *MNRAS*, 348, 369
 Lipari S. L., et al., 2004b, *MNRAS*, submitted (astro-ph 0406515)
 Lipari S. L., et al., 2004c, in T. Storchi Bergmann, L. Ho, H. Schmidt, eds., *The Interplay among Black Hole Stars and IGM in Galactic Nucleus*, IAU Symp. No. 222, (Cambridge: Cambridge Univ. Press), in press
 Lipari S. L., Diaz R., Alonso-Herrero A., 2004d, *IAUC*, 8353 (http://cfa-www.harvard.edu/iau/CBAT_PSN.html)
 Lira P., Ward M., Zezas A., Alonso-Herrero A., Ueno S., 2002, *MNRAS*, 330, 259
 Ne S., Ulvestad J., Campion S., 2003, *ApJ*, 599, 1043
 Nomman C., & Ikeuchi S. 1989, *ApJ* 395, 372

- Sargent A., Sanders D., Phillips T., 1989, *ApJ*, 346, L9
- Schweizer F., 2002, in D. Geisler, E. Geisler, and D. Merrifield, eds., *Extragalactic Star Clusters*, IAU Symposium No. 207 (ASP Conference Series), 630
- Taniguchi Y., Shiota K., 2000, *ApJ*, 532, L12
- Toomre A., 1977, in B. Tinsley, R. Larson eds., *The Evolution of Galaxies and Stellar Population*, (Yale University Observatory, New Haven) 401
- Whitmore B., 2001, in M. Livio ed., *ST ScI Symposium Series 14*
- Zepf S., Ashman K., English J., Freeman K., Shapley R., 1999, *AJ*, 118, 752
- Zhang Q., Fall M., Whitmore B., 2001, *ApJ*, 561, 727

Figure 1. ESO NTT (SUSI) and HST W FPC 2 (W F2 H) images, of the central region of NGC 3256. Letter colour Plate 1.

Figure 1. Continued.

Figure 2. HST W FPC2{PC F435W (B) in age of the central region of NGC 3256, superimposed to: (a) the total sample of young stellar cluster (Y SC) candidates, (b) the sub-sample of very red Y SC candidates, (c) the sub-sample of blue and very blue Y SC candidates (with blue and black colour, respectively). For details see the text. The crosses mark the positions of the Y SC candidates. Letter colour Plate 2.

Figure 2. Continued.

Figure 3. (a) Velocity values of the ionized gas obtained for the core of the main optical nucleus. They were obtained from long slit HST STIS spectra, at PA = 91°, i.e. close to the kinematics major axis. The solid line shows the best Plummer-Kuzmin potential fit, and the grey line corresponds to a point central mass of $2 \times 10^7 M_{\odot}$, with a beam smearing of $0''.15$; (b) the mass to HST WFC2 F814W (I) luminosity ratio.

Figure 3. Continued.

Figure 4. Long slit HST STIS spectrum of the main optical nucleus core of NGC 3256 (at PA = 91°). It shows the H α + [N II] emission line blend, with a clear blue outflow component (in each emission line).