# The X-ray Jet of 3C 120: Evidence for a Non-standard Synchrotron Spectrum

D.E.Harris, A.E.Mossman

Sm ithsonian A strophysical O bservatory, H arvard- Sm ithsonian C enter for A strophysics 60 G arden St., C am bridge, M A 02138 harris@cfa.harvard.edu

amossman@cfa.harvard.edu

and

R.C.W alker<sup>1</sup> NationalRadio A stronom y Observatory, Box 0, Socorro, NM 87801 cwalker@nrao.edu

# ABSTRACT

We report on archival data from the Chandra X -ray O bservatory for the radio jet of 3C 120. We consider the emission process responsible for the X -rays from 4 knots using spectra constructed from radio, optical, and X -ray intensities. While a simple synchrotron model is adequate for three of the knots, the fourth ('k25'), which was previously detected by ROSAT and is now well resolved with Chandra, still represents a problem for the conventional emission processes. If, as we argue, the at X -ray spectra from two parts of k25 are synchrotron emission, then it appears that either the emission comes from an electron distribution spectrally distinct from that responsible for the radio emission, or at the highest electron energies, there is a signi cant deviation from the power law describing the electron distribution at low er energies.

Subject headings: galaxies: active { galaxies: individual (3C 120) { galaxies: jets { radiation m echanism s: nontherm al

1. Introduction

3C 120 is a nearby (z= 0.033) radio galaxy which has variously been classi ed as a Seyfert 1, a broad line radio galaxy, and a late type spiral. The optical appearance is com plex, with multiple dust lanes and two suggestive spiralarm swhich become e radial features at a projected distance of 6.3 kpc from the core (g.1). The radio morphology and lum inosity are more sim ilar to the FRI class radio galaxy (Fanaro & Riley 1974) than to the usual Seyfert. A one sided pc scale jet has been extensively studied with long baseline interferom etry and superlum inal components have been identi ed (apparent speeds between 3 and 6c, W alker et al. 2001). The kpc scale jet is shown in g.2 and there is additional complex em ission at larger scales (beyond the borders of g.2, W alker, Benson, & Unwin 1987).

At X-ray wavelengths, the nucleus is a strong and variable source (e.g. Halpern 1985), but it was the detection with ROSAT of a rather inconspicuous radio knot 25<sup>00</sup> from the core (16 kpc, projected), which presented new problems for the conventionalm odels of X-ray emission from radio jets (Harriset al. 1999; HHSSV hereafter). Therefore we obtained new radio data at higher frequencies to see if this knot was peculiar or unique either in shock structure (as manifest by gradients in ra-

 $<sup>^1\,{\</sup>rm The}$  N ational R adio A stronom y O bservatory is operated by A ssociated U niversities, Inc., under contract with the N ational Science Foundation.

dio surface brightness) or in radio spectra. Now that C handra data are in the archive, we have additionally studied the inner jet as well as being able to resolve the  $25^{00}$  knot, so that we separate X -ray components by their morphology, and then test the various em ission models.

The following sections describe the observations, data reduction, and photom etry (2), the radio results (3), the X-ray emission processes (4), and an evaluation of each detected X-ray feature (5). We label the knots in the jet which are of primary interest, by their distance from the core in arcseconds (e.g. 'k4', 'k25'). We take the distance to 3C 120 to be 140 M pc ( $H_o$ =71 km s<sup>1</sup> M pc<sup>1</sup>; Spergelet al. 2003) so that 1" corresponds to 0.64 kpc. We use the standard de nition of spectral index, : ux density, S /

- The observations and reduction procedures
- 2.1. The Radio Data

D etails of the observations used in this paper are presented in Table 1. A comprehensive set of VLA and VLBI in ages between 1.4 and 15 G H z w as published by W alker, B enson, and U nw in (1987). Here we use the 1983 O ctober 2 VLA Acon guration in age from that paper as the source of our 1.4 G H z ux densities. An extrem ely high dynam ic range in age of 3C 120 at 5 G H z w as produced by W alker (1997) as a byproduct of an e ort to measure the proper motion of the  $4^{\circ\circ}$  knot. W e use that in age, which is actually the weighted sum of 3 in ages widely separated in tim e (no variability in the jet was detected), to measure our 5 G H z ux densities.

To obtain good in ages at appropriate resolution at higher frequencies, we observed 3C120 at 14.9 G Hz using the VLA B-con guration (second largest) and at 22.2 and 43.3 G Hz using the VLA D-con guration (sm allest). Both 22.2 and 43.3 G Hz were observed with subarrays because 43.3 G Hz was only available on 12 antennas. The 22.2 G Hz in age has inadequate resolution for use in this project and will not be presented. Reference pointing was used to keep 3C120 positioned properly in the VLA prim ary beam s at 43 G Hz. Because we were interested in the knot o set by  $25^{\circ0}$  from the core, and the VLA prim ary antenna beam is only about an arcm inute FW HM at 43 GHz, we o set the pointing position by  $12^{00}$  in the direction of the knot at that frequency. For all in ages, baseline based corrections for closure errors were made using scans on the bright, com pact source 3C 84. The procedure is described in the above references.

All of the VLA im ages were calibrated following the norm alprocedures as described in the VLA Calibrator M anual. For all observations, the VLA continuum system was used with 50 M Hz in each of two polarizations in each of two channels (200 M Hz total). All data reduction was done using the NRAO A stronom ical Im aging P rocessing System (A IPS). A prim ary beam correction wasm ade for the 15 and 43 G Hz im ages. At 43 G Hz, that correction took into account the o set pointing position.

The measured ux densities for each of the features of interest are presented in Table 2 along with the optical and X-ray values. We estimate that the errors in our measured ux densities of strong features are 10, 5, 6, and 10% at 1.4, 5, 15, and 43 GHz, respectively. To this we add the in age noise times the square root of the number of beam areas over which a given ux density is m easured. The radio ux densities were m easured in exactly the same boxes from each of the radio im ages although these rectangles, aligned with RA and DEC, are not identical to those used for optical and X-ray photom etry. This is not thought to be a signi cant source of error since we ensured that the sam e features were being m easured on the basis of radio overlays.

The resolutions of the images fallbetween  $0.36^{00}$ and about  $2^{00}$ . Norm ally it is a poor idea to use interferom etric in ages with unm atched resolutions for the measurement of ux densities from which spectra are to be derived. But in this case, there is no large scale structure in the regions of interest that is likely to show up in some images and not in others. The high resolution im ages are of high quality so all the ux density that would be seen at lower resolution should be present. As a result, we do not consider the m ism atched resolutions to be a signi cant lim itation. In fact, the norm al effect of such a m ism atch would be to have too little ux density measured in the higher resolution im ages. An examination of the spectra shows that, if anything, the opposite situation exists here.

# 2.2. The optical data

W e retrieved an H ST in age of 3C 120 from the W FPC 2 Associations website. The selected le was taken from the F 675W lter with a pivot wavelength of 6717 A.W e registered the data so as to align the optical and radio peak intensities  $(0.72^{00} \text{ shift})$ . For the H ST data, the peak position was found by the intersection of the di raction spikes. A fler constructing a ux map (g. 1) we m easured the features of interest: k4, k7, s2, s3, and k25 inner. The other parts of k25 were o the edge of the im age, as was k80.

Uncertainties were estimated by using the same regions on the events in age. For both sets ofm easurements we used 'funtools'<sup>2</sup>. For k4, which lies on a relatively bright optical spiral like feature, we used a circular aperture signicantly smaller than that used in the radio and X -ray bands ( $r= 0.273^{(0)}$ ), chosen so as to include all of the obvious optical emission of k4. For the background, we chose the same size circle, positioned just to the N orth of k4 where the brightness was judged to be about the same as that surrounding k4. N ote how ever, that this subjective choice means that the actual uncertainty in the ux of k4 is larger than the statistical error quoted in Table 2.

For the remaining features we used the same regions as de ned for the X-ray photom etry, except that we used a di erent background region for k25inner since the original background region was o the image. The resulting ux densities are given in Table 2.

For k25in which is close to the edge of the image, we found -58 counts in the source region (the background was oversubtracted), but + 40 and + 58 counts in two equally sized rectangles above and below the source. Thus we take to be about the measured ux value:  $0.34 \ 10^{29}$  (cgs). For an upper limit, we take 3 or 1 Jy.

# 2.3. The X-ray Data

The archival C handra data analyzed consist of an AC IS-S imaging observation (obsid 1613) and the zeroth order of an HETG observation (obsid 3015). The detectors are described in the CHAN - DRA P roposer's O bservatory G uide<sup>3</sup> and the param eters of the observations are given in Table 3.

The imaging data were badly saturated at the core position, whereas the zero order image of the HETG data did not su er from this problem : the so called \level 1" event le had a peak intensity pixel of 4940 counts (0.086 c/s, or 0.28 c/fram e-tim e). Thus there is som e pileup, but no saturation. The imaging observation also was done with a roll angle that m ade the readout streak fall on the inner jet (out to k4 and k7), but k25 was not a ected. For these reasons, m ost of our results are based on the zero order image.

O urbasic data reduction follows the procedures we used for M 87 data (Harris et al. 2003). A ffer checking the background for particle ares (our prim ary dataset was clean, and we decided not to exorcise the high countrate segment of 1613), we rem oved pixel random ization and registered the event le so that the x-ray and radio cores were aligned. This process for 1613 was problem atic since the location of the X-ray core had to be estimated from the readout streak in one dimension, and for the orthogonal direction, an alignment of a circle with the wings of the central count distribution.

W e then created 1024 arrays binned into 1/10 native AC IS pixel for each of four energy bands; constructed m onochrom atic exposure m aps for an energy near the center of each band; and produced ' ux m aps' with units of ergs cm<sup>2</sup> s<sup>1</sup> by dividing the data by the exposure m ap and multiplying by the nom inal h of the band. The exposure m aps were generated using the time dependent e ective area les which deal with the buildup of contam ination on the AC IS detector. An example of a ux m ap is shown in g.3 with the regions overlayed.

F lux m easurem ents consisted of choosing appropriate regions for jet features and corresponding backgrounds (see table 4 and g.3); m easuring the ux; and correcting for the m ean energy (via the sam e source region, but operating on the event

le). Since the ux maps were constructed with a multiplicative factor of h nom in al, the correction factor of  $< E > = E_{nom in al}$  for each m easurem ent is required to recover the actual ux within the band. In Table 5 we give the counts and ux for each jet component of interest.

Conversion of ux to ux density was done for an assumed power law with = 1. A lthough this

<sup>&</sup>lt;sup>2</sup> (http://hea-www.harvard.edu/RD/funtools/)

<sup>&</sup>lt;sup>3</sup>Rev.6.0, December 2003. http://cxc.harvard.edu

procedure is not strictly correct, the error introduced by using an incorrect value of should be of second order since the bands were only a few keV wide. In general, we did not have enough counts to obtain an accurate value of  $_x$ , and for the brighter features such as k4,  $_x$  is not far from 1.

Quoted uncertainties in uxes and ux densities are purely statistical. They were derived by using the source and background regions on count m aps (i.e. before dividing by the exposure m aps). The fractional error in each ux was then carried forward to the ux densities.

Since most features do not have su cient counts to provide an accurate estimate of the spectral parameters using the normal tting procedures in the X-ray band, we use our measurements of ux densities (Table 2) to construct spectra (see sec. 5 for examples). Except for the uncertainty in the absorption correction for the softest band, we believe that our technique of deriving spectral parameters from the ux maps is reliable. For the case of k4 there are su cient counts to allow the usual X-ray method of spectral tting. We did that with Ciao/Sherpa and nd agreement between the two methods, both in values of the amplitude and x.

# 3. Results from the new radio data

We planned and obtained our new VLA radio data on the basis of the RO SAT observations of 3C 120. At that time a persistent question was W hy are a few knots and hotspots in radio jets detected with current X -ray sensitivities while the vast maprity are not?" We now know that this question has been mostly answered because of the large num ber of jet detections with Chandra<sup>4</sup> : with sufcient sensitivity and angular resolution, X-ray detections of radio jets are quite common and no longer drive us to search for 'special conditions'. However, it would still be useful to identify any peculiarities in shock regions containing electrons which produce signi cant X -ray em ission. For k25 in particular, the di culty of nding a reasonable em ission mechanism for the X-rays led HHSSV to suggest the possibility of a atter spectrum (than deduced from the radio data) population of relativistic electrons extending up to Lorentz energies of  $10^7$ .

For these reasons, we undertook high frequency VLA observations to better determ ine the radio spectrum and to evaluate the behavior of the sharp gradients in radio surface brightness at two locations in k25. Both of these parameters m ay be observational signatures of physical attributes of shocks.

#### 3.1. Surface B rightness gradients

W e used the A IPS toolN INER to measure the rst derivative of the radio brightness. In order to measure comparative gradients at dierent frequencies, maps were convolved to the largest beam size. However, when comparing dierent features on the same map, this convolution was not necessary.

For all comparisons we computed the normalized gradient by dividing the actual value of Jy/beam /pix by the intensity of the peak brightness of the feature being m easured. When comparing m aps with di erent pixel size, we also divided by the factor arcsec/pixel. In addition to measuring the gradient for the outer edge of k25, we m easured gradients for the core (which serves to de ne the value for an unresolved source), k4, and the leading edge of k25 (which was undetected in our ROSAT data but is clearly present in the Chandra in ages).

# 3.2. Flux densities for spectra

To compare the spectra of the inner and outer shocks of k25, we measured ux densities on the Walker et al. (1987) maps constructed to have the same beam size of 1.25" FW HM (1.4, 5, and 15 GHz) and then repeated the measurements on m aps sm oothed to m atch the resolution of the 43 GHz map  $(2.2^{00} \ 1.6^{00})$  in order to include the 43 GHz ux density. We de ned a rectangle of  $3^{00}$  2:4<sup>00</sup> (RA DEC) for the internal (southern) shock and  $2.4^{00}$   $3^{00}$  for the outer (western) shock. A Ithough these integration areas are sm all, we corrected for background level by running IMEAN on a nearby region  $15.6^{\circ\circ}$   $132^{\circ\circ}$ . These corrections were < 6% in all cases. The results are given in Table 2. The resulting spectral indices are  $_{out} = 0.66 \ 0.03 \text{ and } _{in} = 0.74 \ 0.06$ .

<sup>&</sup>lt;sup>4</sup> http://hea-www.harvard.edu/XJET/

# 3.3. Evaluation of R adio R esults

There is no clear evidence for a signi cant difference in the spectral indices of the two parts of k25. The values are very close to each other and there are no data that dem and a spectral break between 1.4 and 43 G H z. The outer shock structure is brighter than the inner feature at all frequencies we have measured.

O ur attem pts to m easure the m agnitudes of the gradients in the surface brightness have been less than rewarding. To make progress with this approach, it would be necessary to have a resolution and sensitivity at least equal to those of g. 3. O ur only indicative conclusion is that the ratio of the norm alized gradients of the outer to the inner shock structures is greater than one at and below 5 G H z, and less than one above 10 G H z. H ow ever, the beam sizes em ployed even for this com parison are judged to be too large for any reasonable accuracy.

W e conclude that there is very little or no difference between the two shocks (except for the total ux density) and neither is discernibly exotic. G iven the ever increasing number of X -ray detections of radio jets, we take the absence of evidence for peculiarity in these putative shock features to be consistent with the idea that X -ray emission is relatively common for radio jets with a reasonable velocity towards us (i.e. two-sided X -ray jets are rare; we cannot cite even a single convincing example).

#### 4. X -ray Em ission Processes

In this section we discuss general aspects of the possible emission processes and for each jet feature tabulate the key parameters for each process. In section 5 we com pare the processes for each feature in turn.

There are two basic problems: how to estimate the emitting volume (which in uences the calculated value of the equipartition m agnetic eld strength) and what constraints m ight there be on the D oppler beam ing factor, (which governs the IC calculations). In general, we have elected to use the largest reasonable volume, with the X-ray and highest resolution radio m aps as a guide. Table 4 provides the dimensions and shapes of our assum ptions. In some or possibly all cases the actual volumes m ay be sm aller than that assumed, and that would lead to larger equipartition magnetic eld strengths for synchrotron models or higher electron densities for therm albrem sstrahlung models.

To accom m odate relativistic beam ing from bulk jet uid velocities ( =  $(1 \ ^2)^{1=2}$ ; = v/c), we will derive synchrotron parameters for a few representative values of = 1=  $(1 \ \cos)$  ( is the angle between the jet direction and the line of sight). For therm alm odels we assume there is no beam ing and for IC /CM B m odels we solve for the characteristic value of for which = .

# 4.1. Synchrotron m odels

W e found no evidence that the outer shock in k25 is discernibly exotic in the radio (section 3), but from the X-ray morphology of k25, it is evident that the X-ray and radio em issions are co-spatial. This dem onstrates that the suggestion of HHSSV, that the X-ray em ission m ight originate in a much smaller volum e than the radio em ission, is not supported by the new data.

The synchrotron parameters listed in Table 6 are typical for FRI jets. With beaming, the intrinsic power of the source drops. Since we posit no change in emitting volume, the eld drops as well. The hal ife for electrons responsible for 2 keV emission  $(\frac{1}{2})$  increases as the eld drops, but eventually the IC losses dominate and so  $\frac{1}{2}$  decreases. Since we have chosen to use the largest allow able emitting volume in each case, the actual value of the magnetic eld strength may be larger than indicated, and this would decrease  $\frac{1}{2}$ .

The classical objection to the synchrotron m odel arises for those features which have an obvious in ection in their spectra to accommodate a atter X-ray spectrum than would be anticipated by the segment connecting to optical or radio data. Two possible escapes from this objection are: (a) the existence of a distinct spectral component which is characterized by a atter electron distribution so that its emission is below detection limits at frequencies below the X-ray band (as suggested for k25 by HHSSV), and (b) a scheme that leads to a 'pileup' of electrons near the cuto energy (e.g. Dermer & Atoyan 2002).

# 4.2. Beam ing models

Inverse Compton emission from scattering on CMB photons by jet features with relativistic beam ing has been suggested as a method to increase the energy produced via the IC channel relative to that in the synchrotron channel (Tavechhio et al. 2000; Celotti et al. 2001). Harris and K raw czynski (2002) calculated beam ing param eters in plied from the 3C 120 RO SAT data (HHSSV) if the jet components are in equipartition (energy density, u (particles) u (B)). With our revised values for the spectral indices and ux densities for several other features, we have recalculated the beam ing param eters (Table 7).

The beam ing param eters given in Table 7 are not entirely unreasonable if the of the jet does not decrease signi cantly out to k25. However, as noted in the table, for several features the spectral index observed in the X-ray band disagrees with that expected from the radio spectrum . A dditionally, the sm all angles of the jet to the line of sight (of order 5 ) for m ost of the knots, leads one to infer a large projection e ect so that the physical size of the jet becom es rather long for an FRI.Furtherm ore, the beam ing param eters required (;; ) are m ore extrem e than those generally applied to the pc scale jet (G om ez et al. 2000; W alker et al. 1987).

A more general problem for the IC /CMB model is why the jet is not continuous in the X -rays once the emission 'starts' with the generation of particles at a shock. Since we know the jet uid continues downstream from a knot, the only way to quench the IC emission from the low energy electrons responsible for the X-rays, is to de ect the jet so that it is no longer pointing towards us, or to drastically reduce its velocity so that drops back close to one (in which case it would have to increase again at the next knot). For the synchrotron m odel, X -ray knots end quickly because of the large  $E^2$  losses to the extrem ely high energy electrons required, but the very low energy electrons responsible for X -rays in the IC /CM B m odel have a very long half-life and thus should propagate undim inished for sizable segments of the jet. These and other di culties of the IC /CM B m odels are discussed by Atoyan & Dermer (2004) and Stawarz et al. (2004). We believe the weight of the evidence argues against the IC/CMB model for the X-ray em ission of the 3C 120 jet.

## 4.3. Therm al B rem sstrahlung

G iven the at spectra found for a few features, we consider the possibility that the X-ray em ission comes from hot gas (c.f. the interpretation for NGC 6251 by M ack, K emp, & K lein 1997). For rough estimates of the parameters, we calculate the (uniform) density required to produce the observed X-ray ux. The results are given in Table 8.

The standard arguments against the thermal brem sstrahlung em ission process include:

if the X-ray em ission occupies the same volume as the radio em ission, there would be a departure from the  $^{2}$  law for the radio polarization position angle;

if the therm al region surrounds the em itting region, the predicted large rotation m easures (RM 's) are not seen; and

the therm all pressure of the em itting region generally exceeds the estim ate of the ambient gas pressure.

At least for the inner jet, the observed RM's are small (0 to -10 radians m<sup>2</sup>, W alker et al. 1987), so that the large RM's of Table 8 would appear to preclude therm al X -ray em ission. Reliable RM's are not available for k25, but from polarization position angles at two frequencies for the brighter parts of k25, it seems likely that RM 100 rad m<sup>2</sup>. No polarization data are available for k25new because of the very low radio brightness of this region so part, or conceivably all, of the X-ray em ission from k25new could be from hot gas.

# 5. Evaluation of Features

#### 5.1. The inner jet

As noted earlier, the X-ray in age from the archival AC IS data without the grating is heavily saturated at the core position so that it is in possible to study the innerm ost region of the jet. Even the HETG zero order in age has some degree of pileup, which means that the peak of the PSF is depressed relative to the wings. In spite of these di culties, we constructed a PSF with ChaRT (the Chandra ray tracer) and subtracted it from the im age. From a series of trial runs using di erent norm alizations, we are convinced that there is substantial excess emission to the SW of the nucleus (and here we refer to the position of the radio nucleus since we registered allour les so as to align the radio and X-ray peaks). This emission lies between  $0.5^{00}$  and  $0.9^{00}$  from the nucleus at a PA of 225; i.e. projected on the southerm edge of the radio jet. An example of one of the residual maps is shown in g. 4. G iven the m ism atch between the generated PSF and the data, we will not pursue the attributes of this feature here.

The X-ray emission from k4 is well resolved from the core region and aligns well with the corresponding optical and radio emissions (gs. 1 and 4). The values of the X-ray spectral index,  $_{\rm x}$ , from spectral tting (0.95 0.45) and from our 4 band uxes (0.90 0.20) agree within the errors and are also consistent with the radio to X-ray index,  $r_x = 0.90$ . Furthermore, the optical ux density falls on the power law de ned by the radio and X-ray data. Thus we favor a synchrotron explanation (perhaps with modest beaming, e.g. = 3) for the X-ray emission. If the X-rays were to be beam ed IC /CMB radiation, the apparently straight power law t from radio and optical to X-rays would be a coincidence and would have to be > 10 (Table 7). We note that while it appears that k4 lies in a spiral arm feature of the galaxy, we consider it highly probable that this is a coincidence. At the distance of 3C 120,  $4^{00}$  corresponds to 2.5 kpc, a reasonable distance between the nucleus and a feature de ned by stars. However, from VLBI studies of the superlum inal radio jet, the angle between the jet and our line of sight is of order 5, so k4 would be 29 kpc from the nucleus, and thus starlight would not be a viable source of photons for IC scattering.

The next knot, k7, has only 30 net counts so no spectral analysis was attempted. However, as shown in g.5, there is a signi cant di erence between  $_{rx}$ = 0.86 and  $_x$ = 2.4 0.6, as derived from ux densities for 3 of our energy bands. This sort of spectrum is consistent with a power law distribution of relativistic electrons with a high energy cuto (i.e. su ering E<sup>2</sup> energy losses); so again, we favor the synchrotron model. The 2 optical detection falls on the power law connecting the radio and X-ray data.

The next section of the jet ("s2" in g.3) is very weak in the X-rays and the follow ing section, "s3",

was not detected. However, the measurements are reported in Tables 2 and 5.

# 5.2. K nots k25 and k80

The X-ray emission of k25 continues to be a problem insofar as the standard emission models are concerned (HHSSV). The new data make matters worse since we now nd extended X-ray emission in a region of very low radio brightness. W e call this region 'new', and use the term s 'inner' and 'outer' for the two relatively bright radio edges which show large gradients in surface brightness: 'inner' refers to the upstream edge and 'outer' refers to the western edge. In g.6 we show the events com prising k25.

The only optical data we have is the upper lim it to k25in, and this ux density lies well above the power law connecting the radio and X-ray ux densities. Note however that the optical upper lim its quoted in HHSSV for the area then known to be associated with X-ray emission fall below the spectrum of the total emission, so the suspicion remains that it is not feasible to argue for a single power law from the radio to the X-ray. As shown in g.7, both the inner and outer regions have quite at X-ray spectra, whereas  $_{\rm X}$  for the 'new' region is consistent, within the errors, with rax.

A at power law ( <sub>x</sub> close to zero) is norm ally not considered feasible as a synchrotron spectrum associated with standard shock acceleration expectations, but what are the alternatives?

A side from the large beam ing factor required for the IC/CMB model ( >10, Table 7), one would need to explain why the emission dies away moving downstream since the very low energy electrons responsible for IC/CMB should have very long hal ives. The major unknowns for the IC/CMB model are the validity of extrapolating the electron spectrum down to the required low values of , and the actual value of the bulk relativistic velocity, i.e.

As shown in Table 8, therm all emission models encounter a number of problem s and thus we do not believe that therm all brem sstrahlung is a viable alternative to synchrotron emission for the brighter regions, k25in and k25out.

For the synchrotron m odel, aside from the requirem ent for a separate spectral com ponent with a at spectrum, we face the di culty of propagating high energy electrons from the acceleration sites (i.e. the inner and outer large-gradient radio features) to 11 the enclosed volum e ('new'). The required distance is of order  $3^{00}$  (1.9 kpc in projection); a light travel time of some 6000 years. This may be compared with typical lifetimes of the highest energy electrons against  $E^2$ losses (with equipartition magnetic eld strengths) of 200 years (Table 6). If the X-ray emission from k25new is thermal (the anonym ous referee suggested shocked gas), the travel time problem for high energy electrons would be moot.

An alternative explanation for the presence of high energy electrons in the 'new' region would be distributed acceleration (i.e. not directly associated with obvious shocks), as might be the case for reconnection scenarios (Birk & Lesch 2000, Birk, C rusius-W atzel, & Lesch 2001), or acceleration associated with turbulence along the jet edge (Staw arz et al. 2004, and references therein).

K not k80 (g.2) is of low radio brightness, but we nd associated X -ray emission. The morphological correspondence between radio and X -rays is not close as in the other 3 knots. In g.8 we show the relevant maps. Insofar as the detected structures are concerned, the situation of the X -rays being slightly displaced upstream of the bulk of the radio emission is very similar to that in the outer knot in the jet of PK S1127-145 (Siemiginow ska et al. 2002). The three values of  $S_x$  in Table 2 are consistent with  $_{rx}$  = 0.8. As for the inner knots, the results of section 4 leave little doubt that the x-rays come from synchrotron emission.

## 6. Conclusions

The X-ray emission from k25 remains a puzzle. The only ad-hoc models which we can suggest involve distinct spectral populations of relativistic electrons. For synchrotron emission, this would be a at spectrum component extending to electron Lorentz factors of  $= 10^7$  to  $10^8$  (see table 5 of HHSSV). Note how ever that \a distinct spectral population "need not indicate a separate shock region for its genesis. Derm er & Atoyan (2002) have described  $E^2$  loss conditions which can produce spectral hardening at high energies near the cuto , and Niemiec & O strow ski (2004) have produced num erical simulations for what they call

'realistic' m agnetic eld structures in relativistic shocks that can provide both relatively at particle distributions and even atter high energy tails for energies below the cuto . For IC/CMB with beam ing, the param eters derived above could be relaxed if an additional, steep spectrum distribution of electrons existed below = 2000 (see section 5.2 of H arris and K raw czynski 2002). A 60ks observation with C handra has been approved for A 0 6. W hen these data are available we should obtain m ore accurate X -ray spectral param eters.

A lthough one of us (DEH) has been a long and ardent supporter of the classical view: \If the spectrum is not concave downward, it is not synchrotron emission", it is now our view that there is strong evidence form arked deviations from this scenario. It should be remembered at this point, that there are many other knots in jets for which  $_{\rm X}$  <  $_{\rm ox}$ , (e.g. NGC 6251, Sambruna et al. 2004), and we believe it would be more fruitful to re-interpret these occurrences rather than to cite them as proof that synchrotron models are in possible.

The work at the C fA has been partially supported by NASA contract NAS8-39073 and NASA grant GO 3-4124A. We thank the creators of the W FPC 2 A speciations

(http://archive.stsciedu/hst/w fpc2) for providing public access to useful data. Tahir Yaqoob generously shared the zero order in age of his proprietary data (we joined him in a Chandra AO-4 proposal for an additional observation of 3C 120, but were unsuccessful). We thank C. Cheung for his careful reading of the manuscript and his resulting suggestions. We also thank the anonym ous referee for his useful comments. 7. R eferences

Atoyan, A. & Dermer, C. D. 2004, (submitted to ApJ) Birk, G.T. & Lesch, H. 2000 ApJ 530, L77 Birk, G.T., Crusius-W atzel, A.R., & Lesch, H. 2001 ApJ 559, 96 Celotti, A., Ghisellini, G., & Chiaberge, M. 2001 MNRAS 321, L1 Dermer, C.D. & Atoyan, A.M. 2002 ApJ 586, L81 Gomez, J. L., Marscher, A. P., Alberdi, A., Jorstad, S.G., & Garcia-Miro, C. 2000, Science 289,2317 Halpern, J.P. 1985 ApJ 290, 130. Harris, D.E. and Krawczynski, H. 2002, ApJ 565, 244 Harris, D.E., Hjorth, J., Sadun, A.C., Silverman, JD. & Vestergaard, M. 1999 ApJ 518, 213 (HHSSV) Harris, D.E., Biretta, J.A., Junor, W., Perlman, E.S., Sparks, W.B., and Wilson, A.S. 2003 ApJ 586,L41. Hjorth, J., Vestergaard, M., S rensen, A. N., & G rundahl, F. 1995 ApJ 452, L17 Mack, K. H., Kerp, J., & Klein, U. 1997 A&A 324, 870 Niemiec, J. & Ostrowski, M. 2004 ApJ (in press) Sam bruna, R M ., G liozzi, M ., D onato, D ., Tavecchio, F., Cheung, C.C., & Mushotzky, R.F. 2004, A&A,414,885 Siem iginowska, A., Bechtold, J., Aldcroft, T.L., Elvis, M., Harris, D.E., and Dobrzycki, A. 2002 ApJ 570, 543 Spergel, D. N. et al. 2003 ApJS 148, 175 Stawarz, L., Sikora, M., Ostrowski, M., & Begelman, M.C. 2004 ApJ (in press) http://arX iv.org/abs/astro-ph/0401356 Tavecchio, F., Maraschi, L., Sambruna, R.M., & Urry, C M . 2000 ApJL 544, L23-26 Walker, R.C., Benson, J.M., and Unwin, S.C. 1987 ApJ 316, 546 Walker, R.C. 1997 ApJ 488, 675 Walker, R.C., Benson, JM., Unwin, S.C., Lystrup, M B., Hunter, T R., Pilbratt, G., & Hardee, P E. 2001 ApJ 556, 756

Fig. 1. The HST in age used for photom etry, with radio contours overlaid. K not k4 is towards the right hand edge in this gure. The bright emission along the southern edge of the jet is part of the di raction spike from the nucleus. The radio contours are the same as used in g.3. The W FPC2 in age was taken with the F675W lter.

Fig. 2. Radio contours at 1.6 G Hz. This map from 1982 data (W alker et al. 1987) shows the section of the jet of interest, with several knots labelled. There are additional radio features on larger scales (see W alker et al. 1987). The beam width is  $3.5^{00}$  FW HM; the rst contour level is 0.25 m Jy/beam and successive contours increase by factors of 2.

Fig. 3. The 0.5 to 10 keV m ap of 3C 120 with overlayed regions for m easuring the ux. The X-ray data have been smoothed with a Gaussian of FW HM =  $0.5^{00}$ . D ashed regions are background regions. Radio contours from the VLA at 5 GHz (CLEAN beam width = 0.365") are overlayed. Contour levels start at 0.05 m Jy/beam and increase by factors of two. The regions are labelled by their distance from the core in arcsec for knots, and by 's2' and 's3' for sm ooth sections of the jet detected in the optical by H jorth et al. (1995).

Fig. 4. A residual X-ray m ap after subtraction of the PSF showing the excess emission to the southwest of the nucleus. The white contours are linear: 0.5, 1.0, 1.5, ... 4.0 counts per  $0.049^{00}$  pixel. An additional contour at -8 shows the location of the PSF subtraction. The black area in the center is negative, most likely caused by pileup suppressing the true peak intensity. The image was generated with an energy liter from 0.5 to 10 keV and smoothed with a Gaussian of FW HM =  $0.25^{00}$ . O verified are light m agenta radio contours from a 6cm VLA m ap with the lowest contour level at 0.2 m Jy/beam and successive contours increase by factors of two. The beam is circular with FW HM =  $0.365^{00}$ .



Fig. 5. Spectra for components of the inner jet. The solid line (circles) is for k4; the dashed line (crosses), k7; and the dotted line (squares), s2. The down triangle is a 2 (optical) measurement for k7. The optical values for k4 and s2 are so close to each other that their sym bols overlap. The left panel shows the radio to X-ray spectrum; the right panel shows the X-ray data.



Fig. 6. The events in the 0.5-10 keV band for k25. The radio contours are the same as used in g.3. The regions are shown as well.



Fig. 7. The spectra for the components of k25. The left panel shows the radio to X-ray spectrum; the right panel shows the X-ray data. The strongest feature is the outer region, crosses and solid line; the inner region is marked by circles and a dashed line; and the 'new' region is denoted by squares and a dotted line. The up-triangle is an upper lim it for k25 in from HST, and the two down-triangles are upper lim its reported by HHSSV for the total k25 optical em ission.

Fig. 8. The outer knot, k80. The X-ray in ages come from obsid 1613, consist of counts between 0.4 and 6 keV, and have been regrided to 1/4 native AC IS pixel. The left panel has been sm oothed with a G aussian of FW HM = 1°°; the right panel with a  $2^{\circ0}$  G aussian. R adio contours are overlayed. Those on the left show just the brighter, m ore compact levels from a 5 G H z V LA observation with the B array: with a beam size of  $1.36^{\circ0}$   $1:13^{\circ0}$  in PA = 44, and contours at 0.15, 0.20, and 0.25 m Jy/beam. The contours on the right panel are from a 1.5 G H z (B array) observation with a  $3.8^{\circ0}$  beam with the rst contour at 0.2m Jy/beam and successive contours increase by factors of two. There are between 6 and 9 counts in the two weak X-ray features upstream of k80. In both panels, the colors span a factor of 4 in X-ray brightness. The peak brightnesses are 0.038 (right) and 0.06 (left) counts per 0.123" pixel.



Fig. 9. The spectrum of k80. The left panel shows the radio to X -ray spectrum; the right panel shows the X-ray data.

		E	eam Size	Calibrators					
Frequency (GHz)	D ate (s)	МајАх ( <sup>00</sup> )	M in A x ( <sup>00</sup> )	PA (Deg)	Peak (Jy/beam)	mms (mJy)	3C 286 (Jy)	3C 48 (Jy)	
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	
1.45	1983 Oct.2	1.26	1.25	47	2.76	1.45	14.51		
4.86	Multiple <sup>a</sup>	0.365	0.365		3.	15 0.0	12 7.	55	
14.94	2000 Jan.4	0.54	0.51	24	3.36	0.033	3.43	1.80	
43.34	M ultiple <sup>b</sup>	2,21	1.60	-2	2.07	0.25	1.45	0.575	

Table 1 VLA Observations

 $^{\rm a}{\rm W}$  eighted average of in ages from 1987 Jul. 5, 1991 Aug. 29, and 1996 Nov. 2

<sup>b</sup> Im age m ade from com bined uv data from 1999 Apr. 17 and 1999 M ay 30

Note. The columns in the table give (1) the observing frequency, (2) the date(s) of the observations, the CLEAN restoring beam (3) major and (4) minor axis size (FW HM) and (5) position angle, (6) the peak ux density in the image (always on the 3C 120 core source which is highly variable), (7) the image o -source rms noise level, (8) the assumed ux density of 3C 286 used in the ux density calibration, and (9) the ux density of 3C 48 used in the calibration, when it was used.

Table 2 Flux Densities (ergs cm  $^2\,$  s  $^1$  Hz  $^1$  ) and Spectral Indices

Q uantity	k 4	k7	s2	s3	k25in	k25out	k25new	k80
S(1.5GHz)	1.4 0.1E-24	3.2 0.5E-25	1.5 0.4E-25	9.0 3.9E-26	1.2 0.4E-25	1.5 0.4E-25		6.0 0.6E-26
S (5G H z)	6.6 0.3E-25	1.6 0.1E-25	8.4 0.4E-26	5.1 0.3E-26	4.3 0.2E-26	6.9 0.4E-26	9.1 1.5E-27	
S (15G H z)	2.8 0.2E-25	7.2 0.5E-26	3.9 0.3E-26	2.6 0.2E-26	2.5 0.2E-26	3.5 0.3E-26		
S(43GHz)	1.2 0.1E-25	3.2 0.5E-26	1.6 0.5E-26	8.1 5.0E-27	0.9 0.5E-26	1.6 0.5E-26		
S(4.47E14Hz)	2.7 0.3E-29	0.9 0.4E-29	2.6 0.4E-29	1.0 0.1E-29				
S(2.42E17Hz)	1.0 0.2E-31	6.3 1.6E-32		0.8 0.6E-32	1.7 0.9E-32	6.0 1.5E-32	5.5 1.4E-32	2.4 0.6E-32
S(6.05E17Hz)	5.8 1.0E-32	0.5 0.4E-32	0.5 0.3E-32		1.3 0.4E-32	4.7 0.8E-32	2.5 0.6E-32	1.6 0.5E-32 <sup>a</sup>
S(9.67E17Hz)	3.8 0.8E-32	0.2 0.2E-32			1.4 0.5E-3	32 5.8 0.9E-32	2 2.4 0.5E-32	5.1 4.7E-33
S(1.69E18Hz)	2.7 1.1E-32		0.2 0.1E-32		1.5 0.7E-3	32 3.5 1.2E-32	2 1.0 0.6E-32	
				Spectral Index	es			
r	0.74 0.04	0.68 0.04	0.67 0.06	0.69 0.10	0.74 0.06	0.66 0.03		
rx	0.90	0.86	0.97		0.82	0.81	0.70	0.79
x	0.90 0.20 <sup>b</sup>	2.4 0.6	0.2 0.6		0.0 0.4	0.3 0.3	0.6 0.2	0.7 0.4

<sup>a</sup>This ux density refers to a frequency of 4.8E17 Hz (2 keV).

<sup>b</sup> The value in the table comes from the 4 band ux densities with the soft bands corrected for excess absorption (N h= 2.4 10<sup>21</sup> cm<sup>2</sup>). The galactic column density in this direction is 1.08 10<sup>21</sup> cm<sup>2</sup>. Sherpa spectral tting gives x = 0.95 0.45 (N h= (1.9 2.2) 10<sup>21</sup> cm<sup>2</sup>) for no background subtraction and 1.01 0.6 (N h= (2.4 3.1) 10<sup>21</sup> cm<sup>2</sup>) with background subtraction.

0 bsid	D ate	D etector	G rating	livetim e	CALDB
			<u>)</u>	(æc)	-
1613	2001-09-18	ACIS-235678	none	12,881	2.8
3015	2001-12-21	AC IS-456789	HETG	57 <b>,</b> 218	2.10

Table 3: Chandra observations

Note. Both observations were made with the standard CCD readout time of 3.2s.

Table 4: Speci cs of the regions and assum ed sizes for volum es

		R egion		Assum ed Volum es			
region	shape	RA (J2000)	DEC (J2000)	size <sup>a</sup>	shape	dim ensions <sup>b</sup>	pathlength
				( <sup>00</sup> )		( <sup>00</sup> )	(kpc)
k4	circle	04:33:10.843	+05:21:15.09	0.738	sphere	0.33	0.42
k7	rotbox	04:33:10.628	+ 05:21:15.44	2.165,1.427	sphere	0.33	0.42
s2	rotbox	04:33:10.450	+05:21:16.70	3.542,1.230	cyl.	0.33;3.0	0.42
ടാ	rotbox	04:33:10.232	+ 05:21:18.35	3.444,1.230	cyl.	0.33;2.8	0.42
k25in	rotbox	04:33:09.752	+05:21:25.14	2.706,1.107	cyl.	0.33;2.1	0.42
k25out	rotbox	04:33:09.571	+05:21:26.55	3 <b>.</b> 132,1.447	cyl.	0.6 <b>;</b> 1.8	0.76
k25new	circle	04:33:09.718	+05:21:27.31	1.427	sphere	1.43	1.82
k80	circle	04:33:07.937	+ 05:22:21:22	4.797	sphere	3.0	3.82

<sup>a</sup>A single entry denotes radius of a circle; two entries are sides of the rectangle.

 ${}^{\mathrm{b}}\mathrm{R}$  adius of sphere or cylinder. W hen a second entry is present, it is the length of the cylinder.

Note. | Columns to the left describe the regions used for photom etry except where noted in the text. The last 3 columns give the assumed geometry of the emitting volumes. The background regions are shown in g.3. All coordinates refer to the radio maps.

	Table 5									
X-ray	fluxes	(in	units of 10	14	erg cm	2	s <sup>1</sup> )	and	source	counts

Energy <sup>a</sup>	k4	k7	s2	ടി	k25in		k25out		k25new		k25tot		k80 <sup>b</sup>		
0.5-1.5	1.85 0.39	1.16 0.27		0.15	0.31	0.16	1.11	0.28	1.02	0.25	3.15	0.47	0.49	0.12	
1.5-3	3.35 0.57	0.26 0.22	0.27 0.16		0.77	0.25	2.72	0.49	1.43	0.36	6.03	0.72	0.50	0.17	
3–6	2.55 0.56	0.15 0.12		0.14	0.93	0.31	3.90	0.62	1.60	0.37	7.80	0.86	0.34	0.32	
6–10	2.35 0.96		0.16 0.1	0	1	.29 0	.65 3	.02 1	1.03 0	.90 0	.54 6	.10 1	.40		
0.5-10	10.1 0.91	1.42 0.36	0.43 0.26		3.30	0.99	10.7	1.07	4.95	0.64	23.08	1.62	1.33	0.39	
Source C ounts															
Total counts	124	34	14	7		27	ç	98		55	22	25	2	18	

 $^{\rm a}{\rm T}\,{\rm he}$  energy band is given in keV .

N et counts

 $^{b}$  T he m easurem ents for k80 com e from obsid 1615 and the 3 bands are 0.4-1.5keV, 1.5-3keV, and 3-6 keV; with the broad band being the sum of these 3.

7 0

					-	=	= 3 <sup>a</sup>	=	= 6 <sup>a</sup>		
R egion	B(1) <sup>b</sup> (G)	r	rx <sup>d</sup>	logL <sub>s</sub> (erg/s)	(years)	logL <sub>s</sub> (erg/s)	(years)	logL <sub>s</sub> (erg/s)	(years)		
k4	116	0.74	0.9	41.83	23	39.92	56	38.72	27		
k7	72	0.68	0.84	41.47	47	39.56	86	38.36	47		
s2	33	0.67	0.97	40.62	152	38.73	108	37.53	17		
k25inner	36	0.74	0.82	41.14	131	39.23	108	38.03	18		
k25outer	27	0.66	0.81	41.45	195	39.54	105	38.34	16		
k25new	9		0.	70 41.24	6	69 39.	33	74 38	.13 9	9	
k80	8.3		0.	79 41.14	l 7	62 39.	24	69 38	.03 9	9	

Table 6 Parameters for a Synchrotron M odel

<sup>a</sup> is the beam ing factor

<sup>b</sup>B (1) is the equipartition m agnetic eld strength for the case = 1. For other values of , it is  $B(1)/(H \operatorname{arris} \& K \operatorname{raw} \operatorname{czynski}, 2002)$ .

 $^{\rm c}~_{\rm r}$  is the radio spectral index.

 $^{\rm d}$   $_{\rm rx}$  is the spectral index connecting the radio to the X -ray bands.

 $e_{\frac{1}{2}}$  is the hal ife, as observed at the earth, for the electrons responsible for the observed 2keV em ission (eq B 7 of H arris & K raw czynski, 2002, assuming = ).

Note. The emission spectrum is assumed to cover the range E6 to 5E18 Hz in the observed frame.

R egion	a	B (1) <sup>b</sup>	R (1) <sup>c</sup>	d		R <sup>0e</sup>	D istance <sup>f</sup>
		G			(deg)		(kpc)
k4	0.74	116	0.0917	16	3.5	65	42
k7	0.68	72	0.0176	12	4.5	56	57
s2	0.67	33	0.0104	7.7	7.5	40	49
k25in	0.74	36	0.3967	13	4.5	352	184
k25out	0.66	27	0.1939	18	32	1840	269
k25new	0.70	10	1.3454	14	4.1	5060	215
k80	0.79	8.3	0.6289	53	10.5	137	281

Table 7: Param eters for an Inverse C om pton M odel with Bulk Relativistic Velocities

<sup>a</sup> is the spectral index for both the IC and synchrotron spectra. For most features, it is  $_{\rm r}$  from Table 2. For k25new and k80, it is  $_{\rm rx}$ .

 $^{\rm b}{\rm B}$  (1) is the equipartition m agnetic  $\,$  eld strength for no beam ing.

 $^{\rm c}R$  (1) is the ratio of am plitudes of the IC to synchrotron (observed) spectra.

 $^{\rm d} T \, he\,$  /  $\,$  column gives their values for  $\,$  =  $\,$  , and the relevant angle to the line of sight is  $\,$  .

 ${}^{e}R^{0}$  is the ratio of E  ${}^{2}$  losses in the jet fram e (IC and synchrotron losses).

 $^{\rm f}$  D istance' is the de-projected distance from the core for the particular feature (projected distance/sin ).

Note. For k7, the X-ray spectrum is much steeper than the radio spectrum so the IC/CMB model is not applicable. For both k25in and k25out, the actual X-ray spectrum is significantly atter than the radio spectrum, so also for these features, the IC/CMB beam ing model is not tenable.

Table 8: Therm al Brem sstrahlung Param eters

R egion	ne <sup>a</sup>	Cooling T in e	M ass	P ressure <sup>b</sup>	RM <sup>c</sup>
	(cm <sup>3</sup> )	(years)	(M)	(dyne am $^2$ )	(m <sup>2</sup> )
k4	6.6	1.5E5	6.3E 6	2.0E-8	22 <b>,</b> 450
k7	2.5	4.0E5	2.4E6	7.5E-9	8 <b>,</b> 500
s2	0.6	1.8E6	3.7E6	1.7E-9	2,040
k25inner	1.7	5.6E 5	8.0E6	5.6E-9	5 <b>,</b> 780
k25outer	1.9	5.2E 5	2.4E7	5.6E-9	11 <b>,</b> 450
k25new	0.5	1.9E 6	4.0E7	1.5E-9	7 <b>,</b> 520
k80	80.0	1.2E7	5.9E7	2.5E-10	2 <b>,</b> 500

<sup>&</sup>lt;sup>a</sup>The electron density is that necessary for a uniform plasm a with the volum e described in Table 4.

 $<sup>^{\</sup>mathrm{b}}\mathrm{T}\,\mathrm{he}\,\mathrm{pressure}\,\mathrm{values}\,\mathrm{assum}\,\mathrm{e}\,\mathrm{a}\,\mathrm{tem}\,\mathrm{perature}\,\mathrm{of}\,1\,\mathrm{keV}$  .

<sup>&</sup>lt;sup>c</sup>RM gives the predicted rotation measure for a eld of 10 G and a pathlength from table 4. RM (radians m<sup>2</sup>)=810 B (G) dL (kpc)  $n_e$  (cm<sup>3</sup>).

This figure "f1.jpg" is available in "jpg" format from:

This figure "f2.jpg" is available in "jpg" format from:

This figure "f3.jpg" is available in "jpg" format from:

This figure "f4.jpg" is available in "jpg" format from:

This figure "f8.jpg" is available in "jpg" format from: