Cosm ic Filam ents in Superclusters

Joel N. Bregman, Renato A. Dupke, and Eric D. Miller

Department of Astronomy, University of Michigan, Ann Arbor, MI 48109

jbregman@umich.edu, rdupke@umich.edu, milleric@umich.edu

ABSTRACT

Large-scale structure calculations show that modest overdensity laments will connect clusters of galaxies and these laments are reservoirs of baryons, mainly in gaseous form. To determ ine whether such laments exist, we have examined the UV absorption line properties of three AGNs projected behind possible laments in superclusters of galaxies; the AGNs lie within 3 M pc of the centerlines of loci connecting clusters. All three lines of sight show absorption in Ly , Ly , or/and 0 VI at redshifts within about 1300 km s¹ of the nearby galaxy clusters that would de ne the closest laments. For one AGN, the absorption line redshifts are close to the emission line redshift of the AGN, so we cannot rule out self-absorption for this object. These absorption line associations with superclusters are unlikely to have occurred by chance, a result consistent with the presence of cosm ic laments within superclusters.

Subject headings: galaxies: clusters: general | large-scale structure of universe | quasars: absorption lines

1. Introduction

Big-bang nucleosynthesis and quasar absorption line studies (at z 3) determ ine the same mass fraction of baryons in the universe: $_{\rm b} = 0.04h_{75}^2$ (e.g., Fukugita, Hogan, and Peebles 1998). It was expected that this gas would become the galaxies of today, but a census of stars and gas in galaxies in the local universe accounts for only about 20% of the baryonic mass. This is the \m issing" baryon problem that has drawn much attention, for which the leading solution is that the baryons have remained as dilute gas (e.g., Fukugita, Hogan, and Peebles 1998; C en and O striker 1999a; D av e et al. 2001). However, the Ly forest becomes

less dense in the local universe by about a factor of 3-4 com pared to the num ber of absorption system s at xed equivalent width at z = 3. This decrease in the baryons is not balanced by the increase in the stellarm ass, leading to the conclusion that a signi cant m ass of gas m ust be hot, above 10^5 K.

There is a prediction for the properties of the m issing baryons from hierarchicalm odels of structure form ation. The most visible large structures are clusters of galaxies, which are deep potential wells of high overdensity, but these clusters are relatively rare. The more common collapsed systems are groups of galaxies and the large lam ents that connect the groups and clusters. G roups of galaxies are weak em itters of X-rays, indicating gas at 0.3{ $2 \ 10^7$ K but with extended gaseous m asses that are very poorly known because their extent is broad and their surface brightness fades below the detectability of X-ray instrum ents at a fraction of the virial radius.

It is the laments that contain most of the volume in collapsed structures and here the temperature is generally lower, typically $10^5 \{10^7 \text{ K}, \text{according to theoretical calculations} (e.g., Cen et al. 1995).$ There have been no convincing detections of these laments in emission, but there is good evidence that gas is present at these temperatures, as O VI absorption is detected by both HST and FUSE in observations toward bright AGNs (e.g., Savage et al. 2002). G as bearing O VI occurs when T $10^{5.5}$ K, a temperature expected in laments, and because this doublet is strong and oxygen is the most abundant heavy element, this is an excellent tracer of such gas. These observations show that O VI absorption is common, dN = dz 20 for W > 50 m A (Tripp and Savage 2000). It is predicted that the O VI absorption lies in a network of laments with modest overdensities of 10{40} (Cen et al. 2001) as well as in the galaxy groups (Hellsten et al. 1998). When O VI is present, a sm all fraction of hydrogen is still neutral, so the Lyman series lines of H I accompany O VI as a tracer.

The goal of our e ort is to identify the likely locations of these laments and then to determ inewhether OVI and H I absorption is associated with them. Theoretical simulations guide us to the most likely location of laments (Evrard et al. 2002), which shows that they generally connect rich clusters, so the objects with the highest frequency of laments (and the largest covering factor) are the superclusters of galaxies. These are generally several galaxy clusters near each other on the sky and with only modest velocity di erences, so they are physically close in space. In this study, we exam ine three AGNs that have already been observed in the ultraviolet either by the Far Ultraviolet Spectroscopic Explorer (FUSE), or by the Hubble Space Telescope (HST), and each of these show absorption near the redshift of the supercluster behind which they are projected. We analyze the properties of these absorption features and discuss the implication for the detection of the \cosm ic web" of

lam ents.

2. Sam ple Selection and Results

Practical aspects of observing, involving the identi cation of suitably bright AGNs, drive us toward using the nearby superclusters as the optimum absorption sites. Nearby superclusters subtend a large solid angle (with known redshift), which is very helpful for nding su ciently bright background AGNs. Also, most of the brightest AGNs are nearby, typically Seyfert galaxies at z < 0.1, so the superclusters must be at lower redshift, which are the closest superclusters in a near-linear fashion, so the background AGN should lie close to loci connecting clusters. The tangential dimensions of the laments (at an overdensity of 10) are predicted to be the same size as the virial radii of the galaxy clusters (3 M pc; C en and O striker 1999b), so background AGNs should be within this distance of the line connecting clusters, although, as this is a theoretical prediction, we will include AGNs that are not so ideally aligned.

Superclusters have been cataloged out to a distance of z = 0.12 (E inasto et al. 1997). They have catalogued 220 superclusters using a friends-of-friends algorithm with a neighborhood radius of 24h¹ M pc. W e visually identi ed the most likely locations of laments connecting galaxy clusters within superclusters and searched for UV-bright AGNs projected behind them, using the updated V eron-Cetty & V eron (2001) Catalog and at low Galactic E (B V) as given by Schlegel, Finkbeiner, and D avis (1998). About ten objects satisfy our criteria and are bright enough to be observed by HST or FUSE and we have an observing program to obtain spectra of these targets. However, three targets have already been observed by FUSE and HST (PHL 1811, PG 1402+261 and TON S180), which we report upon here.

The locations of these three targets, relative to the foreground superclusters, are shown in F igure 1, where we indicate the position of clusters of galaxies and the size of the crosses represent the richness of the cluster as de ned in D ekel & O striker (1999). The value for the cluster with m axim um richness is given and the values for the other clusters can be estim ated by the size of the crosses (in unit steps). A loo shown are the AGNs (empty diam ond) and the estimated 2-D location of the relevant laments, where the thickness of the laments is approximately $3h^{-1}$ Mpc. We eliminated m isclassi ed clusters from the E inasto et al. (1997) catalog (those with velocities signi cantly di erent from the rest of the system s in the supercluster; they comprise less than 10% of the clusters). In Table 1 we list the name of the AGN, redshift, visual magnitude, the supercluster projected in the line-of-sight, its

redshift, the full redshift range of the supercluster, the standard deviation of one galaxy cluster about the supercluster redshift, and the estim ated projected distance from the AGN to the nearest candidate lament.

The FUSE observations were reduced in-house with version 2.2.1 of the pipeline and reprocessed most recently with version 2.4 of CALFUSE, but there was no di erence in the resulting spectra. For analysis purposes, the FUSE spectra were binned by a factor of 5 (0.034 A width) and then smoothed with a seven point FFT liter, yielding a resolution of about 4,000 (we exam ined the spectra at full resolution as well before adopting this procedure). For the HST data, we used the standard data products provided by the MAST archive.

PG1402+261

This Seyfert 1 galaxy, at an emission line redshift of 0.164, lies behind the Bootes supercluster, which is composed of 11 galaxy clusters in a redshift range of 0.059-0.079 (there is an additional cluster at z = 0.0948 that could be considered an outlier; Figure 1a). This object lies at high Galactic latitude (73.4), with a low extinction (E (B V) = 0.016; Schlegel et al. 1998) and a Galactic neutral hydrogen column of 1.50 10^{20} cm² (Lockman et al. 2002, Wakker et al. 2003). There are spectra available for the 900-1180 A region (FUSE, a 8 ksec observation taken in June 2000, using the LW RS aperture) as well as for the 1200-1600 A region (HST, with the G130H, G190H, and G270H on the FOS in August 1996), both of which have been discussed in the literature (Bechtold et al. 2002; Wakker et al. 2003).

The FUSE spectrum shows all the strong Galactic atom ic lines, including the high ionization systems (O V I) and the low ionization systems (S III, Si II, C II, C III, O I, A r I, Fe II, Fe III, and N I). These Galactic lines are blueshifted by about 40 km s¹, which is probably due to the target not being located in the center of the slit, but this shift has

Target Name	Z _{AGN}	m _v	Supercluster N am e	z_{SC}	z_{sc}	(z _{SC})	R _{proj}
PG 1402+261	0.164	15.5	Bootes	0.070	0.0151	0.007	< 1 M pc
Ton S180	0.062	14.4	Pisces-Cetus	0.060	0.0187	0.0059	< 3 M pc
PHL 1811	0.192	13.9	Aquarius-Cetus	0.058	0.0089	0.0028	< 1.5 M pc
PHL 1811	0.192	13.9	Aquarius B	0.084	0.0043	0.0034	2 M pc

Table 1. Supercluster and AGN Properties



Fig. 1. Spatial conguration of the background AGNs (open diamonds) projected onto the superclusters. The crosses are galaxy clusters (many with Abell names), where the size of the cross indicates the relative population richness. The linear grey stripes are connections between galaxy clusters that are possible laments projected near the background AGN, based upon three-dimensional proximity and cluster richness; these laments are 3 Mpc wide. We also indicate the redshift of the clusters in parenthesis (in units of 10²), the average redshift of the supercluster, and the standard deviation of cluster redshifts in the supercluster.

no in pact on our analysis. There is very little molecular hydrogen evident, with an upper limit of about $10^{17.5}$ cm². In our analysis of the spectrum, we nd no unidentied lines in the FUSE spectral range, aside from the expected number of 2{3 features. In the HST spectrum, obtained with the Faint Object Spectrograph, using the G130H spectral element (resolution of 1300, or 230 km s¹), there are a few unidentied lines, such as those noted by Bechtold et al. (2002) at 1383.11 A and 1405.36 A. These are most likely Ly absorption features at redshifts of 0.137 and 0.156, respectively.

In addition to the features cataloged by Bechtold et al. (2002), there is a weak feature at 1285.1 A that does not correspond to any Galactic features. It is most likely a Ly absorption line at a redshift of 0.0571 and with an equivalent width of 150-200 m A and an uncertainty of about 50 m A (Figure 2). Con mation of this feature through the detection of some other Lym an series lines is a commonly used tool when the wavelength coverage is broad. Unfortunately, nearly all of the strong lines lie in places in which it is di cult to detect a line uniquely. That is, the Ly line would lie at 1084.3 A, but it would be confused with the strong Galactic N II 1084.0 line, so a unique iden tication is not possible. A lso, the feature does not fall in the Lifla channel, but the Sicla channel, where the S/N is signi cantly lower. We estimate that the 3 upper limit to the Ly line is about 0.25 A, which is much larger than the expected value of 0.03 A if Ly were optically thin. The Ly line would be at 1028.1 A, in the shoulder of the Galactic Ly absorption line, but also coincident with an airglow line, that could ll in a weak absorption feature. The line is not detected, with a 3 upper limit of 0.1 A, which is consistent with the predicted equivalent width of 0.01 A if the Lym an lines are thin. The Ly line, at 1004.0 A, is not detected but the 3 upper lim it, 0.2 A, is considerably larger than the expected line strength. We searched for several atom ic lines but detected none (e.g., C III 977, C II 1036). For the low ionization lines, this is the expected result, as the Lym an lines are always stronger, but O VI can be stronger than the Lym an lines at high tem peratures (5 10^5 K). The stronger OVI line (1031.9 A) would fall at 1090.9 A, and no line is seen, with a 3 upper lim it of 0.1 A equivalent width.

If the line at 1285.1 A is a Ly absorption line at z = 0.0571, it is close to the redshift of the supercluster. O f the clusters in the Bootes supercluster with known redshifts, the one closest to PG 1402+261 is A bell 1831 (2.5 away, z = 0.0615), which has the sm allest redshift di erence with respect to the absorption system, a di erence of 1300 km s¹, com parable to the velocity dispersion of the cluster. Consequently, it is possible that this absorption line system is associated with a supercluster lam ent that connects to A bell 1831, as it lies beyond the virial radius of the cluster (a radius of about 0.5). There are other galaxy clusters nearby, but they do not have measured redshifts, such as A bell 1861, which is closer on the sky to PG 1402+261 than A bell 1831. Two galaxy clusters lie within 1.5



Fig. 2. | HST /FOS spectrum of PG 1402+261, showing the strong Galactic lines of SiII and O I, as well as an unidentied absorption feature, which is most likely Ly near the redshift of the Bootes supercluster. The dotted line indicates the region of expected absorption within

1300 km s 1 of the redshift of the nearest cluster.

of PG 1402+261 and are presumably more distant than the Bootes cluster: A bell 1797 and A bell 1982. The redshifts of these should be measured in order to determ ine whether some of the other unidentied lines could be Ly absorption related to the clusters.

Ton S180

This Seyfert 1.2 galaxy lies near the South G alactic Pole (l, b = 139, -85), it has relatively little G alactic absorption (E (B-V) = 0.014), and it has a redshift of 0.06198. In projection, it lies behind the P isces-C etus supercluster, near two lines of clusters that connect to A bell 133 (z = 0.0566; 82⁰ away). A long the m ore northern system are A bell 114 (z = 0.0587; the closest cluster, 66⁰ away), A bell 2824 (z = 0.049; 136⁰ away), A bell 86 (z = 0.061; 209⁰ away), and A bell 74 (z = 0.0651; 256⁰ away). Toward the south, A bell 133 m ay connect to A bell 2800 (z = 0.0636; 311⁰ away; F igure 1b). If a lam ent connects these clusters both spatially and in velocity, then this Seyfert galaxy is about at the redshift of the lam ent, of which it m ay be part. Being part of a lam ent increases the likelihood that our line of sight toward it would pass through overdense regions, but it also m akes it in possible to distinguish between absorption intrinsic to the AGN or host galaxy and absorption by the lam ent.

The FUSE data were obtained in December 1999, using the LWRS aperture, and with an exposure time of 132 ksec (Shull et al. 2000; Wakker et al. 2003). There were several ultraviolet spectra taken with STIS on HST, using the G140L (January 2000 observation for 1260 seconds) and using the G140M covering the 1194–1300 A region (taken July 1999) and discussed by Shull et al. (2000), Turner et al. (2002), and Penton et al. (2004).

The system has the usual set of G alactic atom ic absorption lines, including O VI (W akker et al. 2003), but it has very little H_2 (< 10^{17} cm²). There are several low redshift Ly and Ly \forest" lines (Shull et al. 2000), but in addition, there are absorption features near z = 0.06 not previously reported. At least three absorption features are seen in both of the O VI lines in the redshift range 0.0614 - 0.0625. These redshifts probably need a sm all correction upward by about 0.00018 since the G alactic atom ic lines are shifted by that am ount (e.g., the nearby Fe II line, and others), an e lines and 0.0625 (Figure 3). The z = 0.0616 system appears to be two lines that are blended together, but the S/N is insu cient to be certain. The other lines appear single and have the same line width, a FW HM of 150 40 mA.

These absorption systems also appear in the ST IS medium resolution spectrum of this object with a resolution of about 1000 (Turner et al. 2002), where three Ly absorption systems are seen at z = 0.0615, 0.0620, and 0.0623, with equivalent widths of 80 m A, 30 m A, and 100 m A. If these lines are optically thin, the Ly lines would be 13 m A, 5 m A,



Fig. 3. FUSE spectrum of Ton S180 shows three O VI absorption line systems near the redshift of the Pisces-Cetus supercluster as well as the emission line redshift of Ton S180. The lines have a FW HM of 0.15 A, which is close to the therm all oppler width for gas at 10^5 K (0.11 A). The dotted line indicates the region of expected absorption within 1300 km s¹ of the redshift of the nearest cluster.

and 16 mA, but the 1 uncertainty for a line in this part of the Lif2a spectrum is about 30 mA, so the failure to detect the Ly line is suggestive that the lines are of low optical depth. These Lya lines are similar to but not exactly at the same redshift, with di erences up to 60 km s¹ for the highest and lowest redshift system s.

The origin of these four redshift system s is am biguous because we cannot distinguish between absorption intrinsic to the Seyfert and hot material within a cosm ological lam ent. If the absorption is associated with a lam ent from a cluster, the lines are 800-1050 km s¹ away from the redshift of the nearest cluster (Abell 114) and may be associated with this system (the line of sight passes about two virial radii from the cluster center). The properties of the absorption system s are consistent with a hot but quiescent medium. The equivalent widths of the 0 VI lines exceed that of the Ly lines, typically by factors of 2-3, which can occur in collisionally ionized gas in the 2{10 10^5 K range (Tripp and Savage 2000), the type of material predicted to lie in lam ents. The line widths of the 0 VI lines are close to the therm al width of oxygen at 3 10^5 K (110 m A), so the gas does not lie in a turbulent medium, as the sound speed of a gas at this tem perature is about 80 km s¹ and if the turbulent speed were equal to the sound speed, it would lead to a FW HM of 500 m A in all lines.

PHL 1811 (J2155-0922)

This narrow-line Seyfert 1 galaxy, at a redshift of 0.192, is a a very bright object, although at a galactic latitude of -44, there is more Galactic extinction along this line of sight, with E(B-V) = 0.046. It lies behind the Aquarius B supercluster as well as the Aquarius-Cetus supercluster, which are distinguished by their redshifts. In the Aquarius B supercluster, the cluster closest to PHL 1811 is Abell 2402 (50°; z = 0.0809), followed by Abell 2410 (109° ; z = 0.0809), Abell 2376 (136° ; z = 0.0896), Abell 2377 (139° ; z = 0.0808), and Abell 2400 (126° ; z = 0.0882). If lam ents connect the clusters and pass in front of PHL 1811, the m ost natural connections would be between Abell 2376/2377 and Abell 2402/2410 (Figure 1c). It has been observed with HST on several occassions with ST IS (with the G 140L in December 2001, and in a current program with the E140M), and with FUSE (in June 2001, and for longer exposures in a current program).

This system has been studied exhaustively by Jenkins et al. (2003), where they combine FUSE and HST data, permitting them to determ ine the strength of many Galactic lines as well as intervening systems. Four intervening absorption line systems are identified at a redshift below 0.1, and they are at z = 0.07344, 0.07779, 0.07900, and 0.08093 (Figure 4; the highest redshift system is a Lyman limit system with 17.5 < logN (HI) < 19.5). Of these systems, they identify many Lyman lines and a few metal lines, including 0 VI in the z = 0.07779 and z = 0.08093 systems at or above the 3 level, several times weaker than Ly . Jenkins et al. (2003) argue that the Lym an limit system is associated with an L galaxy that is 34 kpc from the line of sight and at the same redshift. The origin of the other absorption systems is less clear, as the velocity dimensional errors are great enough (580, 940, and 2250 km s¹) that the gas is unlikely to be gravitationally bound to the galaxy. They suggest that some of this absorbing material is due to tidal interactions between galaxies.

We have little to add to this analysis except to point out that it is possible that three system s are due to absorption within the supercluster. A side from the strongest stystem at z = 0.08093, the next two systems closest in redshift space are within 940 km s¹, while the most distant system is 2250 km s¹ away, which is still within the velocity range of a supercluster (or possibly even a cluster). The line of sight toward PHL 1811 is at least twice the virial radius from the nearest cluster (projected separation of 50°; the virial radius would be about 25°), so the absorbing material is not gravitationally bound to any of the clusters. D espite an extensive study by Jenkins et al., they do not identify any galaxies with redshifts that m ight be associated with these other redshift system s, nor are there tidal features evident with the Lym an lim it galaxy.

A lso, this line of sight passes through the A quarius-C etus supercluster and it lies closest to A bell 2399 (z = 0.0579; 102^{0} away) and A bell 2366 (z = 0.0529; 236^{0}), with a few clusters to the south, such as A bell 2361 and A bell 2362 (z = 0.0608), although these are m ore than 6 away (Figure 1d). Jenkins et al. (2003) report a Ly absorption line system at a redshift of 0.05119 (equivalent width of 0.60 0.05 A), a redshift di erence of 0.0067 (2000 km s¹) for A bell 2399 and 0.0017 (500 km s¹) for A bell 2366 (Figure 4). This is consistent with absorption by m aterial in this second supercluster along the line of sight.

3. Sum m ary and Concluding R em arks

We identi ed three bright AGNs that have been observed by FUSE and HST and lie near possible supercluster laments. Absorption lines, most of which were reported previously, are found to exist at redshifts within about 1300 km sec¹ of the supercluster, often being closest in redshift to the nearest galaxy cluster. These coincidences between the supercluster redshifts (or the redshifts of the nearest clusters) and the absorption line system s are important only if they are statistically signi cant, so we need to calculate the probability that this has occurred by chance. The redshift frequency of Ly absorption lines was the topic of an HST K ey project as well as more recent work (D obrzycki et al. 2002 and references therein), where at low redshift, the frequency of Ly absorption lines with equivalent widths greater than 0.24 A is dN/dz = 28 (for EW > 0.36 A, dN/dz = 15). We do not have Ly equivalent widths for all of the system s discussed above because in som e,



Fig. 4. FUSE spectrum of PHL 1811, showing four Ly systems near the redshift of the Aquarius B supercluster (plus two Galactic H₂ lines). The dotted line indicates the region of expected absorption within 1300 km s¹ of the redshift of the nearest cluster.

only Ly was detected without blending, as for the $z = 0.077\{0.081 \text{ absorption line systems}$ in PHL1811. However, if the ratio of Ly /Ly is about 0.5, as found for other systems of sim ilar equivalent widths (Shull et al. 2000), then the Ly equivalent widths are above the 0.24 A threshold (probably about 0.4 A) and the sum of the three lines have an equivalent width of 1.73 A, consistent with this estimation (Jenkins et al. 2003). For Ton S180, the absorption is due to 0 VI but it is challenging to detect Ly absorption as it would lie in the Ly emission line, but regardless of where one might choose a sensible continuum, there is no Ly absorption line with an equivalent width above 0.05 A. However, as the 0 VI clearly shows evidence for absorption, we will consider the elect of including this system in the analysis. Finally, the absorption line from PG 1402+261 is slightly below the threshold, so we will consider the results of the analysis with and without this system.

The next issue is how one counts a system and over what velocity width. The most conservative approach is to count multiple absorption line systems as a single absorption system (i.e., the 0.077-0.081 systems in PHL1811 and the 0.0616 - 0.0630 systems in Ton S180) even though the determ ination for dN /dz includes all redshift systems. The absorption line systems discussed above fall within 1300 km s¹ (0.0043 in redshift space) of the expected redshift of the lament or nearby cluster, so we should expect 0.24 absorption system s per supercluster. If we use only PHL 1811, there are absorption systems a sasociated with two superclusters (excluding the z = 0.08093 system and counting the systems at z = 0.07779 and 0.07900 as a single system), which should occur by chance 4.6% of the time (for dN /dz 28). U sing the rate for absorption system equivalent widths above 0.36 A (appropriate for these absorption system from PG 1402+ 261, the probability of this occurring by chance falls to 1.4% (for dN /dz 28; if we were to include the 0 VI absorption from Ton S180 as a single system , the chance probability would becom e 0.3%).

W hen we include all of the absorption systems separately, the association between superclusters and absorption features becomes more likely, as there would be three absorption systems in the redshift range 0.077-0.081 in PHL1811 and three in the redshift range 0.0616 – 0.0630 in Ton S180. Excluding Ton S180, the probability that the association with the superclusters occurred by chance is 9 10 5 (1.1 10 3 excluding the z = 0.08093 system), and when Ton S180 is included, the probability becomes 1.7 10 7 (2.2 10 6 excluding the z = 0.08093 system).

It is di cult at this point to dem onstrate that the absorption is associated with regions ofm ild overdensities (lam ents) rather than m aterial that has been ejected or torn away from a galaxy. One would need to be able to determ ine if absorbing m aterial is gravitationally bound to the galaxy, which would involve assum ing a model for the galaxy as well as for the transverse velocity of the absorbing gas relative to the galaxy. A loo, the distinction between these two associations may be unim portant, as material expelled in a galactic wind is unbound to that galaxy and becomes part of the lament once it mixes with this surrounding medium.

An alternative approach to boking for evidence of absorption from galaxy clusters and superclusters is to choose bright but random ly located AGNs and identify the absorption sites by obtaining in aging and spectroscopy of nearby objects. A long these lines, Penton, Stocke, and Shull (2002) found that about 22% of absorbers are not associated with galaxies (also, M cL in et al. 2002). They nd support that absorption line systems a lign with largescale laments of galaxies, which may be similar in nature to the laments that we have tried to probe.

The association of absorption systems with supercluster laments or galaxy clusters may seem surprising since a dedicated search to nd absorbing gas in galaxy clusters failed to produce detections (M iller, B regm an, and K nezek 2002). However, that study was sensitive only to low ionization m etal lines, not to the more common Ly absorption line systems (or to 0 V I). G iven the results here, a search to identify Ly systems around galaxy clusters would be worthwhile.

The relationship between superclusters and Ly absorption systems is enticing, but it is based on the study of just three systems, and it should be possible to improve the statistical situation quite readily. In the near future, we will be investigating several other low redshift superclusters (z < 0.12) to search for Ly absorption systems. Another observation that is badly needed is to obtain redshifts for the clusters that are near the lines of sight. We hope that studies such as those will lead to a deeper understanding of whetherm odest overdensity cosm ological laments can be identi ed through Ly absorption line studies.

A fler this paper was submitted, it was pointed out to us that there is evidence of a Ly absorber associated with the supercluster recently discovered from the RO SAT N orth E cliptic Pole Survey (NEP). The NEP supercluster (M ullis et al. 2001) is at an average redshift of 0.087 and is X-ray selected. It subtends a large angular region, where about 200 AGNs are found. One of them is bright enough for absorption line studies and was previously observed by HST, H1821+ 643 (NEP 5340). Spectral analyses of H1821+ 643 (z= 0.297) revealed the presence of a Ly system at z = 0.0891 (Tripp, Lu & Savage 1998). The nearest galaxy cluster is at a projected distance of 18 h⁻¹ M pc, well beyond the virial radius; no galaxy is identi ed near the absorption site. These observations are consistent with the absorber being associated with di use gas connecting the NEP clusters. As our selection criteria did not include this supercluster, we do not include it in our statistical analysis, but it supports our notings for the other superclusters.

We would like to acknow ledge support for this work from NASA grants NAG 5-10765 and NAG 5-10806, which support research on archival data analysis. A lso, we would like to thank B-G Andersen, Ken Sem bach, Bart Wakker, Jimmy Irwin, Chris Mullis, and Edward Lloyd-Davies for their advice and encouragement.

REFERENCES

- Bechtold, J., Dobrzycki, A., Wilden, B., Morita, M., Scott, J., Dobrzycka, D., Tran, K.-V, & Aldcroft, T.L. 2002, ApJS, 140, 143
- Cen, R., Kang, H., Ostriker, J.P., & Ryu, D. 1995, ApJ, 451, 436
- Cen, R., & Ostriker, J.P. 1999a, ApJ, 514, 1
- Cen, R., & Ostriker, J.P. 1999b, ApJ, 519, L109
- Cen, R., Tripp, T.M., Ostriker, J.P., & Jenkins, E.B. 2001, ApJ, 559, L5
- Dave, R., Cen, R., Ostriker, JP., Bryan, G.L., Hemquist, L., Katz, N., Weinberg, D.H., Noman, M.L., & O'Shea, B. 2001, ApJ, 552, 473
- Dekel, A., & Ostriker, J.P. 1999 Form ation of Structure in the Universe (Cambridge: Cambridge Univ. Press)
- Dobrzycki, A., Bechtold, J., Scott, J., & Morita, M. 2002, ApJ, 571, 654
- Einasto, M., Tago, E., Jaaniste, J., Einasto. J., & Andemach, H. 1997, A& AS, 123, 119

Evrard, A E., et al. 2002, ApJ, 573, 7

- Francis, P.J., Hewett, P.C., Foltz, C.B., Cha ee, F.H., Weymann, R.J., & Morris, S.L. 1991, ApJ, 373, 465
- Fukugita, M, Hogan, CJ., & Peebles, PJE. 1998, ApJ, 503, 518
- Hellsten, U., Gnedin, N.Y., & Miralda-Escude, J. 1998, ApJ, 509, 56
- Jenkins, E.B., Bowen, D.V., Tripp, T.M., Sembach, K.R., Leighly, K.M., Halpern, J.P., & Lauroesch, J.T. 2003, AJ, 125, 2824

Lockman, F.J., Murphy, E.M., Petty-Powell, S., & Urick, V.J. 2002, ApJS, 140, 331

- McLin, K M ., Stocke, J.T ., W eym ann, R J., Penton, S.V ., & Shull, J.M . 2002, ApJ, 574, L115
- Miller, E D ., B regm an, J N ., & K nezek, P M . 2002, A pJ, 569, 134
- Mullis, C.R. et al. 2001, ApJ, 553, L115
- Penton, S.V., Stocke, J.T., & Shull, JM. 2002, ApJ, 565, 720
- Penton, S.V., Stocke, J.T., & Shull, J.M. 2004, ApJS, 152, 29
- Savage, B D ., Sem bach, K R ., Tripp, T M ., & Richter, P. 2002 ApJ, 564, 631
- Schlegel, D.J., Finkbeiner, D.P., & Davis, M 1998, ApJ, 500, 525
- Shull, JM ., et al. 2000, ApJ, 538, L13
- Tripp, T.M., & Savage, B.D. 2000, ApJ, 542, 42
- Tumer, T J., et al. 2002, ApJ, 568, 120
- V eron-Cetty, M.-P., & V eron, P. 2001, Catalogue of Quasars and Active Nuclei (10th ed.; Garching: ESO)
- W akker, B P., et al. 2003, ApJS, 146, 1

This preprint was prepared with the AAS ${\mathbb I}{\mathbb P} T_E X$ m acros v5.0.