# H igh-spatial-resolution CN and CS observation of two regions of massive star form ation

H.Beuther

Harvard-Sm ithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138

hbeuther@cfa.harvard.edu

P.Schilke, F.W yrowski

Max-Planck-Institute for Radioastronomy, Aufdem Hugel 69, 53121 Bonn, Germany

#### ABSTRACT

M olecular line CN, CS and mm continuum observations of two interm ediateto high-m ass star-form ing regions { IRAS 20293+ 3952 and IRAS 19410+ 2336 { obtained with the P lateau de B ure Interferom eter at high spatial resolution reveal interesting characteristics of the gas and dust em ission. In spite of the expectation that the CN and CS m orphology m ight closely follow the dense gas traced by the dust continuum, both m olecules avoid the m ost central cores. C om paring the relative line strengths of various CN hyper ne components, this appears not to be an opacity e ect but to be due to chem ical and physical e ects. The CN data also indicate enhanced em ission toward the di erent m olecular out ows in the region. Regarding CS, avoiding the central cores can be due to high optical depth, but the data also show that the CS em ission is nearly always associated with the out ows of the region. Therefore, neither CS nor CN appear well suited for dense gas and disk studies in these two sources, and we recom m end the use of di erent m olecules for future m assive disk studies. An analysis of the 1 and 3m m continuum uxes toward IRAS 20293+ 3952 reveals that the dust opacity index

is lower than the canonical value of 2. Tentatively, we identify a decreasing gradient of from the edge of the core to the core center. This could be due to increasing optical depth toward the core center and/or grain grow th within the densest cores and potential central disks. We detect 3 mm continuum em ission toward the collimated out ow emanating from IRAS 20293+ 3952. The spectral index of 0.8 in this region is consistent with standard models for collimated ionized winds.

Subject headings: star: form ation { techniques: interferom etric { astrochem istry { ISM : individual (IRAS 20293+ 395; IRAS 19217+ 2336)

## 1. Introduction

M assive star-form ing regions consist of hundreds to thousands of M of dense gas from which future clusters are form ing. The local densities and chem istries vary signi cantly throughout the regions, and if one wants to observe the densest regions, and in particular potential disks at the core center, the right m olecular transitions need to be chosen to study the gas in detail. Especially the question of disks is still open in spite of large e orts over recent years. Pure continuum observations lack the necessary velocity information to prove real disk em ission, maser em ission offen only allows ambiguous interpretations, and therm al line em ission of dense gas tracers is offen contam inated by em ission from the ambient gas (e.g., C esaroni et al. 1997; N orris et al. 1998; W alsh et al. 1998; Zhang et al. 2002; Sandell et al. 2003). N evertheless, we think that observing therm al em ission from the right m olecules { which trace disk-like structures and are weak in ambient gas em ission in high-m ass star formation (e.g., C esaroni et al. 1997; Zhang et al. 2002; Beltran et al. 2004). The question remains: which m olecular transitions are best suited to pursue this case?

M any single-dish m olecular-line studies of m assive star-form ing regions have been undertaken (e.g., Schilke et al. 1997; H atchell et al. 1998; T hom pson & M acdonald 2003), but only very few regions have been observed at the high spatial resolution necessary to disentangle the densest m olecular clum ps from the potential disks (C esaroni et al. 1999; Sandell et al. 2003). Here, we present high-spatial-resolution P lateau de B ure Interferom eter (P dB I<sup>1</sup>) observations of two interm ediate to high-m ass star-form ing regions in two di erent potential dense-gas and disk-tracing m olecules, CN and CS.

O bservations toward low-mass T Tauri stars have shown that CN is enhanced in their disks (D utrey et al. 1997), and theoretical work by A ikawa & Herbst (2001) suggested that CN could be a good tracer of gaseous disks in deeply embedded sources as well. Therefore, we observed the interm ediate-mass region IRAS 20293+ 3952 with the PdBI in several di erent transitions of CN in the 1 and the 3mm band to test the suitability of CN as a potential massive disk tracer. The mm sources we are interested in are part of an interm ediate-mass sub-cluster in the near vicinity of an ultracom pact H ii region which provides most of the lum inosity within the region. For more details on the source and its multiple out ow

<sup>&</sup>lt;sup>1</sup> IRAM is supported by INSU/CNRS (France), MPG (Germany), and IGN (Spain).

properties we refer to Beuther et al. (2004b) and Table 1.

The other PdB I observations aim ed at the high-m ass star-form ing region IRAS 19410+ 2336 in two CS transitions and the continuum at 1 and 3 mm. CS has high critical densities  $(> 10^5 \text{ cm}^3)$  and is expected to follow the dust emission well (Shirley et al. 2003). These observations were meant to study the large scale CS gas and velocity distribution as well as tracing the innerm ost cores and possible disks. IRAS 19410+ 2336 has been studied intensively over recent years, and the complex out ow and mm continuum properties are outlined in Beuther et al. (2004b) and Beuther & Schilke (2004). Further source param eters are listed in Table 1.

Section 2 describes the technical details of the observations, and in section 3 we present the results obtained for both sources. Finally, section 4 discusses these results in the fram ework of future dense gas and disk studies in high-m ass star form ation.

## 2. Observations

W e observed  $\mathbb{R}AS20293+3952$  and  $\mathbb{R}AS19410+2336$  in several runs over the two w inters 2001/2002 and 2002/2003 with the Plateau de Bure Interferom eter (PdBI) at 1 and 3mm. The observing parameters for the various observations are listed in Table 1.

The typical system temperature at 1.3mm was about 300K and at 3mm about 135K. The phase noise at both wavelengths was mostly below 20 and always below 30. Atmospheric phase correction based on the 1.3mm total power was applied. For continuum measurements, we placed two 320M Hz correlator units in each band to cover the largest possible bandwidths. The nal continuum images were produced using only the line-free parts of the spectrum. We estimate the nal ux density accuracy to be 15%. Synthesized beam s at the di erent wavelengths are listed in Table 1.

## 3. Results

One of the goals of both observations { CN in IRAS 20293+ 3952 and CS in IRAS 19410+ 2336 { was to trace the densest gas and in particular the massive disks at the very centers of the evolving protoclusters. Before going into details, we note that at the highest spatial resolution, in both cases CN and CS do not trace the central condensations, but mainly gas in their near vicinity. Thus, the data show that CN and CS are not disk tracers in these two sources. This is distinctively di erent from the observations of low-mass T Tauri disks (D utrey et al. 1997).

## 3.1. M illim eter continuum em ission in IRAS 20293+ 3952

The three mm peaks (mm1 to mm3) observed with the PdBI at 2.6mm (Figs. 1, 2) correspond wellw ith the three peaks previously observed by Beuther et al. (2004b) at 1.3mm. The only di erences are that the previous 1.3mm data showed a small sub-com ponent northeast of the main peak mm1, whereas this new data show an elongation to the south-west of it (mm1a, Fig. 2). The north-eastern 1.3mm sub-com ponent of the previous dataset could be due to dust emission which m ight be litered out by the larger con gurations used to observe the source this time (AB of PdBI com pared to CD in the previous observation). Furtherm ore, at 3mm the sensitivity for dust emission decreases substantially (S /  $^{2+}$  with being the dust opacity index), and therefore we simply might not detect this sub-com ponent at the lower frequency.

Contrary to this, the sub-component mm 1a and its elongation in this new dataset is spatially correlated with the collim ated out ow in this direction (arrows in Fig. 1, Beuther et al. 2004b). This is similar to the recent observations of L1157 where G ueth et al. (2003) distinguish dust emission due to the out ow from the protostellar dust emission. It is likely that the sub-component mm 1a is caused by the collim ated out ow. We will discuss the scenario of a collim ated ionized wind with low spectral index in x42.

A ssum ing optically thin dust em ission for the mm continuum em ission, we can estim ate the dust m asses of mm 1 to mm 3. Setting the dust opacity index to 1 (see the follow ing paragraph) and using a dust temperature of 56K (Sridharan et al. 2002), the derived core m asses from the 3mm data are between 2.8 and 0.4M (Table 2), close to the values derived previously by Beuther et al. (2004b). For mm 1 we can also derive a m ass from the 1.3mm data which is about a factor 2 lower. This di erence can be attributed to the stronger spatial ltering e ects of the interferom eter at shorter wavelengths.

Spectral indices: The high spatial resolution and good signal-to-noise ratio of the continuum data allow us to analyze the spectral index in the mm regime. However, due to the di erent wavelengths, the uv-coverages and thus spatial resolution of the datasets are di erent (Fig. 2). To accurately compare the continuum data it is crucial to employ the same similarly sampled uv-range (in k ) in both bands. Therefore, we agged all data outside the uv-range 13 150 k (covered by both frequency bands) and re-im aged both datasets with the same beam of  $1:5^{00}$ . These new datasets are now well suited to determ ine spectral indices within the central mm core.

The right panel in Figure 2 presents the spectral index m ap of the central core m m 1 assuming a power law S / in the Rayleigh-Jeans regime with = 2 + where is the dust opacity index. Since the calibration uncertainty is 15%, we estimate the error in

and to be 0.3. We nd that the spectral index decreases from 3.4 at the outer edge of the core to values as low as 2.1 at the very center of m m 1. A sum ing a constant tem perature throughout the core, this corresponds to dust opacity indices decreasing from 1.4 at the core edge to 0.1 at the center. The variation of ism ore pronounced in north-west south-eastern direction perpendicular to the out ow than in the direction of the out ow. It is likely that the out ow a ects the mm continuum emission of the core in such a way that the observed in the out ow direction is lower than that perpendicular to it (see also the end of this section and 4.2).

Figure 3 outlines a di erent approach analyzing the data directly in the uv-plane. The 1 and 3 m m data can both be tted with a Gaussian plus a point source (excluding the data-points < 20 k). This way, we derive spectral indices of the Gaussian and point source com ponents of 2.7 and 1.6, respectively. A ssum ing that the Gaussian component is from a dust core with constant tem perature this results in a dust opacity index of 0.7. The central point source could be either a small optically thick disk or an unresolved hypercom pact H ii region.

In addition to the main core mm1, we can estimate the spectral index for the out ow associated emission feature (Table 2). We nd a lower spectral index toward mm1a with (mm1a) 0.8. Thus, is lower in the out ow region than within the central core.

## 3.2. CN in IRAS 20293+3952

Figure 1 presents the CN (1{0) and 2.6mm continuum emission as contours overlaid on the single-dish large-scale 1.2 mm image obtained with the MAMBO-array (Beuther et al. 2002b). The small o set between the single-dish and the interferom etric mm peaks is likely due to pointing inaccuracy of the single-dish data estimated to be  $5^{00}$  (Beuther et al. 2002b).

On large scales, the CN (1{0) emission follows the mm dust emission outlined by the MAMBO 12mm data. But at smaller spatial scales we nd that the CN peaks are always o set from the mm peaks. Because the interferom etric CN line and mm continuum data are taken simultaneously, the relative positional accuracy between the various maps is extremely high and the o sets are real. We observe similar morphologies in the 1.3mm band (Fig. 4). It appears that the CN (2{1) emission forms nearly a ring-like structure around the continuum source. While in the 1.3mm band only one line is strong enough to be imaged in these observations, in the 2.6mm band we have the main line plus 6 satellite lines due to the hyper ne splitting of CN (e.g., Sim on 1997). Figure 5 shows a spectrum of the main line and one hyper ne component (integrated over a region of R A: [5;1]<sup>(0)</sup> and Dec:[5;1]<sup>(0)</sup>.

The ratio of the peak intensities of the two lines is of the order of the theoretical value of

2:7 (Sim on 1997). To rst order, this indicates that all lines are optically thin and trace the whole CN column densities. Therefore, the weak CN emission toward the mm peaks is a realphysical e ect and not due to the optical depth of the observed lines.

One would also like to derive excitation temperature and column density estimates from the CN (2{1) and (1{0) transitions. However, missing ux strongly deteriorating the temperature and column density estimates is a problem. Therefore, we refrain from further analysis of physical parameters analysis based on this data. Short spacing information is crucial to proceed on this path.

# 3.3. CS in IRAS 19410+2336

The mm continuum data and their interpretation are presented in a separate paper (Beuther & Schilke 2004). Although we liter out about 80-90% percent of the continuum emission, the spatial litering is even more severe for the line emission. Comparing the interferom etric measured uxes (see example spectra in Figure 6) with the values derived from single-dish observations (Beuther et al. 2002b), we nd that of the order of only a few percent of the total CS line emission is recovered in the interferom eter data. Therefore, the dust continuum emission appears to be more strongly associated with the dense cores than CS.

Figure 7 (left panel) presents the integrated PdBICS( $2\{1\}$ ) observations (v [17,27]) km/s) as a contours overlaid on the single-dish 1.2mm dust em ission (Beuther et al. 2002b). To associate the CS emission with the many out ows in the regions, we show with arrows, ellipses and markers the 7 (or even 9?) out ows in the region, as presented in Beuther et al. (2003). Additionally, the right panel of F igure 7 presents the shocked  $H_2$  em ission in IRAS 19410+2336 (also taken from Beuther et al. 2003). We ndm any di erent CS em ission features throughout the regions (CS1 to CS9 in Fig. 7). However, the overall morphology diers signi cantly from the single-dish dust continuum data. The CS emission peaks fall into two classes, those in the near vicinity of the mm peaks (triangles in Figure 7; CS1, CS6 and CS7), and the rest of the emission peaks which are allo set from the main mm cores. C on paring these o set C S peaks with the known out ows in the region, it appears that they are all associated with one or other out ow feature: CS2 is at the wake of out ow (B), CS3 and CS4 appear to be associated with the out ows (G) and (H), CS8 is spatially associated with out ow (J) and also with a strong  $H_2$  feature. Finally, CS5 and CS9 both seem to be associated with out ow (F). While CS5 is again at the wake of the out ow, CS9 is within the out ow lobe where there is only very weak CO out ow emission (Beuther et al. 2003).

Zoom ing into the main peak regions in the south (Fig. 8) and in the north (Fig. 9, left) with the higher resolution of the CS (5{4) data, we nd that, although there is emission in the near vicinity of the mm peaks, the CS emission is o set from the main peaks. A spointed out in x32, the positional accuracy between the variousm aps is extrem ely reliable because of the simultaneous interferom etric observations of the CS line and the mm continuum emission. W hile we nd emission at all velocities in the north, the southern CS (5{4) emission is mainly found at blue velocities and the velocity of rest (v [14,22]km/s). There is barely any red CS (5{4) emission in the southern core at all (Fig. 6). Com paring it again with the out ow emission (Beuther et al. 2003), we see that the blue out ow wing (C) follows the same direction as the blue CS emission. The CS spectra observed toward CS1 (Fig. 6) som ewhat resem ble the infall spectra observed in CS toward the massive star-form ing region W 51 (Fig. 8 in Zhang et al. 1998). However, as our data su er strongly from missing ux problem s and the negative dip at 23 km/s is observed toward positions o set from the peak CS1 as well, we attribute this feature to observational artifacts due to missing short spacings and refrain from a physical interpretation.

The image is less clear in the north, the CS peaks are o set from the mm peak emission (Fig. 9), and three out ows emanate from that region. Nevertheless we cannot unam biguously attribute the CS em ission to any of these three out ows. So, it is very possible that other reasons contribute to the CS-mm o sets in the north as well, e.g., optical depth and/or freeze out of CS onto grains in cold core centers (see, e.g., Bergin & Langer 1997; Tafalla et al. 2004). In order to better understand the di erences between the  $CS(5{4})$  and  $(2{1})$ transitions, we con ned both datasets to the sam e uv-range and re-im aged these data. The m issing ux problem s are again severe (see x32) but we could produce integrated CS m aps of both transitions (Fig. 9, right). Both transitions peak o set from the continuum, but it appears that the (2{1) transition is even further o set from the mm continuum peak than the  $(5{4})$  transitions. D i erent explanations for these observations are possible: it could be an opacity e ext with CS (5{4) turning optically thick closer to the center than the CS (2{1) observations. However, physical reasons like higher densities and/or temperatures toward the core-center can produce this observational e ect as well. Unfortunately, based on this data we cannot di erentiate between the di erent scenarios { independent observations of tem perature and density tracers as well as of optically thin isotopom ers are necessary to get a hand on the di erent processes taking place in this region.

Similarly as in x32, we refrain from temperature and column density estimates based on the CS data because the missing ux deteriorates every such an estimate too strongly.

## 4. Discussion

#### 4.1. Line em ission

The search for and investigation of disks in m assive star form ation still rem ains one of the m issing links for a m ore detailed understanding of high-m ass star form ation. Theories predict disks in m assive star form ation (e.g., Yorke & Sonnhalter 2002), and other observations claim indirectly that disks need to exist because m ore and m ore collim ated m assive out ow s have been found (e.g., Beuther et al. 2002a). However, direct disk evidence in high-m ass star form ation is very scarce (C esaroni et al. 1997; Zhang et al. 2002) and yet not com pletely unam biguous. Therefore, it is necessary to establish a procedure to study the dense central cores and potential disks in m assive star form ation. W e believe that therm alm olecular line em ission from dense gas tracers which are not prom inent in the am bient m edium is the m ost prom ising approach.

O ne of the results of this paper is that C S and C N are not well suited for disk and dense gas studies in these sources. In spite of being a rather negative result, it needs to be stressed and we recommend the use of di erent molecules for future massive disk studies.

H igh optical depth probably plays a role for CS avoiding the dense centers of the regions. It remains to be tested whether rarer isotopom ers like  $C^{34}S$  or  $^{13}CS$  m ight trace the dense core centers and potential disks in a better way. In addition, the observations demonstrate that the CS emission is dominated to a large degree by the various out ows in IRAS 19410+2336. This CS emission is still associated with the large scale dust emission observed with single-dish instrum ents which likely rejects the similar range of gas densities needed to produce dust continuum and CS line emission.

The situation in slightly di erent regarding the CN emission in IRAS 20393+3952: it also avoids the core centers but optical depth e ects appear to be negligible. UV radiation is necessary to produce signi cant CN emission (e.g., Sim on 1997), and CN is detected toward low -m ass disks due to photo-dissociation by the star on the surface of the disk (D utrey et al. 1997; A ikawa & Herbst 2001). Two m ain reasons are likely to cause the lack of CN emission toward IRAS 20293+3952: (a) the source is too deeply embedded to excite the surface layers of the potential disk, or (b) the source is in such an early evolutionary stage that it does not produce enough photons. W ith the data available so far, we cannot properly distinguish between the two possibilities.

These observations outline the di culties of studying potential disks in massive star form ation. We have to look for other molecular transitions to properly investigate massive disks. While  $NH_3$  sometimes does a good job (Zhang et al. 2002) it is often confused by

the am bient gas. The sulphur-bearing species SO was observed toward the central driving source of HH 80-81 (G om ez et al. 2003) but it is also found toward the out ows in massive star-form ing regions (e.g., IRAS 18089-1732, Beuther et al. 2004c) in plying that SO can be am biguous as well. The most promising approach might be to focus on real hot-core-tracing molecules like  $CH_3CN$  in IRAS 20126+ 4104 (C esaroni et al. 1997, 1999), or  $HCOOCH_3$  observed toward IRAS 18089 1732 with the Submillimeter A may (SMA) (Beuther et al. 2004a). From an observational point of view, the SMA with a broad bandwidth of 2GHz in two sidebands (4GHz of spectral data) o ers a unique opportunity to observe many interesting lines simultaneously and thus compare di erent molecular transitions within the same observation (H o et al. 2004). This way, we should soon get a better census on the disk-tracing molecules in massive star form ation.

## 4.2. Continuum em ission

The dust opacity indices we derive from the imaging as well as from uv-ts are signi cantly lower than the canonical value of 2 (H ildebrand 1983). These results resemble sim ilar recent ndings of low in other massive star-form ing regions (Beuther et al. 2004a; K um ar et al. 2003). However, due to insu cient signal-to-noise ratio and uv-coverage, the previous study toward IRAS 18089 1732 was only capable of determ ining decreasing with increasing baseline length (Beuther et al. 2004a). Thus, their data could not tell whether the decreasing is due to unresolved clum py structure throughout the core or an actual decrease in toward the core center. The new data toward IRAS 20293+ 3952 now resolve di erent spectral indices for the outer regions as well as the core center. It is not yet clear whether we simply have two di erent components { a dust core with constant plus either an optically thick disk or an unresolved hypercom pact H ii region { or a gradient of varying

from the outside to the core center. M aybe both scenarios contribute to the observed spectral index variation. A gradient in would be similar to results that have been reported in low -m ass systems by H ogenheijde & Sandell (2000). Such a decrease of could either be due to increasing opacity toward the core center or grain grow th within circum -protostellar disks (see also Beckwith et al. 2000). Future work in that eld would be required to cover an even broader range of spatial scales, and it will be interesting to determ ine at what distance from the core center reaches the canonical value of 2, if it reaches it at all.

The nding that is lower toward the out ow than toward the core center is di erent from the recent results from G ueth et al. (2003) in the low-m assout ow L1157, who reported an average spectral index of 5 toward the out ow. Their analysis is based on bolom eter data over very broad band-passes (80 G H z) and thus line contam ination can introduce

arti cial ux increases in one or both bands. But even taking such artifacts into account, Gueth et al. (2003) conclude that in L1157 is higher toward the out ow than toward the core. Their observations point toward sm aller dust grains and/or a di erent chem istry, which is consistent with models predicting the evaporation of icy dust grains during shock interactions of the out ow with the surrounding dust.

These models obviously fail toward mmla. However, thermal ionized wind models predict spectral indices between 2 and 0:1 (Reynolds 1986). A typical spherical wind results in 0:6 whereas highly collimated jets have even lower values around 0.25, and in the optically thin case 0:1 (Reynolds 1986). Therefore, the derived (mmla) 0:8 within the out ow region of IRAS 20393+3952 is consistent with standard models of collimated ionized winds.

We very much appreciate the comments from an anonymous referee which helped to improve the manuscript. We would like to thank the sta of IRAM in Grenoble for help during the observations and the data reduction process. H.B. adknowledges nancial support by the Emmy-Noether-Program of the Deutsche Forschungsgemeinschaft (DFG, grant BE2578/1).

#### REFERENCES

- Aikawa, Y. & Herbst, E. 2001, A & A, 371, 1107
- Beckwith, S.V.W., Henning, T., & Nakagawa, Y. 2000, Protostars and Planets IV, 533
- Beltran, M. T., Cesaroni, R., Neri, R., et al. 2004, ApJ, 601, L187
- Bergin, E.A. & Langer, W.D. 1997, ApJ, 486, 316
- Beuther, H., Hunter, T.R., Zhang, Q., et al. 2004a, ApJ Letter accepted
- Beuther, H. & Schilke, P. 2004, Science, 303, 1167
- Beuther, H., Schilke, P., & Gueth, F. 2004b, ApJ, 608, 330
- Beuther, H., Schilke, P., Gueth, F., et al. 2002a, A & A, 387, 931
- Beuther, H., Schilke, P., Menten, K.M., et al. 2002b, ApJ, 566, 945
- Beuther, H., Schilke, P., & Stanke, T. 2003, A & A, 408, 601

- Beuther, H., Zhang, Q., Hunter, T.R., et al. 2004c, ApJ Letter accepted
- Cesaroni, R., Felli, M., Jenness, T., et al. 1999, A & A, 345, 949
- Cesaroni, R., Felli, M., Testi, L., Walmsley, C.M., & Olmi, L. 1997, A&A, 325, 725
- Dutrey, A., Guilloteau, S., & Guelin, M. 1997, A&A, 317, L55
- Gom ez, Y., Rodr guez, L.F., Girart, J.M., Garay, G., & Mart, J.2003, ApJ, 597, 414
- Gueth, F., Bachiller, R., & Tafalla, M. 2003, A&A, 401, L5
- Hatchell, J., Thompson, M.A., Millar, T.J., & MadDonald, G.H. 1998, A&AS, 133, 29
- Hildebrand, R.H. 1983, QJRAS, 24, 267
- Ho, P., Moran, J., & Kwok, Y.L. 2004, ApJ Letter accepted
- Hogenheijde, M.R. & Sandell, G. 2000, ApJ, 534, 880
- Kumar, M.S.N., Fernandes, A.J.L., Hunter, T.R., Davis, C.J., & Kurtz, S.2003, A&A, 412, 175
- Norris, R.P., Byleveld, S.E., Diamond, P.J., et al. 1998, ApJ, 508, 275
- Reynolds, S.P. 1986, ApJ, 304, 713
- Sandell, G., Wright, M., & Forster, J.R. 2003, ApJ, 590, L45
- Schilke, P., Groesbeck, T.D., Blake, G.A., & Phillips, T.G. 1997, ApJS, 108, 301
- Shirley, Y.L., Evans, N.J., Young, K.E., Knez, C., & Ja e, D.T. 2003, ApJS, 149, 375
- Simon, R. 1997, Ph.D. Thesis
- Sridharan, T.K., Beuther, H., Schilke, P., Menten, K.M., & Wyrowski, F. 2002, ApJ, 566, 931
- Tafalla, M., M. yers, P.C., Caselli, P., & W. alm sley, C.M. 2004, A&A, 416, 191
- Thom pson, M.A. & Macdonald, G.H. 2003, A & A, 407, 237
- Walsh, A.J., Burton, M.G., Hyland, A.R., & Robinson, G. 1998, MNRAS, 301, 640
- Yorke, H.W. & Sonnhalter, C. 2002, ApJ, 569, 846
- Zhang, Q., Ho, P.T.P., & Ohashi, N. 1998, ApJ, 494, 636

Zhang, Q., Hunter, T.R., Sridharan, T.K., & Ho, P.T.P. 2002, ApJ, 566, 982

This preprint was prepared with the AAS  ${\rm I\!AT}_E X$  m acros v5.2.

	<b>IRAS 20293+ 3952</b>		IRAS19410+2336
Date	W	inter 2002/2003	w inter 2001/2002 & 2003/2004
R A .[J2000]	20:31:12.9		19:43:11.4
Dec.[J2000]	+ 40:03:22.9		+ 23:44:06
Vlsr	6.3 km /s		22.4 km /s
Distance <sup>a</sup> [kpc]	2.0		2.0
Cong.	AB		BCD
Lines	$CN N = 1{0 \& N = 2{1}$		CSJ=5{4&J=2{1
Freq.[GHz]	113.385 & 226.656		97.981 & 244.936
Freq. res.		0.312MHz	0.039MHz
Velo.res.	0.83 & 0.41 km/s		0.12 & 0.05 km /s
(1 m m )	0 <b>:</b> 76 <sup>00</sup>	0:61 <sup>00</sup> (PA.37)	$1:52^{\circ\circ}$ $1:03^{\circ\circ}$ (PA.63)
(3mm)	1:4600	121°(PA32)	5:31 <sup>0</sup> 3:42 <sup>0</sup> (PA.77)
Mosaico sets [ $^{0}$ ]			-2/-2 -12/-8 9/2 -1 5/9 5
			-9.5/55 -19.5/49 -9.5/43
			-19.5/37 -19.5/25 -10/17
Phase calib.	2013+ 370 <b>,</b> 2023+ 336		2023+ 336 <b>,</b> 1923+ 210
Amp.calib.	MWC349		M W C 349
3 ms(1.3mm)	3 <b>.</b> 6m Jy		9m Jy <sup>b</sup>
3 ms(2.6mm)		1.2m Jy	0 <b>.</b> 75m Jy <sup>b</sup>

Table 1. Observation parameters

 $^{a}$ T he kinem atic distance am biguities are solved by associating the regions via the nearand m id-infrared surveys 2M ASS and M SX on larger scales with sources of known distance (B ontem ps, priv. com m .).

<sup>b</sup>D ata presented in Beuther & Schilke (2004)

 $S_{2:6m\,m}^{peak}$  $S_{1:3m\,m}^{peak}$  $S_{2:6m m}^{int}$  $S_{1:3m\,m}^{int}$ Source 0 set M <sub>3m m</sub> M <sub>lm m</sub> [\*\*\*] [m Jy] [m Jy/beam ] [m Jy] [m Jy/beam ] [M] [M] 42.3 < 3**:**4<sup>b</sup> m m 1 0/0 28.0 16.1 118.0 2.8 1.6 2**.**9ª 2.9 4.8 mm1a 2:7= 1:3 4.8 0:8 { { m m 210:2= 10:7 8.6 5.6 { { { 0.9 { 5:6= 12:4 3**.**6ª 3.6 { { { { mm3 0.4

Table 2. Continuum data in IRAS 20293+3952

<sup>a</sup>Source is unresolved.

<sup>b</sup>See the main text.

Fig. 1. | IRAS 20293+3952: the grey-scale shows the 12mm large-scale emission obtained with the MAMBO-array (Beuther et al. 2002b), the grey-scale levels are 18 to  $336 \text{ mJy}/(11^{\circ0}\text{beam})$  in  $36 \text{ mJy}/(11^{\circ0}\text{beam})$  steps  $(18(36)336 \text{ mJy}/(11^{\circ0}\text{beam}))$ . The contours present the PdBI observations: white contours the 2.6mm continuum (1.2(1.2)15.6 mJy/beam), and black contours the integrated CN(1{0}) emission (v [4,12]km/s, 29(42)407 \text{ mJy/beam}). The arrows sketch the four out ows previously observed by Beuther et al. (2004c), and the beam is shown at the bottom left. The sm allbox outlines the region presented in Figure 4.

Fig. 2. Millimeter Continuum emission in IRAS 20293+3952. The left and middle panels show the 2.6mm and 1.3mm emission, respectively. The contouring is done in 3 steps (1.2m Jy and 3.6m Jy, respectively). The right panel presents in grey-scale the spectral index m ap between the two bands and in contours the 1.3mm continuum emission. For the right panel, the data of both bands were selected to cover the same uv-range (in k).

Fig. 3. The 1.3 and 2.6mm continuum data in the uv-plane and the corresponding twocomponent ts. The top data-points (triangles) are the 1.3 mm data and the bottom datapoints are the measurements at 3 mm (squares). The dotted lines show the tted point source and the solid lines the combined ts consisting of a Gaussian plus a point source. Fig. 4. IRAS 20293+ 3952: the grey-scale with dotted contours shows the 1.3 mm continuum emission (3.6(3.6)39.6m Jy/beam), and the lined contours present the integrated CN (2(1) data (v [4,9]km/s, 111(74)703m Jy/beam). The beam is shown at the bottom right.

Fig. 5. | CN (1{0) spectrum in IRAS 20293+ 3952 integrated over a region of R A:  $[5;1]^{00}$  and Dec:  $[5;1]^{00}$ , the two di erent transitions are labeled at the top.

Fig. 6. [IRAS 19410+2336:  $CS(2\{1)$  and  $CS(5\{4)$  spectra in full and dashed lines, respectively. The spectra are taken toward the positions CS1 (within 6 arc-sec<sup>2</sup>) and CS6 (within 9 arc-sec<sup>2</sup>) (Fig. 7). The  $CS(5\{4)$  spectrum toward CS1 is scaled up by a factor 2.

Fig. 7. [RAS19410+2336: Left panel: the grey-scale shows the 1.2mm large-scale em ission obtained with the MAMBO-array (Beuther et al. 2002b), the grey-scale levels are  $85(85)765 \text{ mJy}/(11^{\circ0}\text{beam})$ . The contours present the integrated CS(2{1) PdBI em ission (v [17,27]km/s, 0.2(0.2)1.8 Jy/beam). The arrow s, ellipses, triangles and labels mark the outows and mm sources as outlined in Beuther et al. (2003). The 3mm beam is shown at the bottom right. R ight panel: the grey-scale now presents the infrared shocked H<sub>2</sub> em ission taken from Beuther et al. (2003). The contours again show the integrated CS(2{1) em ission, and the numbers label the di erent CS em ission features discussed in the main body.

Fig. 8. IRAS 19410+2336: the grey-scale shows the 1.3mm continuum data already presented in (Beuther et al. 2004c), the grey-scale levels are 7(7)63m Jy/beam. The contours outline the blue CS (5{4) em ission in that region (v [14,19]km/s, 0.5(0.5)4.5 Jy/beam). The arrows sketch out ow C from Beuther et al. (2003), the synthesized beam is shown at the bottom right.

Fig. 9. IRAS19410+2336: (left): the grey-scale shows the 1.3mm continuum data already presented in Beuther et al. (2004c), the grey-scale levels are 9(3)50 mJy/beam. The contours outline the integrated CS(5{4}) emission in that region (v [17,24]km/s, 0.7(0.7)6.3 Jy/beam). In the right image, we present the CS and 1.3mm continuum data imaged all with the same uv-range for better comparisons. The grey-scale shows the continuum, full contours CS(5{4}) and dashed contours (CS(2{1}), the contouring for all three is from 5 to 95% (step 10%) of their peak emissions, respectively. The synthesized beam s are drawn at the bottom left of each panel.

This figure "beutherf1.jpg" is available in "jpg" format from:

This figure "beutherf2.jpg" is available in "jpg" format from:

This figure "beutherf3.jpg" is available in "jpg" format from:

This figure "beutherf4.jpg" is available in "jpg" format from:

This figure "beutherf5.jpg" is available in "jpg" format from:

This figure "beutherf6a.jpg" is available in "jpg" format from:

This figure "beutherf6b.jpg" is available in "jpg" format from:

This figure "beutherf7.jpg" is available in "jpg" format from:

This figure "beutherf8.jpg" is available in "jpg" format from:

This figure "beutherf9.jpg" is available in "jpg" format from: