

Exploring the X-ray Sky with the XMM-Newton Bright Serendipitous Survey[?]

R. Della Ceca¹, T. Maccacaro¹, A. Acciari¹, P. Severgnini¹, V. Braito¹, X. Barcons², F.J. Carrera², M.G. Watson³, J.A. Tedds³, H. Brunner⁴, I. Lehmann⁴, M.J. Page⁵, G. Lammer⁶ and A. Schweppe⁶

¹ INAF-Osservatorio Astronomico di Brera, via Brera 28, I-20121 Milano, Italy. e-mail: rdc@brera.mi.astro.it, tommaso@brera.mi.astro.it, caccia@brera.mi.astro.it, paola@brera.mi.astro.it, braito@brera.mi.astro.it

² Instituto de Fisica de Cantabria (CSIC-UC), Avenida de los Castros, 39005 Santander, Spain. e-mail: barcons@ifca.unican.es, carreraj@ifca.unican.es

³ X-ray Astronomy Group, Department of Physics and Astronomy, Leicester University, Leicester LE1 7RH, UK. e-mail: mgw@star.le.ac.uk, jat@star.le.ac.uk

⁴ Max-Planck-Institut für Extraterrestrische Physik, Postfach 1312, 85741 Garching, Germany. e-mail: hbrunnerl@mpe.mpg.de, ile@mpe.mpg.de

⁵ Mullard Space Science Laboratory, University College London, Holmbury St. Mary, Dorking, Surrey RH5 6NT, UK. e-mail: mjp@mssl.ucl.ac.uk

⁶ Astrophysikalisches Institut Potsdam (AIP), An der Sternwarte 16, 14482 Potsdam, Germany. e-mail: glamer@aip.de, aschwope@aip.de

Abstract. We present here "The XMM-Newton Bright Serendipitous Survey", composed of two flux-limited samples: the XMM-Newton Bright Source Sample (BSS, hereafter) and the XMM-Newton "Hard" Bright Source Sample (HBSS, hereafter) having a flux limit of $f_x \approx 7 \times 10^{-14}$ erg cm⁻² s⁻¹ in the 0.5-4.5 keV and 4.5-7.5 keV energy band, respectively. After discussing the main goals of this project and the survey strategy, we present the basic data on a complete sample of 400 X-ray sources (389 of them belong to the BSS, 67 to the HBSS with 56 X-ray sources in common) derived from the analysis of 237 suitable XMM-Newton fields (211 for the HBSS). At the flux limit of the survey we cover a survey area of 28.10 (25.17 for the HBSS) sq. deg. The extragalactic number-flux relationships (in the 0.5-4.5 keV and in the 4.5-7.5 keV energy bands) are in good agreement with previous and new results making us confident about the correctness of data selection and analysis. Up to now

71% (~90%) of the sources have been spectroscopically identified making the BSS (HBSS) the sample with the highest number of identified XMM-Newton sources published so far. At the X-ray flux limits of the sources studied here we found that: a) the optical counterpart in the majority (~90%) of cases has a magnitude brighter than the POSS II limit ($R \approx 21^{mag}$); b) the majority of the objects identified so far are broad-line AGN both in the BSS and in the HBSS. No obvious trend of the source spectra (as deduced from the Hardness Ratios analysis) as a function of the count rate is measured and the average spectra of the "extragalactic" population corresponds to a (0.5-4.5 keV) energy spectral index of 0.8 (~0.64) for the BSS (HBSS) sample. Based on the hardness ratios we infer that about 13% (40%) of the sources in the BSS (HBSS) sample are described by an energy spectral index steeper than that of the cosmic X-ray background. Based on previous X-ray spectral results on a small subsample of objects we speculate that all these sources are indeed absorbed AGN with the N_{H} ranging from a few times 10^{21} up to few times 10^{23} cm⁻². We do not find strong evidence that the 4.5-7.5 keV survey is sampling a completely different source population if compared with the 0.5-4.5 keV survey; rather we find that, as expected from the CXB synthesis models, the hard survey is simply picking up a larger fraction of absorbed AGN. At the flux limit of the HBSS sample we measure surface densities of optically type 1 and type 2 AGN of 1.63 ~ 0.25 deg⁻² and 0.83 ~ 0.18 deg⁻², respectively; optically type 2 AGN represent 34 ~ 9% of the total AGN population. Finally, we have found a clear separation, in the hardness ratio diagram and in the (hardness ratio) vs. (X-ray to optical flux ratio) diagram, between Galactic "coronae fitting" stars and extragalactic sources. The information and "calibration" reported in this paper will make the existing and incoming XMM-Newton catalogs a unique resource for astrophysical studies.

1. Introduction

Deep Chandra and XMM-Newton observations (Brandt et al., 2001; Rosati et al., 2002; Moretti et al., 2003; Haasinger et al., 2001; Alexander et al., 2003) have recently resolved > 80% of the 2-10 keV cosmic X-ray background (CXB) into discrete sources down to $f_x < 3 \times 10^{-16} \text{ erg cm}^{-2} \text{ s}^{-1}$.

The statistical analysis (stacked spectra and hardness ratios) performed on these faint samples provide information on the X-ray spectral properties of the sources making up most of the CXB. The X-ray data are consistent with AGN being the dominant contributors of the CXB (see Brandt et al., 2004 and reference therein) and, as inferred by the X-ray colors, a significant fraction of these sources have hard, presumably obscured, X-ray spectra, in agreement with the predictions of CXB synthesis models (see Setti and Woltjer, 1989; Madau et al., 1994; Comastri et al., 1995, 2001; Gilli et al. 2001; Ueda et al., 2003).

However the majority of the sources found in these medium to deep fields are too faint to provide good X-ray spectral information. Furthermore, the extremely faint magnitude of a large number of their optical counterparts makes the spectroscopic identifications very difficult, or even impossible, with the present day ground-based optical telescopes.

Thus, notwithstanding the remarkable results obtained by reaching very faint X-ray fluxes, the broad-band physical properties (e.g. the relationship between optical absorption and X-ray obscuration and the reason why AGN with similar X-ray properties have completely different optical appearance) are not yet completely understood. A step forward toward the solution of these problems has been recently obtained by Mainieri et al., 2002; Piconcelli et al., 2002; 2003; Georgantopoulos et al., 2004; Acciari et al., 2004 and Perola et al., 2004 using samples of serendipitous sources for which medium /good quality XMM-Newton and optical data are available.

With the aim of complementing the results obtained by medium to deep X-ray surveys, the XMM-Newton Survey Science Centre¹ (SSC) has conceived the "XMM-Newton Bright Serendipitous Survey". This survey comprises two high galactic latitude ($|b| > 20^\circ$), flux limited samples of serendipitous XMM-Newton sources: the XMM Bright Source Sample (hereafter BSS) and the XMM Hard Bright

Send offprint requests to: R.Della Ceca

¹ Based on observations obtained with XMM-Newton, an ESA science mission with instruments and contributions directly funded by ESA Member States and the USA (NASA). The majority of the new optical spectroscopy data used here have been obtained using the facilities of the Italian "Telescopio Nazionale Galileo" (TNG) and of the European Southern Observatory (ESO).

¹ The XMM-Newton Survey Science Centre is an international collaboration involving a consortium of 10 institutions appointed by ESA to help the SOC in developing the software analysis system, to pipeline process all the XMM-Newton data, and to exploit the XMM-Newton serendipitous detections, see <http://xmmsc.ssc-www.star.le.ac.uk>

Source Sample (hereafter HBSS) having a flux limit of $f_x > 7 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$ in the 0.5-4.5 keV and 4.5-7.5 keV energy bands, respectively. In addition to the issues related to the CXB, where it is now largely accepted that X-ray obscured AGNs play a significant (and perhaps major) role, the use of the 4.5-7.5 keV energy band partially reduces the strong bias against absorbed sources which occurs when selecting at softer energies (or when selecting in the optical domain), and is therefore fundamental to study the accretion history in the Universe (see e.g. Fiore et al., 2003). A similar energy selection band (i.e. 5-10 keV) was pioneered by the BeppoSAX-HELLAS (Fiore et al., 2001) and the ASCA-SHEEP (Nandra et al., 2003) surveys.

The well defined criteria (completeness, representativeness, etc..) of this sample will allow both a detailed study of individual sources of high interest, and statistical studies of populations. In particular, the BSS and HBSS samples will be fundamental to complement other medium and deep XMM-Newton and Chandra survey programs (having fluxes 10 to 100 times fainter and covering a smaller area of the sky) and will provide a larger baseline for all evolutionary studies. Moreover, the high X-ray statistics which characterize most of the sources in the "XMM-Newton Bright Serendipitous Survey", combined with the relative brightness of their optical counterparts, will allow us to investigate their physical properties in detail. Indeed this sample is already contributing to the solution of some critical open (and "hot") questions like the relationship between optical absorption and X-ray obscuration (Acciari et al., 2004) and the physical nature of the "X-ray bright optically normal galaxies" (Severgnini et al., 2003). Many of these issues are investigated with difficulty using the fainter X-ray samples because of their typical poor counts statistics for each source.

The spectroscopic identifications together with the X-ray (spectral, morphological and variability) parameters will be made available to the community and can be used to define statistical identification procedures to select rare and interesting classes of X-ray sources, enabling the application of these procedures to the vast amount of XMM-Newton serendipitous data that will be accumulated during the lifetime of the mission².

² One of the responsibilities of the XMM-Newton SSC is the production of the XMM-Newton Source Catalogue. This catalogue will provide a rich and unique resource for generating well-defined samples for specific studies, using the fact that X-ray selection is a highly efficient way of selecting certain types of objects, like for instance AGN, clusters of galaxies and active stars. The first XMM-Newton Serendipitous Source Catalogue (1XMM), released on 2003 April 7, contains source detections drawn from 585 XMM-Newton EPIC observations and a total of 30000 individual X-ray sources having a likelihood value above 8 and good quality σ . The median flux (in the total photon energy band 0.2-12 keV) of the catalogue sources is $3 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$, with 12% of them having fluxes below $1 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$, see <http://xmmsc.ssc-www.star.le.ac.uk/>

In this paper we discuss the BSS and the HBSS survey strategy, we present a complete sample of 400 sources extracted from the analysis of 237 XMM-Newton fields and we discuss some preliminary statistical results based on the spectroscopic identification done so far.

This paper is organized as follows. In section 2 we discuss the survey strategy (e.g. energy selection bands, primary selection camera and criteria for field and source selection), we present basic information on the XMM-Newton fields used and on the sources belonging to the BSS and HBSS samples and we discuss the completeness of the "XMM-Newton Bright Serendipitous Survey". In section 3 we discuss the number-flux relationship, the identification work done so far, the broad-band X-ray spectral properties of the sample, the position of the sources in the diagram obtained using the X-ray spectral information (provided by the hardness ratio) and the X-ray to optical flux ratio as well as the surface densities ($\log(N > S) - \log S$) of type 1 and type 2 AGN in the HBSS sample. Finally, the summary and the conclusions are reported in section 4. In the appendices we discuss our approach to evaluate the background quality of the data used and to deal with the X-ray sources falling close to the gaps between the CCDs or close to the edge of the CCDs. Throughout this paper $H_0 = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $\Omega = 0.7$, $M = 0.3$ are assumed; the energy spectral index, α , quoted in this paper refers to a power-law spectral model having $S_E \propto E^{-\alpha}$.

2. Survey Strategy and Sample(s) Selection

2.1. Selection Energy Band(s)

We have decided to survey the bright X-ray sky in two complementary energy bands: the 0.5–4.5 keV and the 4.5–7.5 keV energy bands.

The choice of the 0.5–4.5 keV energy band is mainly motivated by the desire to avoid the very soft photons (minimizing non-uniformities introduced by the different values of Galactic absorbing column densities along the line of sight) and by the need to compromise between a broad passband (to favor throughput) and a narrow passband (to minimize non-uniformities in the selection function due to different source spectra). Furthermore in the 0.5–4.5 keV band XMM-Newton has the highest throughput.

The choice of the 4.5–7.5 keV energy band (one of the energy bands used in the standard pipeline processing system of the XMM-Newton data) was instead dictated by the need to study the composition of the source population (in terms of observed and intrinsic energy distribution and absorption properties) as a function of the energy selection band, comparing the sources selected in this band with those selected in the softer 0.5–4.5 keV energy range. Moreover this energy band reduces the strong bias against absorbed sources which occurs when selecting at softer energies.

2.2. Primary Selection Camera

The source sample has been defined using the data from the EPIC MOS2 detector only. The main reasons for this choice are:

1. unlike the EPIC pn, the EPIC MOS cameras have a detector pattern that simplifies the analysis of the field. For example, in the case of the EPIC MOS detectors the source target, in the large majority of the observations, is fully contained in the central chip;
2. the PSF in the 2 EPIC MOSs is narrower than in the EPIC pn. In particular, the EPIC MOS2 has the "best" PSF (FWHM 4.4⁰⁰ and HEW 13.0⁰⁰ at 1.5 keV, see Ehle et al., 2003). As a comparison, the EPIC pn PSF has FWHM 6.6⁰⁰ and HEW 15.2⁰⁰, while the EPIC MOS1 PSF has FWHM 4.3⁰⁰ and HEW 13.8⁰⁰.
3. the gaps between the EPIC MOS chips are narrower than the gaps in the EPIC pn detector, simplifying source detection and analysis and maximizing the survey area;
4. unlike with the EPIC pn camera, we can still use part of the EPIC MOS2 observations in large- and small-window mode by only excluding the area occupied by the central chip. Since 25% of the observations have been performed in window mode, retaining these observations will maximize the searched area, speeding up the creation and definition of the source sample.

The major disadvantage of the EPIC MOS2 camera when compared to the EPIC pn camera is the reduced sensitivity, because of its smaller effective area. However, since the BSS and HBSS samples contain relatively bright sources, and considering the minimum exposure times used here (see section 2.5) this lower efficiency does not affect the source selection of the samples presented here. Obviously, once a source is detected and included in the sample, additional information using data from the EPIC MOS1 and pn detectors are collected to increase the statistics for the X-ray spectra, timing and morphology analysis.

2.3. Source Detection

Each EPIC MOS2 observation used here (see Table 2) has been processed through the pipeline processing system used for the production of the XMM-Newton Serendipitous Source Catalogue, based on tasks from the XMM-Newton Science Analysis Software. Full details about the processing system, the pipeline products as well as the source searching procedures, flux measurements, source likelihood parameter, corrections for vignetting and PSF, etc.. can be found in <http://xmm.ssc-www.star.le.ac.uk>. We note that the count rate(s) reported in this paper have been already corrected for vignetting and PSF.

2.4. Criteria for Source Selection

Since the BSS and HBSS samples have been designed to contain relatively bright X-ray sources not all the sources detected in each individual MOS2 field are adequate to be included in these samples. We discuss here the criteria for the BSS and HBSS source selection within each EPIC MOS2 field:

1. BSS sample: $0.5 \{ 4.5 \text{ keV count-rate } 1 \cdot 10^{-2} \text{ cts/s.}$

At this count rate limit, and given the considered range of MOS2 exposure times (see Table 2 and figure 1), all the selected sources have a likelihood parameter in the $0.5\{4.5 \text{ keV}$ energy band greater than 18 (corresponding to a probability for a random Poissonian fluctuation to have caused the observed source counts of $1.5 \cdot 10^{-8}$). No further constraint is thus needed to ensure the source reliability.

HBSS sample: $4.5 \{ 7.5 \text{ keV count-rate } 2 \cdot 10^{-3} \text{ cts/s}$ and likelihood parameter in the $4.5\{7.5 \text{ keV}$ energy band greater than 12 (corresponding to a probability of $6 \cdot 10^{-6}$ for a spurious detection).

The combination of count-rate limit(s) and likelihood parameter(s) of the sources in the BSS and HBSS samples is such that none of them are expected to be spurious.

The count rate to flux conversion factors (CR2F) depend on the source spectra and in Table 1 we report some CR2F as a function of the input source spectra for a fixed N_H value of $3 \cdot 10^{20} \text{ cm}^{-2}$ (corresponding to the median value for the XMM-Newton fields used here).

For a source with a power-law spectrum with energy spectral index ϵ between 0.7 and 0.8 the count rate limit in the two chosen bands corresponds to a flux limit of $7 \cdot 10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2}$.

2. sources with a distance from the EPIC MOS2 center between an inner radius (R_{in}) and an outer radius (R_{out}).

R_{in} depends on the actual size and brightness of the target and on the window mode. R_{in} ranges between 0 (e.g. survey fields with no "target") and 8 arcmin (e.g. bright/extended X-ray sources or large- and small-window mode). In this way the area of the detector "obscured" by the presence of the target or not exposed is excluded from the analysis. To guarantee that all the sources in the catalogue are truly serendipitous, the size of R_{in} has been adapted in order to exclude the target and the sources physically related to the target. R_{out} is, for the large majority of the fields, equal to 13 arcmin. In the few overlapping fields we have excluded from the analysis the outer region of one of the overlapping fields in order to obtain a mosaic of separate and independent regions on the sky. The values of R_{in} and R_{out} used for each MOS2 image are listed in Table 2.

3. we have also excluded the sources too close to the edges of the field of view or to the gaps between the CCDs. These sources could have either the flux and/or the

Table 1. MOS2 Count Rate to Flux Conversion Factors

E	HR2	Flux (0.5-4.5 keV)		Flux (4.5-7.5 keV) $10^{-11} \text{ ergs cm}^{-2} \text{ s}^{-1}$
		$10^{-12} \text{ ergs cm}^{-2} \text{ s}^{-1}$	(3)	
(1)	(2)	(3)	(4)	(4)
3.0	0.77	14.3; 14.4; 15.0		4.53; 4.54; 4.59
2.0	0.56	12.8; 12.9; 13.6		4.24; 4.25; 4.30
1.0	0.22	10.6; 10.8; 11.7		3.97; 3.97; 4.03
0.5	0.01	9.47; 9.61; 10.6		3.84; 3.84; 3.90
0.0	0.22	8.35; 8.49; 9.55		3.71; 3.72; 3.77
0.4	0.38	7.56; 7.70; 8.82		3.62; 3.62; 3.68
0.5	0.42	7.38; 7.53; 8.66		3.59; 3.60; 3.65
0.7	0.49	7.06; 7.20; 8.37		3.55; 3.55; 3.61
0.9	0.57	6.78; 6.92; 8.11		3.50; 3.51; 3.56
1.0	0.60	6.65; 6.79; 8.00		3.48; 3.49; 3.54
1.5	0.74	6.16; 6.31; 7.61		3.38; 3.38; 3.43
2.0	0.83	5.89; 6.04; 7.46		3.28; 3.28; 3.33
3.0	0.94	5.75; 5.94; 7.62		3.10; 3.10; 3.15

Columns are as follows: (1) Energy spectral index (S_E / E^{α}); (2) corresponding hardness ratio HR2 computed as discussed in section 3.3; (3) Flux (corrected for Galactic absorption) in the 0.5-4.5 keV energy band corresponding to an observed count rate of 1 cts/s in the same energy band. The three numbers refer to thin, medium and thick filters; (4) Flux (corrected for Galactic absorption) in the 4.5-7.5 keV energy band corresponding to an observed count rate of 1 cts/s in the same energy band. The three numbers refer to thin, medium and thick filters. NOTE { A Galactic absorbing column density of $3 \cdot 10^{20} \text{ cm}^{-2}$, the median value for the XMM-Newton fields used here, has been assumed. Given the range of the Galactic absorbing column density along the line of sight (from $5 \cdot 10^{19}$ to 10^{21} cm^{-2}) the CR2F in the 0.5-4.5 keV energy range are accurate to 18%; the CR2F in the 4.5-7.5 keV energy range are independent of the Galactic N_H . Please note that in the case of sources having extreme HR2 values the observed spectra could be much more complex than a simple power law; for these sources the conversion factor reported in the table should be considered only as indicative and a proper X-ray spectral analysis is needed.

source centroid poorly determined (due to the proximity to the edges and/or the gaps), and therefore could degrade the quality of the data, and would require uncertainty corrections thus representing a problem in the subsequent analysis and interpretation of the data. In Appendix A we discuss the procedure used to take into account this problem in an objective way. Obviously, the excluded area has been taken into account in the computation of the sky coverage.

2.5. Criteria for Field Selection

Not all the available EPIC MOS2 pointings are adequate for producing the BSS and HBSS samples. We have defined a set of selection criteria to avoid problematic regions of the sky, to maximize the availability of ancillary information at other frequencies (i.e. optical and radio) and to speed up the optical identification process. The majority

of the fields selection criteria are common to the BSS and HBSS; however, as discussed below, we have been more conservative on the minimum exposure time and background properties for the fields used to define the HBSS sample. The criteria adopted for field selection are:

1. availability to SSC before March 2003 (XM M -Newton fields that are public or with PI granted permission);
2. high Galactic latitude ($|bj| > 20^\circ$) to avoid crowded fields, to obtain a relatively "clean" extragalactic sample and to have magnitude information for the optical counterparts from the Digital Sky Survey material (the Automated Plate Machine - APM - catalogue³ is almost complete for $|bj| > 20^\circ$);
3. Galactic absorbing column density along the line of sight less than 10^{21} cm^{-2} , to minimize non-uniformities introduced by large values of the Galactic N_H ;
4. exclusion of fields centered on bright and/or extended X-ray or optical targets and those containing very bright stars in the optical band. In the first two cases the effective area of sky covered and the actual flux limit are difficult to estimate correctly, making the derivation of the sky-coverage more uncertain; in the latter case the search for the optical counterpart of the X-ray sources could be very difficult or even impossible due to the presence of the bright star;
5. exclusion of fields south of DEC = -80° deg since it could be very difficult to obtain good quality spectroscopy given the location of the optical facilities available to us;
6. good-time interval⁴ exposure $> 5 \text{ ks}$ for the BSS and $> 7 \text{ ks}$ for the HBSS. According to the results presented and discussed below, with these constraints all the sources in the two samples are detectable across the whole field of view considered, ensuring a "flat" sensitivity and therefore a flat sky coverage at the sampling rates.
7. finally, we have also excluded EPIC MOS2 pointings suffering from a high background rate (i.e. accumulated during particle background ares). The background restriction has been more conservative for the set of fields that have been used to define the HBSS sample since the overall background is more critical given the faintness of the sources in the $4.5\text{-}7.5 \text{ keV}$ energy band. We have defined and computed in an automatic way a Background Estimator Parameter (see Appendix B) which is roughly proportional to the "real background" in the MOS2 images used. The set of fields that have been used to define the HBSS sample must have the Background Estimator Parameter less than 100.

³ <http://www.ast.cam.ac.uk/apm/cat/>

⁴ The good-time interval is defined as the on-axis exposure time taken from the exposure map produced in the XM M pipeline processing system.

Note that we have also considered the EPIC MOS2 observations in large- and small-field mode satisfying the criteria discussed above; in these cases we have excluded from the analysis a circular area of 8 arcmin radius enclosing the central chip. No restrictions on the blocking filter in front of the MOS2 camera have been applied since, as shown in section 2.6, the filter used does not affect the statistical properties of the sample⁵.

The complete BSS sample reported here is based on the analysis of 237 XM M -Newton fields, while the complete HBSS sample is based on a "restricted" data set of 211 XM M -Newton pointings.

In Table 2 we report basic information on the XM M -Newton MOS2 fields used for the sample selection; in particular we list the XM M -Newton observation number, the blocking filter in front of the MOS2 instrument, the Right ascension and Declination of the MOS2 image center, the on-axis good-time exposure for the MOS2 detector, the logarithm of the Galactic Hydrogen column density along the line of sight (fromDickey and Lockman, 1990), the inner and outer radius of the part of the MOS2 image used in the survey, and the total number of BSS and HBSS sources found in the surveyed area of each MOS2 image. In Table 2 we have also marked the 26 MOS2 fields not used for the production of the HBSS sample.

In Figure 1 we show the distribution of the MOS2 on-axis good-time exposure, the distribution of the Galactic hydrogen column density along the line of sight and the distribution of the Background Estimator Parameter for the XM M -Newton MOS2 data-set used.

2.6. The XM M -Newton BSS and HBSS samples.

Applying the source selection criteria discussed in section 2.4 to the MOS2 fields reported in Table 2 we have selected 400 XM M -Newton sources: 389 sources belong to the BSS sample and 67 sources to the HBSS sample with 56 sources in common. Basic information on the sources are reported in Table 3 (BSS) and in Table 4 (HBSS); in particular we report the source name, the XM M -Newton observation number, Right Ascension and Declination (J2000) of the X-ray source position, the angular distance (in arcmin) between the source and the MOS2 image center, the source count rate in the $0.5\text{-}4.5 \text{ keV}$ energy band (BSS sample) or in the $4.5\text{-}7.5 \text{ keV}$ energy band (HBSS sample), the hardness ratios computed as described in section 3.3, and the optical spectroscopic classification (see section 3.2 for details). In Table 4 we have also marked the 11 sources belonging to the HBSS sample but not to the BSS sample.

In Figure 2 we show the surface density of the sources belonging to the BSS and to the HBSS as a function of: the MOS2 on-axis good-time exposure (panel a); the Background Estimator Parameter (panel b); the off-axis angle (panel c); and the blocking filter in front of the MOS2 detector (panel d). The appropriate area covered in

⁵ The fraction of MOS2 images with a thin, medium and thick filter used here are 48%, 46% and 6%, respectively.

Table 2. Basic information on the XM M -Newton M O S2 fields used for the sample selection. We have reported here just few lines of Table 2 for illustration purpose. The complete Table 2 will be published only in the electronic version of the Journal.

Obs. ID	Filter	RA ; DEC (J2000)	Exposure s	Log N h cm ⁻²	R _{in} arcmin	R _{out} arcmin	BSS srcs	H BSS srcs
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
0125310101	M edium	00 00 30.4 25 06 43.4	19162.6	20.27	1	13	4	0
0101040101	M edium	00 06 19.7 + 20 12 22.8	34044.7	20.60	8	13	1	0
0127110201	Thin1	00 10 31.2 + 10 58 40.7	7558.0	20.76	2	13	2	0
...
...

Columns are as follows: (1) XM M -Newton Observation number; (2) Filter used; (3) Right ascension and Declination (J2000) of the M O S2 image center; (4) On-Axis good-time exposure; (5) Logarithm of the Galactic hydrogen column density along the line of sight fromDickey and Lockman, 1990; (6) Inner radius of the part of the M O S2 image used; (7) Outer radius of the part of the M O S2 image used; (8) total number of BSS sources found in the surveyed area of each M O S2 image; (9) total number of H BSS sources found in the surveyed area of each M O S2 image. See section 2.4 for details.

each bin has been considered and errors have been computed using Poisson statistic. The dashed lines reported in Figure 2 correspond to the mean surface density obtained considering the whole sample. As can be seen there is no significant trend of the source surface density with respect to the plotted parameters concerning a at sensitivity across the field (i.e. at sky coverage at the sampled fluxes). The only point which seems to be a factor 2.5 above the other is the bin at the highest exposure time in the BSS sample (see panel a). This excess is due to 5 sources found in the field 0022740101 (centered on the Lockman hole), the only pointing in the bin considered; however the error bars are large and so the reported surface density is not significantly different from the mean value. The absolute source surface density as a function of the flux (LogN - LogS) is also in very good agreement with previous and new measurements (see section 3.1) making us confident of the correctness of the data analysis and source selection.

3. First Results

3.1. The Number Counts Relationship (s)

In Figure 3 we show (filled circles) a binned representation of the extragalactic⁶ number-flux relationships in the 0.5-4.5 keV energy band (panel a) and in the 4.5-7.5 keV energy band (panel b). As already shown in section 2.6, the sky coverage of this survey at the flux limit used to define the BSS and the H BSS samples is at and equal to 28.10 sq.deg and 25.17 sq.deg, respectively; given the at sky coverage the errors in the binned representation are Poissonian errors on the total number of sources having a flux greater than any fixed flux. A conversion

⁶ Since we are primarily interested in the extragalactic number-flux relationship we have excluded from the computation the sources classified as stars (see section 3.2). Based on the results presented in section 3.5 we are confident that the large majority of the unidentified sources are associated to extragalactic objects.

factor appropriate for a power-law spectral model with energy index equal to 0.8 (0.7) in the 0.5-4.5 keV (4.5-7.5 keV) energy band, itered by an $N_{\text{H Gal}} = 3 \times 10^{20} \text{ cm}^{-2}$ (the median value of the $N_{\text{H Gal}}$ of the survey), has been used in the conversion between the count rate and the flux. The energy spectral index used in the 0.5-4.5 keV energy band corresponds to the "average" one of the extragalactic BSS population in the same energy selection band (see section 3.3). In the 4.5-7.5 keV energy band we have used $\alpha = 0.7$, the same energy spectral index assumed from other recent surveys in the 5-10 keV band (e.g. the H ELLA S2XMM survey, Baldi et al., 2002) and very close to the median energy spectral index in the 0.5-4.5 keV band of the extragalactic H BSS population (see below for details). We recall that, given the median $N_{\text{H Gal}}$ of the survey, the count rate to flux conversion factor in the 0.5-4.5 keV (4.5-7.5 keV) energy band are accurate 20 (-8)% for an energy spectral index in the range between 0 to 2.

Both LogN > S - LogS distributions can be well described by a power-law model $N > S / S^{\alpha}$; their best fit spectral parameters, obtained applying the maximum likelihood method to the unbinned data (see Macacaro et al., 1982 for details), are reported in Table 5.

The fits have been performed from a flux of 7×10^{14} ergs $\text{cm}^{-2} \text{s}^{-1}$ (the faintest flux) to a flux of 1×10^{12} ergs $\text{cm}^{-2} \text{s}^{-1}$ (we have excluded from the fit extragalactic BSS sources brighter than this flux limit). For fluxes brighter than this limit we may not be complete since "bright" X-ray sources were chosen as targets of observations and then excluded, by definition, from the survey. However we note that the surface density of sources with flux greater than 1×10^{12} ergs $\text{cm}^{-2} \text{s}^{-1}$ is such that about 2.7 (0.9) sources in the BSS (H BSS) sample are expected given the covered sky area; these numbers are fully consistent with what is observed.

Both Log(N > S) - LogS derived here have been compared with a number of representative Log(N > S) - LogS reported in the literature. In particular, in Figure 3 panel

Table 3. Basic information on the XM M -Newton BSS sample. We have reported here just few lines of Table 3 for illustration purpose. The complete Table 3 will be published only in the electronic version of the Journal.

Name XBS ...	Obs. ID	RA; DEC (J2000)	O Axis arcmin	Rate 10^{-2} cts/s	HR 2	HR 3	Class			
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)			
J000027.7	250442	0125310101	00 00 27.8	25 04 42.6	2.10	1.13	0.08	$0.48^{+0.06}_{-0.06}$	$0.37^{+0.11}_{-0.11}$	AGN 1
J000031.7	245502	0125310101	00 00 31.8	24 55 02.9	11.68	1.41	0.13	$0.67^{+0.07}_{-0.06}$	$0.37^{+0.17}_{-0.17}$	AGN 1
J001002.2	250501	0125310101	00 01 00.2	25 05 01.2	6.97	1.38	0.10	$0.37^{+0.07}_{-0.07}$	$0.47^{+0.10}_{-0.10}$	AGN 1
...
...

Columns are as follows: (1) Source name; (2) XM M -Newton Observation number; (3) Right Ascension and Declination (J2000) of the source (X-ray position); (4) Angular distance (in arcmin) between the source and the MOS2 image center; (5) Source count rate, and 1 error, in the 0.5-4.5 keV energy band (units of 10^{-2} cts/s). In Table 1 we have reported the MOS2 conversion factors between the 0.5-4.5 keV count rate and the flux as a function of the energy spectral index, the hardness ratio HR 2 and the blocking filter; (6) and (7) Hardness ratios computed as described in section 3.3. The errors on the hardness ratios have been evaluated using simulations and correspond to 1 ; (8) Optical spectroscopic classification (AGN 1: broad line AGN; AGN 2:narrow line AGN; GAL: Optically Normal Galaxy; CL: Cluster of Galaxies; BL: BL Lac Object; star: star; ?: Tentative classification; see section 3.2 for details).

Table 4. Basic information on the XM M -Newton HBSS sample. We have reported here just few lines of Table 4 for illustration purpose. The complete Table 4 will be published only in the electronic version of the Journal.

Name XBS ...	Obs. ID	RA; DEC (J2000)	O Axis arcmin	Rate 10^{-3} cts/s	HR 2	HR 3	Class	
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	
J002618.5+105019	0001930101	00 26 18.5 + 10 50 19.3	9.56	2.35	0.55	$-0.53^{+0.04}_{-0.04}$	$-0.67^{+0.06}_{-0.06}$	AGN 1
J013240.1+133307	0084230301	01 32 40.1 + 13 33 07.8	11.90	3.23	0.71	$-0.02^{+0.10}_{-0.10}$	$-0.37^{+0.11}_{-0.11}$	AGN 2
J013944.0+674909	0032140401	01 39 44.0 + 67 49 09.4	2.69	2.05	0.46	$-0.50^{+0.07}_{-0.07}$	$-0.43^{+0.13}_{-0.13}$	AGN 1?
...
...

Columns are as follows: (1) Source name; (2) XM M -Newton Observation number; (3) Right Ascension and Declination (J2000) of the source (X-ray position); (4) Angular distance (in arcmin) between the source and the MOS2 image center; (5) Source count rate, and 1 error, in the 4.5-7.5 keV energy band (units of 10^{-3} cts/s). In Table 1 we have reported the MOS2 conversion factors between the 4.5-7.5 keV count rate and the flux as a function of the energy spectral index, the hardness ratio HR 2 and the blocking filter; (6) and (7) Hardness ratios computed as described in section 3.3. The errors on the hardness ratios have been evaluated using simulations and correspond to 1 ; (8) Optical spectroscopic classification (AGN 1: broad line AGN; AGN 2:narrow line AGN; GAL: Optically Normal Galaxy; CL: Cluster of Galaxies; BL: BL Lac Object; star: star; ?: Tentative classification; see section 3.2 for details).

a, we have reported: a) the extragalactic ROSAT (0.5-2.0 keV) Log(N > S)-LogS from Hasinger et al., 1998 (dashed line); b) the EMSS (0.3-3.5 keV) extragalactic number density at 10^{13} ergs cm $^{-2}$ s $^{-1}$ (open triangle; Gioia et al., 1990) and c) the extragalactic surface density obtained from the XM M -Newton AXIS Medium Survey team in the 0.5-4.5 keV energy range (Barcons et al., 2002; open square at $S' \approx 2 \times 10^{14}$ ergs cm $^{-2}$ s $^{-1}$). To convert the ROSAT 0.5-2.0 keV band fluxes and the EMSS 0.3-3.5 keV band fluxes into 0.5-4.5 keV fluxes we have used a power-law spectral model having $\Gamma = 1.0$, corresponding to the mean spectral index of the ROSAT and the EMSS sources (see Hasinger et al., 1993 and MacAccarone et al., 1988, respectively).

In panel b) we have compared our result with: a) the extragalactic XM M -Newton (5-10 keV) Log(N > S)-LogS from the HELLA2XM M survey (area inside the thick

solid lines; Baldi et al., 2002); b) the BeppoSAX -HELLAS (5-10 keV) Log(N > S)-LogS (open squares; Fiore et al., 2001); c) the ASCA-SHEEP (5-10 keV) Log(N > S)-LogS (dashed line; Nandra et al., 2003). Finally, the open circle at $S' \approx 2 \times 10^{15}$ ergs cm $^{-2}$ s $^{-1}$ represents the extragalactic surface density in the (5-10 keV) energy band from the XM M -Newton observation of the Lockman hole field (Hasinger et al., 2001). For consistency with previous hard survey both these latter number densities have been converted from their original (5-10 keV) band to the 4.5-7.5 keV band using a power-law spectral model having $\Gamma = 0.7$. We found that our results are fully consistent with those obtained from the other XM M -Newton related survey (e.g. the HELLA2XM M 5-10 keV survey) and, moreover, our better statistics above 7×10^{14} ergs cm $^{-2}$ s $^{-1}$ allow us to significantly constrain the 4.5-7.5 keV extragalactic number densities above this flux.




Fig. 1. Histograms of some basic properties of the XMM-Newton MOS2 fields used for the sample selection. Normal histograms refer to the XMM-Newton fields used to define the BSS sample, while shaded histograms refer to the subset of XMM-Newton fields used to define the HBSS sample. Panel a: histogram of the MOS2 on-axis good-time exposure; Panel b: histogram of the Galactic hydrogen column density along the line of sight; Panel c: histogram of the Background Estimator Parameter (see Appendix B for details).

Table 5. The Extragalactic Log($N > S$) - LogS maximum likelihood best-fit parameters: $N > S = K \cdot (S/10^{13})^{\alpha}$, where $N > S$ is the surface densities of sources having a flux greater than S in deg^{-2}

Sample	Objects	K	
(1)	(2)	(3)	(4)
BSS ^a	330	$1.80_{-1.69}^{+1.91}$	$6.1_{-6.4}^{+5.8}$
HBSS	65	$1.64_{-1.41}^{+1.89}$	$1.5_{-1.6}^{+1.4}$
HBSS AGN 1	41	$1.72_{-1.42}^{+2.03}$	$0.92_{-1.0}^{+0.82}$
HBSS AGN 2	21	$1.57_{-1.18}^{+1.99}$	$0.50_{-0.58}^{+0.42}$

Columns are as follows: (1) Sample; (2) Number of sources used in the fit; (3) Best-fit power-law slope and 68% confidence intervals; (4) Log($N > S$) - LogS normalization and 68% confidence intervals. Note that the normalization K is not a parameter of the fit but is determined by re-scaling the model to the number of objects in the sample. The normalization K reported here corresponds to the surface densities at $1 \cdot 10^{-13}$ ergs $\text{cm}^{-2} \text{s}^{-1}$. NOTE {^a} As explained in section 3.1 we have excluded from the fit the 3 BSS extragalactic sources having a flux greater than $1 \cdot 10^{-12}$ ergs $\text{cm}^{-2} \text{s}^{-1}$.

On the other hand the extragalactic HBSS Log($N > S$) - LogS falls below both the BeppoSAX - HELLAS and the ASCA - SHEEP determinations. Given the results discussed in section 2.6 and the very similar slope between our Log($N > S$) - LogS and the BeppoSAX / ASCA Log($N > S$) - LogS we have checked if this problem could be related to an offset of the absolute flux scale in the 4.5 - 7.5 keV energy range between XMM-Newton and BeppoSAX / ASCA. To this purpose we have cross-correlated the HELLAS and the SHEEP sources with the total catalogue of XMM-Newton sources obtained from the analysis of the 237 XMM-Newton fields reported in

Table 2. Using a search radius of 90^{arcmin} and considering the point-like XMM-Newton sources with a 4.5 - 7.5 keV likelihood parameter greater than 12 and with an "Illumination Factor" (see Appendix A for details) greater than 0.8, we have found 6 "bona fide" HELLAS-XMM coincidences and 2 "bona fide" SHEEP-XMM coincidences. The ratio between the 4.5 - 7.5 keV XMM-Newton fluxes and the 5-10 keV BeppoSAX fluxes in the case of the HELLAS sources ranges between 0.09 and 0.98 with a mean value of 0.47, while in the case of the two SHEEP sources the ratio between the 4.5 - 7.5 keV XMM-Newton fluxes and the 5-10 keV ASCA fluxes is equal to 0.63 for both objects. Although these small numbers do not allow us to draw firm conclusions we note that using a conversion factor between the fluxes in the 5-10 keV energy range and the fluxes in the 4.5 - 7.5 keV energy range equal to 0.47 (instead of 0.69 as expected for $E = 0.7$ and as assumed in figure 3) the HBSS Log($N > S$) - LogS and the BeppoSAX - HELLAS Log($N > S$) - LogS turn out to be in perfect agreement. This suggests that an offset in the absolute flux scale could easily explain the disagreement in the number densities discussed above; this possible discrepancy in the flux scale has to be further investigated.

3.2. Optical identification and classification

One of the main characteristics of the X-ray sources presented here is that the majority (~90%) of them have an optical counterpart above the POSSUM limit ($R < 21^{\text{mag}}$), thus allowing spectroscopic identification even on 2-4 meter class telescopes. Furthermore, given the good accuracy of the X-ray positions⁷ and the magnitude of the optical

⁷ Using the optical position of the sources classified as Type 1 AGN we have evaluated that the 90% confidence level error circle has a radius equal to 4^{arcsec} ; about 99% of the Type 1




Fig. 2. Panel a: Source surface density as a function of the MOS2 on-axis good-time exposure for the sources belonging to the BSS (open circles) and to the H BSS (filled circles) sample. Panel b: Source surface density as a function of the Background Estimator Parameter; symbols as in panel a. Panel c: Source surface density as a function of the off-axis angle; the bin size has been adapted in order to have similar areas in each bin; symbols as in panel a. Panel d: Source surface density as a function of the blocking filter in front of the MOS2 detector; symbols as in panel a. For all the panels the dashed lines correspond to the mean surface density considering the whole BSS or H BSS sample. Errors have been computed using Poisson statistic.

counterparts there is no ambiguity in the optical identification for the large majority of cases.

AGN are within 6^{arcmin} from the X-ray position. This is consistent with what found in other XMM-Newton surveys (Barcons et al. 2002; Fiore et al. 2003).

Up to now 285 X-ray sources have been spectroscopically identified (either from the literature or from our own observations mainly at the Italian "Telescopio Nazionale Galileo" -TNG, at the ESO 3.6m, at the Calar Alto 2.2




Fig. 3. The extragalactic number–flux relationship in the 0.5–4.5 keV energy band (panel a) and in the 4.5–7.5 keV energy band (panel b) obtained using the BSS and HBSS samples (binned representation: black filled circles). In the 0.5–4.5 keV $\text{Log}(N > S)$ – $\text{Log}S$ (panel a) we have also reported the ROSAT (0.5–2.0 keV) $\text{Log}(N > S)$ – $\text{Log}S$ (dashed line) and the EMSS (0.3–3.5 keV) extragalactic number density at 10^{13} $\text{ergs cm}^{-2} \text{s}^{-1}$ (open triangle) both converted to the 0.5–4.5 keV energy band. The open square at $S \approx 2 \times 10^{14}$ $\text{ergs cm}^{-2} \text{s}^{-1}$ represents the extragalactic surface density (0.5–4.5 keV) obtained by the XMM-Newton AXIS Medium Survey project. In panel b (4.5–7.5 keV $\text{Log}(N > S)$ – $\text{Log}S$) we have also reported the HELLA2XMM $\text{Log}(N > S)$ – $\text{Log}S$ interval (area inside the thick solid lines), the HELLA $\text{Log}(N > S)$ – $\text{Log}S$ (open squares) and the SHEEP $\text{Log}(N > S)$ – $\text{Log}S$ (dashed line). The open circle at $S \approx 2 \times 10^{15}$ $\text{ergs cm}^{-2} \text{s}^{-1}$ represents the extragalactic surface density (5–10 keV band) in the Lockman hole field. Both these latter number densities have been converted from their original (5–10 keV) band to the 4.5–7.5 keV band.

m or at the NOT 2.6m⁸ telescopes) leading to a 71% and 90% identification rate for the BSS and HBSS samples respectively.

The optical breakdown of the sources identified so far is reported in Table 6. We stress that the source detection algorithm is optimized for point-like sources, so the sample of clusters of galaxies is not statistically complete nor representative of the cluster population.

To our present knowledge all but one (XBS J014100.6 675328⁹) of the sources classified as stars are coronalemitters. If we consider the BSS sources with Right Ascension below 5^h or above 17^h (spectroscopic identification rate of 88%, see Table 6), the X-ray coronalemitters represent 14% of the $\delta\beta > 20^\circ$ (0.5–4.5 keV) population at the sampled fluxes. This fraction must be compared with 15% of coronalemitters in

the HBSS sample¹⁰; this smaller fraction of stars in the HBSS sample, compared with that in the BSS sample, is entirely consistent with their low temperature coronal emission. Note that in the softer (0.5–2.0 keV) ROSAT Bright Survey Catalog (RBS, Schwope et al., 2000) the fraction of coronalemitters is around 37%.

The largemajority (~90%) of the extragalactic X-ray sources are emission line objects, i.e. sources for which at least one strong ($\text{EW} >> 5 \text{\AA}$ in the source rest frame) emission line is present in the optical spectrum. As a comparison in the RBS the fraction of emission line AGN amongst the extragalactic sources is around 55% (Schwope et al., 2000). The few remaining non-emission line objects have been classified as "Normal galaxies" or BL Lacs objects according to the measured Calcium break discontinuity at the rest frame wavelength of 4000 \AA (see e.g. Landt et al., 2002). We stress that some of the sources classified as "Normal galaxies" could indeed host an AGN. As a

⁸ As part of the AXIS (An XMM-Newton International Program) project, see <http://www.ifca.unican.es/~xray/AXIS>

⁹ XBS J014100.6 675328 (BL Hyi), also belonging to the HBSS sample, is a well known AM Herculis object (a polar) i.e. a binary system composed of a magnetized white dwarf and a low-mass star (see Caccianiga et al., 2004 and references therein).

¹⁰ On the basis of the results presented in section 3.5 we are confident that the bulk of unidentified sources both in the BSS and HBSS samples are associated with extragalactic objects and we know that XBS J014100.6 675328 is an accreting binary system (see Caccianiga et al., 2004).

Table 6. The current optical breakdown of the BSS and HBSS samples

	BSS ¹	HBSS
Objects ²	389 (166)	67
Identified:	278 (146)	60
Identification rate	71% (88%)	90%
AGN-1	180 (100)	39
AGN-2	26 (15)	16
Galaxies ³	7 (3)	1
Clusters of galaxies	4 (1)	1
BL Lacs	5 (3)	1
Stars ⁴	56 (24)	2

¹ In brackets we have reported the optical breakdown for the BSS sources with Right Ascension below 5^{h} or above 17^{h} ; ² Note that 56 sources are in common between the BSS and HBSS samples; ³ We stress that some of the sources classified as "Optically Normal Galaxy" could indeed host an optically elusive AGN (see e.g. Severgnini et al., 2003); ⁴ To our knowledge all but one (XBS J014100.6 675328) of the sources classified as stars are coronal emitters.

already discussed by Severgnini et al. (2003) using X-ray and optical spectral data from this project, the lack of significant emission lines in the optical spectra can be explained by an adequate combination of the absorption associated with the AGN and of the optical faintness of the active nucleus with respect to the host galaxy. Furthermore for some of the sources classified as "Normal Galaxies" the H line (in some case the only spectroscopic evidence of the presence of an AGN in the optical domain) is not sampled. Although the presence of an AGN in the nucleus of some of these sources is highly probable (e.g. observed L_x well in excess of 10^{42} ergs s⁻¹) we prefer to wait for a confirmation also from optical/infrared follow-ups; for the moment these objects are classified as "Normal Galaxies".

To classify the emission line objects we have used the criteria presented for instance in Veron-Cetty & Veron (2001) which are based on the line width and the line flux ratios. Type 1 AGN are those sources showing broad (FWHM > 1000 km s⁻¹) permitted lines, while Type 2 AGN are those sources showing only narrow lines (FWHM < 1000 km s⁻¹) and, when detected, [O III] 5007/H > 3 .

A few sources show permitted lines with 1000 km s⁻¹ $<$ FWHM $<$ 2000 km s⁻¹ and [O III] 5007/H below 3. These sources are probably narrow line Seyfert 1 candidates and, according to our classification, have been included in the Type 1 AGN group. For some sources classified as Type 2 AGN we have indication of the presence of a broad component at the bottom of the narrow H α and/or H β lines. These sources should be properly classified as Seyfert 1.8 or Seyfert 1.9 objects; for the purpose of the present paper these sources have been included in Type 2 AGN group. Finally for 26 objects a better S/N optical

spectrum and/or a more appropriate set-up for the spectroscopic observations are needed to truly classify them as Type 1 or Type 2 AGN. These 26 sources have been marked in column 8 of Table 3 and 4.

3.3. 0.5-4.5 keV spectral properties

A "complete" spectral analysis for all the sources in the BSS and HBSS samples (using data from the two EPIC MOSs and the EPIC pn) is in progress; first results on selected sub-samples of sources have been already discussed in Severgnini et al. (2003) and Caccianiga et al. (2004). In the meantime, and in order to extract first order X-ray spectral information we present here a "Hardness Ratio" analysis of the single sources using only EPIC MOS2 data; this latter method is equivalent to the "color-color" analysis largely used at optical wavelengths. The use here of the "Hardness Ratio" analysis is twofold. First of all it is much faster than a complete spectral analysis with the combined use of three different instruments. Second, a "Hardness Ratio" is often the only X-ray spectral information available for the faintest sources in the XMM-Newton catalogue, and thus, a "calibration" in the parameter space is needed to select "clean" and well-defined samples. On the other hand, in Caccianiga et al. (2004) we have already shown and discussed a tight correlation between X-ray absorption, as deduced from a complete X-ray spectral analysis, and "Hardness Ratio" properties.

We have used the hardness ratios as defined from the XMM-Newton pipeline processing¹¹:

$$HR2 = \frac{C(2-4.5 \text{ keV}) - C(0.5-2 \text{ keV})}{C(2-4.5 \text{ keV}) + C(0.5-2 \text{ keV})}$$

and

$$HR3 = \frac{C(4.5-7.5 \text{ keV}) - C(2-4.5 \text{ keV})}{C(4.5-7.5 \text{ keV}) + C(2-4.5 \text{ keV})}$$

where $C(0.5-2 \text{ keV})$, $C(2-4.5 \text{ keV})$ and $C(4.5-7.5 \text{ keV})$ are the "PSF and vignetting corrected" count rates in the 0.5-2, 2-4.5 and 4.5-7.5 keV energy bands, respectively.

In Figure 4 (panel a and panel b) we plot HR2 as a function of the 0.5-4.5 keV count rate for the extragalactic (and unidentified) sources of the BSS sample. In particular, in panel a) we show the position of the optically classified Type 1 and Type 2 AGN, while in panel b) we have shown the unidentified sources, the "Optically Normal Galaxies", the Clusters of galaxies and the BL Lac objects. On the top, we have also reported the

¹¹ We have not used here the "pipeline processing product" HR1 which is defined using the corrected count rate in the (0.15-0.5) keV and in the (0.5-2) keV energy band since the measured count rate in the (0.15-0.5) keV band is a strong function of the Galactic N_H along the line of sight. Note that the effect on HR2 and HR3 due to the different N_{H,gal} for the objects in the sample (which ranges between 10^{20} to 10^{21} cm⁻²) is negligible.

scale computed assuming a conversion factor appropriate for E' = 0.8, which is the "mean" energy spectral index of the "extragalactic" sample in the 0.5–4.5 keV energy band (see below).

No obvious trend in the source spectra as a function of the count rate is measured. If we split the extragalactic BSS sample in two different bins of count rate (below or above a count rate of 1.41×10^2 cts/s in the 0.5–4.5 keV energy band) which includes a similar number of objects (163 and 170 sources, respectively) we found that the HR2 distributions of the two sub-samples are consistent with being extracted from the same distribution with a probability of 61% according to a KS test.

The weighted average of the extragalactic population is $\text{HR2} = 0.51 \pm 0.01$ corresponding to $E = 0.74 \pm 0.03$ (the relation between the spectral index and HR2 has been tabulated for few representative spectral indices in Table 1). For comparison the weighted averages of the sources classified as Type 1 AGN and unidentified objects are $\text{HR2} = 0.54 \pm 0.01$ and $\text{HR2} = 0.52 \pm 0.02$, respectively. The HR2 distribution of the sources identified as Type 1 AGN appears to be "narrow" with 90% of the sources inside the HR2 range 0.75 to 0.35. On the contrary the Type 2 AGN are characterized by a broader distribution with 42% of the objects having an observed energy spectral index apparently steeper than that of the cosmic X-ray background ($E = 0.4$ corresponding to $\text{HR2} = 0.38$). For comparison only 10% of the unidentified X-ray sources or of the X-ray sources identified as broad line AGN seem to have spectra apparently steeper than $E = 0.4$.

Please note that two broad line AGN, which are clearly separated from the majority of the other broad line AGN, seem to be characterized by an extremely hard spectrum ($E < 1$). These two objects, belonging both to the BSS and HBSS samples, are XBS J091828.4+513931 ($\text{HR2} = 0.31$) and XBS J143835.1+642928 ($\text{HR2} = 0.18$). For the first object the optical spectra show broad H and H α emission lines without any obvious sign of peculiarity, while in the case of XBS J143835.1+642928 the optical spectrum in hand is very noisy and therefore the optical classification is at the moment tentative. The X-ray spectra of both sources are described by an absorbed power-law model having an intrinsic N_{H} in excess of 10^{22} cm $^{-2}$. At the moment these two sources are the only broad line AGN displaying intrinsic absorption (as derived from a complete X-ray spectral analysis) above 10^{22} cm $^{-2}$ but the completion of the X-ray spectral work for the total sample is needed to evaluate correctly the fraction of X-ray absorbed broad line AGN in this survey¹².

¹² We note that in the HBSS28 sample discussed in Caccianiga et al., 2004 none of the 19 Type 1 AGN is absorbed with N_{H} in excess to 10^{22} cm $^{-2}$, implying that the fraction of X-ray absorbed Type 1 AGN is less than 10% of the total Type 1 AGN population. This result is also confirmed by the present HBSS sample since at most 4 (out of 39) broad line AGN are characterized by HR2 values typical of absorbed ($N_{\text{H}} > 10^{22}$ cm $^{-2}$) AGN. See also section 3.4.

It is now worth comparing the HR2 properties of the extragalactic BSS and HBSS sources; in figure 4 (panel c and d) we plot HR2 as a function of the 4.5–7.5 keV count rate for the extragalactic (and unidentified) sources of the HBSS sample. As a class, the Type 1 AGN in the HBSS sample seem to have the same HR2, and thus 0.5–4.5 keV spectral properties, of the Type 1 AGN in the BSS; about 87% of them reside in the HR2 range between 0.75 and 0.35, with a median HR2 value of 0.53. On the contrary the Type 2 AGN in the HBSS seem to be characterized by more extreme spectral properties if compared with the Type 2 AGN in the BSS sample; 81% of them seem to have an energy spectral index steeper than 0.4 and 23 seem to have inverted spectra ($E < 0$). It is worth noting that 2 of the 3 type 2 AGNs having HR2 around 0.55 in the HBSS sample are Seyfert 1.9 galaxies.

The HR2 statistic for some relevant BSS and HBSS sub-sample(s) have been summarized in Table 7.

We note that in the case of the extragalactic population in the BSS sample the use of the weighted average, the unweighted average or the median value of HR2 give consistent results on the underlying spectral index ($E = 0.74$, $E = 0.83$ and $E = 0.91$, respectively). On the contrary in the case of the extragalactic population in the HBSS sample the use of the weighted average, the unweighted average or the median value of HR2 gives completely different results ($E = 0.27$, $E = 0.11$ and $E = 0.64$, respectively). Since the HR2 distribution of the extragalactic population in the HBSS sample is significantly different from a Gaussian we prefer to use as "average" spectral index of this sample that related to the median value ($E = 0.64$).

A similar result has been pointed out by Nandra et al., 2003 studying the spectral properties of the sources in the SHEEP (5–10 keV) survey. These authors prefer the use of the unweighted average over the weighted average (they did not consider median values) and reach the result that $< E > < 0$ and the conclusion that the 5–10 keV surveys are sampling a completely different population compared with the 2–10 keV surveys.

On the contrary we find strong evidence that the 0.5–4.5 spectral properties of the class of broad line AGN in the BSS and in the HBSS are very similar. Moreover, since the majority of the objects in the BSS and HBSS samples are in common we do not find compelling evidence that the surveys in the two bands are selecting completely different populations. The HBSS survey is simply more efficient than the BSS survey in selecting the hard part of the intrinsic source spectral distribution.

The eleven objects belonging only to the HBSS sample (enclosed inside empty squares in figure 4, 5 and 6 below) are amongst the hardest X-ray sources in the sample; all but one (XBS J140113.4+024016, still optically unidentified) of them seem to be characterized by an apparently inverted spectrum ($E < 0.0$). Among these sources there are 5 Type 2 AGN, one optically normal galaxy, one BLLac object and 4 unidentified objects.

Table 7. HR 2 statistic for some relevant BSS and HBSS sub-sample(s).

Sample (1)	Objects (2)	weighted (3)	unweighted (4)	median (5)	st.dev. (6)	intrins. (7)
BSS Sample:						
Extragalactic	333	-0.51	0.01	-0.54	0.01	-0.57
AGN 1	180	-0.54	0.01	-0.56	0.01	-0.58
AGN 2	26	-0.32	0.07	-0.34	0.07	-0.49
Unidentified	111	-0.52	0.02	-0.54	0.02	-0.56
Stars	56	-0.83	0.01	-0.88	0.01	-0.90
HBSS Sample:						
Extragalactic	65	-0.10	0.05	-0.27	0.05	-0.47
AGN 1	39	-0.44	0.03	-0.49	0.03	-0.53
AGN 2	16	0.14	0.11	0.05	0.11	-0.02
						0.45
						0.44

Columns are as follows: (1) Sample; (2) Number of sources; (3) HR 2 weighted average; (4) HR 2 unweighted average; (5) HR 2 median value; (6) observed standard deviation of the HR 2 distribution; (7) intrinsic dispersion of the HR 2 distribution computed using the maximum likelihood method as described in Macacaro et al., 1988.

3.4. Broad-band X-ray Spectral Properties

Combining the information on HR 2 and HR 3 we can now investigate in more detail the broad band spectral properties of the sample(s) as well as the selection function(s) of the BSS and HBSS.

The comparison is made in figure 5 where we show the position of the BSS (panels a, b and c) and HBSS (panel d) sources in the HR 2-HR 3 plane. We have used different symbols and panels to mark the spectroscopically identified and unidentified objects. Useful information can be extracted by cross-comparing the position of the different optical types of sources as well as by comparing the position of the sources in the BSS and HBSS samples.

The "bulk" of the sources optically identified as broad line AGNs are strongly clustered in the region between $\text{HR } 2 = 0.75$ and $\text{HR } 2 = 0.35$; this is true both for the broad line AGNs belonging to the BSS sample (by definition) and for those belonging to the HBSS sample (only 5 of 39 HBSS broad line AGN are outside these limits). The spread on HR 3 is much larger than the spread on HR 2 but note that HR 3 is much noisier than HR 2 since many of the sources are detected with poor statistics (or even undetected) in the 4.5-7.5 keV energy band.

All but 2 (~96%) of the sources classified as stars¹³ in the BSS sample have an HR 2 less than 0.75. If we assume a simple Rayleigh-Jeans thermal model, $\text{HR } 2 = 0.75$ corresponds to temperatures below 1.5 keV, in very good agreement with the identification as coronalemitting stars.

¹³ The two sources that are outside the "stellar" HR 2 locus are XBS J014100.6-675328 and XBS J215323.7+173018. The first object, XBS J014100.6-675328, has been already discussed in footnote 9. The XM M spectrum, discussed in Caccianiga et al., 2004, is well fitted by a unabsorbed power-law model with $kT = 1.53$ plus a thermal component having a temperature $kT = 55$ eV and an emission line at $E = 6.7$ keV. The second object (XBS J215323.7+173018) is a star having an XM M spectrum described by a thermal component with $kT = 1.6$ keV and a possible hard tail.

In Caccianiga et al., 2004, using a "pilot" 4.5-7.5 keV sample composed of 28 X-ray sources (26 of which in common with the current version of the HBSS sample reported in Table 4), we have already discussed the correlation between X-ray absorption, as deduced from a complete X-ray spectral analysis, and "Hardness Ratio" properties. In particular we have found that a) at the sampled fluxes, the 4.5-7.5 keV selection is picking up AGN having an intrinsic N_{H} up to few times 10^{23} cm^{-2} ; b) all the AGN having $\text{HR } 2 > 0.35$ are X-ray absorbed with N_{H} ranging from few times 10^{21} up to few times 10^{23} cm^{-2} . Assuming that this result is valid also for the sources presented here we can conjecture that all the sources having HR 2 greater than 0.35 are absorbed AGN; in this part of the diagram, besides narrow line AGN, we also note a few broad line AGN and 2 sources (one in the BSS sample and one in the HBSS sample) optically identified as normal galaxies. However also in these sources their point-like X-ray emission, their X-ray spectra (a preliminary spectral analysis shows that they are described by an absorbed power-law model having $N_{\text{H}} > 10^{22} \text{ cm}^{-2}$) and their intrinsic luminosity (in excess of $7 \times 10^{43} \text{ ergs s}^{-1}$ in the 2-10 keV energy range) strongly suggest the presence of an AGN. The existence of relatively luminous X-ray sources, optically identified with "normal galaxies", has been reported since the Einstein Observatory era in the early eighties (Elvis et al., 1981); this kind of sources were called in a variety of names such as optically dull galaxies (Elvis et al., 1981), passive galaxies (Gritths et al., 1995) and X-ray bright optically normal galaxies (XBONG, Comastri et al., 2002). We have already discussed in Severgnini et al. (2003) that detailed and specific optical-infrared follow-ups or higher-quality optical spectra are needed to unveil the AGN also in the optical domain. An advection-dominated accretion flow model has been recently used by Yuan and Narayan (2004) to explain their broad band properties.

Contrary to broad line AGNs and stars, narrow line (Type 2) AGNs seem to be distributed over a larger area




Fig. 4. HR2 vs. EPIC MOS2 count rate in the selection band for the sources in the BSS (panel a and panel b) and HBSS sample (panel c and panel d). We have also reported the HR2 expected from a unabsorbed power-law model with α_E ranging from -1 to 2 ($S_E / E^{-\alpha_E}$). The flux scale on the top has been computed assuming a conversion factor appropriate for $\alpha_E = 0.8$ (BSS sample) or $\alpha_E = 0.7$ (HBSS sample). We have used different symbols to mark the different kinds of objects. The eleven sources belonging only to the HBSS sample are enclosed inside empty squares in panels c and d.

in the HR2–HR3 plane with a well visible difference in the source position between the Type 2 AGNs in BSS and those in the HBSS sample. Although many of them have the hardest spectra amongst the identified objects, highly suggestive of intrinsic absorption (see also

Fig. 4), a new fact which seems to emerge from this investigation is the large number of narrow line AGN in the BSS sample occupying the locus typical of X-ray unabsorbed broad line AGN. Taking into account the still incomplete spectroscopic identification work and that some sources




Fig. 5. HR 2 vs. HR 3 for the sources belonging to the B SS sample (panels a, b and c) and to the H B SS sample (panel d). The dotted lines at constant HR 2 correspond to the locus enclosing 90% of the Type 1 AGN in the B SS sample; these lines have been reported in all panels to assist with the comparison(s). We have used different symbols to mark the spectroscopically identified and unidentified objects in the two samples. The eleven sources belonging only to the H B SS sample are enclosed inside empty squares in panel d. In the lower right corner of each panel we have also reported the median error on HR 2 and HR 3 for the total B SS and H B SS sample (solid line) and the 90% percentile on these errors (dotted line).

need a better quality optical spectrum we estimate that the relative fraction of these objects over the entire Type 2 AGN population may range between 50 and 75% in the case of the B SS sample and around 20% in the case of

the H B SS sample¹⁴. It is also worth noting that 2 out of 4 of the Type 2 AGN belonging to the (2–10 keV)

¹⁴ This latter estimate is consistent with the fact that in the H B S28 sample discussed in Caccianiga et al., 2004 there is only

HELLAS2XM M survey and having a good X-ray statistic (sample S1 in Perola et al., 2004) are characterized by an "observed" intrinsic N_{H} well below 10^{22} cm^{-2} .

To our knowledge two kinds of narrow line AGN could populate this zone, and thus could have X-ray spectral properties similar to those expected from unabsorbed AGN: a) "Compton thick" absorbed AGN (see e.g. the results presented in Della Ceca et al., 1999 from ASCA data); b) objects similar to the class of unabsorbed Seyfert 2 discussed in Pappa et al. (2001); Panessa and Bassani (2002) and in Barcons, Carrera and Ceballos (2003). Also the variability could play some role if the nucleus was bright at the time of the XM M -Newton observation but was turned off at the time of the optical spectroscopy. A detailed and exhaustive analysis of these sources is beyond the scope of the present paper. A deeper investigation of their optical (e.g. their optical classification between Seyfert 1.8, Seyfert 1.9 and Seyfert 2, analysis of the O [III] to 2-10 keV flux ratio) and X-ray (e.g. presence of Fe K emission lines to evaluate the Compton-thickness of the source) properties, as well as an assessment of the role played by selection effects, is in progress and will be presented elsewhere.

3.5. X-ray to optical flux ratio

A useful parameter to discriminate between different classes of X-ray sources is the X-ray to optical flux ratio (X/O flux ratio hereafter; see Maccacaro et al., 1988).

Previous investigations (see Fiore et al., 2003 and references therein) have shown that standard X-ray selected AGN (both type 1 and type 2) have typical X/O flux ratios in the range between 0.1 and 10 (for comparison standard optically selected QSOs and Seyfert 1 galaxies have X/O flux ratio > 1). X/O flux ratios below 0.1 are typical of coronal emitting stars, normal galaxies (both early type and star-forming) and nearby heavily absorbed (Compton thick) AGN. Finally at high X/O flux ratios (well above 10) we can find broad and narrow line AGN as well as high-z high-luminosity obscured AGN (type 2 QSOs), high-z clusters of galaxies and extreme BL Lac objects.

In this paper we have defined the X-ray to optical flux ratio using the observed X-ray flux in the 0.5-4.5 keV energy range and the optical red-band flux (see Fukugita et al., 1995 for the appropriate conversion factors).

For the large majority (73%) of the objects we have found and used APM red magnitudes. For 19% of the objects red magnitudes have been found from the literature, measured by us during spectroscopic runs or estimated using magnitudes measured in other optical bands. Finally for the unidentified objects with optical magnitudes below the POSSUM limit (30 sources) we have used an upper limit of $m_{\text{R}} > 21.0$. We note here that, for a fixed X-ray flux, an error of 1 magnitude corresponds to an uncertainty

one Type 2 AGN with HR2 between 0.75 and 0.35 out of 8 Type 2 objects.

of 60% on the X/O flux ratio; this uncertainty does not affect any of the general conclusions we discuss below.

In figure 6 we have plotted the X/O flux ratio versus the HR2 value for each source (BSS sample in panels a and b; HBSS sample in panel c). The boxes defined by the dot-dashed lines (dashed line) indicate the locus of coronalem itting stars (broad line AGN) in the BSS sample and have been reported in all panels to assist with the comparisons.

As is clearly visible in figure 6 the bulk of coronalem itting stars is well separated from the bulk of extragalactic sources; some of the AGN (both broad and narrow line) have X/O flux ratio typical of stars but HR2 values typical of AGN; similarly some AGN with an HR2 typical of stars can be distinguished from stars thanks to their X/O flux ratio. Therefore the combined use of X/O flux ratio and HR2 allows us to distinguish almost unequivocally galactic sources from the extragalactic ones.

A round 10% of the extragalactic population have a X/O flux ratio greater than 10. If we consider the 2-10 keV fluxes instead of the 0.5-4.5 keV fluxes this fraction increases to 15%, in good agreement with the results obtained by Fiore et al., 2003 at fainter fluxes. Amongst the sources identified so far at X/O flux ratio > 10 there are some broad line and a few narrow line AGN. These results are consistent with those reported in Fiore et al., 2003 who also discuss the observational evidence that at X/O flux ratio greater than 10 is where to look for type 2 QSOs. Since the large majority of the X-ray sources at X/O flux ratio > 10 presented here is still unidentified we can not comment further on this; we can only note that many of the unidentified objects with high or very high X/O flux ratio seem to have rather standard hardness ratios.

The opposite side of the extragalactic X/O distribution (X/O flux ratio < 0.1) is populated by optically normal galaxies, Type 2 AGNs and a few broad line AGN. As discussed above this part of the X/O distribution could be also populated by nearby Compton thick AGN; in this respect it is worth noting that a few of the type 2 AGN in the BSS sample populating the same HR2 region of the broad line AGN population have X/O flux ratios < 0.1 . In the HBSS sample there are also a few Type 2 AGN having apparently hard spectra but low X/O flux ratios (< 0.1).

3.6. The number densities of optically broad and narrow line AGN

The measured X-ray $\log(N > S) - \log S$ of broad and narrow line AGN is a fundamental observable for cosmological investigations and, in particular, provides very strong constraints on the CXB synthesis models and on the history of the accretion in the Universe.

As discussed above the spectroscopic identification rate of the HBSS sample is 90%, allowing us to investigate for the first time the X-ray $\log(N > S) - \log S$ of optically broad and narrow line AGN in the same sample.




Fig. 6. HR2 vs. X-ray (0.5–4.5 keV) to optical flux ratio for the sources belonging to the BSS sample (panels a and b) and to the HBSS sample (panel c). The X-ray (0.5–4.5 keV) to optical flux ratio for each source has been computed as discussed in section 3.5. We have used different symbols to mark the spectroscopically identified and unidentified objects in the two samples. The box defined by the dot-dashed lines (dashed line) enclose about 95% (85%) of the sources optically identified as coronalemmitting stars (broad-line AGNs) in the BSS sample; these boxes have been reported in all panels to assist with the comparison(s) discussed in the text. The eleven sources belonging only to the HBSS sample are enclosed inside empty squares in panel c.

In order to deal with the 7 unidentified HBSS sources we have made the reasonable assumption that the 5 unidentified sources having $HR2 > 0.35$ are Type 2 AGN, while the 2 unidentified sources having $HR2$ in the range from 0.75 to 0.35 are Type 1 AGN (see section 3.3 for details).

The 4.5{7.5 keV Log(N > S)-LogS of optically broad and narrow line AGN are reported in figure 7 (AGN 1: open circles; AGN 2: filled circles). A conversion factor appropriate for a power-law spectral model with $\alpha = -0.7$ has been used to obtain the fluxes. Both Log(N > S)/LogS can be well described by a power-law model and their linear

minimum likelihood best fit parameters have been reported in Table 5.

At the 4.5{7.5 keV flux limit of $f_{\text{lim}} = 7 \times 10^{-14}$ erg cm $^{-2}$ the surface densities of optically type 1 AGN and type 2 AGN are 1.63 ± 0.25 deg $^{-2}$ and 0.83 ± 0.18 deg $^{-2}$, respectively. Optically type 2 AGN represent 34 ± 9% of the AGN population shining in the 4.5{7.5 keV energy selection band at the flux limit of the HBSS. In Table 8 we compare the optically type 2 AGN fraction from the HBSS sample with that found in a few representative X-ray surveys. As expected from the CXB synthesis models the fraction of optically type 2 AGN decreases going to




Fig. 7. The number-flux relationship in the 4.5–7.5 keV energy band for optically type 1 (open circles) and type 2 (filled circles) AGN obtained using the sources in the HBSS sample.

softer X-ray surveys (e.g. by a factor 2.6 in the XMM-BSS survey and by a factor 34 in the ROSAT Bright Survey) while remaining almost constant (around 35%) for surveys above 2 keV down to a flux limit of 10^{14} ergs $\text{cm}^{-2} \text{s}^{-1}$. In the flux range between 10^{15} and 10^{14} ergs $\text{cm}^{-2} \text{s}^{-1}$ there is apparently an increase of the fraction of optically type 2 AGN but the spectroscopic identification rate in this flux range is still low (~50%) preventing us from speculating further.

The optically type 2 AGN fraction from the HBSS sample is also in very good agreement with the fraction of X-ray absorbed ($N_{\text{H}} > 10^{22} \text{ cm}^{-2}$) AGN found from several studies (e.g. Piconcelli et al., 2003; Perola et al., 2004) which ranges between 25% to 40% for hard X-ray fluxes spanning four orders of magnitude, from 10^{10} to 10^{14} ergs $\text{cm}^{-2} \text{s}^{-1}$. While we anticipate that the Mardioli et al. (2003) prediction scheme for the synthesis of the CXB (Ueda et al., 2003) predicts a fraction of absorbed AGN around 41% for the fluxes (and energy band) covered by the HBSS, a proper comparison will have to wait for the completion of the X-ray spectroscopic work.

4. Summary and Conclusions

In this paper we have discussed the scientific goals and the strategy of the 'The XMM-Newton Bright Serendipitous Survey'. This survey comprises two flux-limited samples: the BSS sample and the HBSS sample having a flux limit

of 7×10^{14} erg cm^{-2} in the 0.5–4.5 keV and 4.5–7.5 keV energy band, respectively.

From the analysis of 237 suitable XMM-Newton fields (211 for the BSS), covering a useful survey area of 28.10 (25.17 for the HBSS) sq. deg of sky, we have defined (and presented here) a sample of 400 X-ray sources: 389 of them belong to the BSS, 67 to the HBSS with 56 X-ray sources in common. Up to now 71% (90%) of the sources in the BSS (HBSS) sample have been spectroscopically identified, either from the literature or from our spectroscopic observations. The main results from this study can be summarized:

a) the extragalactic number-flux relationship in the 0.5–4.5 keV and 4.5–7.5 keV band are in good agreement with previous and new results. They are well described by a power law model, $N \propto S^{\alpha}$, with best-fit values for the slope of 1.80 ± 0.11 and 1.64 ± 0.24 in the 0.5–4.5 keV and 4.5–7.5 keV bands, respectively;

b) at the X-ray flux limits of the survey presented here we found that the optical counterpart in them a majority (~90%) of cases has an optical magnitude brighter than the POSSUM limit ($R = 21^m$ mag). Galactic counterparts represent about 14% of the sources in the BSS sample and less than 3% of the sources in the HBSS sample. The majority of the extragalactic objects identified so far are broad-line AGN both in the BSS (~80%) and in the HBSS (~60%);

c) we have investigated the broad-band spectral properties of the selected sources using hardness ratios. No obvious trend of the source spectra as a function of the count rate is measured. The average spectrum of the 'extragalactic' population corresponds to a (0.5–4.5 keV) energy spectral index of 0.8 for the BSS sample and 0.64 for the HBSS sample. About 13% (40%) of the sources in the BSS (HBSS) sample seem to be described by an energy spectral index steeper than that of the CXB; following the results presented in Caccianiga et al., 2004 we speculate that these sources are absorbed AGN with N_{H} ranging from few times 10^{21} up to few times 10^{23} cm^{-2} . There are hints from this study of a significant number of narrow-line AGN in the BSS sample occupying the locus typical of X-ray unabsorbed broad-line AGN. A deeper investigation of their optical and X-ray properties, of the source selection function, as well as a complete X-ray spectral analysis for all the BSS and HBSS sources using data from all the EPIC instruments (a major goal of this project) is in progress and will be presented elsewhere;

d) we do not find a compelling evidence that the HBSS 4.5–7.5 keV survey is sampling a completely different source population compared with the BSS 0.5–4.5 keV survey. Rather we find that the HBSS survey is simply picking up a larger fraction of absorbed AGN, consistent with CXB synthesis models based on the unification scheme of the AGN;

e) at the flux limit of the HBSS sample we measure surface densities of optically classified type 1 and type 2 AGN of $1.63 \pm 0.25 \text{ deg}^{-2}$ and $0.83 \pm 0.18 \text{ deg}^{-2}$, respectively. The AGN optically classified as Type 2 represent (34–9)% of the total AGN population shining in the

Table 8. Optically Type 2 AGN fraction in few representative X-ray surveys.

Sample (1)	Energy Range (2)	Flux Range (3)	Type 2 Fraction (4)	Ref./Note (5)
ROSAT Bright Survey	0.5{2.0	> 2 10 ⁻¹²	1 0.5%	(a)
XMM-BSS	0.5{4.5	> 7 10 ⁻¹⁴	13 4%	(b)
XMM-HBSS	4.5{7.5	> 7 10 ⁻¹⁴	34 9%	(c)
HELLAS2XMM + Other Surveys	2{10 10 ⁻¹⁵	10 ⁻¹⁴	57 12%	(d)
HELLAS2XMM + Other Surveys	2{10 10 ⁻¹⁴	10 ⁻¹³	37 6%	(e)

Columns are as follows: (1) Sample; (2) Energy Selection Band; (3) Investigated flux range; (4) Type 2 AGN fraction over the total AGN density in the specific survey; (5) References and Note as follows: (a) From Schwope et al., 2000; (b) From this paper using the BSS sources with Right Ascension below 5^h or above 17^h. For this BSS subsample the spectroscopic identification rate is around 88%; (c) From this paper using the HBSS sample and taking into account the small number of sources still unidentified as discussed in section 3.6; (d) Fraction of optically obscured AGN from Fiore et al., 2003. The spectroscopic ID rate of the sample in this flux range is 50%; (e) Fraction of optically obscured AGN from Fiore et al., 2003. The spectroscopic ID rate of the sample in this flux range is 81%.

4.5–7.5 keV energy band at the sampled fluxes. A proper comparison with X-ray absorbed/unabsorbed AGN have to wait for the completion of the ongoing X-ray spectroscopic work.

Finally, we found a clear separation between Galactic "coronalemitting" stars and extragalactic sources in the hardness ratio diagram and in the (hardness ratio) vs. (X/O flux ratio) diagram. Since the investigated sample is a fair representation of the high Galactic latitude X-ray sky, this result will help with the selection of defined classes of sources from the XMM-Newton catalogue prior to spectroscopic observations making the existing and incoming XMM-Newton catalogs an unique resource for astrophysical studies.

Acknowledgments. This research has made use of the NASA/IPAC Extragalactic Database (NED; which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration) and of the SIMBAD database (operated at CDS, Strasbourg, France). We thanks Piero Rosati for providing us someIDL routines used in the data analysis and preparation. We thanks L. Maraschi, G. Trinchieri, A. Wolter, D. Worrall, N. Webb and the referee for a careful reading of the paper and for useful comments which have improved the paper. We thanks Y. Ueda to have provided us with the predictions from the Modelled Unification Model in a tabular form. RDC, TM, AC, PS, VB acknowledge partial financial support by the Italian Space Agency (ASI grants: I/R/037/01, I/R/062/02 and I/R/071/02), by the MURST (Co n-03-02-23) and INAF. XB and FJC acknowledge financial support from the Spanish Ministerio de Ciencia y Tecnología, under the project ESP2003-00812. Based on observations collected at the Italian Telescopio Nazionale Galileo" (TNG), at the German-Spanish Astronomical Center, Calar Alto (operated jointly by Max-Planck Institut für Astronomie and Instituto de Astrofísica de Andalucía, CSIC), at the European Southern Observatory (ESO) and at the Nordic Optical Telescope (NOT). We would like to thank the members of the TNG, ESO, Calar Alto and NOT for their support during the observations. The TNG and NOT telescopes are operated in the island of La Palma by the Nordic Optical Telescope Scientific Association and the Centro Galileo

Galilei of the INAF, respectively, in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. We finally thank the APM team for maintaining this facility.

References

- Alexander, D. M., et al. 2003, *AJ*, 126, 539
- Baldi, A., Molendi, S., Comastri, A., et al. 2002, *ApJ*, 564, 190
- Barcons, X., Carrera, F. J., & Ceballos, M. T. 2003, *MNRAS*, 339, 757
- Barcons, X., Carrera, F. J., Watson, M. G., et al. 2002, *A&A*, 382, 522
- Brandt, W. N., Alexander, D. M., Homscheid, A. E., et al. 2001, *AJ*, 122, 2810
- Brandt, W. N., Alexander, D. M., Bauer, F. E., & Vignali, C. 2004, in Physics of Active Galactic Nuclei at All Scales, eds. Alloin D., Johnson R., Lira P. (Springer-Verlag, Berlin), astro-ph/0403646
- Caccianiga, A., Severgnini, P., Braito, V., et al. 2004, *A&A*, 416, 901
- Comastri, A., Setti, G., Zamorani, G., Hasinger, G. 1995, *A&A*, 296, 1
- Comastri, A., Fiore, F., Vignali, C., Matt, G., Perola, G. C., La Franca, F. 2001, *MNRAS*, 327, 781
- Comastri, A., et al. 2002, *ApJ*, 571, 771
- Della Ceca, R., Castelli, G., Bratto, V., Cagnoni, I., & Macacaro, T. 1999, *ApJ*, 524, 674
- Dickey, J. M. & Lockman, F. J. 1990, *ARA&A*, 28, 215
- Ehle, M., Breitfellner, M., Gonzales Riestra, R., et al. 2003, XMM-Newton Users' Handbook, Issue 2.1
- Elvis, M., Schreier, E. J., Tonry, J., Davis, M., & Huchra, J. P. 1981, *ApJ*, 246, 20
- Fiore, F., Giommi, P., Vignali, C., Comastri, A., Matt, G., Perola, G. C., La Franca, F., Molendi, S., Tamburelli, F., Antonelli, L. A. 2001, *MNRAS*, 327, 771
- Fiore, F., Brusa, M., Cocchia, F., et al. 2003, *A&A*, 409, 79
- Fukugita, M., Shimasaki, K., & Ichikawa, T. 1995, *PASP*, 107, 945
- Galbiati, E., Caccianiga, A., Macacaro, T., et al. 2004, *A&A*, submitted.
- Georgantopoulos, I., Georgakakis, A., Akylas, A., Stewart, G. C., Giannakis, O., Shanks, T., & Kitsonas, S. 2004, *MNRAS*, 352, 91.

- Gilli, R., Salvati, M., Hasinger, G. 2001, A & A, 366, 407
- Gioia, I. M., Maccacaro, T., Schild, R. E., Wolter, A., Stocke, J. T., Morris, S. L., & Henry, J. P. 1990, ApJS, 72, 567
- Grihs, R. E., Georgantopoulos, I., Boyle, B. J., Stewart, G. C., Shanks, T., & della Ceca, R. 1995, MNRAS, 275, 77
- Hasinger, G., Burg, R., Giacconi, R., Hartner, G., Schmidt, M., Trumper, J., & Zamorani, G. 1993, A & A, 275, 1
- Hasinger, G., Burg, R., Giacconi, R., Schmidt, M., Trumper, J., & Zamorani, G. 1998, A & A, 329, 482
- Hasinger, G., Almaini, B., Arnaud, M., et al. 2001, A & A, 365, L45
- Landt, H., Padovani, P., & Giommi, P. 2002, MNRAS, 336, 945
- Maccacaro, T. et al. 1982, ApJ, 253, 504
- Maccacaro, T., Gioia, I. M., Wolter, A., Zamorani, G., & Stocke, J. T. 1988, ApJ, 326, 680
- Madau, P., Ghisellini, G., Fabian, A. C. 1994, MNRAS, 270, 17
- Malinieri, V., Bergeron, J., Hasinger, et al. 2002, A & A, 393, 425
- Maretti, A., Campana, S., Lazzati, D., Tagliaferri, G. 2003, ApJ, 588, 696
- Nandra, K., Georgantopoulos, I., Ptak, A., Turner, T. J. 2003, ApJ, 582, 615
- Panessa, F. & Bassani, L. 2002, A & A, 394, 435
- Pappa, A., Georgantopoulos, I., Stewart, G. C., & Zelaz, A. L. 2001, MNRAS, 326, 995
- Perola, G. C., et al. 2004, A & A, 421, 491.
- Piconcelli, E., Cappi, M., Bassani, L., DiCocco, G., Dadina, M. 2003, A & A, 412, 689
- Piconcelli, E., Cappi, M., Bassani, L., Fiore, F., DiCocco, G., Stephen, J. B. 2002, A & A, 394, 835
- Rosati, P., Tozzi, P., Giacconi, R., et al. 2002, ApJ, 566, 667
- Schwope, A. et al. 2000, Astronomische Nachrichten, 321, 1
- Severgnini, P., Caccianiga, A., Bracco, et al. 2003, A & A, 406, 483
- Setti, G. & Wolter, L. 1989, A & A, 224, 21
- Ueda, Y., Akiyama, M., Ohta, K., & Miyaji, T. 2003, ApJ, 598, 886
- Veron-Cetty, M.-P. & Veron, P. 2001, A & A, 374, 92
- Yuan, F., & Narayan, R. 2004, ApJ, in press, astro-ph/0401117

Online Material

Table 2. Basic information on the XMM-Newton MOS2 fields used for the sample selection

Obs. ID	Filter	RA ; DEC (J2000)		Exposure s	Log N h cm ⁻²	R _{in} arcmin	R _{out} arcmin	BSS srcts	H BSS srcts
(1)	(2)	(3)		(4)	(5)	(6)	(7)	(8)	(9)
0125310101	Medium	00 00 30.4	25 06 43.4	19162.6	20.27	1	13	4	0
0101040101	Medium	00 06 19.7	+20 12 22.8	34044.7	20.60	8	13	1	0
0127110201	Thin1	00 10 31.2	+10 58 40.7	7558.0	20.76	2	13	2	0
0111000101	Medium	00 18 33.0	+16 26 08.0	31362.1	20.61	3	13	2	0
0001930101	Thin1	00 26 06.8	+10 41 12.5	18032.3	20.71	1	13	1	1
0050140201	Thin1	00 26 35.9	+17 09 37.8	50380.4	20.63	3	13	2	0
0112320101	Thin1	00 30 27.6	+04 51 47.4	24577.8	20.48	1	13	1	0
0065770101	Medium	00 32 47.0	+39 34 33.3	7360.0	20.82	1	13	1	0
0125920201	Thin1	00 33 54.5	12 07 10.8	9709.9	20.40	1	13	3	0
0125320701	Thin1	00 50 03.0	52 07 36.7	16524.9	20.51	1	13	3	—
0110890401	Medium	00 57 20.0	22 23 04.1	29741.8	20.19	8	13	0	0
0111280301	Thin1	00 58 51.3	27 35 47.1	7975.6	20.28	0	12	1	0
0112650401	Medium	01 04 24.0	06 24 10.6	23695.9	20.71	0	13	1	0
0103861601	Thick	01 05 16.4	58 26 10.4	7127.1	20.44	8	13	1	0
0025540101	Medium	01 07 46.5	17 30 14.0	7678.2	20.18	2	13	2	0
0113040801	Thin1	01 20 39.7	10 56 24.5	5692.0	20.51	1	13	4	—
0101040201	Medium	01 23 46.0	58 48 25.8	28994.1	20.50	8	13	0	0
0109860101	Thin1	01 25 33.4	+01 45 38.2	38533.6	20.49	4	13	2	0
0112600601	Medium	01 27 32.1	+19 10 32.3	3994.2	20.68	8	13	3	—
0084230301	Medium	01 31 53.7	13 36 56.4	20446.4	20.21	8	13	1	1
0112630201	Thin1	01 33 01.9	40 06 23.5	31094.2	20.28	1	13	1	0
0111430101	Medium	01 39 01.2	17 57 06.3	33440.3	20.16	8	13	1	0
0032140401	Medium	01 40 12.1	67 48 40.3	7610.7	20.40	0	13	4	2
0093641001	Medium	01 43 01.7	+13 38 24.4	9329.1	20.69	4	13	2	0
0101640201	Medium	01 59 50.2	+00 23 46.8	7543.3	20.41	8	13	3	1
0084140101	Medium	02 08 38.2	+35 23 00.2	38001.6	20.80	2	13	3	0
0112371701	Thin1	02 17 12.4	04 39 04.2	20001.1	20.38	0	11	2	1
0112371001	Thin1	02 18 00.3	04 59 47.8	50800.3	20.40	0	11	4	3
0112371501	Thin1	02 18 48.3	04 39 02.9	6684.2	20.40	0	11	0	—
0112370301	Thin1	02 19 36.3	04 59 47.7	50384.1	20.41	0	11	1	0
0111110501	Thin1	02 22 40.0	05 09 49.0	24185.8	20.41	0	10	0	0
0109520501	Thin1	02 23 20.0	04 49 49.3	24107.8	20.42	0	10	1	0
0111110401	Thin1	02 24 00.1	05 09 53.6	28276.2	20.42	0	10	1	0
0111110301	Thin1	02 25 20.3	05 09 52.5	24158.7	20.43	0	10	0	0
0111110201	Thin1	02 26 40.1	05 09 47.1	8229.8	20.42	0	10	1	0
0111110101	Thin1	02 27 60.0	05 09 48.4	23809.9	20.42	0	10	0	0
0103861101	Medium	02 35 12.8	29 36 09.9	7409.2	20.25	8	13	1	0
0098810101	Thin1	02 36 12.3	52 19 55.6	23450.3	20.52	2	13	2	1
0075940301	Medium	02 36 57.9	+24 38 53.7	47426.5	20.96	8	13	0	0
0067190101	Medium	02 38 19.4	52 11 34.1	25627.4	20.50	3	10	1	0
0111200101	Medium	02 42 41.1	00 00 53.7	35485.0	20.55	8	13	4	0
0111490401	Medium	02 48 43.8	+31 06 59.2	31028.6	20.97	8	13	0	0
0056020301	Thin1	02 56 32.8	+00 06 01.7	17250.5	20.81	2	13	2	0
0041170101	Thin1	03 02 38.5	+00 07 31.9	47275.7	20.85	0	13	2	1
0042340501	Thin1	03 07 03.8	28 40 24.1	13285.0	20.13	4	13	2	1
0122520201	Thick	03 11 59.3	76 51 53.0	28295.0	20.91	2	13	2	1
0110970101	Thin1	03 13 09.6	55 03 48.4	10239.4	20.41	0	13	2	1
0110970401	Thin1	03 14 15.9	55 21 33.5	8457.5	20.42	0	8	0	0
0129320901	Thin1	03 16 03.7	55 06 08.5	9948.2	20.43	0	13	1	0
0105660101	Medium	03 17 56.0	44 14 15.1	23150.7	20.41	6	13	2	1
0108060501	Thin1	03 32 29.1	27 48 27.1	46833.4	19.95	0	13	2	0
0099010101	Thin1	03 35 27.6	25 44 54.5	18776.9	20.01	8	13	3	0

Obs. ID	Filter	RA ; DEC (J2000)	Exposure s	Log N h cm ⁻²	R _{in} arcmin	R _{out} arcmin	BSS srcts	H BSS srcts
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
0055140101	Thin1	03 39 35.0 35 25 58.1	47634.6	20.12	1	11	3	1
0111970301	Thin1	04 09 07.2 71 17 43.9	18740.3	20.88	8	13	3	3
0112600401	Medium	04 25 44.2 57 13 34.4	7627.0	20.31	8	13	1	0
0103861701	Medium	04 35 17.1 78 01 54.3	8010.5	20.88	8	13	1	0
0112880401	Thick	04 59 35.4 + 01 47 16.0	18914.3	20.89	2	13	2	0
0111160201	Thin1	05 05 20.0 28 48 57.6	47984.4	20.17	2	13	4	1
0103860701	Medium	05 16 21.3 10 33 34.5	4740.3	20.97	8	13	1	—
0094400101	Medium	05 16 40.1 + 79 40 16.7	31727.2	20.90	2	13	4	0
0090050701	Thin1	05 19 49.4 45 46 48.9	13426.7	20.62	8	13	2	0
0085640101	Thin1	05 21 00.9 25 21 52.6	9720.6	20.28	1	13	5	2
0050150101	Thin1	05 25 05.9 33 43 13.3	13440.2	20.35	1	13	2	0
0123720301	Medium	05 28 44.1 65 26 58.6	4762.6	20.73	8	13	0	—
0009220601	Medium	06 15 35.2 + 71 02 12.2	11376.9	20.92	2	13	1	0
0110930101	Medium	06 18 41.9 + 78 21 25.7	19578.2	20.86	5	13	0	0
0103860101	Medium	06 23 08.8 64 36 15.8	8862.8	20.73	8	13	2	0
0029340101	Medium	06 41 18.4 + 82 15 59.3	22288.1	20.70	8	13	1	0
0061540101	Medium	06 52 11.4 + 74 25 30.1	21080.3	20.81	2	13	3	0
0112980201	Thin1	06 58 17.6 55 57 30.6	41042.1	20.81	8	13	2	0
0083000101	Medium	07 35 19.9 + 43 44 27.3	13589.8	20.77	2	13	0	0
0103862101	Medium	07 42 32.8 + 49 48 30.4	5909.3	20.75	8	13	1	—
0123100101	Thin1	07 44 03.3 + 74 33 42.9	39171.0	20.55	3	13	4	2
0111100301	Thin1	07 51 09.3 + 18 07 43.0	32111.2	20.61	1	13	1	0
0110070401	Thin1	07 55 05.0 + 22 00 04.6	22449.2	20.75	8	13	0	0
0094400301	Medium	08 04 34.3 + 64 59 55.6	19077.9	20.64	2	13	2	1
0094530401	Thin1	08 05 35.2 + 24 49 56.7	8709.6	20.59	0	13	2	0
0092800101	Medium	08 31 42.4 + 52 45 20.8	16808.2	20.59	2	13	1	0
0025540301	Medium	08 38 22.8 + 25 45 13.9	11803.5	20.56	1	13	4	2
0111400101	Thick	08 39 11.6 + 65 01 12.1	24224.9	20.63	8	13	1	0
0112620101	Medium	08 41 22.9 + 70 53 37.8	4301.9	20.47	8	13	0	—
0103660201	Medium	08 47 42.4 + 34 45 07.3	16223.3	20.52	8	13	1	0
0085030101	Thin1	08 55 35.7 + 58 44 05.8	6545.6	20.62	5	13	2	—
0110660201	Medium	09 06 02.5 + 62 03 02.2	8671.8	20.65	0	13	1	0
0083240201	Thin1	09 11 28.2 + 05 50 57.3	19349.7	20.56	2	13	1	0
0084230601	Medium	09 17 52.1 + 51 43 33.1	16347.4	20.16	6	13	1	1
0021740101	Medium	09 39 53.7 + 35 54 06.4	10323.5	20.16	2	13	0	0
0106460101	Thin1	09 43 01.2 + 46 59 34.4	49423.9	20.10	3	13	0	0
0017540101	Medium	09 45 25.9 08 39 03.2	7573.9	20.56	8	13	2	0
0111290101	Thin1	09 50 48.7 + 39 26 59.8	21359.3	20.20	8	13	1	0
0065790101	Thin1	09 52 09.1 01 48 14.6	9322.6	20.60	2	13	2	1
0070940101	Thin1	09 53 41.5 + 01 34 55.9	6021.1	20.55	5	13	2	—
0112850101	Thin1	09 54 56.3 + 17 43 29.7	17345.1	20.50	1	13	2	0
0111290201	Thin1	09 56 52.6 + 41 15 24.9	12896.7	19.90	8	13	1	0
0041170201	Medium	10 00 40.8 + 25 14 24.8	46751.9	20.44	0	13	3	0
0110930201	Medium	10 01 56.9 + 55 40 41.1	7317.1	19.90	2	13	2	0
0070340201	Thin1	10 08 47.1 + 53 42 00.8	21851.2	19.87	1	13	3	0
0085170101	Thin1	10 11 13.5 + 55 44 55.0	31755.9	19.89	2	13	0	0
0086750101	Thin1	10 16 15.4 + 52 09 24.1	6574.0	19.88	2	13	3	—
0028740301	Thin1	10 18 40.5 + 41 25 15.2	28448.5	20.05	8	13	4	2
0093640301	Medium	10 19 59.8 + 08 13 32.3	12804.9	20.48	1	13	2	0
0101040301	Medium	10 23 30.7 + 19 51 53.1	36611.7	20.33	4	13	1	0
0108670101	Thin1	10 23 39.9 + 04 11 25.6	50918.8	20.48	8	13	2	0
0102040301	Thick	10 30 59.0 + 31 02 51.8	26293.5	20.29	8	13	2	0
0110950101	Thin1	10 31 54.2 14 16 52.5	7308.6	20.82	8	13	0	0
0112810301	Thin1	10 38 44.4 + 53 30 08.4	4917.1	20.05	3	13	2	—

Obs. ID	Filter	RA ; DEC (J2000)	Exposure s	Log N h cm ⁻²	R _{in} arcmin	R _{out} arcmin	BSS srcts	H BSS srcts
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
0059800101	Thin1	10 39 53.7 + 20 50 52.2	14293.2	20.31	2	13	5	1
0125300101	Thin1	10 44 32.4 01 25 05.3	32227.5	20.62	2	13	4	1
0055990201	Thin1	10 49 53.6 + 32 58 58.1	19840.1	20.30	4	13	2	1
0022740101	Medium	10 52 44.1 + 57 28 48.7	82880.4	19.75	0	13	5	0
0094800101	Thin1	10 56 59.5 03 37 29.8	31924.0	20.55	2	13	1	0
0110660301	Thin1	11 01 08.4 + 86 09 59.2	6216.2	20.83	1	13	0	—
0112880201	Medium	11 01 52.3 34 42 27.4	28570.1	20.70	2	13	3	1
0070340301	Medium	11 03 36.6 + 35 55 11.8	8386.3	20.30	1	13	1	—
0112630101	Thin1	11 06 33.6 18 21 35.1	12627.6	20.66	1	13	1	0
0099030101	Thin1	11 17 18.6 + 17 57 50.3	20091.0	20.18	1	13	1	0
0111290401	Thin1	11 19 08.8 + 21 19 17.4	8525.8	20.11	8	13	2	0
0107860201	Thin1	11 20 06.3 + 43 17 59.9	22162.9	20.32	1	13	0	1
0093641101	Medium	11 20 15.0 + 12 59 29.1	10566.2	20.39	5	13	3	0
0112270301	Thin1	11 22 21.7 + 24 17 39.2	14296.9	20.10	6	13	0	0
0110660401	Medium	11 26 00.6 + 42 53 05.1	13085.5	20.32	1	13	0	0
0112810101	Thin1	11 28 29.1 + 58 33 43.7	20593.3	20.00	3	13	0	0
0102040201	Thick	11 31 09.2 + 31 14 08.0	23552.4	20.31	8	13	3	3
0042341001	Thin1	11 31 56.2 19 55 32.8	12683.8	20.65	6	13	1	0
0112210101	Medium	11 39 02.3 37 44 17.2	37360.0	20.93	4	13	1	0
0094800201	Thin1	11 40 23.9 + 66 08 37.6	24709.2	20.09	1	13	0	0
0112551401	Thin1	11 52 54.0 + 36 59 06.6	6471.1	20.28	4	13	1	—
0090020101	Thin1	11 57 55.3 + 55 27 07.8	10362.7	20.09	3	13	1	0
0109141401	Medium	12 03 09.4 + 44 31 52.4	62156.3	20.12	8	13	2	0
0112310101	Medium	12 10 32.3 + 39 24 19.8	30192.1	20.30	8	13	0	0
0112610101	Medium	12 14 17.0 + 14 03 16.9	51088.9	20.44	8	13	1	0
0111840101	Medium	12 21 22.2 + 30 10 38.1	9637.0	20.24	4	13	0	—
0124110101	Medium	12 21 41.5 + 75 18 38.0	17009.1	20.46	2	13	2	0
0108860101	Thin1	12 23 07.2 + 10 37 15.1	20960.4	20.35	4	13	0	0
0070340501	Thin1	12 27 15.6 + 33 32 18.9	15390.5	20.14	1	13	4	0
0110990201	Thin1	12 27 18.4 + 01 29 20.2	11954.5	20.26	1	13	2	1
0126700301	Medium	12 29 06.6 + 02 03 08.0	56647.6	20.25	8	13	2	0
0112650101	Medium	12 31 11.4 + 20 47 52.2	23562.4	20.30	0	13	0	0
0124900101	Thin1	12 31 30.9 + 64 14 20.0	33644.1	20.30	2	13	4	0
0006220201	Thick	12 35 34.9 39 54 55.9	42741.5	20.86	1	13	2	1
0111550101	Thin1	12 36 48.5 + 62 12 56.4	43468.9	20.17	0	13	3	0
0109970101	Medium	12 39 39.1 05 20 40.3	8087.8	20.36	4	13	0	0
0136950201	Thin1	12 42 37.9 11 19 22.4	29166.8	20.56	1	13	1	0
0111190701	Medium	12 42 48.1 + 02 41 17.8	59825.8	20.26	8	13	0	0
0051760101	Thin1	12 46 34.9 + 02 22 01.1	12244.0	20.24	2	13	4	1
0060370201	Thin1	12 49 13.5 05 59 27.5	40156.8	20.33	1	13	4	0
0008220201	Thick	12 51 41.5 + 27 32 25.3	39747.7	19.95	8	13	0	0
0111020101	Medium	12 53 02.9 29 23 41.4	29008.5	20.78	8	13	0	0
0081340201	Medium	12 56 13.6 + 56 52 24.0	21493.1	20.10	1	13	2	0
0017940101	Thin1	13 05 43.5 + 18 00 59.0	50660.2	20.32	8	13	1	0
0002940101	Medium	13 07 06.6 23 40 27.2	7263.6	20.96	1	13	2	0
0093640401	Medium	13 20 35.4 + 34 08 21.4	19663.3	20	1	13	4	1
0100240201	Medium	13 30 47.4 + 24 13 59.1	35954.0	20.06	3	13	3	0
0061940101	Medium	13 31 52.9 + 11 16 43.9	7011.7	20.29	1	13	1	0
0111160101	Thin1	13 34 19.2 + 50 30 55.6	47144.5	20.04	2	13	1	0
0111570201	Medium	13 35 53.9 34 17 44.3	50435.3	20.61	6	13	1	0
0096010101	Thin1	13 37 18.8 + 24 23 02.3	32367.8	20.06	8	13	1	0
0035940301	Medium	13 39 55.8 31 38 30.3	42378.9	20.59	5	13	1	1
0112250201	Thin1	13 47 40.0 + 58 12 45.2	33008.0	20.11	1	13	2	2
0098010101	Thin1	14 01 02.0 + 02 52 44.3	32776.7	20.37	5	13	1	2

Obs. ID	Filter	RA ; DEC (J2000)	Exposure s	Log N h cm ⁻²	R _{in} arcmin	R _{out} arcmin	BSS srcts	H BSS srcts
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
0109910101	Thin1	14 01 34.5 11 07 40.3	48892.6	20.63	6	13	3	1
0092850501	Thin1	14 09 23.9 + 26 18 21.4	41167.0	20.17	2	13	2	0
0013140101	Medium	14 13 14.9 03 12 25.5	17838.3	20.58	8	13	1	0
0112250301	Thin1	14 15 45.6 + 11 29 35.7	25593.4	20.26	2	13	2	1
0127921001	Thin1	14 17 10.3 + 52 23 54.1	54477.7	20.11	0	13	3	0
0109960101	Medium	14 17 59.2 + 25 08 11.7	23142.3	20.23	4	13	3	1
0111850201	Medium	14 28 32.5 + 42 40 16.4	39671.3	20.14	4	13	2	1
0111530101	Thick	14 38 59.9 + 64 17 25.5	50080.5	20.23	8	13	3	2
0057560301	Thin1	14 49 29.2 + 08 59 51.1	38812.8	20.31	0	13	2	0
0056030101	Medium	14 51 22.8 + 19 06 02.7	58940.0	20.39	8	13	0	0
0067750101	Thin1	14 58 21.3 - 31 40 09.0	34424.0	20.93	2	13	1	0
0070740301	Thin1	15 04 01.1 + 10 26 15.0	10625.1	20.37	8	13	1	0
0018741001	Thin1	15 18 46.4 + 06 14 10.4	7406.8	20.51	4	13	2	0
0100240801	Medium	15 32 23.0 08 32 01.3	23738.7	20.96	3	13	2	0
0091140401	Thin1	15 32 28.6 + 04 40 48.6	34556.1	20.62	5	13	0	0
0112190401	Medium	15 35 01.4 + 01 20 45.4	13512.1	20.69	2	13	3	1
0094380801	Medium	15 55 43.2 + 11 11 16.3	5805.5	20.56	8	13	0	-
0067340601	Medium	16 07 13.8 + 08 04 51.5	14215.2	20.60	0	13	3	1
0103460801	Thin1	16 16 28.7 + 12 12 22.1	13130.2	20.66	1	13	2	0
0103461001	Thin1	16 17 36.8 + 12 36 57.7	11734.5	20.66	1	13	1	1
0061940301	Medium	16 30 53.6 + 78 10 54.5	4965.0	20.61	2	13	6	-
0112230301	Medium	16 32 45.7 + 05 34 39.6	20662.8	20.75	8	13	1	0
0049540101	Medium	16 34 20.8 + 57 09 43.1	10855.0	20.27	2	13	2	-
0067340501	Medium	16 42 17.3 + 03 10 57.5	14421.1	20.74	0	13	1	0
0101640101	Medium	16 52 58.4 + 02 24 00.4	16112.3	20.76	8	13	1	0
0113070101	Medium	16 53 59.8 + 14 17 52.6	7298.1	20.74	0	13	4	2
0111060101	Medium	16 57 49.6 + 35 20 35.8	9421.0	20.25	3	13	2	0
0102040101	Thick	17 23 20.8 + 34 18 00.6	6893.5	20.49	8	13	1	-
0067340101	Medium	18 55 37.1 46 30 57.6	10652.7	20.72	0	11	2	-
0081341001	Thin1	19 31 21.7 72 39 13.3	15287.3	20.77	1	13	2	1
0081340501	Medium	20 13 30.0 41 47 25.7	19195.9	20.69	0	13	0	0
0111180201	Medium	20 40 10.0 00 52 14.7	16494.9	20.83	8	13	2	1
0111510101	Medium	20 41 50.9 32 26 19.8	14918.6	20.71	8	13	3	0
0130720201	Thin1	20 44 09.0 10 43 11.2	5311.7	20.61	8	13	0	-
0111420101	Medium	20 45 09.4 31 20 36.6	42956.6	20.74	8	13	0	0
0112600501	Medium	20 46 20.0 02 48 48.4	10549.7	20.70	8	13	1	0
0083210101	Thin1	20 54 18.8 15 55 38.2	10434.9	20.64	2	13	2	0
0112190601	Medium	20 56 21.6 04 37 59.4	16639.7	20.70	2	13	1	1
0081340401	Thin1	20 58 27.1 42 38 56.9	15001.6	20.59	1	13	2	1
0041150101	Medium	21 04 11.1 11 21 40.0	38734.2	20.67	1	13	1	0
0038540301	Thin1	21 04 40.2 12 20 05.6	14697.9	20.65	2	13	1	0
0088020201	Thin1	21 27 38.1 44 48 38.5	16093.9	20.54	1	13	2	0
0103060101	Medium	21 29 12.2 15 38 34.7	22041.8	20.70	3	13	1	1
0092850201	Medium	21 37 45.3 14 32 56.4	16014.8	20.67	2	13	1	1
0109463501	Thin1	21 37 56.5 43 42 19.8	7577.3	20.46	2	13	3	0
0061940201	Medium	21 38 07.9 42 36 06.6	4867.9	20.43	1	13	5	-
0008830101	Thin1	21 40 15.5 23 39 32.1	13986.0	20.55	2	13	1	1
0103060401	Medium	21 51 55.9 30 27 43.5	24071.7	20.33	3	13	2	0
0111270101	Thin1	21 53 37.3 + 17 41 57.7	13650.1	20.83	6	13	1	0
0124930201	Medium	21 58 52.9 30 13 28.8	36244.5	20.23	8	13	0	0
0130920101	Thick	22 03 09.3 + 18 52 27.9	16490.5	20.78	1	13	1	0
0012440301	Thin1	22 05 10.1 01 55 11.2	30935.9	20.79	2	13	2	1
0106660101	Thin1	22 15 31.9 17 44 02.6	56908.9	20.37	1	13	1	0
0009650201	Medium	22 17 55.4 08 20 58.0	21026.5	20.73	8	13	4	0

Obs. ID	Filter	RA; DEC (J2000)	Exposure s	Log N h cm ⁻²	R _{in} arcmin	R _{out} arcmin	BSS srcts	H BSS srcts
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
0103861201	M edium	22 19 19.2 + 12 08 00.1	12172.6	20.73	8	13	1	0
0049340201	M edium	22 20 45.1 24 40 58.1	26840.9	20.29	6	13	0	0
0100440101	Thick	22 28 30.4 05 18 49.5	45013.1	20.72	2	13	1	0
0018741701	Thin1	22 34 32.9 37 43 48.1	7080.0	20.09	4	13	0	0
0111790101	M edium	22 35 45.9 26 03 00.4	41302.3	20.16	4	13	2	0
0103860201	M edium	22 36 55.9 22 13 10.1	8489.4	20.33	8	13	0	0
0103860301	M edium	22 40 18.1 + 08 03 23.2	5879.2	20.82	8	13	1	—
0006810101	M edium	22 42 39.5 + 29 43 35.6	7009.1	20.81	8	13	0	0
0109070401	M edium	22 48 41.5 51 09 57.9	14721.9	20.13	8	13	2	0
0112240101	M edium	22 49 48.3 64 23 11.2	30606.8	20.47	8	13	3	0
0081340901	M edium	22 51 49.4 17 52 17.0	22397.4	20.43	1	13	1	0
0112910301	Thin1	22 53 58.8 17 33 55.8	5301.0	20.43	8	13	1	—
0009650101	M edium	22 55 18.1 03 10 35.6	33943.5	20.66	8	13	0	0
0112170301	M edium	23 03 15.8 + 08 52 25.9	23348.5	20.69	4	13	0	0
0109130701	Thin1	23 04 43.6 08 41 14.5	10615.1	20.56	8	13	1	0
0033541001	Thin1	23 04 45.0 + 03 11 35.6	12460.9	20.72	2	13	2	0
0025541001	M edium	23 04 56.8 + 12 19 32.0	9954.8	20.70	3	13	5	0
0123900101	Thin1	23 13 58.9 42 43 28.3	37399.4	20.25	6	13	1	0
0109463601	Thin1	23 15 18.7 59 10 31.7	5403.3	20.46	8	13	1	—
0093640701	M edium	23 16 11.0 42 35 01.8	14442.9	20.27	3	13	4	0
0112880301	Thick	23 31 49.9 + 19 56 28.5	14317.3	20.63	3	13	0	0
0093550401	M edium	23 33 40.0 15 17 12.2	22256.9	20.30	1	13	2	0
0100241001	Thick	23 49 40.6 + 36 25 30.6	8738.5	20.91	3	13	2	0

Columns are as follows: (1) XM-M-Newton Observation number; (2) Filter used; (3) Right ascension and Declination (J2000) of the MOS2 image center; (4) On-Axis good-time exposure; (5) Logarithm of the Galactic hydrogen column density along the line of sight fromDickey and Lockman, 1990; (6) Inner radius of the part of the MOS2 image used; (7) Outer radius of the part of the MOS2 image used; (8) total number of BSS sources found in the surveyed area of each MOS2 image; (9) total number of H BSS sources found in the surveyed area of each MOS2 image. See section 2.4 for details.

NOTE { These 26 fields have not been considered for the definition of the H BSS sample.

Table 3. Basic information on the XM M -Newton BSS sample

Name XBS...	Obs. ID	RA ; DEC (J2000)	O Axis arm in	Rate 10^{-2} cts/s	HR 2	HR 3	Class
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
J000027.7	250442	0125310101	00 00 27.8 25 04 42.6	2.10	1:13 0:08	0.48 ^{+0.06} _{-0.06}	0.37 ^{+0.11} _{-0.11} AGN 1
J000031.7	245502	0125310101	00 00 31.8 24 55 02.9	11.68	1:41 0:13	0.67 ^{+0.07} _{-0.06}	0.37 ^{+0.17} _{-0.17} AGN 1
J000100.2	250501 ^d	0125310101	00 01 00.2 25 05 01.2	6.97	1:38 0:10	0.37 ^{+0.07} _{-0.07}	0.47 ^{+0.10} _{-0.10} AGN 1
J000102.4	245850	0125310101	00 01 02.5 24 58 50.7	10.72	1:48 0:12	0.54 ^{+0.07} _{-0.07}	0.71 ^{+0.11} _{-0.10} AGN 1
J000532.7+	200716	0101040101	00 05 32.8 + 20 07 16.2	12.15	2:47 0:12	0.81 ^{+0.03} _{-0.03}	0.70 ^{+0.11} _{-0.10} GAL
J001002.4+	110831	0127110201	00 10 02.4 + 11 08 31.0	12.12	13:12 0:71	0.79 ^{+0.03} _{-0.03}	0.88 ^{+0.07} _{-0.06} star
J001051.6+	105140	0127110201	00 10 51.6 + 10 51 40.2	8.61	1:01 0:15	0.76 ^{+0.09} _{-0.10}	1.00 ^{+0.76} _{-0.00} star
J001749.7+	161952	0111000101	00 17 49.7 + 16 19 52.5	12.11	3:35 0:15	1.00 ^{+0.01} _{-0.00}	1.00 ^{+0.00} _{-0.96} star
J001831.6+	162925	0111000101	00 18 31.7 + 16 29 25.6	3.31	1:82 0:08	0.67 ^{+0.03} _{-0.03}	0.56 ^{+0.08} _{-0.08} AGN 1
J002618.5+	105019 ^a	0001930101	00 26 18.5 + 10 50 19.3	9.56	5:19 0:24	0.53 ^{+0.04} _{-0.04}	0.67 ^{+0.06} _{-0.06} AGN 1
J002637.4+	165953	0050140201	00 26 37.4 + 16 59 53.5	9.74	2:78 0:10	0.61 ^{+0.03} _{-0.03}	0.62 ^{+0.06} _{-0.05} AGN 1
J002707.5+	170748	0050140201	00 27 07.6 + 17 07 48.5	7.78	1:58 0:08	0.58 ^{+0.04} _{-0.04}	0.39 ^{+0.08} _{-0.08} AGN 1
J002953.1+	044524	0112320101	00 29 53.2 + 04 45 24.6	10.69	1:33 0:12	0.91 ^{+0.04} _{-0.04}	1.00 ^{+0.83} _{-0.00} star
J003255.9+	394619 ^d	0065770101	00 32 55.9 + 39 46 19.9	11.90	1:11 0:21	0.76 ^{+0.12} _{-0.11}	0.24 ^{+0.32} _{-0.44} AGN 1
J003315.5	120700	0125920201	00 33 15.6 12 07 00.9	9.52	1:62 0:20	0.75 ^{+0.08} _{-0.08}	0.05 ^{+0.25} _{-0.26} AGN 1
J003316.0	120456	0125920201	00 33 16.1 12 04 56.6	9.66	1:03 0:15	0.90 ^{+0.06} _{-0.06}	0.97 ^{+1.27} _{-0.03} AGN 1
J003418.9	115940	0125920201	00 34 19.0 11 59 40.1	9.60	1:05 0:14	0.69 ^{+0.09} _{-0.09}	0.39 ^{+0.24} _{-0.26} AGN 1
J005009.9	515934	0125320701	00 50 09.9 51 59 34.7	8.10	1:38 0:11	0.56 ^{+0.07} _{-0.07}	0.57 ^{+0.13} _{-0.13} AGN 1
J005031.1	520012	0125320701	00 50 31.1 52 00 12.8	8.57	1:89 0:14	0.65 ^{+0.05} _{-0.05}	0.47 ^{+0.13} _{-0.13} AGN 1
J005032.3	521543	0125320701	00 50 32.3 52 15 43.5	9.28	1:12 0:13	0.45 ^{+0.09} _{-0.09}	0.53 ^{+0.14} _{-0.15} AGN 1
J005822.9	274016	0111280301	00 58 22.9 27 40 16.1	7.71	2:07 0:21	0.91 ^{+0.04} _{-0.05}	0.19 ^{+0.42} _{-0.38} star
J010421.4	061418	0112650401	01 04 21.4 06 14 18.8	9.89	1:01 0:09	0.49 ^{+0.08} _{-0.08}	0.81 ^{+0.10} _{-0.11} AGN 1
J010432.8	583712	0103861601	01 04 32.8 58 37 12.2	12.41	1:61 0:26	0.53 ^{+0.14} _{-0.13}	0.07 ^{+0.26} _{-0.27} AGN 1
J010701.5	172748	0025540101	01 07 01.5 17 27 48.4	10.99	3:56 0:33	0.47 ^{+0.08} _{-0.08}	0.84 ^{+0.09} _{-0.09} AGN 1
J010747.2	172044	0025540101	01 07 47.3 17 20 44.6	9.49	2:27 0:23	0.62 ^{+0.08} _{-0.08}	0.48 ^{+0.19} _{-0.17} AGN 1
J012000.0	110429	0113040801	01 20 00.1 11 04 29.3	12.66	1:18 0:25	0.77 ^{+0.14} _{-0.14}	0.01 ^{+0.50} _{-0.53} AGN 1
J012025.2	105441	0113040801	01 20 25.3 10 54 41.6	3.95	1:39 0:18	0.50 ^{+0.11} _{-0.11}	0.68 ^{+0.16} _{-0.18} N
J012057.4	110444	0113040801	01 20 57.5 11 04 44.6	9.40	2:34 0:28	0.61 ^{+0.09} _{-0.09}	0.48 ^{+0.22} _{-0.21} AGN 2
J012119.9	110418	0113040801	01 21 20.0 11 04 18.6	12.65	5:85 0:48	0.71 ^{+0.06} _{-0.06}	0.67 ^{+0.15} _{-0.16} AGN 1
J012505.4+	014624 ^d	0109860101	01 25 05.4 + 01 46 24.4	7.04	1:94 0:09	0.46 ^{+0.04} _{-0.04}	0.51 ^{+0.06} _{-0.06} AGN 1
J012540.2+	015752	0109860101	01 25 40.3 + 01 57 52.1	12.35	1:41 0:10	0.56 ^{+0.06} _{-0.06}	0.29 ^{+0.11} _{-0.12} AGN 2
J012654.3+	191246 ^d	0112600601	01 26 54.3 + 19 12 47.0	9.20	1:02 0:26	0.64 ^{+0.17} _{-0.17}	0.07 ^{+0.41} _{-0.39} GAL
J012757.3+	185923	0112600601	01 27 57.4 + 18 59 23.9	12.64	34:37 1:49	0.92 ^{+0.02} _{-0.02}	0.34 ^{+0.17} _{-0.16} star
J012757.2+	190000	0112600601	01 27 57.2 + 19 00 00.4	12.09	10:69 0:81	0.81 ^{+0.04} _{-0.04}	1.00 ^{+0.27} _{-0.00} star
J013204.9	400050	0112630201	01 32 04.9 40 00 50.9	12.23	1:29 0:12	0.67 ^{+0.06} _{-0.06}	0.32 ^{+0.16} _{-0.16} AGN 1
J013240.1	133307 ^{a,d}	0084230301	01 32 40.1 13 33 07.8	11.90	1:42 0:14	0.02 ^{+0.10} _{-0.10}	0.37 ^{+0.11} _{-0.11} AGN 2

Name XBS...	Obs. ID	RA; DEC (J2000)	O Axis arcmin	Rate 10^{-2} cts/s	HR 2	HR 3	Class
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
J013811.7	175416 ^d	01 38 11.8 17 54 16.3	12.09	3.09 0.17	$0.73^{+0.04}_{-0.04}$	$0.77^{+0.08}_{-0.08}$	BL?
J013924.8	674126	0032140401	01 39 24.8 67 41 26.5	8.50 1.02	$0.21^{+0.14}_{-0.14}$	$0.54^{+0.13}_{-0.13}$	
J013944.0	674909 ^a	0032140401	01 39 44.0 67 49 09.4	2.69 2.04	0.17 0.07	$0.43^{+0.13}_{-0.13}$	AGN?
J014100.6	675328 ^a	0032140401	01 41 00.7 67 53 29.0	6.64 57.52	1.03 0.02	$0.38^{+0.03}_{-0.03}$	star
J014109.9	675639	0032140401	01 41 09.9 67 56 39.6	9.67 1.97	0.23 0.10	$0.45^{+0.16}_{-0.15}$	AGN?
J014227.0+	133453	0093641001	01 42 27.1 +13 34 53.7	9.12 1.29	0.14 0.10	$0.51^{+0.18}_{-0.18}$	AGN?
J014251.5+	133352	0093641001	01 42 51.6 +13 33 52.8	5.15 1.43	0.13 0.08	$0.49^{+0.14}_{-0.14}$	AGN
J015916.9+	003010 ^c	0101640201	01 59 17.0 +00 30 10.4	10.49 5.74	0.54 0.06	$0.71^{+0.20}_{-0.20}$	CL ^b BL?
J015957.5+	003309 ^a	0101640201	01 59 57.5 +00 33 09.7	9.56 6.50	0.39 0.05	$0.51^{+0.10}_{-0.10}$	AGN
J020029.0+	002846	0101640201	02 00 29.0 +00 28 46.1	10.90 3.39	0.30 0.07	$0.36^{+0.15}_{-0.15}$	AGN
J020757.3+	351828	0084140101	02 07 57.3 +35 18 28.9	9.48 1.19	0.07 0.05	$0.55^{+0.09}_{-0.09}$	AGN
J020845.1+	351438	0084140101	02 08 45.2 +35 14 38.0	8.49 1.80	0.09 0.04	$0.52^{+0.07}_{-0.07}$	AGN
J020916.4+	351850	0084140101	02 09 16.5 +35 18 50.4	8.85 1.30	0.08 0.05	$0.66^{+0.08}_{-0.08}$	AGN
J021640.7	044404 ^{a,d}	0112371701	02 16 40.7 04 44 04.9	9.35 2.56	0.17 0.05	$0.26^{+0.13}_{-0.13}$	AGN
J021642.3	043553	0112371701	02 16 42.4 04 35 53.0	8.14 1.32	0.12 0.08	$0.61^{+0.11}_{-0.11}$	AGN
J021808.3	045845 ^a	0112371001	02 18 08.3 04 58 45.7	2.26 4.87	0.10 0.02	$0.52^{+0.04}_{-0.04}$	AGN
J021817.4	045113 ^a	0112371001	02 18 17.4 04 51 13.3	9.58 4.47	0.13 0.03	$0.56^{+0.04}_{-0.04}$	AGN
J021820.6	050427	0112371001	02 18 20.7 05 04 27.6	6.90 1.22	0.06 0.04	$0.57^{+0.07}_{-0.07}$	AGN
J021830.0	045514	0112371001	02 18 30.0 04 55 14.8	8.69 3.36	0.10 0.01	$0.69^{+0.09}_{-0.09}$	star
J021923.2	045148	0112370301	02 19 23.2 04 51 48.1	8.63 1.40	0.07 0.04	$0.42^{+0.09}_{-0.10}$	AGN
J022253.0	044515	0109520501	02 22 53.0 04 45 16.0	8.12 1.15	0.09 0.07	$0.60^{+0.12}_{-0.12}$	AGN
J022339.6	050728	0111110401	02 23 39.6 05 07 28.2	5.66 1.45	0.08 0.05	$0.76^{+0.07}_{-0.07}$	AGN
J022707.7	050819	0111110201	02 27 07.7 05 08 19.8	7.05 1.89	0.19 0.09	$0.67^{+0.08}_{-0.08}$	AGN
J023459.7	294436	0103861101	02 34 59.7 29 44 36.9	8.91 1.18	0.23 0.14	$0.48^{+0.14}_{-0.14}$	AGN
J023530.2	523045	0098810101	02 35 30.2 52 30 45.2	12.59 1.88	0.14 0.05	$0.38^{+0.14}_{-0.14}$	AGN
J023713.5	522734 ^a	0098810101	02 37 13.5 52 27 34.4	12.07 5.60	0.24 0.03	$0.58^{+0.07}_{-0.06}$	AGN
J023853.2	521911	0067190101	02 38 53.3 52 19 11.1	9.22 1.45	0.11 0.06	$0.42^{+0.11}_{-0.11}$	AGN
J024200.9+	0000020	0111200101	02 42 01.0 +00 00 20.1	10.11 2.13	0.12 0.04	$0.52^{+0.09}_{-0.09}$	AGN
J024204.7+	000814	0111200101	02 42 04.7 +00 08 14.3	12.90 1.01	0.10 0.07	$0.81^{+0.10}_{-0.10}$	AGN
J024207.3+	0000037	0111200101	02 42 07.3 +00 00 37.6	8.59 1.26	0.09 0.05	$0.77^{+0.09}_{-0.09}$	AGN
J024325.6	000413	0111200101	02 43 25.6 00 04 13.3	11.61 1.04	0.08 0.07	$0.40^{+0.10}_{-0.10}$	AGN
J025606.1+	001635	0056020301	02 56 06.2 +00 16 35.5	12.49 1.28	0.15 0.09	$0.93^{+0.06}_{-0.06}$	AGN
J025645.4+	0000031	0056020301	02 56 45.5 +00 00 31.3	6.35 1.01	0.10 0.08	$0.65^{+0.12}_{-0.13}$	AGN?
J030204.2+	001625	0041170101	03 02 04.2 +00 16 25.6	12.36 1.01	0.07 0.06	$0.50^{+0.10}_{-0.10}$	
J030206.8	000121 ^a	0041170101	03 02 06.9 00 01 21.2	11.90 3.56	0.13 0.03	$0.46^{+0.05}_{-0.06}$	AGN
J030614.1	284019 ^a	0042340501	03 06 14.2 28 40 19.9	10.87 3.85	0.23 0.05	$0.35^{+0.09}_{-0.09}$	AGN
J030641.0	283559	0042340501	03 06 41.0 28 35 59.5	6.66 1.25	0.11 0.07	$0.81^{+0.11}_{-0.11}$	AGN
J031015.5	765131 ^a	0122520201	03 10 15.6 76 51 31.5	5.90 5.36	0.15 0.03	$0.53^{+0.04}_{-0.04}$	AGN

Name XBS...	Obs. ID	RA; DEC (J2000)	O Axis arcmin	Rate 10^{-2} cts/s	HR 2	HR 3	Class			
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)			
J031146.1	550702 ^a	0110970101	03 11 46.1	55 07 02.5	12.38	6.55	0.45	$0.58^{+0.05}_{-0.05}$	$0.39^{+0.11}_{-0.11}$	AGN 2
J031311.7	765428	0122520201	03 13 11.7	76 54 28.8	4.86	1.10	0.07	$0.43^{+0.06}_{-0.06}$	$0.30^{+0.09}_{-0.09}$	AGN 1
J031401.3	545959	0110970101	03 14 01.3	54 59 59.5	8.34	1.29	0.14	$0.51^{+0.09}_{-0.09}$	$0.68^{+0.14}_{-0.14}$	AGN 1
J031549.4	551811	0129320901	03 15 49.4	55 18 11.5	12.22	1.04	0.16	$0.83^{+0.09}_{-0.08}$	$0.85^{+0.21}_{-0.13}$	AGN 1
J031851.9	441815	0105660101	03 18 51.9	44 18 15.3	10.78	1.15	0.12	$0.59^{+0.08}_{-0.07}$	$0.40^{+0.16}_{-0.15}$	AGN 1
J031859.2	441627 ^{a,b}	0105660101	03 18 59.3	44 16 27.6	11.55	1.54	0.14	$0.21^{+0.08}_{-0.08}$	$0.47^{+0.10}_{-0.10}$	AGN 1
J033208.7	274735	0108060501	03 32 08.7	27 47 35.7	4.60	1.13	0.06	$0.70^{+0.03}_{-0.04}$	$0.40^{+0.10}_{-0.09}$	AGN 2?
J033226.9	274107 ^d	0108060501	03 32 27.0	27 41 07.3	7.35	1.42	0.07	$0.63^{+0.04}_{-0.04}$	$0.59^{+0.08}_{-0.07}$	AGN 1
J033435.5	254259	0099010101	03 34 35.6	25 42 59.3	11.87	1.10	0.12	$0.50^{+0.09}_{-0.09}$	$1.00^{+0.18}_{-0.00}$	AGN 1
J033453.9	254154	0099010101	03 34 54.0	25 41 54.6	8.14	1.89	0.13	$0.62^{+0.05}_{-0.05}$	$0.38^{+0.12}_{-0.12}$	AGN 1
J033506.0	255619	0099010101	03 35 06.1	25 56 19.9	12.41	1.18	0.12	$0.63^{+0.08}_{-0.08}$	$0.82^{+0.13}_{-0.16}$	AGN 1
J033851.4	352646	0055140101	03 38 51.5	35 26 46.3	8.89	1.38	0.08	$0.55^{+0.04}_{-0.04}$	$0.68^{+0.07}_{-0.07}$	AGN 1
J033912.1	352813	0055140101	03 39 12.1	35 28 13.8	5.18	1.23	0.06	$0.40^{+0.04}_{-0.04}$	$0.52^{+0.06}_{-0.06}$	N
J033942.8	352411	0055140101	03 39 42.8	35 24 11.3	2.40	1.33	0.06	$0.75^{+0.03}_{-0.03}$	$0.56^{+0.09}_{-0.09}$	AGN 1
J040744.6	710846	0111970301	04 07 44.6	71 08 47.0	11.15	1.27	0.15	$0.77^{+0.07}_{-0.07}$	$0.75^{+0.18}_{-0.22}$	Newton
J040807.2	712702	0111970301	04 08 07.2	71 27 02.5	10.47	6.46	0.27	$0.91^{+0.02}_{-0.02}$	$0.74^{+0.12}_{-0.11}$	star
J041108.1	711341 ^a	0111970301	04 11 08.1	71 13 41.1	10.52	1.03	0.11	$0.53^{+0.08}_{-0.08}$	$0.04^{+0.16}_{-0.16}$	AGN 1
J042417.9	571539	0112600401	04 24 17.9	57 15 39.2	11.86	1.04	0.18	$0.64^{+0.13}_{-0.13}$	$0.46^{+0.32}_{-0.30}$	Right Survey
J043448.3	775329	0103861701	04 34 48.4	77 53 29.2	8.55	2.03	0.22	$0.35^{+0.10}_{-0.10}$	$0.61^{+0.14}_{-0.13}$	AGN 1
J045942.4+	015843	0112880401	04 59 42.5 + 01 58 43.5	11.59	1.38	0.13	$0.35^{+0.09}_{-0.08}$	$0.49^{+0.11}_{-0.12}$	AGN 1	
J050011.7+	013948	0112880401	05 00 11.8 + 01 39 48.0	11.76	1.19	0.11	$0.35^{+0.09}_{-0.09}$	$0.38^{+0.13}_{-0.13}$	AGN 1?	
J050446.3	283821	0111160201	05 04 46.3	28 38 21.8	12.92	1.36	0.10	$0.57^{+0.06}_{-0.05}$	$0.67^{+0.10}_{-0.10}$	AGN 1
J050453.4	284532	0111160201	05 04 53.5	28 45 32.1	6.75	1.06	0.06	$0.39^{+0.05}_{-0.05}$	$0.41^{+0.08}_{-0.08}$	AGN 1
J050501.8	284149	0111160201	05 05 01.9	28 41 49.1	8.18	1.41	0.08	$0.65^{+0.04}_{-0.04}$	$0.51^{+0.09}_{-0.09}$	AGN 1
J050536.6	290050	0111160201	05 05 36.7	29 00 50.8	12.43	1.68	0.10	$0.27^{+0.05}_{-0.05}$	$0.47^{+0.07}_{-0.07}$	AGN 1
J051413.5+	794345	0094400101	05 14 13.6 + 79 43 45.8	7.42	1.43	0.09	$0.29^{+0.06}_{-0.06}$	$0.53^{+0.08}_{-0.08}$	AGN 1	
J051617.1+	794408	0094400101	05 16 17.2 + 79 44 08.7	4.00	4.05	0.12	$0.85^{+0.02}_{-0.02}$	$0.73^{+0.07}_{-0.07}$	star	
J051651.9+	794314	0094400101	05 16 51.9 + 79 43 14.6	3.01	1.77	0.08	$0.28^{+0.04}_{-0.04}$	$0.55^{+0.06}_{-0.05}$	AGN 1	
J051655.3	104104	0103860701	05 16 55.3	10 41 04.0	11.23	1.97	0.31	$0.33^{+0.14}_{-0.14}$	$1.00^{+0.29}_{-0.00}$	AGN 1
J051822.6+	793208	0094400101	05 18 22.7 + 79 32 08.0	9.37	1.27	0.09	$0.54^{+0.06}_{-0.06}$	$0.41^{+0.10}_{-0.11}$	AGN 1	
J051955.5	455727	0090050701	05 19 55.5	45 57 27.4	10.69	3.27	0.24	$0.51^{+0.06}_{-0.06}$	$0.77^{+0.08}_{-0.08}$	AGN 1
J052022.0	252309	0085640101	05 20 22.0	25 23 09.5	8.87	1.21	0.14	$0.66^{+0.09}_{-0.09}$	$0.37^{+0.21}_{-0.22}$	AGN 1
J052048.9	454128	0090050701	05 20 48.9	45 41 28.5	11.67	8.35	0.45	$0.95^{+0.02}_{-0.02}$	$0.81^{+0.17}_{-0.17}$	star
J052108.5	251913 ^d	0085640101	05 21 08.5	25 19 13.1	3.17	3.54	0.20	$0.59^{+0.05}_{-0.05}$	$0.55^{+0.09}_{-0.09}$	AGN 1
J052116.2	252957	0085640101	05 21 16.2	25 29 57.8	8.80	1.00	0.15	$0.50^{+0.11}_{-0.12}$	$0.70^{+0.18}_{-0.17}$	AGN 1
J052144.1	251518	0085640101	05 21 44.1	25 15 18.8	11.77	1.25	0.19	$0.60^{+0.11}_{-0.12}$	$0.87^{+0.17}_{-0.12}$	AGN 1
J052155.0	252200	0085640101	05 21 55.0	25 22 00.7	12.24	1.82	0.24	$1.00^{+0.07}_{-0.00}$	{	star
J052509.3	333051	0050150101	05 25 09.3	33 30 51.9	12.38	1.38	0.18	$0.56^{+0.10}_{-0.10}$	$0.95^{+0.09}_{-0.05}$	star

Name XBS...	Obs. ID	RA; DEC (J2000)	O Axis arcmin	Rate 10^{-2} cts/s	HR 2	HR 3	Class
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
J052543.6 334856	0050150101	05 25 43.6 33 48 56.4	9.70	1:01 0:15	0.53 ^{+0.11} _{-0.10}	0.76 ^{+0.15} _{-0.14}	
J061342.7+ 710725 ^d	0009220601	06 13 42.8 + 71 07 25.9	10.51	6:44 0:34	0.67 ^{+0.04} _{-0.04}	0.75 ^{+0.08} _{-0.07}	BL
J062134.8 643150	0103860101	06 21 34.8 64 31 50.5	11.02	1:04 0:20	0.41 ^{+0.15} _{-0.16}	0.22 ^{+0.23} _{-0.25}	D star
J062425.7 642958	0103860101	06 24 25.7 64 29 58.3	10.39	1:99 0:22	1.00 ^{+0.04} _{-0.00}	1.00 ^{+0.00} _{-1.30}	star
J063709.1+ 820949	0029340101	06 37 09.2 + 82 09 49.1	10.46	1:28 0:11	0.58 ^{+0.06} _{-0.06}	0.52 ^{+0.13} _{-0.13}	AGN?
J065214.1+ 743230	0061540101	06 52 14.1 + 74 32 30.6	7.01	2:33 0:14	0.49 ^{+0.05} _{-0.05}	0.58 ^{+0.08} _{-0.08}	AGN?
J065237.4+ 742421	0061540101	06 52 37.5 + 74 24 21.9	2.09	1:08 0:09	0.62 ^{+0.05} _{-0.06}	0.35 ^{+0.12} _{-0.12}	AGN?
J065400.0+ 742045	0061540101	06 54 00.1 + 74 20 45.1	8.72	1:82 0:15	0.49 ^{+0.06} _{-0.06}	0.65 ^{+0.10} _{-0.10}	AGN?
J065744.3 560817	0112980201	06 57 44.3 56 08 17.5	11.74	1:40 0:10	0.56 ^{+0.05} _{-0.05}	0.78 ^{+0.08} _{-0.08}	The
J065839.5 560813	0112980201	06 58 39.6 56 08 13.3	11.14	1:60 0:10	0.65 ^{+0.04} _{-0.04}	0.45 ^{+0.10} _{-0.10}	
J074202.7+ 742625	0123100101	07 42 02.7 + 74 26 25.8	10.86	3:02 0:13	0.55 ^{+0.03} _{-0.03}	0.28 ^{+0.07} _{-0.07}	AGN?
J074312.1+ 742937	0123100101	07 43 12.1 + 74 29 37.4	5.33	19:59 0:25	0.61 ^{+0.01} _{-0.01}	0.56 ^{+0.02} _{-0.02}	AGN?
J074338.7+ 495431	0103862101	07 43 38.7 + 49 54 31.1	12.22	1:27 0:23	0.14 ^{+0.20} _{-0.20}	0.47 ^{+0.24} _{-0.24}	AGN?
J074352.0+ 744258	0123100101	07 43 52.1 + 74 42 59.0	9.30	1:99 0:10	0.56 ^{+0.04} _{-0.04}	0.49 ^{+0.08} _{-0.08}	AGN?
J074359.7+ 744057	0123100101	07 43 59.7 + 74 40 57.5	7.25	1:47 0:08	0.85 ^{+0.03} _{-0.03}	0.62 ^{+0.13} _{-0.13}	star
J075117.9+ 180856 ^b	0111100301	07 51 17.9 + 18 08 56.1	2.37	1:65 0:08	0.35 ^{+0.04} _{-0.04}	0.46 ^{+0.06} _{-0.06}	AGN?
J080309.8+ 650807	0094400301	08 03 09.8 + 65 08 07.6	12.10	2:91 0:22	0.97 ^{+0.02} _{-0.01}	1.00 ^{+1.30} _{-0.00}	star
J080411.3+ 650906	0094400301	08 04 11.4 + 65 09 06.2	9.49	1:39 0:12	0.06 ^{+0.09} _{-0.09}	0.51 ^{+0.09} _{-0.09}	Serendipitous
J080504.6+ 245156	0094530401	08 05 04.6 + 24 51 56.4	7.23	1:07 0:13	0.53 ^{+0.11} _{-0.11}	0.56 ^{+0.20} _{-0.20}	
J080608.1+ 244420	0094530401	08 06 08.2 + 24 44 20.4	9.34	4:49 0:30	0.74 ^{+0.05} _{-0.04}	0.48 ^{+0.14} _{-0.13}	AGN?
J083049.8+ 524908	0092800101	08 30 49.8 + 52 49 08.2	8.81	1:40 0:13	0.48 ^{+0.08} _{-0.08}	0.85 ^{+0.09} _{-0.08}	
J083737.1+ 254751	0025540301	08 37 37.2 + 25 47 51.1	10.61	9:08 0:37	0.47 ^{+0.04} _{-0.04}	0.53 ^{+0.06} _{-0.06}	AGN?
J083737.0+ 255151	0025540301	08 37 37.1 + 25 51 51.2	12.25	2:89 0:23	0.34 ^{+0.07} _{-0.08}	0.52 ^{+0.10} _{-0.10}	AGN?
J083838.6+ 253616	0025540301	08 38 38.6 + 25 36 16.8	9.63	1:32 0:16	0.57 ^{+0.09} _{-0.09}	0.45 ^{+0.20} _{-0.19}	AGN?
J083905.9+ 255010	0025540301	08 39 05.9 + 25 50 10.5	10.89	1:02 0:15	0.69 ^{+0.10} _{-0.10}	1.00 ^{+0.54} _{-0.00}	
J084026.2+ 650638 ^d	0111400101	08 40 26.2 + 65 06 38.5	9.56	1:21 0:09	0.50 ^{+0.06} _{-0.07}	0.27 ^{+0.12} _{-0.12}	AGN?
J084651.7+ 344634	0103660201	08 46 51.7 + 34 46 34.1	10.51	1:05 0:15	0.71 ^{+0.08} _{-0.08}	1.00 ^{+0.42} _{-0.00}	AGN?
J085427.8+ 584158	0085030101	08 54 27.8 + 58 41 58.9	9.06	2:92 0:32	0.83 ^{+0.06} _{-0.06}	0.71 ^{+0.23} _{-0.27}	star
J085530.7+ 585129	0085030101	08 55 30.8 + 58 51 29.1	7.42	1:21 0:19	0.64 ^{+0.11} _{-0.11}	1.00 ^{+0.47} _{-0.00}	star
J090729.1+ 620824	0110660201	09 07 29.2 + 62 08 24.6	11.48	1:00 0:17	0.26 ^{+0.15} _{-0.15}	0.41 ^{+0.20} _{-0.20}	Giant
J091043.4+ 054757	0083240201	09 10 43.4 + 05 47 57.8	11.54	2:40 0:19	0.80 ^{+0.04} _{-0.04}	1.00 ^{+0.25} _{-0.00}	star
J091828.4+ 513931	0084230601	09 18 28.5 + 51 39 31.3	6.92	1:09 0:11	0.31 ^{+0.09} _{-0.10}	0.39 ^{+0.09} _{-0.09}	AGN?
J094526.2 085006	0017540101	09 45 26.2 08 50 06.5	11.06	1:39 0:20	0.50 ^{+0.12} _{-0.12}	0.80 ^{+0.16} _{-0.18}	AGN?
J094548.3 084824	0017540101	09 45 48.3 08 48 24.6	10.87	1:02 0:16	0.58 ^{+0.13} _{-0.13}	0.24 ^{+0.30} _{-0.27}	
J095054.5+ 393924	0111290101	09 50 54.5 + 39 39 24.3	12.46	1:19 0:13	0.59 ^{+0.08} _{-0.09}	0.32 ^{+0.19} _{-0.18}	AGN?
J095134.6 015451	0065790101	09 51 34.6 01 54 51.6	10.87	1:05 0:16	0.50 ^{+0.13} _{-0.13}	0.72 ^{+0.19} _{-0.25}	
J095218.9 013643	0065790101	09 52 19.0 01 36 43.1	11.78	9:02 0:51	0.77 ^{+0.03} _{-0.03}	0.41 ^{+0.09} _{-0.09}	AGN?
J095309.7+ 013558	0070940101	09 53 09.8 + 01 35 58.6	8.00	1:15 0:19	0.64 ^{+0.12} _{-0.13}	0.54 ^{+0.30} _{-0.43}	AGN?

Name XBS...	Obs. ID	RA; DEC (J2000)	O Axis arcmin	Rate 10^{-2} cts/s	HR 2	HR 3	Class
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
J095341.1+ 014204 ^c	0070940101	09 53 41.2 + 01 42 04.4	7.14	16.29	0.72	$0.62^{+0.04}_{-0.04}$	$0.83^{+0.09}_{-0.10}$ CL
J095416.9+ 173627	0112850101	09 54 16.9 + 17 36 27.8	11.71	4.01	0.23	$0.72^{+0.04}_{-0.04}$	$0.62^{+0.10}_{-0.11}$
J095509.6+ 174124	0112850101	09 55 09.6 + 17 41 24.5	3.80	1.41	0.10	$0.55^{+0.06}_{-0.06}$	$0.51^{+0.11}_{-0.11}$
J095606.4+ 411814	0111290201	09 56 06.5 + 41 18 14.7	9.12	1.23	0.15	$0.57^{+0.09}_{-0.10}$	$0.62^{+0.18}_{-0.17}$
J095955.2+ 251549	0041170201	09 59 55.3 + 25 15 49.1	10.40	1.62	0.10	$0.77^{+0.04}_{-0.04}$	$0.84^{+0.09}_{-0.09}$
J100020.6+ 252046	0041170201	10 00 20.6 + 25 20 46.7	7.83	2.37	0.10	$0.78^{+0.02}_{-0.03}$	$0.45^{+0.09}_{-0.09}$
J100032.5+ 553626	0110930201	10 00 32.5 + 55 36 26.4	12.64	1.90	0.27	$0.48^{+0.11}_{-0.11}$	$0.30^{+0.21}_{-0.21}$
J100100.0+ 252103	0041170201	10 01 00.1 + 25 21 03.4	7.94	1.37	0.07	$0.65^{+0.04}_{-0.04}$	$0.60^{+0.08}_{-0.09}$
J100309.4+ 554135	0110930201	10 03 09.5 + 55 41 35.3	10.26	1.48	0.22	$0.47^{+0.13}_{-0.13}$	$0.62^{+0.20}_{-0.20}$
J100828.8+ 535408	0070340201	10 08 28.8 + 53 54 08.3	12.42	1.09	0.11	$0.60^{+0.08}_{-0.08}$	$0.38^{+0.17}_{-0.17}$
J100921.7+ 534926	0070340201	10 09 21.8 + 53 49 26.7	9.03	1.82	0.13	$0.66^{+0.05}_{-0.05}$	$0.59^{+0.11}_{-0.12}$
J100926.5+ 533426	0070340201	10 09 26.6 + 53 34 26.1	9.57	1.36	0.12	$0.45^{+0.08}_{-0.08}$	$0.46^{+0.12}_{-0.13}$
J101506.0+ 520157	0086750101	10 15 06.0 + 52 01 57.9	13.00	1.41	0.27	$0.59^{+0.14}_{-0.14}$	$1.00^{+0.63}_{-0.00}$
J101511.8+ 520708 ^d	0086750101	10 15 11.8 + 52 07 08.0	10.02	1.72	0.27	$0.65^{+0.10}_{-0.10}$	$0.71^{+0.20}_{-0.26}$
J101706.5+ 520245 ^d	0086750101	10 17 06.5 + 52 02 45.6	10.28	1.83	0.24	$0.66^{+0.09}_{-0.09}$	$0.99^{+0.36}_{-0.01}$
J101838.0+ 411635	0028740301	10 18 38.0 + 41 16 35.6	8.67	1.43	0.10	$0.71^{+0.04}_{-0.04}$	$0.73^{+0.11}_{-0.11}$
J101843.0+ 413515	0028740301	10 18 43.1 + 41 35 15.5	10.02	1.93	0.13	$0.60^{+0.05}_{-0.05}$	$0.52^{+0.10}_{-0.11}$
J101850.5+ 411506	0028740301	10 18 50.6 + 41 15 06.6	10.32	4.33	0.18	$0.69^{+0.03}_{-0.03}$	$0.51^{+0.07}_{-0.08}$
J101922.6+ 412049	0028740301	10 19 22.6 + 41 20 49.7	9.06	3.36	0.14	$0.44^{+0.04}_{-0.04}$	$0.58^{+0.06}_{-0.05}$
J102016.1+ 082143 ^d	0093640301	10 20 16.1 + 08 21 43.9	9.13	1.14	0.14	$0.40^{+0.10}_{-0.10}$	$0.47^{+0.15}_{-0.15}$
J102044.1+ 081424	0093640301	10 20 44.1 + 08 14 24.7	10.99	3.35	0.26	$0.94^{+0.03}_{-0.03}$	$1.00^{+0.76}_{-0.00}$
J102252.0+ 194837	0101040301	10 22 52.1 + 19 48 37.9	9.65	1.42	0.10	$0.48^{+0.06}_{-0.06}$	$0.51^{+0.09}_{-0.09}$
J102412.3+ 042023	0108670101	10 24 12.3 + 04 20 23.9	12.07	1.08	0.07	$0.48^{+0.06}_{-0.06}$	$0.68^{+0.08}_{-0.08}$
J102417.5+ 041656	0108670101	10 24 17.5 + 04 16 56.4	10.88	1.60	0.08	$0.55^{+0.04}_{-0.04}$	$0.44^{+0.08}_{-0.08}$
J103120.0+ 311404	0102040301	10 31 20.0 + 31 14 04.1	12.08	1.68	0.13	$0.69^{+0.05}_{-0.05}$	$0.28^{+0.14}_{-0.14}$
J103154.1+ 310732	0102040301	10 31 54.1 + 31 07 32.2	12.70	1.03	0.10	$0.34^{+0.08}_{-0.08}$	$0.46^{+0.12}_{-0.12}$
J103745.7+ 532353	0112810301	10 37 45.8 + 53 23 53.8	10.73	1.03	0.18	$0.71^{+0.14}_{-0.14}$	$0.18^{+0.44}_{-0.43}$
J103909.4+ 205222	0059800101	10 39 09.4 + 20 52 22.0	10.44	2.25	0.20	$0.55^{+0.07}_{-0.07}$	$0.47^{+0.14}_{-0.14}$
J103932.7+ 205426	0059800101	10 39 32.8 + 20 54 26.0	6.04	1.94	0.15	$0.51^{+0.06}_{-0.06}$	$0.67^{+0.10}_{-0.10}$
J103935.8+ 533036	0112810301	10 39 35.9 + 53 30 36.9	7.67	2.72	0.31	$0.55^{+0.09}_{-0.09}$	$0.71^{+0.16}_{-0.16}$
J103958.3+ 203848	0059800101	10 39 58.3 + 20 38 48.2	12.11	1.11	0.14	$0.43^{+0.10}_{-0.11}$	$0.42^{+0.17}_{-0.17}$
J104026.9+ 204542	0059800101	10 40 26.9 + 20 45 43.0	9.32	11.76	0.37	$0.58^{+0.03}_{-0.03}$	$0.48^{+0.05}_{-0.05}$
J104034.3+ 205110	0059800101	10 40 34.4 + 20 51 10.4	9.51	1.16	0.12	$0.45^{+0.09}_{-0.09}$	$0.76^{+0.11}_{-0.12}$
J104425.0 013521	0125300101	10 44 25.0 01 35 21.2	10.43	1.19	0.09	$0.56^{+0.06}_{-0.06}$	$0.52^{+0.12}_{-0.12}$
J104451.5 012227	0125300101	10 44 51.5 01 22 27.7	5.45	1.23	0.07	$0.41^{+0.05}_{-0.05}$	$0.57^{+0.08}_{-0.08}$
J104509.3 012442	0125300101	10 45 09.3 01 24 42.3	9.23	1.02	0.08	$0.73^{+0.05}_{-0.05}$	$0.68^{+0.14}_{-0.14}$
J104522.1 012843	0125300101	10 45 22.1 01 28 43.3	12.94	4.45	0.22	$0.63^{+0.04}_{-0.04}$	$0.46^{+0.08}_{-0.08}$
J104912.8+ 330459	0055990201	10 49 12.8 + 33 04 59.8	10.46	2.57	0.18	$0.45^{+0.06}_{-0.06}$	$0.54^{+0.09}_{-0.10}$

Name XBS...	Obs. ID	RA; DEC (J2000)	O Axis arcmin	Rate 10^{-2} cts/s	HR 2	HR 3	Class
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
J105014.9+ 331013	0055990201	10 50 14.9 + 33 10 13.2	12.11	1:51 0:15	0.55 ^{+0.08} _{-0.08}	0.61 ^{+0.14} _{-0.14}	
J105131.1+ 573439	0022740101	10 51 31.1 + 57 34 39.1	11.41	2:06 0:10	0.89 ^{+0.02} _{-0.02}	0.33 ^{+0.13} _{-0.13}	star
J105239.7+ 572431	0022740101	10 52 39.8 + 57 24 31.1	4.33	1:68 0:06	0.65 ^{+0.02} _{-0.02}	0.62 ^{+0.05} _{-0.05}	AGN 1
J105316.9+ 573551	0022740101	10 53 17.0 + 57 35 51.6	8.31	2:48 0:09	0.59 ^{+0.02} _{-0.02}	0.44 ^{+0.05} _{-0.05}	AGN 1
J105335.0+ 572540	0022740101	10 53 35.1 + 57 25 40.7	7.53	1:77 0:07	0.56 ^{+0.03} _{-0.03}	0.54 ^{+0.05} _{-0.05}	AGN 1
J105339.7+ 573104	0022740101	10 53 39.7 + 57 31 04.3	7.80	1:11 0:06	0.68 ^{+0.03} _{-0.03}	0.26 ^{+0.08} _{-0.08}	AGN 1
J105624.2 033522	0094800101	10 56 24.3 03 35 22.9	9.05	1:34 0:09	0.57 ^{+0.05} _{-0.05}	0.73 ^{+0.08} _{-0.08}	
J110050.6 344331	0112880201	11 00 50.6 34 43 32.0	12.71	3:97 0:18	0.48 ^{+0.04} _{-0.04}	0.33 ^{+0.07} _{-0.07}	
J110119.0 345303	0112880201	11 01 19.1 34 53 03.8	12.61	1:33 0:13	0.73 ^{+0.06} _{-0.06}	0.40 ^{+0.17} _{-0.18}	
J110244.1 344604	0112880201	11 02 44.2 34 46 04.7	11.26	1:17 0:11	0.49 ^{+0.08} _{-0.08}	0.71 ^{+0.11} _{-0.11}	
J110320.1+ 355803	0070340301	11 03 20.1 + 35 58 03.7	4.40	7:76 0:32	0.98 ^{+0.01} _{-0.01}	1.00 ^{+0.76} _{-0.00}	stack
J110652.0 182738	0112630101	11 06 52.0 18 27 38.9	7.47	1:14 0:12	0.52 ^{+0.09} _{-0.09}	0.99 ^{+0.20} _{-0.01}	MM
J111654.8+ 180304	0099030101	11 16 54.8 + 18 03 04.6	7.71	1:29 0:13	0.76 ^{+0.05} _{-0.05}	0.66 ^{+0.16} _{-0.17}	AGN 2
J111928.5+ 130250 ^c	0093641101	11 19 28.6 + 13 02 50.7	11.81	1:74 0:18	0.60 ^{+0.08} _{-0.08}	0.29 ^{+0.19} _{-0.18}	AGN 1
J111933.0+ 212756	0111290401	11 19 33.0 + 21 27 56.7	10.33	1:59 0:18	0.42 ^{+0.11} _{-0.11}	0.70 ^{+0.15} _{-0.15}	AGN 1
J111942.1+ 211516	0111290401	11 19 42.2 + 21 15 16.2	8.75	1:01 0:14	0.76 ^{+0.09} _{-0.09}	0.17 ^{+0.26} _{-0.27}	AGN 1
J112022.3+ 125252	0093641101	11 20 22.3 + 12 52 52.7	6.84	2:56 0:19	0.66 ^{+0.06} _{-0.06}	0.61 ^{+0.13} _{-0.13}	AGN 1
J112046.7+ 125429	0093641101	11 20 46.8 + 12 54 30.0	9.21	3:07 0:24	0.67 ^{+0.06} _{-0.06}	0.66 ^{+0.13} _{-0.14}	AGN 1
J113106.9+ 312518	0102040201	11 31 07.0 + 31 25 18.2	11.18	1:33 0:12	0.53 ^{+0.08} _{-0.08}	0.15 ^{+0.14} _{-0.14}	AGN 1
J113121.8+ 310252	0102040201	11 31 21.8 + 31 02 52.6	11.58	1:67 0:12	0.93 ^{+0.07} _{-0.07}	0.35 ^{+0.08} _{-0.08}	AGN 2
J113128.6 195903	0042341001	11 31 28.7 19 59 03.4	7.35	1:50 0:13	0.67 ^{+0.07} _{-0.07}	0.73 ^{+0.14} _{-0.14}	AGN 2
J113148.7+ 311358	0102040201	11 31 48.7 + 31 13 58.8	8.45	1:54 0:10	0.10 ^{+0.06} _{-0.07}	0.43 ^{+0.07} _{-0.07}	AGN 2
J113837.9 373402	0112210101	11 38 37.9 37 34 02.0	11.33	1:79 0:12	0.60 ^{+0.05} _{-0.05}	0.44 ^{+0.11} _{-0.10}	AGN 1?
J115317.9+ 364712	0112551401	11 53 17.9 + 36 47 12.5	12.82	1:08 0:21	0.46 ^{+0.19} _{-0.19}	1.00 ^{+0.76} _{-0.00}	AGN 1
J115846.9+ 551625	0090020101	11 58 46.9 + 55 16 25.4	12.98	1:74 0:25	0.65 ^{+0.10} _{-0.10}	0.98 ^{+0.45} _{-0.02}	AGN 1
J120359.1+ 443715	0109141401	12 03 59.2 + 44 37 15.1	10.37	1:37 0:07	0.73 ^{+0.03} _{-0.03}	0.63 ^{+0.09} _{-0.09}	
J120413.7+ 443149	0109141401	12 04 13.8 + 44 31 49.7	11.47	1:28 0:08	0.60 ^{+0.05} _{-0.04}	0.70 ^{+0.08} _{-0.08}	
J121501.7+ 140113	0112610101	12 15 01.8 + 14 01 13.7	11.06	1:84 0:10	0.58 ^{+0.04} _{-0.04}	0.68 ^{+0.07} _{-0.08}	
J122017.5+ 752217 ^c	0124110101	12 20 17.5 + 75 22 17.8	6.45	10:78 0:35	0.87 ^{+0.03} _{-0.03}	1.0 ^{+0.30} _{-0.12}	GAL
J122350.4+ 752231	0124110101	12 23 50.5 + 75 22 31.7	9.04	1:49 0:13	0.55 ^{+0.07} _{-0.07}	0.72 ^{+0.11} _{-0.11}	AGN 1
J122628.9+ 333626	0070340501	12 26 28.9 + 33 36 26.6	10.57	1:08 0:13	0.83 ^{+0.06} _{-0.06}	0.64 ^{+0.25} _{-0.32}	
J122655.1+ 012002	0110990201	12 26 55.1 + 01 20 02.8	10.96	1:32 0:16	0.80 ^{+0.07} _{-0.07}	0.78 ^{+0.20} _{-0.20}	
J122656.5+ 013126	0110990201	12 26 56.6 + 01 31 26.2	5.85	1:41 0:12	0.11 ^{+0.09} _{-0.09}	0.34 ^{+0.10} _{-0.10}	AGN 2?
J122658.1+ 333246 ^{c;d}	0070340501	12 26 58.1 + 33 32 46.5	3.68	7:15 0:25	0.51 ^{+0.04} _{-0.04}	0.80 ^{+0.04} _{-0.04}	CL
J122751.2+ 333842	0070340501	12 27 51.3 + 33 38 43.0	9.80	1:87 0:17	0.85 ^{+0.04} _{-0.04}	0.39 ^{+0.23} _{-0.23}	star
J122803.7+ 333957	0070340501	12 28 03.7 + 33 39 57.1	12.60	1:15 0:16	0.58 ^{+0.10} _{-0.10}	1.00 ^{+0.00} _{-0.00}	
J122837.3+ 015720	0126700301	12 28 37.3 + 01 57 20.6	9.33	2:16 0:09	0.88 ^{+0.02} _{-0.02}	0.99 ^{+0.03} _{-0.01}	star
J122942.3+ 015525	0126700301	12 29 42.4 + 01 55 25.4	11.80	2:06 0:11	0.84 ^{+0.03} _{-0.03}	0.83 ^{+0.09} _{-0.09}	star

Name XBS...	Obs. ID	RA; DEC (J2000)	O Axis arcmin	Rate 10^{-2} cts/s	HR 2	HR 3	Class
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
J123036.2+ 642531	0124900101	12 30 36.3 + 64 25 31.8	12.66	1.09 0.09	$0.70^{+0.05}_{-0.06}$	$0.49^{+0.15}_{-0.16}$	
J123116.5+ 641115	0124900101	12 31 16.5 + 64 11 15.1	3.46	1.06 0.06	$0.51^{+0.05}_{-0.05}$	$0.45^{+0.09}_{-0.09}$	
J123208.7+ 640304	0124900101	12 32 08.8 + 64 03 04.3	12.00	1.16 0.11	$0.82^{+0.05}_{-0.05}$	$0.91^{+0.11}_{-0.07}$	star
J123218.5+ 640311	0124900101	12 32 18.6 + 64 03 11.3	12.30	1.48 0.12	$0.58^{+0.06}_{-0.06}$	$0.52^{+0.13}_{-0.12}$	star
J123538.6+ 621644 ^d	0111550101	12 35 38.6 + 62 16 44.7	8.98	2.06 0.09	$0.59^{+0.03}_{-0.03}$	$0.59^{+0.07}_{-0.07}$	AGN 1
J123549.1 395026	0006220201	12 35 49.1 39 50 26.8	5.25	3.23 0.10	$0.94^{+0.01}_{-0.01}$	$1.00^{+0.14}_{-0.00}$	star
J123600.7 395217	0006220201	12 36 00.7 39 52 18.0	5.62	17.71 0.22	$0.77^{+0.01}_{-0.01}$	$0.77^{+0.02}_{-0.02}$	star
J123759.6+ 621102	0111550101	12 37 59.7 + 62 11 02.9	8.51	1.95 0.10	$0.59^{+0.04}_{-0.04}$	$0.55^{+0.08}_{-0.08}$	AGN 1
J123800.9+ 621338	0111550101	12 38 00.9 + 62 13 38.5	8.47	1.48 0.08	$0.78^{+0.03}_{-0.03}$	$0.36^{+0.13}_{-0.12}$	AGN 1
J124214.1 112512	0136950201	12 42 14.1 11 25 12.3	8.24	1.55 0.09	$0.57^{+0.05}_{-0.05}$	$0.33^{+0.10}_{-0.10}$	AGN 1
J124557.6+ 022659	0051760101	12 45 57.7 + 02 26 60.0	10.56	1.34 0.17	$0.55^{+0.09}_{-0.09}$	$0.59^{+0.16}_{-0.17}$	The XMM
J124607.6+ 022153	0051760101	12 46 07.7 + 02 21 53.6	6.81	2.41 0.17	$0.62^{+0.05}_{-0.05}$	$1.00^{+0.10}_{-0.00}$	AGN 1
J124641.8+ 022412	0051760101	12 46 41.8 + 02 24 12.1	2.78	3.00 0.17	$0.58^{+0.05}_{-0.05}$	$0.48^{+0.09}_{-0.09}$	AGN 1
J124647.9+ 020955	0051760101	12 46 48.0 + 02 09 55.8	12.52	1.03 0.19	$0.39^{+0.14}_{-0.14}$	$0.93^{+0.10}_{-0.06}$	Newton
J124903.6 061049 ^d	0060370201	12 49 03.6 06 10 49.2	11.63	2.54 0.13	$0.61^{+0.04}_{-0.04}$	$0.55^{+0.08}_{-0.08}$	AGN 1
J124914.6 060910	0060370201	12 49 14.7 06 09 11.0	9.73	1.06 0.08	$0.50^{+0.06}_{-0.06}$	$0.75^{+0.08}_{-0.08}$	Brown
J124938.7 060444	0060370201	12 49 38.7 06 04 44.9	8.20	35.88 0.41	$0.88^{+0.01}_{-0.01}$	$0.85^{+0.02}_{-0.02}$	star
J124949.4 060722	0060370201	12 49 49.5 06 07 22.0	11.93	1.35 0.11	$0.58^{+0.06}_{-0.06}$	$0.43^{+0.12}_{-0.12}$	Stern
J125457.2+ 564940	0081340201	12 54 57.2 + 56 49 40.7	10.79	1.72 0.13	$0.46^{+0.06}_{-0.06}$	$0.63^{+0.09}_{-0.10}$	AGN 1
J125648.4+ 570349	0081340201	12 56 48.5 + 57 03 49.1	12.37	1.19 0.12	$0.38^{+0.09}_{-0.09}$	$0.72^{+0.11}_{-0.11}$	Surprise
J130616.8+ 175923	0017940101	13 06 16.9 + 17 59 23.7	8.09	1.01 0.06	$0.50^{+0.05}_{-0.05}$	$0.44^{+0.08}_{-0.08}$	Dipper
J130619.7 233857	0002940101	13 06 19.7 23 38 57.9	10.84	2.83 0.32	$0.68^{+0.08}_{-0.08}$	$0.56^{+0.21}_{-0.21}$	Material
J130658.1 234849	0002940101	13 06 58.2 23 48 49.7	8.60	1.39 0.19	$0.52^{+0.11}_{-0.11}$	$0.51^{+0.21}_{-0.19}$	AGN 1
J132038.0+ 341124	0093640401	13 20 38.1 + 34 11 24.3	3.10	2.91 0.13	$0.43^{+0.04}_{-0.04}$	$0.48^{+0.06}_{-0.06}$	AGN 1
J132052.5+ 341742	0093640401	13 20 52.5 + 34 17 42.9	10.01	1.50 0.12	$0.53^{+0.07}_{-0.07}$	$0.58^{+0.12}_{-0.12}$	AGN 1
J132101.6+ 340656	0093640401	13 21 01.6 + 34 06 56.6	5.61	3.88 0.15	$0.71^{+0.03}_{-0.03}$	$0.57^{+0.08}_{-0.07}$	AGN 1
J132105.5+ 341459 ^d	0093640401	13 21 05.5 + 34 14 59.4	9.10	2.74 0.15	$0.56^{+0.05}_{-0.05}$	$0.56^{+0.09}_{-0.09}$	Line
J133023.8+ 241707	0100240201	13 30 23.8 + 24 17 07.3	6.22	1.15 0.07	$0.68^{+0.04}_{-0.05}$	$0.41^{+0.12}_{-0.11}$	Material
J133026.6+ 241520 ^d	0100240201	13 30 26.7 + 24 15 20.5	4.91	2.08 0.09	$0.82^{+0.02}_{-0.03}$	$0.57^{+0.10}_{-0.10}$	Material
J133120.2+ 242305	0100240201	13 31 20.2 + 24 23 05.1	11.78	1.77 0.11	$0.56^{+0.05}_{-0.05}$	$0.60^{+0.09}_{-0.09}$	Material
J133232.6+ 111220 ^d	0061940101	13 32 32.7 + 11 12 20.7	10.70	1.05 0.18	$0.35^{+0.15}_{-0.15}$	$1.00^{+0.31}_{-0.00}$	Material
J133321.2+ 503102	0111160101	13 33 21.3 + 50 31 02.2	9.21	1.79 0.08	$0.91^{+0.02}_{-0.02}$	$0.49^{+0.16}_{-0.16}$	star
J133626.9 342636	0111570201	13 36 26.9 34 26 36.7	11.19	1.00 0.08	$0.90^{+0.03}_{-0.03}$	$1.00^{+0.42}_{-0.00}$	star
J133807.5+ 242411	0096010101	13 38 07.6 + 24 24 11.8	11.17	1.69 0.11	$0.66^{+0.05}_{-0.05}$	$0.65^{+0.10}_{-0.10}$	AGN 1
J133942.6 315004	0035940301	13 39 42.6 31 50 04.8	11.91	1.67 0.11	$0.36^{+0.05}_{-0.06}$	$0.20^{+0.08}_{-0.08}$	AGN 1?
J134732.0+ 582103	0112250201	13 47 32.0 + 58 21 03.8	8.38	3.43 0.13	$0.86^{+0.02}_{-0.02}$	$0.81^{+0.07}_{-0.08}$	star
J134749.9+ 582111	0112250201	13 47 49.9 + 58 21 11.0	8.53	12.44 0.24	$0.60^{+0.02}_{-0.02}$	$0.54^{+0.03}_{-0.03}$	AGN 1
J140100.0 110942	0109910101	14 01 00.0 11 09 42.2	8.70	1.12 0.07	$0.57^{+0.05}_{-0.05}$	$0.66^{+0.09}_{-0.09}$	

Name XBS...	Obs. ID	RA; DEC (J2000)	O Axis arcmin	Rate 10^{-2} cts/s	HR 2	HR 3	Class
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
J140102.0 111224	0109910101	14 01 02.0 11 12 24.3	9.27	8.34 0.18	$0.52^{+0.02}_{-0.02}$	$0.46^{+0.03}_{-0.03}$	AGN 2?
J140127.7+ 025605	0098010101	14 01 27.7 + 02 56 05.4	7.25	7.04 0.19	$0.38^{+0.02}_{-0.02}$	$0.53^{+0.04}_{-0.04}$	AGN 1
J140219.6 110458	0109910101	14 02 19.7 11 04 58.8	11.40	1.16 0.08	$0.94^{+0.02}_{-0.02}$	$0.16^{+0.29}_{-0.27}$	star
J140921.1+ 261336	0092850501	14 09 21.2 + 26 13 36.8	4.78	1.24 0.06	$0.32^{+0.04}_{-0.05}$	$0.53^{+0.06}_{-0.06}$	star
J140936.9+ 261632	0092850501	14 09 36.9 + 26 16 32.1	3.44	1.11 0.06	$0.92^{+0.02}_{-0.02}$	$0.70^{+0.16}_{-0.17}$	star
J141235.8 030909	0013140101	14 12 35.8 03 09 09.3	10.29	1.72 0.15	$0.40^{+0.08}_{-0.08}$	$0.59^{+0.11}_{-0.11}$	star
J141523.8+ 113737	0112250301	14 15 23.8 + 11 37 37.3	9.64	1.16 0.09	$0.70^{+0.05}_{-0.05}$	$0.54^{+0.15}_{-0.15}$	star
J141531.5+ 113156	0112250301	14 15 31.6 + 11 31 56.2	4.15	3.15 0.12	$0.55^{+0.03}_{-0.03}$	$0.47^{+0.06}_{-0.06}$	AGN 1
J141643.8+ 521434	0127921001	14 16 43.8 + 52 14 34.2	10.17	1.41 0.07	$0.58^{+0.04}_{-0.04}$	$0.50^{+0.08}_{-0.08}$	The
J141658.8+ 521202	0127921001	14 16 58.8 + 52 12 02.8	11.98	1.01 0.08	$0.71^{+0.05}_{-0.05}$	$0.28^{+0.14}_{-0.14}$	
J141722.6+ 251335	0109960101	14 17 22.7 + 25 13 35.2	9.86	1.33 0.12	$0.65^{+0.06}_{-0.06}$	$1.00^{+0.21}_{-0.00}$	AGN 1
J141736.3+ 523028	0127921001	14 17 36.3 + 52 30 28.5	7.67	1.15 0.06	$0.65^{+0.04}_{-0.04}$	$0.40^{+0.09}_{-0.09}$	AGN 1
J141809.1+ 250040	0109960101	14 18 09.1 + 25 00 40.3	7.85	1.31 0.09	$0.59^{+0.06}_{-0.05}$	$0.47^{+0.12}_{-0.12}$	AGN 1
J141830.5+ 251052 ^c	0109960101	14 18 30.5 + 25 10 52.6	7.58	14.99 0.37	$0.52^{+0.04}_{-0.04}$	$0.65^{+0.07}_{-0.07}$	C L
J142800.1+ 424409	0111850201	14 28 00.1 + 42 44 09.5	7.10	1.53 0.07	$0.90^{+0.02}_{-0.02}$	$0.48^{+0.17}_{-0.16}$	star
J142901.2+ 423048	0111850201	14 29 01.2 + 42 30 48.8	10.84	1.70 0.11	$0.98^{+0.01}_{-0.01}$	$0.56^{+0.28}_{-0.30}$	star
J143835.1+ 642928	0111530101	14 38 35.1 + 64 29 28.3	12.34	1.31 0.09	$0.18^{+0.07}_{-0.07}$	$0.37^{+0.07}_{-0.06}$	AGN 1?
J143923.1+ 640912	0111530101	14 39 23.2 + 64 09 12.4	8.60	2.36 0.09	$0.99^{+0.01}_{-0.01}$	$0.51^{+0.40}_{-0.46}$	star
J144021.0+ 642144	0111530101	14 40 21.0 + 64 21 44.1	9.79	1.42 0.07	$0.47^{+0.04}_{-0.04}$	$0.37^{+0.07}_{-0.08}$	star
J144937.5+ 090826	0057560301	14 49 37.6 + 09 08 26.0	8.83	1.86 0.10	$0.52^{+0.04}_{-0.04}$	$0.54^{+0.07}_{-0.07}$	
J144945.8+ 085921	0057560301	14 49 45.9 + 08 59 21.8	4.14	1.04 0.06	$0.56^{+0.05}_{-0.04}$	$0.51^{+0.09}_{-0.09}$	dip
J145857.1 313535	0067750101	14 58 57.1 31 35 36.0	8.89	1.03 0.08	$0.55^{+0.06}_{-0.06}$	$0.45^{+0.12}_{-0.12}$	AGN Survey
J150428.3+ 101856	0070740301	15 04 28.4 + 10 18 57.0	9.91	1.67 0.17	$0.37^{+0.09}_{-0.09}$	$1.00^{+0.13}_{-0.00}$	AGN 1
J151815.0+ 060851	0018741001	15 18 15.1 + 06 08 51.5	9.43	1.12 0.19	$0.77^{+0.10}_{-0.10}$	$0.22^{+0.33}_{-0.32}$	
J151832.3+ 062357	0018741001	15 18 32.4 + 06 23 57.6	10.39	1.72 0.27	$0.72^{+0.10}_{-0.09}$	$0.70^{+0.23}_{-0.27}$	GAL
J153156.6 082610	0100240801	15 31 56.6 08 26 11.0	8.75	1.32 0.11	$0.96^{+0.02}_{-0.02}$	$1.00^{+0.96}_{-0.00}$	star
J153205.7 082952	0100240801	15 32 05.8 08 29 52.6	4.77	1.05 0.08	$0.45^{+0.07}_{-0.06}$	$0.79^{+0.08}_{-0.08}$	AGN 1
J153419.0+ 011808	0112190401	15 34 19.1 + 01 18 08.2	10.91	1.37 0.15	$0.62^{+0.08}_{-0.08}$	$0.30^{+0.18}_{-0.18}$	AGN 1
J153452.3+ 013104 ^d	0112190401	15 34 52.4 + 01 31 04.6	10.57	9.44 0.38	$0.39^{+0.04}_{-0.04}$	$0.55^{+0.05}_{-0.05}$	AGN 1
J153456.1+ 013033	0112190401	15 34 56.2 + 01 30 33.4	9.89	3.05 0.23	$0.64^{+0.05}_{-0.05}$	$0.64^{+0.11}_{-0.11}$	AGN 1
J160645.9+ 081525	0067340601	16 06 46.0 + 08 15 25.1	12.61	1.28 0.17	$0.81^{+0.08}_{-0.08}$	$0.23^{+0.10}_{-0.10}$	AGN 2?
J160706.6+ 075709	0067340601	16 07 06.7 + 07 57 09.5	7.90	1.49 0.14	$0.65^{+0.07}_{-0.06}$	$0.34^{+0.16}_{-0.16}$	AGN 1
J160731.5+ 081202	0067340601	16 07 31.6 + 08 12 02.9	8.43	1.67 0.14	$0.75^{+0.05}_{-0.05}$	$0.44^{+0.17}_{-0.18}$	AGN 1?
J161544.2+ 121708	0103460801	16 15 44.2 + 12 17 08.3	11.87	1.41 0.18	$0.64^{+0.09}_{-0.09}$	$0.18^{+0.22}_{-0.24}$	
J161615.1+ 121353	0103460801	16 16 15.1 + 12 13 53.2	3.65	1.11 0.10	$0.65^{+0.07}_{-0.07}$	$1.00^{+0.19}_{-0.00}$	
J161825.4+ 124145	0103461001	16 18 25.5 + 12 41 45.5	12.82	1.14 0.15	$0.76^{+0.09}_{-0.09}$	$0.17^{+0.31}_{-0.31}$	AGN 1
J162813.9+ 780342	0061940301	16 28 14.0 + 78 03 42.7	10.92	2.64 0.34	$0.67^{+0.09}_{-0.09}$	$0.33^{+0.24}_{-0.24}$	AGN 1
J162911.1+ 780442	0061940301	16 29 11.2 + 78 04 42.7	8.13	1.66 0.22	$0.81^{+0.08}_{-0.08}$	$1.00^{+0.76}_{-0.00}$	star

Name XBS...	Obs. ID	RA; DEC (J2000)	O Axis arcmin	Rate 10^{-2} cts/s	HR 2	HR 3	Class
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
J162923.3+ 781306	0061940301	16 29 23.4 + 78 13 06.6	5.11	1.41 0.18	0.19 ^{+0.13} _{-0.13}	0.68 ^{+0.14} _{-0.14}	
J162944.8+ 781128	0061940301	16 29 44.8 + 78 11 28.5	3.57	1.77 0.20	0.99 ^{+0.02} _{-0.01}	1.00 ^{+2.00} _{-0.00}	star
J163141.1+ 781239	0061940301	16 31 41.2 + 78 12 39.3	2.99	1.39 0.17	0.61 ^{+0.10} _{-0.11}	0.84 ^{+0.15} _{-0.16}	AGN 1
J163223.6+ 052547	0112230301	16 32 23.6 + 05 25 47.1	10.44	1.89 0.14	0.71 ^{+0.05} _{-0.05}	0.23 ^{+0.14} _{-0.15}	AGN 1
J163309.8+ 571039	0049540101	16 33 09.8 + 57 10 39.3	9.67	2.87 0.27	0.43 ^{+0.07} _{-0.07}	0.49 ^{+0.10} _{-0.11}	AGN 1
J163332.3+ 570520	0049540101	16 33 32.4 + 57 05 20.2	7.90	1.03 0.16	0.84 ^{+0.06} _{-0.06}	1.00 ^{+0.63} _{-0.00}	AGN 1?
J163427.5+ 781002	0061940301	16 34 27.6 + 78 10 02.5	11.00	1.79 0.30	0.62 ^{+0.13} _{-0.14}	0.71 ^{+0.24} _{-0.26}	AGN 1
J164237.9+ 030014 ^d	0067340501	16 42 38.0 + 03 00 14.3	11.90	1.22 0.14	0.63 ^{+0.09} _{-0.08}	0.38 ^{+0.20} _{-0.20}	galaxy
J165313.3+ 021645	0101640101	16 53 13.3 + 02 16 45.6	8.16	1.36 0.13	0.96 ^{+0.03} _{-0.02}	1.00 ^{+1.30} _{-0.00}	star
J165314.4+ 141943	0113070101	16 53 14.5 + 14 19 43.6	11.13	1.54 0.21	0.42 ^{+0.12} _{-0.12}	0.67 ^{+0.17} _{-0.16}	The X
J165406.6+ 142123	0113070101	16 54 06.6 + 14 21 23.6	3.89	1.12 0.13	0.45 ^{+0.11} _{-0.10}	0.76 ^{+0.14} _{-0.14}	AGN 1
J165425.3+ 142159	0113070101	16 54 25.4 + 14 21 59.3	7.44	11.86 0.53	0.63 ^{+0.03} _{-0.03}	0.61 ^{+0.07} _{-0.07}	AGN 1
J165448.5+ 141311	0113070101	16 54 48.5 + 14 13 11.6	12.71	6.78 0.56	0.54 ^{+0.07} _{-0.07}	0.42 ^{+0.13} _{-0.13}	AGN 2
J165710.5+ 352024	0111060101	16 57 10.5 + 35 20 24.0	7.98	1.93 0.20	0.90 ^{+0.04} _{-0.04}	1.00 ^{+0.76} _{-0.00}	star
J165800.7+ 352333	0111060101	16 58 00.7 + 35 23 33.5	3.73	1.40 0.13	0.51 ^{+0.08} _{-0.08}	0.69 ^{+0.12} _{-0.13}	AGN 2?
J172230.6+ 341344	0102040101	17 22 30.7 + 34 13 44.0	11.20	1.74 0.25	0.65 ^{+0.10} _{-0.11}	0.31 ^{+0.28} _{-0.27}	AGN 1
J185518.7 462504	0067340101	18 55 18.8 46 25 04.0	6.69	1.55 0.20	0.40 ^{+0.09} _{-0.09}	1.00 ^{+0.13} _{-0.00}	AGN 1
J185613.7 462239 ^a	0067340101	18 56 13.8 46 22 39.2	10.44	3.02 0.30	0.47 ^{+0.07} _{-0.07}	0.01 ^{+0.11} _{-0.10}	AGN 1
J193138.9 725115	0081341001	19 31 39.0 72 51 15.0	12.10	1.48 0.16	0.42 ^{+0.09} _{-0.09}	0.62 ^{+0.13} _{-0.14}	AGN 1
J193248.8 723355 ^a	0081341001	19 32 48.8 72 33 55.2	8.39	1.37 0.12	0.23 ^{+0.08} _{-0.08}	0.08 ^{+0.10} _{-0.10}	AGN 2?
J204043.4 004548 ^a	0111180201	20 40 43.5 00 45 48.2	10.55	1.29 0.15	0.04 ^{+0.11} _{-0.11}	0.35 ^{+0.12} _{-0.12}	AGN 2
J204047.5 005853	0111180201	20 40 47.6 00 58 53.8	11.50	1.05 0.14	0.33 ^{+0.12} _{-0.12}	0.64 ^{+0.14} _{-0.15}	BLS
J204159.2 321439	0111510101	20 41 59.3 32 14 40.0	11.80	3.36 0.23	0.54 ^{+0.05} _{-0.05}	0.79 ^{+0.08} _{-0.07}	AGN 1
J204204.1 321601	0111510101	20 42 04.1 32 16 01.5	10.68	1.39 0.14	0.52 ^{+0.08} _{-0.08}	0.80 ^{+0.10} _{-0.11}	AGN 1
J204208.2 323523	0111510101	20 42 08.3 32 35 23.6	9.78	1.16 0.12	0.63 ^{+0.08} _{-0.08}	0.89 ^{+0.09} _{-0.10}	AGN 1
J204548.4 025234	0112600501	20 45 48.4 02 52 34.4	8.74	1.29 0.14	0.38 ^{+0.10} _{-0.10}	0.76 ^{+0.12} _{-0.11}	AGN 1
J205411.9 160804	0083210101	20 54 11.9 16 08 04.8	12.55	1.97 0.23	0.58 ^{+0.09} _{-0.09}	0.69 ^{+0.16} _{-0.16}	AGN 1
J205429.9 154937	0083210101	20 54 30.0 15 49 37.8	6.59	1.14 0.13	0.65 ^{+0.08} _{-0.08}	0.80 ^{+0.14} _{-0.19}	AGN 1
J205635.7 044717 ^a	0112190601	20 56 35.8 04 47 17.9	9.96	3.41 0.21	0.65 ^{+0.04} _{-0.05}	0.48 ^{+0.11} _{-0.11}	AGN 1
J205829.9 423634 ^a	0081340401	20 58 30.0 42 36 35.0	2.42	5.01 0.19	0.57 ^{+0.03} _{-0.03}	0.47 ^{+0.06} _{-0.06}	AGN 1
J205847.0 423704	0081340401	20 58 47.0 42 37 04.7	4.11	1.71 0.12	0.93 ^{+0.03} _{-0.03}	1.00 ^{+0.54} _{-0.00}	star
J210325.4 112011	0041150101	21 03 25.4 11 20 11.5	11.29	1.01 0.09	0.49 ^{+0.07} _{-0.07}	0.22 ^{+0.12} _{-0.13}	AGN 1
J210355.3 121858	0038540301	21 03 55.3 12 18 58.2	11.02	1.56 0.17	0.50 ^{+0.09} _{-0.09}	0.80 ^{+0.12} _{-0.10}	AGN 1
J212635.8 445046	0088020201	21 26 35.8 44 50 46.1	11.25	3.03 0.25	0.92 ^{+0.03} _{-0.03}	0.37 ^{+0.32} _{-0.32}	
J212759.5 443924	0088020201	21 27 59.6 44 39 24.7	9.99	1.08 0.12	0.65 ^{+0.08} _{-0.08}	0.47 ^{+0.18} _{-0.19}	
J213002.3 153414 ^a	0103060101	21 30 02.3 15 34 14.1	12.83	3.95 0.21	0.65 ^{+0.04} _{-0.04}	0.50 ^{+0.09} _{-0.09}	AGN 1
J213719.6 433347	0109463501	21 37 19.6 43 33 47.0	10.85	2.41 0.27	0.27 ^{+0.10} _{-0.10}	0.74 ^{+0.11} _{-0.11}	AGN 1
J213729.7 423601	0061940201	21 37 29.8 42 36 01.6	7.02	1.51 0.23	0.56 ^{+0.12} _{-0.13}	0.33 ^{+0.25} _{-0.27}	

Name XBS...	Obs. ID	RA; DEC (J2000)	O Axis arcmin	Rate 10^{-2} cts/s	HR 2	HR 3	Class			
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)			
J213733.2	434800	0109463501	21 37 33.2	43 48 00.9	7.07	1:03	0:14	$0:91^{+0:06}_{-0:06}$	$0:05^{+0:48}_{-0:52}$	AGN 1
J213757.6	422334	0061940201	21 37 57.6	42 23 34.2	12.68	1:37	0:27	$0:59^{+0:15}_{-0:16}$	$0:15^{+0:36}_{-0:34}$	AGN 1
J213820.2	142536	0092850201	21 38 20.2	14 25 37.0	11.17	2:10	0:17	$0:29^{+0:07}_{-0:07}$	$0:45^{+0:10}_{-0:10}$	AGN 1
J213824.0	423019 ^a	0061940201	21 38 24.0	42 30 19.2	6.50	2:88	0:27	$0:62^{+0:08}_{-0:08}$	$0:36^{+0:18}_{-0:18}$	AGN 1
J213829.8	423958	0061940201	21 38 29.9	42 39 58.9	5.59	1:45	0:19	$0:66^{+0:10}_{-0:10}$	$0:50^{+0:24}_{-0:24}$	AGN 1
J213840.5	424241	0061940201	21 38 40.6	42 42 41.3	8.91	1:89	0:26	$1:00^{+0:07}_{-0:00}$	{ star	
J213852.2	434714	0109463501	21 38 52.3	43 47 14.9	11.21	1:32	0:23	$0:68^{+0:11}_{-0:12}$	$0:36^{+0:31}_{-0:33}$	AGN 1
J214041.4	234720 ^a	0008830101	21 40 41.5	23 47 20.1	9.80	3:46	0:22	$0:57^{+0:05}_{-0:05}$	$0:39^{+0:10}_{-0:10}$	AGN 1
J215218.0	302721	0103060401	21 52 18.1	30 27 21.6	4.80	1:15	0:08	$0:69^{+0:05}_{-0:05}$	$0:55^{+0:12}_{-0:12}$..
J215244.2	302407	0103060401	21 52 44.3	30 24 07.4	11.03	3:28	0:16	$0:65^{+0:04}_{-0:04}$	$0:68^{+0:08}_{-0:07}$	AGN 1
J215323.7+	173018	0111270101	21 53 23.7	+ 17 30 18.6	12.09	1:08	0:16	$0:37^{+0:12}_{-0:13}$	$0:73^{+0:16}_{-0:14}$	star
J220320.8+	184930	0130920101	22 03 20.9	+ 18 49 30.3	4.03	1:58	0:11	$0:49^{+0:06}_{-0:06}$	$0:39^{+0:10}_{-0:10}$	AGN 2?
J220446.8	014535	0012440301	22 04 46.8	01 45 35.6	11.22	1:23	0:10	$0:45^{+0:07}_{-0:07}$	$0:69^{+0:10}_{-0:09}$	AGN 1
J220601.5	015346 ^a	0012440301	22 06 01.5	01 53 47.0	12.92	1:85	0:12	$0:41^{+0:06}_{-0:06}$	$0:41^{+0:09}_{-0:09}$	AGN 1
J221623.3	174317	0106660101	22 16 23.4	17 43 17.9	12.29	1:15	0:08	$0:55^{+0:05}_{-0:06}$	$0:57^{+0:10}_{-0:10}$	AGN 1
J221722.4	082018	0009650201	22 17 22.5	08 20 18.2	8.18	1:76	0:13	$0:38^{+0:07}_{-0:07}$	$0:48^{+0:10}_{-0:10}$	AGN 1
J221729.3	081154	0009650201	22 17 29.4	08 11 54.2	11.12	2:00	0:16	$0:54^{+0:06}_{-0:06}$	$0:53^{+0:12}_{-0:11}$	AGN 1
J221750.4	083210	0009650201	22 17 50.4	08 32 10.8	11.28	1:32	0:12	$0:79^{+0:06}_{-0:06}$	$1:00^{+0:37}_{-0:00}$	star
J221821.9	081332	0009650201	22 18 22.0	08 13 33.0	9.91	2:10	0:13	$0:63^{+0:05}_{-0:05}$	$0:85^{+0:07}_{-0:07}$	AGN 1
J221951.6+	120123 ^d	0103861201	22 19 51.6	+ 12 01 23.1	10.33	1:33	0:15	$0:36^{+0:10}_{-0:10}$	$0:39^{+0:15}_{-0:15}$	AGN 2
J222852.2	050915	0100440101	22 28 52.3	05 09 15.4	11.01	1:33	0:08	$0:97^{+0:01}_{-0:01}$	$1:00^{+0:76}_{-0:00}$	star
J223547.9	255836	0111790101	22 35 48.0	25 58 37.0	4.41	1:21	0:06	$0:55^{+0:04}_{-0:04}$	$0:50^{+0:08}_{-0:08}$	AGN 1
J223555.0	255833	0111790101	22 35 55.0	25 58 33.9	4.89	1:03	0:06	$0:61^{+0:04}_{-0:04}$	$0:66^{+0:08}_{-0:09}$	AGN 1
J223949.8+	080926	0103860301	22 39 49.9	+ 08 09 26.4	9.24	1:07	0:21	$0:65^{+0:14}_{-0:14}$	$0:09^{+0:29}_{-0:31}$	AGN 1
J224756.6	642721	0112240101	22 47 56.6	64 27 21.8	12.76	1:43	0:11	$0:51^{+0:06}_{-0:06}$	$0:62^{+0:10}_{-0:10}$..
J224833.3	511900	0109070401	22 48 33.4	51 19 00.2	9.13	1:18	0:12	$0:91^{+0:04}_{-0:04}$	$0:93^{+0:90}_{-0:06}$	star
J224846.6	505929	0109070401	22 48 46.7	50 59 29.6	10.50	1:03	0:14	$0:91^{+0:05}_{-0:05}$	$1:00^{+1:30}_{-0:00}$	star
J225025.1	643225	0112240101	22 50 25.2	64 32 25.3	10.05	1:00	0:09	$0:66^{+0:06}_{-0:06}$	$0:52^{+0:14}_{-0:14}$	AGN 1
J225050.2	642900	0112240101	22 50 50.3	64 29 00.4	8.87	2:42	0:13	$0:57^{+0:04}_{-0:04}$	$0:45^{+0:08}_{-0:08}$	AGN 1
J225118.0	175951	0081340901	22 51 18.1	17 59 51.3	10.63	3:39	0:20	$0:68^{+0:04}_{-0:04}$	$0:56^{+0:10}_{-0:10}$	AGN 1?
J225349.6	172137	0112910301	22 53 49.6	17 21 37.2	12.50	1:83	0:28	$0:95^{+0:05}_{-0:05}$	$1:00^{+2:00}_{-0:00}$	star
J230400.4	083755	0109130701	23 04 00.4	08 37 55.5	11.19	1:05	0:14	$0:96^{+0:04}_{-0:04}$	$0:33^{+0:39}_{-0:61}$	AGN 1
J230401.0+	031519	0033541001	23 04 01.0	+ 03 15 19.2	11.59	1:06	0:14	$0:69^{+0:09}_{-0:09}$	$0:44^{+0:25}_{-0:25}$	GAL
J230408.2+	031820	0033541001	23 04 08.3	+ 03 18 20.7	11.38	5:66	0:31	$0:92^{+0:02}_{-0:02}$	$1:00^{+0:31}_{-0:00}$	star
J230434.1+	122728	0025541001	23 04 34.1	+ 12 27 28.3	9.68	1:39	0:17	$0:57^{+0:10}_{-0:10}$	$0:31^{+0:21}_{-0:21}$	AGN 2?
J230443.8+	121636	0025541001	23 04 43.9	+ 12 16 36.8	4.30	1:22	0:13	$0:54^{+0:09}_{-0:09}$	$0:77^{+0:12}_{-0:14}$	AGN 1
J230459.6+	121205	0025541001	23 04 59.7	+ 12 12 05.4	7.48	1:35	0:15	$0:49^{+0:10}_{-0:09}$	$0:47^{+0:16}_{-0:16}$	AGN 2?
J230522.1+	122121	0025541001	23 05 22.1	+ 12 21 21.1	6.45	1:28	0:14	$0:37^{+0:09}_{-0:09}$	$0:60^{+0:14}_{-0:13}$	AGN 2

Name XBS...	Obs. ID	RA; DEC (J2000)	O Axis arcmin	Rate 10^{-2} cts/s	HR 2	HR 3	Class
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
J230523.0+121325	0025541001	23 05 23.0 + 12 13 25.4	8.85	1.31	0.15	$0.47^{+0.10}_{-0.10}$	$0.89^{+0.10}_{-0.11}$
J231342.5 423210	0123900101	23 13 42.5 42 32 10.7	11.69	2.65	0.13	$0.64^{+0.04}_{-0.04}$	$0.53^{+0.08}_{-0.08}$
J231541.2 424125	0093640701	23 15 41.3 42 41 25.6	8.42	1.19	0.14	$0.86^{+0.05}_{-0.05}$	$0.40^{+0.33}_{-0.31}$
J231546.5 590313	0109463601	23 15 46.5 59 03 13.6	8.13	1.31	0.22	$0.54^{+0.14}_{-0.14}$	$0.07^{+0.26}_{-0.26}$
J231553.0 423800	0093640701	23 15 53.0 42 38 00.4	4.46	1.17	0.11	$0.84^{+0.05}_{-0.05}$	$0.18^{+0.25}_{-0.23}$
J231601.7 424038	0093640701	23 16 01.7 42 40 38.9	5.87	1.34	0.12	$0.62^{+0.07}_{-0.07}$	$0.62^{+0.13}_{-0.14}$
J231658.8 423853	0093640701	23 16 58.9 42 38 53.6	9.61	2.46	0.18	$0.45^{+0.06}_{-0.06}$	$0.72^{+0.08}_{-0.08}$
J233325.7 152240	0093550401	23 33 25.8 15 22 40.2	6.45	2.87	0.13	$0.87^{+0.02}_{-0.02}$	$0.64^{+0.12}_{-0.13}$
J233421.9 151219	0093550401	23 34 21.9 15 12 19.0	11.23	1.33	0.12	$0.46^{+0.08}_{-0.08}$	$0.29^{+0.14}_{-0.15}$
J235032.3+363156	0100241001	23 50 32.4 + 36 31 56.9	12.23	2.05	0.22	$0.80^{+0.06}_{-0.07}$	$0.38^{+0.28}_{-0.26}$
J235036.9+362204 ^d	0100241001	23 50 37.0 + 36 22 05.0	11.84	2.39	0.24	$0.37^{+0.09}_{-0.09}$	$0.59^{+0.12}_{-0.13}$

Columns are as follows: (1) Source name; (2) XMM-Newton Observation number; (3) Right Ascension and Declination (J2000) of the source (X-ray position); (4) Angular distance (in arcmin) between the source and the MOS2 image center; (5) Source count rate, and 1 error, in the 0.5-4.5 keV energy band (units of 10^{-2} cts/s). In Table 1 we have reported the MOS2 conversion factors between the 0.5-4.5 keV count rate and the flux as a function of the energy spectral index, the hardness ratio HR 2 and the blocking filter; (6) and (7) Hardness ratios computed as described in section 3.3. The errors on the hardness ratios have been evaluated using simulations and correspond to 1 ; (8) Optical spectroscopic classification (AGN 1:broad line AGN; AGN 2:narrow line AGN; GAL:Optically Normal Galaxy; CL: Cluster of Galaxies; BL: BL Lac Object; star: star; ?: Tentative classification; see section 3.2 for details).

NOTE { For two sources (XBS J052155.0 252220 and XBS J213840.5 424241) the hardness ratio HR 3 is undefined since these sources are undetected above 2 keV .

^a A detailed X-ray and optical spectral analysis of these sources have been reported in Caccianiga et al. (2004).

^b A detailed X-ray and optical spectral analysis of these sources have been reported in Severgnini et al. (2003).

^c These sources are more extended than the XMM-Newton EPIC MOS2 Point Spread Function at their off-axis angle. Count rates have been evaluated using aperture photometry.

^d A detailed X-ray spectral analysis of these sources have been reported in Gabici et al. (2004).

Table 4 . Basic information on the XM M -N ewton HBSS sample

Name XBS...	Obs. ID (1)	RA ; DEC (J2000) (2)	O Axis arm in (4)	Rate 10^{-3} cts/s (5)	H R 2 (6)	H R 3 (7)	Class (8)
J002618.5+ 105019 ^b	0001930101	00 26 18.5 + 10 50 19.3	9.56	2:35 0:55 -0.53 + 0:04 0:04 + 0:01	-0.67 + 0:06 0:06 + 0:11	A G N 1	
J013240.1 133307 ^{b,e}	0084230301	01 32 40.1 13 33 07.8	11.90	3:23 0:71 -0.02 + 0:10 0:10 + 0:07	-0.37 + 0:11 0:11 + 0:13	A G N 2	
J013944.0 674909 ^b	0032140401	01 39 44.0 67 49 09.4	2.69	2:05 0:46 -0.50 + 0:07 0:07 + 0:02	-0.43 + 0:13 0:13 + 0:03	A G N 1?	
J014100.6 675328 ^b	0032140401	01 41 00.7 67 53 29.0	6.64	70:95 3:84 -0.45 + 0:02 0:02 + 0:02	-0.38 + 0:03 0:03 + 0:03	star	
J015957.5+ 003309 ^b	0101640201	01 59 57.5 + 00 33 09.7	9.56	3:80 0:92 -0.63 + 0:05 0:05 + 0:04	-0.51 + 0:10 0:10 + 0:07	A G N 1	
J021640.7 044404 ^{b,e}	0112371701	02 16 40.7 04 44 04.9	9.35	2:08 0:49 -0.72 + 0:04 0:05 + 0:02	-0.26 + 0:13 0:13 + 0:04	A G N 1	
J021808.3 045845 ^b	0112371001	02 18 08.3 04 58 45.7	2.26	2:67 0:25 -0.65 + 0:02 0:02 + 0:02	-0.52 + 0:04 0:04 + 0:04	A G N 1	
J021817.4 045113 ^b	0112371001	02 18 17.4 04 51 13.3	9.58	3:30 0:40 -0.48 + 0:02 0:03 + 0:02	-0.56 + 0:04 0:04 + 0:04	A G N 1	
J021822.2 050615 ^{a,b,d}	0112371001	02 18 22.3 05 06 15.7	8.48	4:54 0:43 0.65 + 0:09 0:09 + 0:08	0.40 + 0:07 0:07 + 0:07	A G N 2	
J023713.5 522734 ^b	0098810101	02 37 13.5 52 27 34.4	12.07	3:23 0:61 -0.57 + 0:03 0:03 + 0:03	-0.58 + 0:07 0:06 + 0:05	A G N 1	
J030206.8 000121 ^b	0041170101	03 02 06.9 00 01 21.2	11.90	3:10 0:42 -0.53 + 0:03 0:03 + 0:03	-0.46 + 0:05 0:06 + 0:05	A G N 1	
J030614.1 284019 ^b	0042340501	03 06 14.2 28 40 19.9	10.87	4:61 0:92 -0.49 + 0:05 0:05 + 0:05	-0.35 + 0:09 0:09 + 0:09	A G N 1	
J031015.5 765131 ^b	0122520201	03 10 15.6 76 51 31.5	5.90	4:39 0:46 -0.46 + 0:03 0:03 + 0:03	-0.53 + 0:04 0:04 + 0:04	A G N 1	
J031146.1 550702 ^b	0110970101	03 11 46.1 55 07 02.5	12.38	5:87 1:12 -0.58 + 0:05 0:05 + 0:05	-0.39 + 0:11 0:11 + 0:11	A G N 2	
J031859.2 441627 ^{b,d}	0105660101	03 18 59.3 44 16 27.6	11.55	2:16 0:49 -0.21 + 0:08 0:08 + 0:08	-0.47 + 0:10 0:10 + 0:10	A G N 1	
J033845.7 352253 ^{a,b}	0055140101	03 38 45.8 35 22 53.4	10.49	2:37 0:36 -0.05 + 0:12 0:12 + 0:12	0.27 + 0:11 0:11 + 0:11	A G N 2	
J040658.8 712457 ^{a,b}	0111970301	04 06 58.9 71 24 57.7	12.55	3:41 0:68 0.39 + 0:19 0:19 + 0:19	0.20 + 0:16 0:16 + 0:16	A G N 2	
J040758.9 712833 ^{a,b}	0111970301	04 07 59.0 71 28 33.5	12.12	4:96 0:91 0.49 + 0:17 0:17 + 0:17	0.33 + 0:13 0:13 + 0:13	A G N 2	
J041108.1 711341 ^b	0111970301	04 11 08.1 71 13 41.1	10.52	2:20 0:42 -0.53 + 0:08 0:08 + 0:08	-0.04 + 0:16 0:16 + 0:16	A G N 1	
J050536.6 290050	0111160201	05 05 36.7 29 00 50.8	12.43	2:19 0:44 -0.27 + 0:05 0:05 + 0:05	-0.47 + 0:07 0:07 + 0:07	previous surveys	
J052108.5 251913 ^e	0085640101	05 21 08.5 25 19 13.1	3.17	2:11 0:55 -0.59 + 0:05 0:05 + 0:05	-0.55 + 0:09 0:09 + 0:09	A G N 1	
J052128.9 253032 ^a	0085640101	05 21 28.9 25 30 32.4	10.73	3:08 0:88 1.00 + 0:00 0:17 + 0:00	-0.32 + 0:15 0:15 + 0:15	A G N 1	
J074202.7+ 742625	0123100101	07 42 02.7 + 74 26 25.8	10.86	3:38 0:49 -0.59 + 0:03 0:03 + 0:03	-0.28 + 0:07 0:07 + 0:07	A G N 1	
J074312.1+ 742937	0123100101	07 43 12.1 + 74 29 37.4	5.33	10:92 0:64 -0.61 + 0:01 0:01 + 0:01	-0.56 + 0:02 0:02 + 0:02	A G N 1	
J080411.3+ 650906	0094400301	08 04 11.4 + 65 09 06.2	9.49	2:41 0:50 0.06 + 0:09 0:09 + 0:09	-0.51 + 0:09 0:09 + 0:09	unpublished	
J083737.1+ 254751	0025540301	08 37 37.2 + 25 47 51.1	10.61	7:30 1:12 -0.47 + 0:04 0:04 + 0:04	-0.53 + 0:06 0:06 + 0:06	A G N 1	
J083737.0+ 255151	0025540301	08 37 37.1 + 25 51 51.2	12.25	2:98 0:71 -0.34 + 0:07 0:08 + 0:07	-0.52 + 0:10 0:10 + 0:10	A G N 2	
J091828.4+ 513931	0084230601	09 18 28.5 + 51 39 31.3	6.92	3:03 0:54 0.31 + 0:09 0:10 + 0:09	-0.39 + 0:09 0:09 + 0:09	A G N 1	
J095218.9 013643	0065790101	09 52 19.0 01 36 43.1	11.78	24:50 2:98 -0.77 + 0:03 0:03 + 0:03	0.41 + 0:09 0:09 + 0:09	A G N 1	
J101850.5+ 411506	0028740301	10 18 50.6 + 41 15 06.6	10.32	2:16 0:49 -0.69 + 0:03 0:03 + 0:03	-0.51 + 0:07 0:07 + 0:07	A G N 1	
J101922.6+ 412049	0028740301	10 19 22.6 + 41 20 49.7	9.06	2:46 0:47 -0.44 + 0:04 0:04 + 0:04	-0.58 + 0:06 0:05 + 0:06	A G N 1	
J104026.9+ 204542	0059800101	10 40 26.9 + 20 45 43.0	9.32	8:70 1:15 -0.58 + 0:03 0:03 + 0:03	-0.48 + 0:05 0:05 + 0:05	A G N 1	
J104522.1 012843	0125300101	10 45 22.1 01 28 43.3	12.94	3:00 0:67 -0.63 + 0:04 0:04 + 0:04	-0.46 + 0:08 0:08 + 0:08	A G N 1	
J104912.8+ 330459	0055990201	10 49 12.8 + 33 04 59.8	10.46	2:06 0:51 -0.45 + 0:06 0:06 + 0:06	-0.54 + 0:09 0:10 + 0:09	A G N 1?	
J110050.6 344331	0112880201	11 00 50.6 34 43 32.0	12.71	5:12 0:69 -0.48 + 0:04 0:04 + 0:04	-0.33 + 0:07 0:07 + 0:07	A G N 1	
J112026.7+ 431520 ^a	0107860201	11 20 26.7 + 43 15 20.3	4.58	2:32 0:37 0.67 + 0:08 0:08 + 0:08	-0.32 + 0:09 0:09 + 0:09	A G N 2	

Name XBS...	Obs. ID	RA ; DEC (J2000)	O Axis arcmin	Rate 10^{-3} cts/s	HR 2	HR 3	Class
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
J113106.9+ 312518	0102040201	11 31 07.0 + 31 25 18.2	11.18	2.27 0.44 -0.53 + 0.08 + 0.07	-0.15 + 0.14 + 0.08	-0.15 + 0.14 + 0.08	AGN 1
J113121.8+ 310252	0102040201	11 31 21.8 + 31 02 52.6	11.58	3.88 0.69 -0.03 + 0.07 + 0.07	-0.35 + 0.08 + 0.08	-0.35 + 0.08 + 0.08	AGN 2
J113148.7+ 311358	0102040201	11 31 48.7 + 31 13 58.8	8.45	3.43 0.48 0.10 + 0.06 + 0.07	-0.43 + 0.07 + 0.07	-0.43 + 0.07 + 0.07	AGN 2
J122656.5+ 013126	0110990201	12 26 56.6 + 01 31 26.2	5.85	3.07 0.54 -0.11 + 0.09 + 0.09	-0.34 + 0.10 + 0.10	-0.34 + 0.10 + 0.10	AGN 2?
J123600.7 395217	0006220201	12 36 00.7 39 52 18.0	5.62	2.66 0.29 -0.77 + 0.01 + 0.01	-0.77 + 0.02 + 0.02	-0.77 + 0.02 + 0.02	star
J124641.8+ 022412	0051760101	12 46 41.8 + 02 24 12.1	2.78	2.22 0.49 -0.58 + 0.05 + 0.05	-0.48 + 0.09 + 0.09	-0.48 + 0.09 + 0.09	AGN 1
J132038.0+ 341124	0093640401	13 20 38.1 + 34 11 24.3	3.10	2.89 0.40 -0.43 + 0.04 + 0.04	-0.48 + 0.06 + 0.06	-0.48 + 0.06 + 0.06	AGN 1
J133942.6 315004	0035940301	13 39 42.6 31 50 04.8	11.91	3.52 0.51 -0.36 + 0.05 + 0.06	-0.20 + 0.08 + 0.08	-0.20 + 0.08 + 0.08	AGN 1?
J134656.7+ 580315 ^a	0112250201	13 46 56.7 + 58 03 15.4	11.08	3.33 0.56 0.46 + 0.10 + 0.10	-0.09 + 0.10 + 0.10	-0.09 + 0.10 + 0.10	GAL
J134749.9+ 582111	0112250201	13 47 49.9 + 58 21 11.0	8.53	7.39 0.66 -0.60 + 0.02 + 0.02	-0.54 + 0.03 + 0.03	-0.54 + 0.03 + 0.03	AGN 1
J140102.0 111224	0109910101	14 01 02.0 11 12 24.3	9.27	7.21 0.59 -0.52 + 0.02 + 0.02	-0.46 + 0.03 + 0.03	-0.46 + 0.03 + 0.03	AGN 2?
J140113.4+ 024016 ^a	0098010101	14 01 13.4 + 02 40 17.0	12.78	2.10 0.52 -0.55 + 0.09 + 0.09	0.08 + 0.16 + 0.16	0.08 + 0.16 + 0.16	M
J140127.7+ 025605	0098010101	14 01 27.7 + 02 56 05.4	7.25	6.66 0.62 -0.38 + 0.02 + 0.02	-0.53 + 0.04 + 0.04	-0.53 + 0.04 + 0.04	AGN 1
J141531.5+ 113156	0112250301	14 15 31.6 + 11 31 56.2	4.15	2.58 0.34 -0.55 + 0.03 + 0.03	-0.47 + 0.06 + 0.06	-0.47 + 0.06 + 0.06	AGN 1
J141830.5+ 251052 ^c	0109960101	14 18 30.5 + 25 10 52.6	7.58	7.70 1.20 -0.52 + 0.04 + 0.04	-0.65 + 0.07 + 0.07	-0.65 + 0.07 + 0.07	C
J142741.8+ 423335 ^a	0111850201	14 27 41.9 + 42 33 35.8	11.45	3.62 0.53 0.48 + 0.07 + 0.07	-0.23 + 0.07 + 0.07	-0.23 + 0.07 + 0.07	B
J143835.1+ 642928	0111530101	14 38 35.1 + 64 29 28.3	12.34	3.57 0.54 0.18 + 0.07 + 0.07	-0.37 + 0.06 + 0.06	-0.37 + 0.07 + 0.07	AGN 1?
J143911.2+ 640526 ^a	0111530101	14 39 11.2 + 64 05 26.8	12.04	2.44 0.39 0.86 + 0.09 + 0.09	0.29 + 0.11 + 0.11	0.29 + 0.11 + 0.11	B?
J153452.3+ 013104 ^e	0112190401	15 34 52.4 + 01 31 04.6	10.57	8.36 1.30 -0.39 + 0.04 + 0.04	-0.55 + 0.05 + 0.05	-0.55 + 0.05 + 0.05	AGN 1
J160645.9+ 081525	0067340601	16 06 46.0 + 08 15 25.1	12.61	7.36 1.37 0.81 + 0.08 + 0.08	-0.23 + 0.10 + 0.10	-0.23 + 0.10 + 0.10	AGN 2?
J161820.7+ 124116 ^a	0103461001	16 18 20.7 + 12 41 16.3	11.56	2.09 0.61 0.11 + 0.18 + 0.18	-0.18 + 0.18 + 0.19	-0.18 + 0.18 + 0.19	AGN 2?
J165425.3+ 142159	0113070101	16 54 25.4 + 14 21 59.3	7.44	5.26 1.08 -0.63 + 0.03 + 0.03	-0.61 + 0.07 + 0.07	-0.61 + 0.07 + 0.07	AGN 1
J165448.5+ 141311	0113070101	16 54 48.5 + 14 13 11.6	12.71	6.25 1.79 -0.54 + 0.07 + 0.07	-0.42 + 0.13 + 0.13	-0.42 + 0.13 + 0.13	AGN 2
J193248.8 723355 ^b	0081341001	19 32 48.8 72 33 55.2	8.39	4.47 0.73 -0.23 + 0.08 + 0.08	-0.08 + 0.10 + 0.10	-0.08 + 0.10 + 0.10	AGN 2?
J204043.4 004548 ^b	0111180201	20 40 43.5 00 45 48.2	10.55	3.24 0.80 0.04 + 0.11 + 0.11	-0.35 + 0.12 + 0.12	-0.35 + 0.12 + 0.12	AGN 2
J205635.7 044717 ^b	0112190601	20 56 35.8 04 47 17.9	9.96	2.08 0.50 -0.65 + 0.04 + 0.05	-0.48 + 0.11 + 0.11	-0.48 + 0.11 + 0.11	AGN 1
J205829.9 423634 ^b	0081340401	20 58 30.0 42 36 35.0	2.42	3.91 0.56 -0.57 + 0.03 + 0.03	-0.47 + 0.06 + 0.06	-0.47 + 0.06 + 0.06	AGN 1
J213002.3 153414 ^b	0103060101	21 30 02.3 15 34 14.1	12.83	2.30 0.47 -0.65 + 0.04 + 0.04	-0.50 + 0.09 + 0.09	-0.50 + 0.09 + 0.09	AGN 1
J213820.2 142536	0092850201	21 38 20.2 14 25 37.0	11.17	2.80 0.59 -0.29 + 0.07 + 0.07	-0.45 + 0.10 + 0.10	-0.45 + 0.10 + 0.10	AGN 1
J214041.4 234720 ^b	0008830101	21 40 41.5 23 47 20.1	9.80	3.30 0.68 -0.57 + 0.05 + 0.05	-0.39 + 0.10 + 0.10	-0.39 + 0.10 + 0.10	AGN 1
J220601.5 015346 ^b	0012440301	22 06 01.5 01 53 47.0	12.92	2.26 0.55 -0.41 + 0.06 + 0.06	-0.41 + 0.09 + 0.09	-0.41 + 0.09 + 0.09	AGN 1

Columns are as follows: (1) Source name; (2) XMM-Newton Observation number; (3) Right Ascension and Declination (J2000) of the source (X-ray position); (4) Angular distance (in arcmin) between the source and the MOS2 image center; (5) Source count rate, and 1 σ error, in the 4.5–7.5 keV energy band (units of 10⁻³ cts/s). In Table 1 we have reported the MOS2 conversion factors between the 4.5–7.5 keV count rate and the flux as a function of the energy spectral index, the hardness ratio HR2 and the blocking filter; (6) and (7) Hardness ratios computed as described in section 3.3. The errors on the hardness ratios have been evaluated using simulations and correspond to 1 σ ; (8) Optical spectroscopic classification (AGN1:broad line AGN; AGN2:narrow line AGN; GAL:Optically Normal Galaxy; CL: Cluster of Galaxies; BL: BL Lac Object; star: star; ?: Tentative classification; see section 3.2 for details).

NOTE { ^a These 11 sources belong to the HBSS sample but not to the BSS sample, while the remaining 56 HBSS sources are present also in the BSS sample.

^b A detailed X-ray and optical spectral analysis of these sources have been reported in Caccianiga et al. (2004).

^c These sources are more extended than the XMM-Newton EPIC MOS2 Point Spread Function at their off-axis angle. Count rates have been evaluated using aperture photometry.

^d A detailed X-ray and optical spectral analysis of these sources have been reported in Severgnini et al. (2003).

^e A detailed X-ray spectral analysis of these sources have been reported in Gialbauti et al. (2004).

Appendix A : Illumination Factor

The MOS2 cameras consist of a mosaic of 7 identical, front-illuminated CCDs with a dead space between the different chips. It is clear that serendipitous X-ray sources falling close to the gaps between the CCDs (or to their edges) could have either the flux and/or the source centroid poorly determined. The poor understanding of the corrections to be applied to these sources could represent a problem in the subsequent analysis and/or interpretation of the data. In order to take into account this problem in an objective way we have used the procedure detailed below.

From the exposure map produced from the pipeline processing system we have built a mask image representing the area on the sky which is "effectively" imaged by the CCDs. To produce this mask we have used the SAS task emask with threshold1= 0.25 and threshold2= 0.20 (see http://xmm.vilspa.esa.es/external/xmm_sw_cal/sas.html for specific details).

Using this mask image we have thus defined the "Illumination Factor" of each source as the fraction of sky "effectively" imaged by the CCDs in a circle of 20 arcsec around the source.

The "Illumination Factor" so defined ranges between 0.3 and 1 and the lower the "Illumination Factor" the closer is the source to gaps and/or edges in the CCDs. In the BSS and/or HBSS catalogues we have retained only sources with "Illumination Factor" > 0.8. Given the PSF of the MOS2 detector and its energy and off-axis dependence we have evaluated that less than 10% of the flux is lost in the case of a source with an "Illumination Factor" equal to the lower limit of 0.8. In table B.1 we report the complete list of sources which meet the selection criteria for the BSS and/or HBSS samples (e.g., inside the selected area between the inner and outer radius of each MOS2 image, count rate and likelihood limits, etc..) but that have been excluded from the sample because their "Illumination Factor" is below 0.8.

Finally the produced mask has been also used, in the computation of the sky coverage, to take into account the excluded area because of edges and gaps.

Appendix B : Cleaning procedure

XMM-Newton observations are subject to "holes" in the background rate, probably due to soft protons which are collimated by the X-ray mirrors toward the EPIC cameras and interact with the structure of the detectors and the detectors itself. The current understanding is that soft protons are probably organized in clouds populating the Earth's magnetosphere.

In order to check the background quality of the dataset used we have defined a "Background Estimator Parameter" which is roughly proportional to the "realbackground" accumulated in the MOS2 images.

To set this "Background Estimator Parameter" for each image we have produced an histogram of the total accumulated MOS2 counts in the 10-12 keV energy range as a function of time; the histogram bin size has been set to 100 seconds.

Thus, using this histogram we have:

- evaluated the mean count rate ($\langle bck \rangle$) and its standard deviation (σ_{bck})
- eliminated the time intervals which have a count rate greater than $\langle bck \rangle + 2\sigma_{bck}$
- repeated points a) and b) 10 times

The mean count rate at the end of the loop described above is the "Background Estimator Parameter".

Table A.1. Basic information on the sources excluded from the sample(s) since their "Illumination Factor" is less than 0.8

Obs. ID	RA; DEC (J2000)	O Axis arcmin	Illum.
(1)	(2)	(3)	(4)
0111000101	00 18 41.6 + 16 20 33.4	5.94	0.69
0001930101	00 26 24.9 + 10 31 23.7	10.77	0.68
0112320101	00 30 08.3 + 05 03 40.9	12.83	0.70
0065770101	00 32 41.0 + 39 40 08.2	5.70	0.42
0112600601	01 26 40.3 + 19 12 13.0	12.36	0.60
0112630201	01 34 03.1 40 02 22.8	12.38	0.68
0112371501	02 18 21.9 04 34 51.2	7.80	0.48
0111110501	02 22 49.4 05 14 53.0	5.58	0.27
0098810101	02 36 30.3 52 27 04.3	7.66	0.77
0098810101	02 37 02.1 52 23 48.0	8.53	0.62
0122520201	03 13 14.6 76 55 55.4	5.88	0.71
0110970401	03 13 34.3 55 26 43.6	7.85	0.76
0111970301	04 09 02.1 71 07 54.6	9.83	0.63
0085640101	05 21 00.2 25 28 52.5	7.00	0.49
0050150101	05 25 57.1 33 44 35.7	10.72	0.74
0110930101	06 17 56.8 + 78 16 05.3	5.81	0.66
0103860101	06 24 46.8 64 33 46.0	10.80	0.78
0112980201	06 57 26.2 55 49 53.9	10.49	0.78
0025540301	08 38 27.2 + 25 50 53.7	5.75	0.65
0025540301	08 38 52.5 + 25 37 25.3	10.27	0.58
0110660201	09 07 51.1 + 62 01 58.3	12.77	0.65
0084230601	09 16 45.2 + 51 41 45.0	10.53	0.57
0110930201	10 00 56.7 + 55 41 01.2	8.50	0.69
0101040301	10 23 13.3 + 19 56 50.5	6.43	0.62
0055990201	10 50 29.0 + 33 00 42.8	7.63	0.70
0110660401	11 26 11.9 + 42 52 45.9	2.09	0.78 ^a
0124110101	12 21 34.3 + 75 09 18.1	9.34	0.60
0124110101	12 22 07.3 + 75 26 20.7	7.88	0.53
0092360601	12 52 14.1 83 46 42.6	9.48	0.66
0002940101	13 06 32.8 23 31 13.1	12.05	0.62
0111160101	13 34 16.9 + 50 23 09.6	7.78	0.64
0111160101	13 35 09.6 + 50 39 17.8	11.58	0.73
0111570201	13 35 18.7 34 21 45.5	8.30	0.62
0112250301	14 15 39.9 + 11 24 04.4	5.69	0.37
0109960101	14 18 42.7 + 25 07 09.5	9.90	0.79
0070740301	15 03 39.6 + 10 16 04.9	11.46	0.66
0018741001	15 18 59.7 + 06 18 39.4	5.57	0.73
0061940301	16 32 39.0 + 78 11 54.3	5.48	0.56
0102040101	17 22 54.3 + 34 27 25.3	10.89	0.58
0061940201	21 37 47.7 42 26 14.5	10.55	0.68
0008830101	21 39 57.9 23 45 35.1	7.27	0.76
0111790101	22 36 06.1 26 08 04.6	6.80	0.55
0109070401	22 47 48.1 51 10 18.1	8.38	0.73
0025541001	23 04 43.5 + 12 12 10.2	8.04	0.53
0123900101	23 14 32.2 42 33 01.5	12.11	0.72

Columns are as follows: (1) XMM-Newton Observation number; (2) Right Ascension and Declination (J2000) of the source (X-ray position); (3) Angular distance (in arcmin) between the source and the MOS2 image center; (4) "Illumination factor". NOTE { ^a Although this source is well contained inside the MOS2 central CCD it fall above a bad column; for this reason its "Illumination factor" is below 0.8.