# IA C -star: a code for synthetic color-m agnitude diagram com putation

Antonio Aparicio

Departamento de Astrof sica, Universidad de La Laguna/Instituto de Astrof sica de Canarias.V a Lactea s/n.E38200 - La Laguna, Tenerife, Canary Islands, Spain

Carm e Gallart

Instituto de Astrof sica de Canarias. V a Lactea s/n. E 38200 – La Laguna, Tenerife, Canary Islands, Spain

#### ABSTRACT

The code IAC-star is presented. It generates synthetic HR and colorm agnitude diagrams (CMDs) and is mainly aim ed to star form ation history studies in nearby galaxies. Com posite stellar populations are calculated on a star by star basis, by computing the lum inosity, e ective tem perature and gravity of each star by direct bi-logarithm ic interpolation in the metallicity and age grid of a library of stellar evolution tracks. V isual (broad band and HST) and infrared magnitudes are also provided for each star after applying bolom etric corrections. The Padua (Bertelli et al. 1994, Girardi et al. 2000) and Teram o (Pietrinferni et al. 2004) stellar evolution libraries and various bolom etric corrections libraries are used in the current version. A variety of star form ation rate functions, initial mass functions and chem ical enrichm ent laws are allowed and binary stars can be computed. A lthough the main motivation of the code is the computation of synthetic CMDs, it also provides integrated m asses, lum inosities and m agnitudes as well as surface brightness uctuation lum inosities and magnitudes for the total synthetic stellar population, and therefore it can also be used for population synthesis research.

The code is o ered for free use and can be executed at the site http://iac-star.iac.es, with the only requirement of referencing this paper and crediting as indicated in the site.

Subject headings: HR-diagram , color-m agnitude digram , population synthesis, star form ation history

## 1. Introduction

G alaxies evolve on two main paths: dynam ically, including interactions and merging with external systems and through the process of form ation, evolution and death of stars within them. The latter has the following relevant e ects on the galaxy: (i) the evolution of gas content, (ii) the chemical enrichment and (iii) the form ation of the stellar populations with dierent properties as the gas from which they form evolves. The star form ation history is therefore fundamental to understand the galaxy evolution process.

The color-m agnitude diagram (CM D) is the best tool to study and derive the star form ation history (SFH) of a galaxy. Deep CM Ds display stars born all over the life-time of the galaxy and are indeed fossil records of the SFH. A qualitative sketch of the stellar populations present in a galaxy can be done from a quick look to a good CM D. The presence of stars in characteristic evolutionary phases indicates that star form ation took place in the system in di erent epochs of its history. For exam ple, the presence of RR-Lyre stars is indicative of an old, low m etallicity stellar population; a substantial am ount of red-giant branch (RGB) stars is associated to an interm ediate-age to old star form ation activity; a well developed red-tail of asym ptotic giant branch (AGB) stars show s that interm ediate-age and young stars with relatively high m etallicity are present in the system, and even a few blue, bright stars as well as H II regions are evidences of a very recent star form ation activity.

A higher degree of sophistication is provided by isochrone thing to signi cant features of the CMD. Indeed, this method is simple and powerful enough to determ ine age and metallicity of single stellar populations as the ones present in star clusters.

However, actually deciphering the inform ation contained in a complex CMD and deriving a quantitative, accurate SFH is complicated and requires more sophisticated techniques. The standard procedure involves three main ingredients: (i) good data, from which a deep, ideally reaching the oldest main-sequence turn-o s, observational CMD can be plotted; (ii) a stellar evolution library complemented by a bolom etric correction library providing colors and magnitudes of stars as a function of age, mass and m etallicity, and (iii) a method to relate the number of stars populating di erent regions of the observational CMD with the density distribution of stars in the CMD as a function of age, mass and metallicity as predicted by the stellar evolution theory. The result of this comparison provides the SFH of the system.

U sually, item (iii) involves computation of one or several synthetic CMDs from which the star density distribution predicted from theory is deduced. Indeed, the synthetic CMD technique is the most powerful one available for this kind of analysis and, in general, for the study of the SFH of resolved stellar populations. The rst to compute and use a synthetic CMD using a monte-carlo technique was M aeder (1974). Chiosi, Bertelli, & Bressan (1988) presented the rst application of the Padua synthetic CMD code. Tosi et al. (1991) used, for the rst time, lum inosity functions derived from synthetic CMDs to sketch recent SFHs of nearby galaxies. Bertelli et al. (1992) were the rst to use the global detailed morphology together with number counts and ratios of star counts in dierent areas of the synthetic and observed CMDs to derive the SFH of a galaxy (the LMC).

A more recent version of the Padua code is ZVAR, which distinctive feature is the age and metallicity interpolation that is performed for each single star in the Padua grid of stellar evolution tracks. This has been a heavily used code in the recent past (e.g., Gallart et al. 1996; Vallenari et al. 1996a,b; Aparicio, Gallart, & Bertelli 1997a,b; Hurley-Keller, Mateo, & Nemec 1998; Gallart et al. 1999; Aparicio, Tikhonov, & Karachentsev 2000; Vallenari, Bertelli & Schmidtobreick 2000; Bertelli & Nasi 2001; Ng et al. 2002; Bertelli et al. 2003; see Aparicio et al. 1996 for several exam ples of synthetic CMDs using ZVAR and Aparicio 2003, for a review). O ther codes have been developed by di erent groups for the com putation of synthetic CMDs (see Tolstoy & Saha 1996; Dolphin 1997; Holtzm an et al. 1999; A rdeberg et al. 1997; Stappers et al. 1997; Dohm-Palm er et al. 1997; Hernandez, Valls-Gabaud, & Gilm ore 1999; Harris & Zaritsky 2001).

In this paper we present a new algorithm for the computation of synthetic CMDs. It is intended to be as general as possible. For this purpose, a variety of inputs for the initial mass function (IMF), SFH, metallicity law and binariety are allowed. Stars with age and metallicity following a continuous distribution are computed through interpolation in a stellar evolution library, providing synthetic CMDs with smooth, realistic stellar distributions. This signi cantly improves the accuracy of the comparison with observed CMDs as compared to with methods without such interpolation. The Padua stellar evolution libraries by Bertelli et al. (1994; Bertelli94 from here on) and G irardiet al. (2000; G irardi00; from here on) and Teram o (Pietrinfemi et al. 2004) stellar evolution library are used in the current version of the program, completed by the Cassisi et al. (2000) models for very low mass stars. Together with the lum inosity and temperature of each star, the code provides the absolute magnitudes in Johnson-Cousins UBVRI, HSTF218W, F336W, F 439W m F 450W, F 555W and F 814 and infrared JHKLL<sup>M</sup> Iters. To this purpose the bolom etric correction libraries by Lejeune, Cuisinier, & Buser (1997), Castelli & Kurukz (2002), G irardiet al. (2002) and O riglia & Leitherer (2000) are used, the later for the HST

Iter system. In summary, in the current version of the code, the user can select among three stellar evolution libraries and four bolom etric correction libraries. The code includes the therm ally pulsating AGB (TP-AGB) following the prescription of M arigo, B ressan, & Chiosi (1996).

The code is made available for free use. It can be executed at the internet site http://iac-star.iac.es with the only requirement of crediting as mentioned in that page. The present paper should be considered as the reference for the code. This paper will be complemented by several others presenting, in particular, a method based on a genetic algorithm to compute the SFH from comparison of an observed and a generic synthetic CMD (Aparicio & Hidalgo 2004) and a library of population synthesis based in surface brightness uctuations (Mar n-Franch & Aparicio 2004).

This paper is organized as follows. In x2, the fundam entals of synthetic CMD computation are given, including short descriptions of the stellar evolution and bolom etric correction libraries used. In x3, x4 and x5, the basic stellar evolution, the main features arising in a real CMD and the main aspects of them ical enrichment laws are summarized. These topics are relevant for a better understanding of the computation process and the synthetic results. Section 6 is devoted to explain how the code works as well as some particularities including the computation of mass loss in advanced evolutionary phases. Section 7 is a short outline of the code input and output. A full commented list of input choices and a description of the output le content are available in the internet site from which the program is run (http://iac-star.iac.es). This will be permanently updated, including any future change in the input and output. Finally, a summary of the paper is given in x8.

#### 2. Fundam entals of synthetic C M D

The distribution of stars in a CM D depends on several aspects on the physics of stars and galaxies. Broadly speaking, the density of stars in a given region of the CM D is directly proportional to the tim e that stars of any m ass and m etallicity spend in that region during their evolution. This inform ation is provided prim arily by stellar evolution theory, on which the computation of synthetic CM D s strongly relies, com plem ented by bolom etric corrections. But, since the fraction of stars form ed with di erent m asses vary with m ass, the distribution of the stellar m asses at birth, the so called IM F, provides a second, im portant input characterizing the star density distribution on the CM D. Stars, on the other hand, are not form ed at the same rate during the life-tim e of a galaxy. The function accounting for the star form ation rate (SFR) as a function of tim e is a further im portant input. Besides this, stars are born with the m etallicity of the interstellar m edium from which they form . So, the chem ical enrichm ent law (CEL) of the system holding them is another fundam ental inform ation. The last im portant contribution to the m orphology of the CM D com es from the fraction and m ass ratio distribution of binary stars. O ther physical properties do a ect that m orphology, like the internal, di erential reddening or the m ass distribution (stellar population gradients, depth e ects, etc) within the system . However, these are properties that can be considered as external to the CMD itself. In m ost (or all) cases in which we are interested, they can be added later on to the com puted synthetic CMD.

In sum m ary, the inputs necessary to compute a synthetic CMD are the following:

A stellar evolution library, covering a wide range of stellar evolutionary phases and stellar m asses and m etallicities. It should provide stellar lum inosities, tem peratures and surface gravities as a function of m ass, m etallicity and age.

A bolom etric correction library to transform from stellar lum inosities, tem peratures and surface gravities to magnitudes in a given photom etric system.

An MF, (m).

A SFR function, (t).

A CEL,Z(t).

A function accounting for the fraction, and secondary to primary mass ratio distribution, of binary stars, (f;q).

In the following we will shortly comment on the stellar evolution and bolom etric correction libraries used here and on the commutation algorithm.

2.1. The stellar evolution libraries

Computation of a synthetic CMD relies on a stellar evolutionary library, which should be computed using the most up-to-date physics, well tested against observations, and complete in terms of ages, metallicities and evolutionary phases. These three characteristics are hard to be accomplished and, indeed, many of the accuracy and precision limitations of synthetic CMD based techniques originate in this. In its current version, IAC -star uses the Bertelli94, G irardi00 and Teram o (Pietrinfemi et al. 2004) stellar evolution libraries, completed by the Cassisi et al. (2000) models for very low mass stars. Even though the Bertelli94 library has been basically superseded by the G irardi00 one, we have chosen to include both libraries since many SFH studies in the past have been perform ed using the Bertelli94 one which may be chosen for comparative studies. It is our intention to include additional libraries in the future. In Table 1 the characteristics and main input physics of the three libraries are listed. Figure 1 shows several tracks from the Bertelli94 library as well as two very low mass starm odels computed from Cassisi et al. (2000).

## 2.2. The bolom etric correction libraries

C om parison of synthetic and observed CMDs require transforming the rst from the theoretical plane (the  $T_e$  L plane) to the observational one (the color-m agnitude plane). This is done by adopting a bolom etric correction scale and a color - e ective tem perature relation. It is important to note that, while the stellar evolution library is fundam ental for the computation of the synthetic CMD itself, an equally reliable bolom etric correction library is required if synthetic CMDs are to be compared with observed ones. This also implies that the derivation of accurate SFHs relies on the accuracy of both stellar evolution library and the model atm ospheres set adopted for computing the bolom etric correction scale and color  $-T_e$  relation.

In its current version, IAC-star in plements four choices for the bolometric correction and  $color - T_e$  relation libraries:

The G irardi et al. (2002) library providing bolom etric corrections for visual, broad band UBVRI and infrared JHK lters.

The Castelli & Kurucz (2002) library for visual, broad band UBVRI and infrared JHKL liters, completed with the Flucks et al. (1994) sem i-empirical transform ations for M giants. Care has been taken that the two sets properly m atch.

The Lejeune, Cuisinier, & Buser (1997) library for visual, broad band UBVRI liters and infrared JHK LL<sup>M</sup> liters.

The Origlia & Leitherer (2000) library for F 218W , F 336W , F 439W , F 450W , F 555W and F 814W liters of the HST W FPC2 and ACS.

These libraries full the requirements of completeness and wide metallicity ( $F \in H$ ), temperature ( $T_e$ ) and gravity (g) coverage and sampling, as well as having been empirically calibrated. It is our intention to include other libraries, as well as transformations for additional photometric bands such as the Stromgren ones, in the future as they become available.

The bolom etric correction and the color  $-T_e$  are functions of the following three parameters: BC (F e=H ];  $T_e$ ; g). The used libraries allow obtaining bolom etric corrections

for alm ost all cases, although they fail to fully cover a few, very low temperature cases. Linear extrapolation is used in such cases. Since it is necessary in very few cases, the deviation e ects that could arise in the CMD from that extrapolation should be negligible. However, detailed analysis of color distribution of stars in phases as the extended AGB should be made with caution.

#### 2.3. Computing a synthetic CM D

IAC -star uses a monte-carlo method to compute synthetic CMDs on a star by star basis. A random number generator (RNG) is used to compute, rst the mass, and then the time of birth of the star, according to the distribution functions (m) and (t), respectively. The value of time is introduced in the function Z (t) to obtain the metallicity. If metallicity dispersion is allowed, its exact value is obtained using such a dispersion and a new random number. The binariety of the star is determined in a similar way, using a RNG and the function (f;q), in which f is the fraction of binaries and q the internal mass rate. If the star turns out to be a binary, it is assumed to be the primary (most massive) star of the system. The age and metallicity of the secondary are assumed to be the same as for the primary. The mass of the secondary is computed using again the RNG and according to the secondaries. The lifetime of each star, according to the stellar evolution models, is used at this point to decide whether the star is alive (and to calculate then its remaining parameters), or dead (and to calculate the properties of the remnant). The case in which only the secondary star is alive is allowed.

Once the mass, age and metallicity are known, lum inosity, temperature and surface gravity of the star are computed through interpolation within the chosen stellar evolutionary library. If the star is a binary, the process is repeated for each component. Finally, lum inosity, temperature and gravity are interpolated in the chosen bolom etric correction library to obtain magnitudes and colors in the standard photom etric system.

These are the computation fundam entals. In x7, we provide details about the procedure itself. But before, it is clarifying to outline the aspects of the stellar evolution relevant to the morphology of the CMD (x3); to review such morphology (x4), and to discuss the CEL, its param eterization and its relation to basic physical properties of the system (x5).

#### 3. Basic stellar evolution

Lum inosity, L, and temperature  $T_{\rm e}$  , as a function of stellar age and metallicity, as well as lifetime, are, for our purposes, the main information provided by the stellar evolutionary library. If s is a star of m ass m<sub>s</sub>, m etallicity  $Z_s$  and age  $a_s$ , a rst estimate of its location in the CMD might be obtained just by time interpolation within a single evolutionary track of age and m etallicity closest to the stellar values. However, if a realistic synthetic CMD is to be computed, interpolation must be done between tracks both in age and m etallicity. The fact that stars experience di erent particular physical processes along their evolution that, like the He-ash, abruptly interrupt their sm ooth evolutionary paths, makes this interpolation rather cumbersome. In the following, we will sketch the basic stellar evolution directly a ecting the synthetic CMD computation problem. In this sense, it is useful to divide stars in three groups according to their masses: low, intermediate and high mass stars (see Chiosi et al. 1992 and Chiosi 1998 for a short but clear sum mary of the basic stellar evolution). The critical masses separating these groups depend on the chem ical composition and the input physics. But, roughly, the mass separating the low and intermediate dominions, m  $_{\rm lw}$  , is in the range 1:7 2:0 M and the mass separating the interm ediate and high dom inions,  $m_{up}$ , is in the range 5 7 M . In Fig. 2 the loci of the di erent features that will be introduced in the following are displayed.

Stars of any mass spend a major fraction of their lives in the core H-burning, main sequence (MS) phase. As H is exhausted in the core, the star leaves the MS while the H-burning proceeds in a shell. At this point, a starm ay follow two types of paths in the CMD, depending on its mass. In low mass stars, the H-exhausted, He core is not massive, hot and dense enough to immediately start the Hebuming. The core gravitationally contracts and, as a consequence, degeneracy appears in the electronic gas. This degeneracy stops the core collapse. As H-burning proceeds in a shell, the star climbs the RGB and m ore He is incorporated into the degenerated core. W hen the mass of this core reaches about 0.45 to 0.50 M (the precise mass depends on chem ical composition, stellar mass and input physics), He-burning starts violently, rem oving the electron degeneracy. This is the so called He-ash. The star abruptly term inates the RGB phase and start the horizontal-branch, in which He-burning and H-burning proceed in the core and a shell, respectively. Since the He-ash is produced at essentially identical core mass, the maximum RGB lum inosity is almost constant, independently of the stellar metallicity and initial mass. The same happens to the lum inosity of the horizontal branch (HB). However, the stellar surface e ective tem perature, which determ ines the position of the star in the HB, strongly depends on several factors, am ong which the most important are the chemical com position and total stellar mass. It must be noted that the latter depends not only on the initial stellar mass, but also on the mass loss during the RGB and the He-ash.

In interm ediate and high m ass stars, the H e-core ignites in non-degenerate conditions as soon as the central tem perature and density reach some threshold values, approximately  $10^8$  K and  $10^4$  g cm<sup>-3</sup>, respectively. This requires a minimum core m ass of 0.33 M . Since this m ass depends on the initial stellar m ass, the lum inosity of the star during this phase increases with stellar m ass. In interm ediate m ass stars (including the lower end of high m ass stars), the core H e-burning phase takes place in two regions of the CM D. The rst one is close to the H ayashi line. Subsequently, when the energy produced in the H e core, which is increasing, equals the energy produced in the H shell, which is decreasing, the outer envelope rapidly contracts passing from convective to radiative. Tem perature increases and the star m ignates to a bluer region. This produces the blue-loop (BL) phase. It appears rather red in F ig. 2 as corresponds to the high m etallicity of the com puted young population.

Both in low and intermediate mass stars, a C-O core is formed as a result of core Heburning. Upon the exhaustion of He in the core, the mass of the C-O core is constant for low mass stars, but increases with stellar mass for all the others. The structure of stars is hence form ed by a C-O core, a He-burning shell, a H-burning shell and a H-rich envelope. In low and interm ediate m ass stars, the m ass of the C  $\rightarrow$  core is not high enough to ignite and an electron degenerated core again develops. The star expands, produces a convective envelope and evolves upward running alm ost parallel to the RGB. This is the asymptotic giant branch (AGB) phase. Cooling in the regions external to the C-O core extinguishes the H-burning shell but, eventually, the expansion of the envelope is stopped by its own cooling, the envelope contracts and the H-shell re-ignites. This term inates the early-AGB (E-AGB). A fter that, the He-shell becomes therm ally unstable. Both, the H and He shells alternate as the main sources of energy producing the TP-AGB phase. This process results in e cient m ass loss and term inates with the complete ejection of the envelope and tem porally producing a planetary nebula. The rem nant is a C-O white dwarf of m as lower than the Chandrasekar limit (1.4 M). Intermediate (and low) mass stars fail to ignite the C-O core.

In high m ass stars, the C-O core ignites in non-degenerate conditions. The fraction of high m ass stars form ed in a galaxy is small and their life-times short. For these reasons they do not a ect signi cantly the structure of the CMD and we do not go further in the discussion of their evolution. It is enough to mention that m ass loss plays a fundamental role in determining the position in the CMD of core He-burning stars more massive than about 15 M , and that, after core C-O ignition, they proceed through a series of nuclear burnings and nish theirs lives with a supernova explosion.

#### 4. Main features of the C M D

The main features and the distribution of stars produced by stellar evolution in a CM D can be seen in Fig. 2, where a synthetic CM D, computed for constant SFR from 13 Gyr ago to date, is shown. The CM D is based in the Bertelli94 stellar evolution library and the Lejeune et al. (1997) bolom etric correction library. The IM F obtained by K roupa, Tout & G ilm ore (1993) and a simple CEL with m etallicity Z (t) increasing linearly with time, initial value  $Z_0 = 0.0001$  and nal value  $Z_f = 0.02$  have been used. Colors are associated to age intervals to illustrate the distribution of stars in the CM D according to their age. Labels are included to identify the M S, BL, RGB, AGB, HB and red-clum p (RC) phases. The RC is form ed by a m ix of the reddest HB stars and the bottom part of the core Herburning phase of interm ediate m ass stars. The remaining features have been introduced in x3.

The M S provides the most important and unambiguous age information. But it is interesting to note that, besides it, the BL and the AGB, if well populated, provide useful and complem entary age information, which is poor, however, on the RGB.But the latter gives information on the integrated star formation rate for intermediate and old stellar populations and provides clues to constrain the CEL.

The morphology of the CMD is altered by the presence of binary stars. Figure 3 illustrates the e ect, which is most obvious in the lowest MS but a ects also the overall distribution of stars in the CMD.

To simplify the discussion of stellar populations based on CM D analysis, it is useful to introduce some term inology. We will use the term sold, intermediate-age and young to specify three stellar age ranges which are associated to features clearly recognizable in the CM D. By old we will refer to populations that are not able to produce bright, extended AGBs. This corresponds to ages larger than about 10 G yr. Good examples of old objects are globular clusters. Old populations will always produce a well populated RGB and additionally, they may show a blue extended HB. However, the latter is not a unambiguous age indicator, since it depends on m etallicity and mass loss on the RGB. Interm ediate-age populations are those producing an RGB as well as extended, bright AGBs. They are older than 1 2 G yr. Finally, young populations are those igniting He in the nucleus in non-degenerate conditions, and therefore failing to produce an RGB. The age separating the young from the interm ediate-age ranges is the MS life time of stars of mass equal to m<sub>lw</sub>, the mass separating the low and interm ediate-mass dom inions. A pure young population would not have had time for low-mass stars to evolve from the MS.

## 5. The chem ical enrichm ent law of a stellar system

In its current version, IAC-star o ers two main procedures to compute the CEL, Z (t): (i) by linear interpolation in several, arbitrary Z (t) nodes given by the user, and (ii) from the parameters (yield, infall and out ow rates, etc.) used in physical CEL scenarios. For a better understanding of the second procedure, the CEL fundamentals are outlined in this section. However, it must be noted and kept in m ind that IAC-star is not a code to solve for chemical enrichment law scenarios. It allows a wide range of choices for them and computes stars with the necessary coupling between age, metallicity, SFR and gas fraction provided by the input parameters. But it is the user responsibility to decide what kind of chemical law is to be used. On the other hand, the linear interpolation in Z (t) nodes, which includes a linear law and a constant one, is a simplistic, perhaps physically unrealistic choice, but it may work well form any users, in particular when details about the actual metallicity law are not known. It must be also noted that a simple, linear CEL is not far from the law of a moderate infallm odel.

M etallicity laws for a closed box and for infall, out ow of well mixed material and out ow of rich material can be found in Peimbert, Coln, & Sam iento (1994). In IAC-star the infall and out ow (of well mixed material) scenarios can be used as well as the close box one, which is but a limit case of any of the form er. The unmixed, rich material out ow can be simulated by the well mixed material out ow if the yield is allowed to vary and we are not interested on the Helium to Oxygen ratio, which, in any case, is beyond the scope of this software.

A ssum ing instantaneous recycling, infall and out ow laws are given by (Peimbert et al. 1994)

$$Z(t) = Z_0 + \frac{Y}{2} f1 [ (1)(t)^{-1} ]^{-(1)} g$$
 (Infall) (1)

$$Z(t) = Z_0 + \frac{Y}{+1} \ln [(+1)(t)^{-1}] \qquad (Out ow) \qquad (2)$$

The chem ical enrichm ent theory has been developed elsewhere (see T insley 1980; Peim bert et al. 1994 and references therein) and it is not our aim to discuss it here in any detail. However, the form er equations deserve som e comments in order to understand the CEL computation in IAC-star. The initial metallicity,  $Z_0$  should be 0 for physically realistic models, but it will be useful to have the possibility of assuming some initial enrichment in some cases. The yield, y is de ned as the mass of new ly form ed metals that a generation of stars ejects to the interstellar medium, relative to the mass locked in stars and stellar remnants by the same stellar generation. is the gas mass fraction relative to the total m ass intervening in the chem ical evolution; i.e. including stars, stellar rem nants, low m ass objects, dust and gas itself but excluding dark m atter, which does not intervene in such process.

Parameters and control infall and out ow respectively. They are de ned as  $f_I = (1 \ R)$  and  $f_0 = (1 \ R)$ , where  $f_I$  and  $f_0$  are the infall and out ow rates, respectively and R is the mass fraction returned to the interstellar medium by a generation of stars. In this way,  $(1 \ R)$  is the mass locked into stars and stellar remnants and  $f_I$  and  $f_0$  are given as fractions ( and ) of this mass. A few things must be pointed out. First, since is a function of time, also  $f_I$  and  $f_0$  are so. Second, both and can take values > 1. Finally, the simple closed box model is obtained from any of the former relations if, respectively,  $= 0 \ or = 0$ .

The mass fraction, , appearing in equations 1 and 2 is coupled to the SFR or, rather, to its integral, by

$$(t) = 1 \qquad \frac{(1 \quad R) \quad (t)}{M_0 + (1 \quad R) \quad (t) \quad ()}$$
(3)

where  $M_0$  is the initial mass of the system and  $(t) = {R_t \atop 0} (t^0) dt^0$ . This relation for is valid for equation 1 and 2 just using = 0 in the rst case and = 0 in the second one. Lets assume for simplicity, and only in the context of the chemical law computation, that the time unit is the present age of the system, with t = 0 for the initial instant and t = 1 for the present day time, and that the integral of the SFR is normalized also for the system age  ${R_1 \choose 0} (t^0) dt^0 = 1$ . With this, the initial mass of the system can be written as

$$M_0 = (1 R) ( \frac{1}{f 1} )$$
 (4)

where  $_{f}$  is the nalgas fraction. The yield, which is assumed here to be constant, can also be expressed as a function of  $Z_{0}$  and  $Z_{f}$  (the nalm etallicity):

$$y = \frac{(Z_{f} \quad Z_{0})}{1 \quad [ \quad ( \quad 1)_{f}^{1} ]^{-(1)}}$$
 (Infall) (5)

$$y = \frac{(Z_{f} Z_{0})(+1)}{\ln[(+1)_{f}]^{1}} \qquad (Out ow)$$
(6)

In practice, the most frequent situation is one in which the researcher has observational information about  $_{\rm f}$  and  $Z_{\rm f}$  and can assume that  $_{0} = 1$  and  $Z_{0} = 0$  or small. R can be derived from the stellar evolution models (it is R ' 0.2 for the Padua stellar evolution

m odels, although it depends slightly on the m etallicity) and (t) is the key function of the stellar population m odel (usually what is sought in the analyzed galaxy). The remaining parameters are and , for which reasonable guesses need to be m ade. Of course, in this way y is no longer the physical stellar yield, but an elective yield de ned just by equations 5 and 6. If the physical relevance of the chem ical evolution m odel is seeked, the consistency of all the parameters must be analyzed. However, for the purposes of computing a synthetic CMD, the chem ical law is only an ingredient of the code, which can be imposed from outside. Users of IAC -star are encouraged to check and decide the required chem ical law. We insist that IAC -star only provides the coupling between (t), (t) and Z (t) for the set of input  $Z_0$ ,  $Z_f$ , f and or .

In Fig. 4 several examples of CELs for di erent choices of  $_{\rm f}$ , and (t) are given for xed Z  $_0$  and Z  $_{\rm f}$ . For (t), an exponential law of the form (t) = A exp(t=) is used. In Fig. 5 a law with m etallicity dispersion is shown.

#### 6. IA C -star com putation outline

In this section we will outline how the IAC -star algorithm works paying attention to the general interpolation procedure and to how di erent stellar mass ranges and evolutionary phases are handled. Computation of mass loss in advanced evolutionary phases is explained in the last paragraph. In Fig. 1 several tracks of di erent masses from the Bertelli94 library are plotted as reference for the following discussion.

## 6.1. The general case

As discussed in x2, a monte-carlo algorithm and the corresponding probability distribution functions are used to provide the age,  $a_s$ , m ass, m  $_s$ , and m etallicity,  $Z_s$ , of each synthetic star. In the general, simplest case, m ass and m etallicity are used to compute the evolutionary track of the star by bi-logarithm ic interpolation within the evolutionary track library. It is important to make this simple but important and cum bersom e process clear. Note that one of the di culties is that tracks are not functions, but arbitrary curves, time being the driving parameter. For this, interpolation must be done between hom ologous sections of di erent tracks.

Let rst introduce some notation. Tracks are identied by mass and metallicity. We will name tracks by K<sub>ijj</sub>, the rst index denoting the metallicity and the second one, the mass.  $Z_s$  represents a value of metallicity between two values present in the stellar evolution

library. The algorithm input does not allow extrapolation outside the library limits in Z. Lets call  $Z_{i}^{lb}$  and  $Z_{i+1}^{lb}$  these two values, respectively. So  $Z_{i}^{lb}$   $Z_{s}$   $Z_{i+1}^{lb}$ . A search is then made within  $Z_{i}^{lb}$  and  $Z_{i+1}^{lb}$  track subsets to look for the mass values embracing  $m_{s}$ . Lets call  $m_{ij}^{lb}$  and  $m_{ij+1}^{lb}$  track mass values embracing  $m_{s}$  for the  $Z_{i}^{lb}$  subset and  $m_{i+1,k}^{lb}$  and  $m_{i+1,k}^{lb}$  and  $m_{i+1,k+1}^{lb}$  the track mass values embracing  $m_{s}$  for the  $Z_{i}^{lb}$  subset and  $m_{i+1,k+1}^{lb}$  and  $m_{i+1,k+1}^{lb}$  subset. So  $m_{ij}^{lb}$   $m_{s}$   $m_{ij+1}^{lb}$  and  $m_{i+1,k}^{lb}$   $m_{s}$   $m_{i+1,k+1}^{lb}$ . The track corresponding to stars can now be obtained by logarithm ic interpolation between tracks K  $_{i;j}$ , K  $_{i;j+1}$ , K  $_{i+1,k+1}$ . Lets call this interpolated track K  $_{s}$ . Additionally, lets call  $_{ij}$  the life time of track K  $_{i;j}$ , i.e., the time at which evolution within that track mish, and  $_{s}$  the corresponding life time for the interpolated track.

Once K<sub>s</sub> has been obtained, the photom etric parameters of the star are obtained straightforwardly by logarithm ic interpolation of  $a_s$  within this track. However, before this, whether the star is death or alive must be established. This is done just comparing  $a_s$  with s. If  $a_s$  s, logarithm ic interpolation is performed and the computation is nished for this star. If not, a new dichotom y arises. If the star is of intermediate or high mass,  $a_s > s$  m eans that it is death. The mass of the remnant is then computed as explained below and added to the total mass of the system. If the star is of low mass,  $a_s > s$  means that it is beyond the He- ash. Computation in the HB track subset must be performed in this case. For this, the mass loss of the star during the RGB is subtracted from  $m_s$  (see below) and s is subtracted from  $a_s$ . The resulting mass and age,  $m_{s,HB}$  and  $a_{s,HB}$ , are used in the HB track subset. Computation is similar to that in the general library. If  $a_{s,HB}$  is larger than the life time of the new interpolated HB track, the star is considered death and the remnant mass is computed and stored as before.

We have outlined the general computation. However, a few special cases can occur that need more comments. We will treat them in the following.

#### 6.2. Interpolation and extrapolation between di erent m ass ranges

It may occur that the tracks embracing the mass and metallicity of the synthetic star belong to two dierent mass ranges (see Fig. 1). If some tracks are in the high mass range and some in the intermediate mass range, interpolation can be performed as in the general case up to the end of the tracks. A fler that, high mass stars are considered death while intermediate mass stars continue through the TP-AGB, computed as outlined below.

The case is som ewhat more complicated if som e tracks are in the interm ediate-mass range and som e in the low-mass range. The stellar physical processes are qualitatively similar for both ranges up to the basis of the RGB: core H-burning (MS) or He-core

gravitationally contracting plus shell H-burning (sub-giant phase). Interpolation can hence be performed up to that point but not afterward, when intermediate mass stars are about to start the core He-burning while low mass stars evolve climbing the RGB. In this case, the mass separating the intermediate and low mass ranges is rstly determined by logarithmic interpolation between the four tracks embracing the mass and metallicity of the synthetic star. This separating mass is used to determ ine the range to which the star belongs. A fter this, interpolation is performed up to the RGB basis. Beyond that point, extrapolation is made from tracks of the mass range to which the star belongs. The CMD related features can be seen in Fig. 1

## 6.3. Very high and very low mass stars

The stellar mass interval is controled by the input IM F. Stars of masses beyond the interval covered by the chosen stellar evolution library are handled in di erent ways, depending on the mass being larger than the maximum mass of the library or smaller than the minimum one. In the rst case the star is simply considered death. The case of very low mass stars is more complex.

The smallest mass computed in the used libraries are given in Table 1. Stars less m assive than that have MS life times larger than the age of globular clusters and are certainly in the MS for any realistic case. They are norm ally below the limiting magnitude in most CMD s of galaxies. However, they have a quite signi cant contribution to the total m ass of the system and, also to the integrated lum inosity, m agnitudes and colors. A lthough this contribution can be estimated by integration of the IMF, it may be useful to have the possibility of explicitly computing their colors and magnitudes. To this purpose, the libraries mentioned above have been completed downward in mass using the models by Cassisiet al. (2000). They present MSL and Te of stars with masses between 0:09 M for m etallicity values in the range: 0:0002 0:002. For present purpose these and 0.8 M models have been completed with a very low mass sequence for solar chemical composition (Cassisi 2003, private communication). Since these stars have long evolution times and the analysis of the CMD does not usually rely on their exact distribution in it, the e ect of age on their L and Te can be neglected. A polynom ic t has been performed, for each m etallicity, to the m L and the m T<sub>e</sub> plots of the model stars. This can be used to estimate L and  $T_e$  of the synthetic star. The coe cients of the t for the general equation

$$y = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + a_4 x^4 + a_5 x^5$$
(7)

Figure 1 shows two models computed with these relations for masses 0.4 and 0.6 M and Z = 0.004. A good agreement exists with the Bertelli94 tracks. Consistence is also quite good comparing with the G irardi00 and the Teramo libraries (not shown here). However the fact that the very low mass stars are computed from a dimension of the model set (namely the Cassisi et al. 2000 one) and that no age computation is made for them must be taken into account if accurate analysis of CMD is intended in the region around 0.6 M for the Bertelli94 and the Teramo libraries and 0.15 M for the G irardi00 library; i.e., around the lower mass limits of these libraries.

## 6.4. M ass loss

M ass boss by stellar winds plays an important role in the evolution of massive stars. It is included in the Bertelli94 computation for stars with initial mass larger than 12 M . The reader is referred to Bertelli94 for details. Besides massive stars, mass loss during the RGB and the AGB has dramatic consequences on the evolution of low and intermediate mass stars. In these cases the elects of mass loss are included at the moment of computing synthetic stars in IAC -star. Unfortunately, mass loss during these phases and, in particular, during the AGB depends on poorly known physical parameters that should be ne-tuned. In the modeling of mass loss during these phases we follow the criteria given by Bertelli94 and Marigo et al. (1996).

M ass loss during the RGB does not a ect the internal structure of the star and can be neglected in the computation of L and  $T_e$ . However, the integrated m ass loss during this phase determ ines the m ass envelope of the subsequent HB stars and hence its  $T_e$ . For this reason, m ass loss can be integrated during the RGB and introduced in a single step at the tip of the RGB (TRGB). To this purpose, the empirical relation by Reimers (1975) is used:

$$\underline{m} = 1.27 \quad 10^{5} \text{ m}^{-1} \text{L}^{1:5} \text{T}_{2}^{-2} \text{ M} = \text{yr}$$
 (8)

where L and m are in solar units and is a scaling parameter. The lowest stellar initial m ass which evolution time up to the TRGB is smaller than or equal to the globular cluster age (about 13 Gyr) is about 0:80 0:95 M, depending on the metallicity, lower masses corresponding to lower metallicities. From this, the mass loss during the RGB must be subtracted. Since the core mass at the moment of the HeFlash is about 0.55 M, the envelope mass of these stars can be very small. As a consequence, the exact value adopted for strongly a ects  $T_e$  at the zero-age HB. In the current version of IAC-star, has a

default value of 0.35, but it can be modi ed by the user.

C om puting the m ass bost during the AGB is m ore complicated. It can be neglected during the E-AGB, but it is the key parameter controlling the nal evolution of the star during the TP-AGB. Stellar evolution m odels of low and intermediate-m ass stars are in general computed up to the beginning of the TP-AGB only. From that point on, computation is performed within IAC-star following the prescriptions by M arigo et al. (1996). First, the starting point of the TP-AGB is assumed to be the point in the track just before the rst signi cant He-shell ash. Hence, the evolution through the TP-AGB is followed using several basic relations: (i) the m ass base rate; (ii) a relation connecting the core m ass with the total lum inosity of the star; (iii) the rate at which the core m ass increases as a result of shell H-burning, and (iv) an L  $T_e$  relation for the TP-AGB. D etails on these relations can be found in M arigo et al. (1996) and references therein. We shortly sum marize them here for self-consistency.

The mass loss rate is taken as the minimum of the two following values:

$$\log \underline{m} = 11.4 + 0.0123P$$
(9)  
$$\underline{m} = 6.07023 \quad 10^{-3} Lc^{-1} v_{exp}^{-1}$$
(10)

where m and L are the stellar mass and lum inosity (in solar units), P is the pulsation period (in days) c is the light speed (in km s<sup>1</sup>) and  $v_{exp}$  is the term inal velocity of the stellar wind (in km s<sup>1</sup>). Equations 9 and 10 stand for periods shorter and larger than about 500 days, respectively. The wind expansion velocity is calculated in term s of the period as (Vassiliadis & W ood 1993):

$$v_{exp} = 13.5 + 0.056P$$
 (11)

and the period is derived from the period-m ass-radius relation by Vassiliadis & W ood (1993):

$$\log P = 2.07 + 1.94 \log R_2 \quad 0.9 \log m \tag{12}$$

where  $R_{2}$  is the stellar radius in solar units.

To connect the core mass with the total lum inosity of the star, the following two equations are used:

$$L = 238000 \frac{3}{4} Z_{CNO}^{0.04} \text{ (m}_{c}^{2} = 0.0305 \text{m}_{c} = 0.1802 \text{)}$$
(13)

for stars with core mass in the range 0.5 m<sub>c</sub> 0.66 (Boothroyd & Sackmann 1988), and

$$L = 122585 \frac{2}{w} (m_c = 0.46) m^{0.19}$$
(14)

for stars with 0:95 m<sub>c</sub> (Iben & Truran 1978; G roenew egen & de Jong 1993). For stars in the interval 0:66 < m<sub>c</sub> < 0:95, linear interpolation is performed. In the former equations,  $Z_{CNO}$  is the total abundance (in mass fraction) of carbon, nitrogen and oxygen in the envelope and  $_{W} = 4 = (5X + 3 Z)$  is the average molecular weight for a fully ionized gas, where X and Z are the hydrogen and metal abundances, respectively.

The rate at which the core mass increases as a result of shell H-burning is computed using

$$\frac{\text{@m}_{c}}{\text{@t}} = 9:555 \quad 10^{-12} \frac{\text{L}_{H}}{\text{X}}$$
(15)

where X is the hydrogen abundance (in m ass fraction) in the envelope and  $L_H$  is the lum inosity produced by the H-burning shell (in solar units). In fact, the latter is computed by the relation (Iben 1977):

$$L = L_{H} + 2000 (M = 7)^{0.4} \exp[3.45 (m_{c} 0.96)]$$
(16)

Finally, the L  $T_e$  relation of the TP-AGB is obtained by extrapolating the slope of the E-AGB. In fact, to m inim ize random e ects and possible inconsistencies, the slope is computed using the following relation, obtained by bi-logarithmic tofthe slopes of all the models at the end of the E-AGB:

$$\log(\frac{\theta L}{\theta T_{e}}) = 2.870 + 0.994 \log L \quad 0.110 \log Z$$
(17)

where Z is the metallicity.

U sing the form er relations, the TP-AGB is computed for each HB and interm ediate age evolutionary track with mass m 5M, according to the interm ediate mass limit found by M arigo et al. (1996). The TP-AGB computation is term inate either when the envelope mass is zero or the core mass reaches the Chandrasekhar limit (1.4 M).

## 7. Running IAC-star

IAC-star is made available for free use. It can be executed from the internet site http://iac-star.iac.es. A number of interactive software facilities are intended to be m ade available from this site, together with IAC-star itself. In fact, the content and form at of the page is expected to be up-dated, including new software and taking into account user feed-backs. In the present form, a tem plate is provided in which several input parameters can be entered. For all them, default values as well as a brief option list for quick reference are available. Once all the inputs are provided, the user can request the program to be executed. This is performed by a dedicated computer at the Instituto de A strof sica de C anarias (IAC). The resulting data le is stored upon completion in a public access flp directory at the IAC. The address of this directory is given to the user by em ail, to the address provided by him or herself, together with the output le name and information on the used inputs and the output le content. In the following we will describe the input parameters and the output le content. Since the speci c content of both m ay change in future versions of the software, only a general description of them is provided. D etailed, updated information can be found in the IAC-star internet site http://iac-star.iac.es.

## 7.1. Input param eters

The input parameters are introduced as follows (see the IAC-star internet site, http://iac-star.iac.es, for updated details).

The chosen stellar evolution and bolom etric correction libraries.

A seed for the random number generator. The routine used for random number generation is \ran2", obtained from Numerical Recipes in Fortran (Press et al. 1997).

Total num ber of stars computed or saved into the output le. To prevent too long runs, the maximum allowed total computed and saved stars are limited to some big num bers.

The minimum stellar lum inosity or maximum magnitude in a given lter. Although fainter stars are computed, only those brighter than this value are saved into the output le.

The SFR, (t). It is computed by interpolation between several (t) nodes, de ned by the user arbitrarily.

The chem ical enrichment law. Two alternative approaches are allowed to produce the chem ical enrichment law: (i) just an interpolation between several age-metallicity nodes, de ned by the user and permitting a completely arbitrary metallicity law; (ii) computation from usual parameters involved in physical scenarios of chemical evolution, even though a self-consistent physical formulation is not intended in this case. Metallicity dispersion is allowed in both cases.

The  $\mathbb{I}M$  F. It is assumed to be a power law of the mass, but several mass intervals can be de ned.

B inary star control. Both the fraction of binary stars and the secondary to primary m inim um mass ratio can be supplied.

M ass loss parameter. A lthough the default = 0.35 seems a reasonable choice, the user can modify it here. It must be noted that significantly a ects the extension of the HB for low metallicity stars.

## 7.2. The output le content

In the current version, the content of the output le is organized as follows (see the IAC-star internet site, http://iac-star.iac.es, for updated details):

A head containing inform ation about the input param eters. In particular: the libraries used, the total number of computed and stored stars; the minimum luminosity or maximum magnitude stored; the current SFR, CEL and IMF laws; the fraction of binaries and minimum secondary to primary mass ratio, and a heading line for the column content, including a list of the photom etric bands for which magnitudes have been computed.

Several lines each containing the information for a single or binary star. This information includes physical parameters (lum inosity, tem perature, gravity, mass and metallicity), photometric parameters and age.

C bosing: Integrated quantities are provided in the le closing lines, including the total number of ever form ed, currently alive and stored stars; the total mass ever incorporated into stars (in other words, this is just the time integral of the SFR,  $R_{0}^{R_{T}}$  (t)dt); the mass currently locked into alive stars and into stellar remnants, and the total lum inosity and the integrated magnitudes. Besides these integrated quantities, the logarithm of the sum of squared lum inosities (log  $P_{i}L_{i}^{2}$ , where  $L_{i}$  is the lum inosity

of the ith star if it is single or the total lum inosity of the system, if it is binary) and the magnitudes derived from this are also given. These are the magnitudes associated to surface brightness uctuations (SBF). In this sense, IAC -star can be used as a SBF population synthesis code (see M ar n-Franch & Aparicio 2004).

## 8. Final rem arks and conclusions

The program IAC -star, designed to generate synthetic CMD s is presented in this paper. It calculates full synthetic stellar populations on a star by star basis, by computing the lum inosity, e ective temperature and magnitudes of each star. A variety of stellar evolution and bolom etric correction libraries, SFR, IMF and CEL are allowed and binary stars can be computed. The program provides also integrated masses, lum inosities and magnitudes as well as SBF lum inosity and magnitudes for the total synthetic stellar population. In this way, although it is mainly intended for synthetic CMD computation, it can be also used for traditional and SBF population synthesis research.

Am ong the main characteristics of the program, the following are the most relevant:

Lum inosity and e ective tem peratures of each star are computed by direct bilogarithm ic interpolation in the age-m etallicity grid of a stellar evolution library. Two Padua libraries (Bertelli94 and Girardi00) and the new Teram o library (Pietrinfemi et al. 2004) are used in the current version completed by the Cassisi et al. (2000) m odels for very low mass stars. This produces a smooth distribution of stars in the output synthetic CMD which is necessary if accurate studies of SFH are intended.

M ass loss is computed during the RGB and the AGB phases.

The AGB phase is extended to the TP-AGB, covering in this way all the signi cant stellar evolution phases accurately.

Color and lum inosities in a variety of visual broad band, infrared and HST Iters are provided. Bolom etric correction transform ations by G irardi et al. (2002); C astelli & K unucz (2002); F lucks et al. (1994); Lejeune, C uisinier, & Buser (1997); and O riglia & Leitherer (2000) are used for this purpose.

The program is o ered for free use and can be executed at the site http://iac-star.iac.es, with the only requirement of referencing this paper and adknowledging the IAC in any derived publication. It is intended to produce further improved versions of the program after feed-back by the user community. We are deeply indebted to Cesare Chiosi, G ianpaolo Bertelli and other colleagues of the Padua stellar evolution group for a fruitful, long-term collaboration. Building up IAC-star would not have been possible without this collaboration. We are also indebted to Santi Cassisi for his help in the implementation of the Teram o stellar evolution library and corresponding bolom etric corrections, and form any useful discussions and advice. The Computer D ivision of the IAC is acknow ledged for implementing the hardware and the web page interface of the program. Financial support has be provided by the IAC Research D ivision. The authors are funded by the IAC (grant P3/94) and by the Science and Technology M inistry of the Kingdom of Spain (grant AYA 2001-1661 and AYA 2002-01939).

#### REFERENCES

- A lexander, D R. & Ferguson, JW . 1994, ApJ, 437, 879
- Angulo, C., Amould, M., Rayet, M., et al. (NACRE Collaboration), 1999, Nucl. Phys. A., 656, 3
- A paricio, A. 2003, in O bærved HR diagram s and stellar evolution models: the interplay between observational constraints and theoretical limitations, eds. T. Lejeune & J. Fernandes, ASP Conference Series, in press
- Aparicio, A., Gallart, C., Chiosi, C. & Bertelli, G. 1996, ApJ, 469, L97
- Aparicio, A., Gallart, C., & Bertelli, G. 1997a, AJ, 114, 669
- Aparicio, A., Gallart, C., & Bertelli, G. 1997b, AJ, 114, 680
- Aparicio, A. & Hidalgo, S. 2004, in preparation
- Aparicio, A., Tikhonov, N., & Karachentsev, I. 2000, AJ, 119, 177
- Ardeberg, A., Gustafsson, B., Linde, P., & Nissen, P.E. 1997, A&A, 322, L13
- Bertelli, G., Bressan, A., Chiosi, C., Fagotto, F., & Nasi, E. 1994, A& AS, 106, 275
- Bertelli, G., Mateo, M., Chiosi, C., & Bressan, A. 1992, ApJ, 388, 400
- Bertelli, G. & Nasi, E. 2001, AJ, 121, 1013
- Bertelli, G., Nasi, E., Girardi, L., Chiosi, C., Zoccali, M., Gallart, C. 2003, AJ, 125, 770
- Boothroyd, A. I., & Sackmann, I.-J. 1988, ApJ, 328, 641
- Cassisi, S., Castellani, V., Ciarcelluti, P., Piotto, G. & Zoccali, M. 2000, MNRAS, 315, 679
- Castelli, F., Gratton, R.G. & Kurucz, R.L. 1997, A&A, 318, 841
- Castelli, F. & Kurucz, R.L. 2002, in Modeling of Stellar Atmospheres, IAU Symp. 210 (N.E. Piskunov et al. Eds), in press

Caughlan, G R. & Fowler, W A. 1988, Atom ic Data and Nuclear Data Tab. 40, 283

- Chiosi, C. 1998, in Stellar A strophysics for the Local G roup, V III C anary Islands W inter School of A strophysics, eds. A paricio, A., H errero, A., & Sanchez, F., C am bridge University P ress.
- Chiosi, C., Bertelli, G., & Bressan, A. 1988, A&A, 196, 84
- Chiosi, C., Bertelli, G., & Bressan, A. 1992, ARA & A, 30, 235
- Cox, AN., & Stewart, JN. 1970a, ApJS, 19, 243
- Cox, AN., & Stewart, JN. 1970b, ApJS, 19, 261
- Dohm -P lamer, R.C., Skillman, E.D., Saha, A., et al. 1997, AJ, 114, 2527
- Dolphin, A.E. 1997, New Astronom y, 2, 397
- Fluks, M A., Plez, B., The, P.S. et al. 1994, A&AS, 105, 311
- Gallart, C., Aparicio, A., Bertelli, G. & Chiosi, C. 1996, AJ, 112, 1950
- Gallart, C., Frædman, W. L., Aparicio, A., Bertelli, G., & Chiosi, C. 1999, AJ, 118, 2245
- Girardi, L., Bertelli, G., Bressan, A., Chiosi, C., Groenewegen, M.A.T., Marigo, P., Salasnich, B. & Weiss, A. 2002, A & A, 391, 195
- GirardiL, Bressan A, BertelliG, & ChiosiC.2000, A&AS, 141, 371
- GirardiL, Bressan A, ChiosiC, BertelliG, NasiE, 1996 A&AS, 117, 113
- Groenewegen, M.A.T. & de Jong, T. 1993, A & A, 267, 410
- Harris, J. & Zaritsky, D. 2001, ApJS, 136, 25
- Hernandez, X., Valls-Gabaud, D., & Gilmore, G., 1999, MNRAS, 304, 705
- Holtzman, J.A., Gallagher, J.S., Cole, A.A. et al. 1999, AJ, 118, 2262
- Huebner W F., Merts, A L., Magee, N H. & Argo, M F. 1997, Los A lam os Scienti c Laboratory Report LA-6760-M
- Hurley-Keller, D., Mateo, M., & Nemec, J. 1998, AJ, 115, 1840
- Iben, I. 1977, ApJ, 217, 788
- Iben, I., & Truran, J.W . 1978, ApJ, 220, 980
- Iglesias, C.A. & Rogers, F.J. 1996, ApJ, 464, 943
- Irw in, A W . et al. 2004, in preparation
- Kippenhahn, R., Thomas, H.-C. & Weigert, A. 1965, Z. Astrophys. 61, 241
- Kroupa, P., Tout, C.A., & Gilmore, G. 1993, MNRAS, 262, 545

- Kunz, R., Fey, M., Jaeger, M., Mayer, A., Hammer, J.W., Staudt, G., Harissopulos, S., & Paradellis, T. 2002, ApJ, 567, 643
- Kurucz, R.L. 1992, in Stellar Populations of Galaxies, eds. B.Barbuy and A.Renzini (D ordrecht: Kluwer) p. 225
- Landre, V., Prantzos, N., Aguer, P., Bogaert, G., Lefebre, A., & Thibaud, J.P. 1990, A&A, 240, 85
- Lejeune, Th., Cuisinier, F., & Buser, R. 1997, A&AS, 125, 229
- M aeder, A. 1974, A & A, 32, 177
- Marigo, P., Bressan, A., & Chiosi, C. 1996, A & A, 313, 545
- Marn-Franch, A., & Aparicio, A. 2004, in preparation
- Mihalas, D., Hummer, D.G., Mihalas, B.W. & Dappen, W. 1990, ApJ, 350, 300
- Ng, Y K, Brogt, E, Chiosi, C, & Bertelli, G. 2002, A & A, 392, 1129
- Origlia, L, & Leitherer, C. 2000, AJ, 119, 2018
- Peimbert, M., Coln, P., & Sammiento, A. 1994, in Violent Star Formation: from 30 Doradus to QSOs, ed. Tenorio-Tagle, G., Cambridge University Press
- Pietrinfemi, A., Cassisi, S., Salaris, M. & Castelli, F. 2004, ApJ, submitted.
- Press, W. H., Teukolsky, S. A., Vetterling, W. T., & Flannery, B. P. 1997. Numerical Recipes in Fortran 77. The art of scientic computation, Cambridge University Press
- Reimers, D. 1975, Mem. Soc. R. Sci. Liege 6 (8), 369
- Rogers, F.J., & Iglesias, C.A. 1992, ApJS, 79, 507
- Stappers, B.W., Mould, J.R., Sebo, K.R. et al. 1997, PASP, 109, 292
- Tinsley, B. 1980, Fund. Cosm ic Phys., 5, 287
- Tolstoy, E., & Saha, A. 1996, ApJ, 462, 672
- Tosi, M., Greggio, L., Marconi, G., & Focardi, P. 1991, AJ, 102, 951
- Vallenari, A., Bertelli, G., & Schmidtobreick, L. 2000, A&A, 361, 73
- Vallenari, A., Chiosi, C., Bertelli, G., & Ortolani, S. 1996a, A & A, 309, 358
- Vallenari, A., Chiosi, C., Bertelli, G., Aparicio, A. & Ortolani, S. 1996b, A & A, 309, 367
- Vassiliadis, E., & Wood, P.R. 1993, ApJ, 413, 641
- Weaver, TA.& Woosley, SE. 1993 Phys. Rep. 227, 65

This preprint was prepared with the AAS  $\mathbb{P}T_{E}X$  m acros v4.0.

M odels	Teram o	G irardi00	Bertelli94	
Mass range [M]	0.5–10	015-7	0.6–120	
Z range (Z <b>,</b> n <i>s</i> tep)	0.0001-0.04 (10)	0.0-0.03 (8)	0.0004-0.05 (5)	
= 1=H p	1.913	1.68	1.63	
Y ;Z	0.2734,0.0198	0.273,0.019	0.282,0.02	
0 vershooting	canonical& 0.2H <sub>p</sub> <sup>a</sup>	0.5H <sub>p</sub> <sup>a</sup> ; <sup>b</sup>	0.5H <sub>p</sub> <sup>a;b</sup>	
EOS	Irw in04	К ірр65+G іг96+М НD 90	K ipp65	
Rad.Opacity	0 PA L 96+ A lex 94	0 PAL92+A lex94	0 PAL92+Hueb77+Cox70	
Nucl. Reactions	NACRE+Kunz02	Caughlan88+Landre90+WW 93	Caughlan88	
ModelAtm.	Castelli02	Castelli97 K unicz92+ e		

Table 1. Model grids used in the current version of IAC-star

Note. | Key to references: Teram o: Pietrinferni et al. (2004); GirardiOO: Girardi et al. (2000); Bertelli94: Bertelli et al. (1994); Inw in04: Inw in, A.W. et al. (2004); OPAL96: Iglesias & Rogers (1996); A lex94: A lexander & Ferguson (1994); NACRE: Angub et al. (1999); KunzO2: Kunz et al. (2002); CastelliO2: Castelli & Kunucz (2002); Kipp65: Kippenhahn, Thom as & Weigert (1965); Gir96: Girardi et al. (1996); M HD 90: M ihalas et al. (1990); OPAL92: Rogers & Iglesias (1992); Caughlan88: Caughlan & Fow ler (1988); Landre90: Landre et al. (1990); W W 93: Weaver & Woosley (1993); Castelli97: Castelli, Gratton & Kunucz (1997); Huebb77: Huebner et al. (1997); Cox70: Cox & Stewart (1970a,b); Kunucz92: Kunucz (1992)

<sup>a</sup>The prescription on how overshooting is turned o as a function of mass heavily determines the morphology of the MS-TO at intermediate ages. Teram o: = (M/M - 0.9)/4 for 1.1M M 1:7M, =  $0.2H_p$  for M 1:7M and = 0 for M 1:1M ; G irardiOO: = M = M - 1.0 for 1.0M M 1:5M, with =  $0.5H_p$  for M 1:5M and = 0 for M 1:0M ; Bertelli94: =  $0.5H_p$  for M 1:6M, =  $0.25H_p$  for M 1:5M.

<sup>b</sup>The param eter describing overshooting is its extent across the border of the convective zone, in units of pressure scale height  $(H_p)$ . This param eter, de ned in Bressan et al. (1981) is not equivalent to others present in the literature. E.g. = 0.5 in the Padova form alism approx corresponds to  $0.25H_p$  above the convective border adopted by the Teram o group.

Table 2. Coe cients of the log L  $\,$  log m and log T\_e  $\,$  log m  $\,$  ts for very low mass stars

Relation	a <sub>0</sub>	a <sub>1</sub>	a <sub>2</sub>	a <sub>3</sub>	a4	<b>a</b> <sub>5</sub>
Z = 0.0002, Y = 0.23:						
logL vs. logm :	1.1387	11.3958	15.8615	8.7825	0.1773	
logT <sub>e</sub> vs. logm : Z=0.0006,Y=0.23:	4.2038	4.3333	13.5738	22.1148	18,2446	6.0709
bgL vs. bgm :	1.1457	11.7429	16.6033	9.4539	0.3695	
$\log T_{\rm e}$ vs. $\log m$ :	4.1851	4.1282	12,2825	19.1799	15.4070	5.0853
Z=0.001,Y=0.23:						
logL vs. logm :	1.0342	10.9539	14,2405	6.5840	{0 <b>.</b> 8234	
$\log T_{\rm e}$ vs. $\log m$ :	4,2229	4.7253	14.9449	24.4181	20.0889	6.6370
Z=0.002, Y=0.23:						
logL vs. logm :	0.8190	9.6332	10.7591	2.8464	{2.2331	
$\log T_{\rm e}$ vs. $\log m$ :	4.2593	5.4559	18.3152	31.0882	26.0219	8.5882
Z = 0.02, Y = 0.27:						
logL vs. logm :	0.3668	9.8685	16.8057	14.4196	3.9769	
$\mbox{log} T_e$ vs. logm :	3.8133	1.5630	3,2232	2.8249	0.7851	



Fig. 1. Set of evolutionary tracks for m etallicity Z = 0.004 from the library of Bertelli94 (full and dashed lines) extended by two very low mass models interpolated from Cassisi et al. (2000) of age 10 G yr (starred dots). Colors indicate di erent mass ranges, as follows. Red: high mass tracks; blue: interm ediate mass tracks; magenta: low mass stars up to the He-ash; black: low mass stars HB. In all the cases, dashed lines represent the TP-AGB extensions of the Padua tracks computed by IAC-star. Labels correspond to the track initial masses, in M .



Fig. 2. Synthetic CMD computed using constant SFR from 13 Gyr ago to date and metallicity linearly increasing from  $Z_0 = 0.0001$  to  $Z_f = 0.02$  and a K roupa et al. (1993) IMF. The Bertelli94 stellar evolution library and the Lejeune et al. (1997) bolom etric correction library have been used. Stars in di erent age intervals are plotted in di erent colors. The color code is given in the gure, in Gyr. Labels indicate the evolutionary phase (see text for an explanation).



Fig. 3. Synthetic CMD computed using constant SFR from 13 Gyr ago to date and metallicity linearly increasing from  $Z_0 = 0.0001$  to  $Z_f = 0.001$ , Salpeter IMF, and a 0.3 fraction of binary stars with minimum mass ratio 0.7. The Bertelli94 stellar evolution library and the Lejeune et al. (1997) bolom etric correction library have been used. Colors and magnitudes of binary stars are represented in green. G reen points in the single star locus of the MS correspond to binary stars in which the primary is dead.



Fig. 4. Examples of CELs for di erent choices of  $f_{f}$ , and (t) are given for xed  $Z_{0} = 0$  and  $Z_{f} = 0.02$ . For (t), an exponential law of the form (t) = A exp(t=) has been used, where A is set to normalyze the integral of (t) for the entyre life of the system. Time scale is normalyzed, with the instant t = 0 corresponding to the beginning of the star form ation and the instant t = 1 corresponding to its end. In the upper window the parameter choices are displayed. Bracketed gures correspond to [; f; ; ]. C losed-box m odels are drawn in blue, infallm odels are drawn in red and out ow m odels in m agenta.



Fig. 5. Example of a CEL with m etallicity dispersion. A constant SFR has been used with = 0 and  $_{\rm f} = 0.2$  in both cases, but  $Z_0 = 0$ ,  $Z_{\rm f} = 0.01$  for the lower law and  $Z_0 = 0.004$ ,  $Z_{\rm f} = 0.02$  for the upper one. Time scale is normalyzed, with the instant t = 0 corresponding to the beginning of the star form ation and the instant t = 1 corresponding to its end. For each age, stars would have m etallicities comprised within the shaded area.