

Abstract. We present CCD UBVI photometric and medium/high resolution spectroscopic observations obtained in the field of the previously unstudied dissolving open cluster candidates NGC 5385, NGC 2664 and Collinder 21. Our analysis stands on the discussion of star counts, photometry, radial velocity distribution, and proper motions available from the Tycho 2 catalogue. All the three aggregates clearly emerge from the mean Galactic field, but, regrettably, the close scrutiny of proper motions and radial velocities reveals that we are not facing any physical group. Instead, what we are looking at are just chance alignments of a few bright unrelated stars. Our analysis casts some doubt on the Bica et al. (2001) criterion to look for Possible Open Cluster Remnants. It seems mandatory to define a better criterion to adopt for further investigations.

Key words: open clusters and associations: individual: Collinder 21, NGC 5385 and NGC 2664 – open clusters and associations : general

NGC 5385, NGC 2664 and Collinder 21: three candidate Open Cluster Remnants^{?, ??}

Sandro Villanova¹, Giovanni Carraro^{1,2,3}, Raul de la Fuente Marcos⁴, and Ruggero Stagni¹

¹Dipartimento di Astronomia, Università di Padova, Vicolo Osservatorio 2, I-35122 Padova, Italy

²Dipartimento de Astronomia, Universidad de Chile, Casilla 36-D, Santiago, Chile

³Astronomy Department, Yale University, P.O. Box 208101, New Haven, CT 06520-8101 USA

⁴South University Madrid Campus, C/ Vima 3, E-28003, Madrid, Spain

Received April 2004; accepted

1. Introduction

Bica et al. (2001) identified 34 neglected star clusters (see http://obswww.unige.ch/webda/dissolving_ocl.htm) having relatively high galactic latitude ($|b| > 15^\circ$), being poorly populated and appearing to be candidate objects to be experiencing the late stages of star cluster dynamical evolution. For these objects the acronym PO CR {Possible Open Cluster Remnant} is used. In their study, Bica et al. (2001) basically select these candidates on the basis of star counts. All of them indeed clearly emerge from the background.

The final residue of open star cluster evolution is often called an open cluster remnant (OCR). These ghostly objects are characterized by very low surface brightness and they are hardly distinguishable against the background field stars (de la Fuente Marcos 1998). They consist of a small number of coeval relatively massive stars, most of which are binaries, confined in a core, since due to the evolution the less massive star members evaporated from the cluster and merged with the Galactic disk field (de la Fuente Marcos 1997).

Recently, these objects have started to receive some attention mainly because they play a fundamental role in our understanding of the subject of open cluster evolution and dissolution, and, ultimately, on the origin of the field star population. Photometry, kinematics, binary percentage and membership are basic data to constrain N-body models of cluster dissolution, which aim to reconstruct {for instance} the Initial Mass Function (IMF) of Galactic clusters (e.g. de la Fuente Marcos 1997).

On the other hand, they may be what still remains of old clusters, and therefore once discovered they can allow us to improve on the statistics of old open clusters in the Galactic disk. These objects are widely recognised to be of paramount importance for our understanding of the formation and early evolution of the Galactic disk (Carraro et al. 1998). Some PO CR candidates have been studied recently.

The best discussed case is doubtless NGC 6994 (M 73), studied by Bassino et al. (2000), Carraro (2000) and Odenkirchen and Soubiran (2002). By combining together photometry, astrometry and spectroscopy this object turned out to be just a chance alignment of four stars.

Pavani et al. (2001) discuss new photometry of NGC 1901 and NGC 1252, bringing new evidence that these stellar groups are PO CRs. However, proper motions studies for NGC 1252 seem to point to the opposite conclusion (Baumgardt 1998). On the other hand, the radial velocity survey conducted by Villanova et al. (2003a,b) confirms that NGC 1901 is a genuine PO CR.

Finally, on the basis of only star counts and photometry, Carraro (2002) discusses the cases of NGC 7772 and NGC 7036 and Baume et al. (2003) the case of NGC 1663, favoring the possibility that these two objects be indeed OCRs and providing a list of possible candidate members for spectroscopic follow up.

In this paper we present a photometric, astrometric and spectroscopic investigation of three previously unstudied PO CRs, namely NGC 5385 and NGC 2664 and Collinder 21, with the goal to clarify their real nature. The analysis closely follows the strategy earlier proposed by Odenkirchen and Soubiran (2002).

The layout of this paper is as follows.

In Sect. 2 we briefly present the observations and data reduction. Sect. 3 illustrates star counts analysis and spatial configurations. Sect. 4 is dedicated to the outcomes of the proper motion analysis, whereas in sect. 5 we discuss our spectroscopy. Our conclusions are presented in sect. 6.

Send o print requests to: Sandro Villanova: villanova@pd.astro.it

[?] Based on observations carried out at Mt Ekar Observatory, Asiago, Italy

^{??} Photometry is only available in electronic form at the CDS via anonymous ftp to cdsarc.u-strasbg.fr (130.79.128.5) or via <http://cdsweb.u-strasbg.fr/cgi-bin/qcat?J/A+A/>

Table 1. Basic parameters of the observed objects. Coordinates are for J2000.0 equinox and have been taken from Dias et al. (2002)

Name	l		b	
	hh:mm:ss	°:′:″	°	°
NGC 5385	13:52:27	+76:10:24	118.19	+40.38
NGC 2664	08:47:11	+12:36:06	214.34	+31.31
Collinder 21	01:50:11	+27:04:00	138.73	-33.99

2. Observations and Data Reduction

2.1. Photometry

Observations were carried out with the AFO SC camera at the 1.82 m Copernico telescope of Cima Ekar (Asiago, Italy), in the photometric nights of December 17 and 18, 2001. AFO SC samples a $8^{\circ}:14$ $8^{\circ}:14$ field in a 1K 1K thinned CCD. The typical seeing was between 1.8 and 2.3 arcsec. The basic data of the studied objects are summarized in Table 1, whereas the details of the observations are listed in Table 2.

The data has been reduced by using the IRAF¹ packages CCDRED, DAOPHOT, and PHOTCAL. The calibration equations obtained by observing Landolt (1992) SA 93, PG 1047+003, PG 2331+055 and PG 0231+051 fields along both the nights, are:

$$\begin{aligned}
 u &= U + 3.520 \quad 0.042 + (0.099 \quad 0.030) (U - B) + 0.58 X \\
 b &= B + 1.407 \quad 0.012 \quad (0.004 \quad 0.017) (B - V) + 0.29 X \\
 v &= V + 0.752 \quad 0.009 + (0.036 \quad 0.012) (B - V) + 0.16 X \\
 i &= I + 1.619 \quad 0.017 \quad (0.011 \quad 0.015) (V - I) + 0.08 X
 \end{aligned}
 \tag{1}$$

where $UBVI$ are standard magnitudes, $ubvi$ are the instrumental ones, and X is the air mass. The standard stars in these fields provide a very good color coverage, being 0.329 $(B - V)$ 1.448 . For the extinction coefficients, we assumed the typical values for the Asiago Observatory (Desidera et al. 2001). Photometric global errors have been estimated following Patat and Carraro (2001). For the V filter, they amount to 0.02, 0.04 and 0.08 at $V = 12.0$, 16.0 and 20.0, respectively.

The photometry for the most obvious candidates is reported in Tables 3 to 5, whereas all the photometric catalogues are available upon request to the authors.

2.2. Spectroscopy

Medium ($R = 3;600$) and high resolution ($R = 20;000$) spectra in the field of NGC 5385, NGC 2664 and Collinder 21 have been obtained by

¹ IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

Table 2. Journal of photometric observations in the field of Collinder 21, NGC 5385, and NGC 2664 and standard star fields (December 17-18, 2001).

Field	Filter	Time integration (sec)	Seeing (′′)
Collinder 21	U	15,60	2.1
	B	2,5,15	2.0
	V	1,3,10,10	2.1
	I	1,8	2.0
SA 93	U	120	1.9
	B	60,60	2.0
	V	30,30,30	2.0
	I	30,30	2.0
NGC 5385	U	240	2.2
	B	60,60	2.1
	V	15,15	2.3
	I	15,15	2.3
PG 1047+003	U	120	1.9
	B	60,60	2.0
	V	30,30	2.0
	I	30,30	2.0
PG 0231+051	U	120	1.9
	B	60,60	2.0
	V	30,30	2.0
	I	30,30	2.0
PG 2331+055	U	120	1.9
	B	60,60	2.0
	V	30,30	2.0
	I	30,30	2.1
NGC 2664	U	15,240	1.8
	B	1,10,60	1.7
	V	1,5,30,30	1.8
	I	1,5,30	1.8

using AFO SC in Echelle mode (in the grism # 9 and # 10 combination) and the REOSC Echelle Spectrograph onboard the 1.82 m telescope of Asiago Astronomical Observatory.

The Echelle spectrograph works with a Thomson 1024 1024 CCD and the allowed wavelength coverage is approximately 4140 - 6840 Å. Details on this instrument are given in Munari & Zwitter (1994) and at the Asiago Observatory Homepage².

The exposures times were 45 minutes for the all stars. In order to improve the signal to noise ratio, two exposures were taken for each star reaching at the end $S=N$ values up to 150. The data have been reduced with the IRAF package ECHELLE using thorium lamp spectra for wavelength calibration purposes. By comparing nal

² <http://www.pd.astro.it/Asiago/2000/2300/2320.htm> 1




Fig. 1. Distribution of stars from USNO-B1 catalogue in a $20'' \times 20''$ field around NGC 5385. The size of the dots are proportional to the magnitude of the stars.

known sky lines positions along the spectra we obtained an error measurement of about 0.01 Å.

The AFOSC in Echelle mode spectra cover the wavelength range 4100–6800 Å, and for typical exposure times of 30 to 60 minutes we obtained S/N values up to 300, which allows us to derive radial velocity with error almost always lower than 8 km/s. In this case, the data has been reduced with the IRAF package for one-dimensional spectroscopy CTIO SLIT and by using the HgCd lamp spectra for wavelength calibration purposes. Together with radial velocities we provide also spectral classifications, which have been derived as described in Villanova et al. (2004).

In Tables 6 to 8 we report some details of the spectroscopic observations.

3. Star counts, surface density and spatial distribution

POCRs are identified by Bica et al. (2001) as star overdensities at relatively high galactic latitude. For star count purposes we are going to use USNO-B1 (Monet et al. 2003) catalogue for NGC 5385 and UCAC2 (Zacharias et al. 2003) catalogue for NGC 2664 and Collinder 21, because UCAC2 does not extend to NGC 5385 declination. In the following, we are comparing star counts inside the OCR region with star counts in a *o*-set field, which is aimed to represent the mean Galactic disk field.

A crucial role in star counts is played by binaries. Antici-




Fig. 2. Histogram of the surface density of the stars shown in Fig. 1, measured by counts in $0''.5$ wide annuli centered on $\alpha = 13^{\text{h}}52^{\text{m}}19^{\text{s}}$, $\delta = +76^{\circ}10'56''$. Dots with error bars indicate the mean surface density of stars in a surrounding field centered on $\alpha = 13^{\text{h}}54^{\text{m}}10^{\text{s}}$, $\delta = +76^{\circ}49'30''$, and the expected 1- σ range of statistical fluctuations around this mean for each bin.

pating the results of Section 5, we find that in the OCRs under investigation the binary fraction ranges from 18% to 25%, a value which is not much different from the estimated typical percentage in the Galactic disk field population, which in the the proper F-K spectral range amounts to about 14% (Hallowachs et al. 2003). If one considers also M-type stars, this percentage would increase. Therefore, we do not expect that binary stars significantly affect star counts.

3.1. NGC 5385

Fig. 1 shows a finding chart in a $20'' \times 20''$ region around NGC 5385 down to $B = 21.5$ according to USNO-B1. The center of this image, which we also consider the center of NGC 5385, has been derived as the mean of the position of the 9 most obvious members of the aggregate. Indeed this object seems to consist of 9 stars with magnitudes 11–13 in a field of about $3''.5 \times 3''.5$. The aggregate appears rather sparse, being the mean mutual angular separation larger than $30''$. Out of the central group, there is

a clear deficiency of similar brightness stars, which appear at least 1 mag fainter on the average, if we exclude some obvious case northward NGC 5385.

In Fig. 2 we show the surface density of stars of different magnitudes as a function of angular distance from the center of NGC 5385, (based on USNO-B1). The central group of bright stars produces a significant local enhancement in the surface density of stars with magnitudes $B < 13.5$ (see panel (a)). Fainter stars in the magnitude bins $13.5 < B < 17.0$ (panel (b)) $B < 18.0$ (panel (c)) do not show any signs of a spatial concentration, star counts in the cluster area always agree within the uncertainties with the surrounding fields. In fact in the lower panel the overdensity in the central part lies quite close to the noise level to be considered significant.

In conclusion only the brightest stars indicate the existence of a cluster, and we are here going to look for membership in this magnitude range. This of course does not exclude that also fainter stars could be members, in the case the cluster turns out to be a physical one.

3.2. NGC 2664

Fig. 3 shows the distribution of stars in a field of $30^0 \times 30^0$ around NGC 2664 down to a magnitude UCMag (between V and R) of 16.3 according to UCAC2. This figure reveals that NGC 2664 is defined by 4 stars with magnitude $10.5 < \text{UCMag} < 11.5$ concentrated in $2^0.5 \times 2^0.5 \text{ arcmin}$ around the nominal center, taken as the mean of the 4 most obvious members positions. In the immediate surroundings there is a clear lack of bright stars, although outward about 3^0 from the cluster center the field is quite rich in bright stars, and a few other concentrations of the same kind as NGC 2664 seem to be present.

In Fig. 4 we show the surface density of stars of different magnitudes as a function of angular distance from the center of NGC 2664, (based on UCAC2 as well). The central group of bright stars produces a small but sizeable local enhancement in the surface density of stars with magnitudes $\text{UCMag} < 13.5$ (see panel (a)). The same seems to occur for the magnitude bins $13.0 < \text{UCMag} < 16.0$ (panel (b)), whereas fainter stars, in the magnitude bin $\text{UCMag} < 16.0$ (panel (c)) do not show any clear sign of a spatial concentration. At all distances in fact the cluster density profile is comparable with the field, and only at a distance of 3 arcmin does there seem to be a significant enhancement.

Therefore, the asterism NGC 2664 seems to be mainly identified by four bright stars and a few fainter stars in its immediate vicinity.

3.3. Collinder 21

Fig. 5 shows the distribution of stars in a field of $30^0 \times 30^0$ around Collinder 21 down to a magnitude of 15.3 according to UCAC2. The appearance of Collinder 21




Fig. 3. Distribution of stars from UCAC2 catalogue in a $30^0 \times 30^0$ field around NGC 2664. The size of the dots are proportional to the magnitude of the stars.

(also designated as OCL 371 and C0147+270) on the sky is very impressive. It consists of about 10 stars distributed in a ring-like structure in a very poorly populated field, which makes this structure to emerge very sharply. There are two well known binary stars in this circlet: the visual binary system BD+26 305AB and the binary HD 11142, resolved by speckle interferometry, which presents a separation of $0^0.56$ between its components. All the other stars have much larger angular separation.

In Fig. 6 we show the surface density of stars of different magnitudes as a function of angular distance from the center of Collinder 21, based on USNO-B1. The bright star belonging to the circlet produce a significant local enhancement in the surface density of stars with magnitudes $\text{UCMag} < 13.0$ (see panel (a)) at a distance corresponding to the radius of the circlet. Fainter stars in the magnitude bins $13.0 < \text{UCMag} < 16.0$ (panel (b)) $\text{UCMag} < 16.0$ (panel (c)) do not show any signs of a spatial concentration since star counts in the cluster area always agree within the uncertainties with the surrounding fields.

The only exception is again at a distance of about 3 arcmin , where the enhancement is always significant at 2 level. In conclusion star counts suggests that a clear concentration exist in the region of Collinder 21.

At the end of this section, we would like to stress that the existence of some stars overdensity is a necessary condition




Fig. 4. Histogram of the surface density of the stars shown in Fig. 3, measured by counts in $0.5'$ wide annuli centered on $\alpha = 08^h 47^m 07^s$, $\delta = +12^\circ 36' 30''$. Dots with error bars indicate the mean surface density of stars in a surrounding field centered on $\alpha = 08^h 53^m 05^s$, $\delta = +12^\circ 59' 50''$, and the expected 1 σ range of statistical fluctuations around this mean for each bin.

for a physical cluster to exist, but not a sufficient one. We repeat here the kind of analysis proposed by Odenkirchen and Soubiran (2002), to show what actually number statistics can tell us about these 3 objects.

We use their Equation (1):

$$p(n) = \frac{1}{n!} (\lambda r^2)^n \exp(-\lambda r^2) \quad (2)$$

which gives us the probability of finding n neighbours in a circular field of radius r around an arbitrarily selected star, for a random distribution of stars with mean surface density λ .

The statistics of star counts in the comparison fields for NGC 5385, NGC 2664 and Collinder 21 provide in the magnitude range 10:0 m ag 13:0 a mean surface density of 20, 16 and 64 stars/deg², respectively.

In the case of NGC 2664, we are looking at the probability to find 3 or more neighbours within a radius of 2.5 arcmin. As a consequence the probability of finding 3 or more neighbours turns out to be lower than 1.2×10^{-4} . As for NGC 5385, we must look for the probability to find at least 8 neighbours within about 3 arcmin, which




Fig. 5. δ distribution of stars from UCAC2 catalogue in a $30' \times 30'$ field around Collinder 21. The size of the dots are proportional to the magnitude of the stars.

amounts to less than 1.1×10^{-11} . Finally, in the case of Collinder 21, we need at least 5 neighbours within about 3 arcmin, and the corresponding probability results lower than 4.4×10^{-4} .

These numbers are saying to us that random configurations like those ones we are facing are rare. As a consequence, in all the three cases (and especially for NGC 5385) we cannot exclude these are physical groupings, and therefore only proper motions and radial velocities can settle the question of their real nature.

4. Proper motion analysis

Important information on the kinematics of the luminous stars in and around our targets can be derived from the proper motions available in the Tycho-2 catalogue (Høg et al. 2000). We decided to opt for this catalogue since it provides homogeneous proper motions data for all the targeted stars. The recently released UCAC2 catalogue in fact does not provide data for Collinder 21.

In Figs. 7, 9 and 11 we show the finding charts of our targets, in a field which roughly corresponds to our photometric survey, and indicate there the stars for which we have proper motion measurements at our disposal. Moreover, Tables 3 to 5 list proper motion values, magnitudes and colors for the most obvious candidate members of the aggregates under investigation.




Fig. 8. Photometry and proper motions in the region of NGC 5385. a) Color-Magnitude diagram. Open squares are Tycho 2 stars having both proper motion and radial velocity measurements (see panel b) and Table 6), while filled triangles are stars having only proper motion measurements. b) Vector point plot of Tycho 2 proper motions and proper motion errors. Symbols are as in panel a). As for the identification, we are plotting here Tycho 2 numbering. See text for more details.

Table 3. Photometry and proper motion of the most obvious candidate members in the field of NGC 5385.

ID	TYC 4558-	(J2000.0)	(J2000.0)	V	(B - V)	(U - B)	cos	[mas/yr]	[mas/yr]
1	1641	13:52:54.104	+76:09:59.63	11.29	0.50	0.18	-7.5	1.9	11.7 2.1
2	1385	13:52:08.976	+76:10:55.12	11.52	0.39	0.14	2.5	2.0	-3.0 2.1
3	457	13:51:58.757	+76:13:20.41	11.39	0.58	0.32	-44.6	2.0	-6.6 2.1
4	399	13:52:18.535	+76:10:45.58	11.31	0.68	0.39	-29.0	2.3	13.8 2.3
5	1797	13:51:49.975	+76:12:11.67	11.32	0.72	0.56	-23.6	2.1	17.0 2.2
6	1873	13:52:32.078	+76:08:07.41	11.78	0.64	0.33	-42.2	2.2	26.8 2.3
7	217	13:52:34.850	+76:11:10.84	11.77	1.04	1.03	-12.5	2.3	6.5 2.4

Table 4. Photometry and proper motion of the most obvious candidate members in the field of NGC 2664.

ID	TYC 816-	Name	(J2000.0)	(J2000.0)	V	(B - V)	(U - B)	(V - I)	cos	[mas/yr]	[mas/yr]
1	2354	BD + 13 1898	08:47:11.173	+12:37:58.63	10.75	0.43	0.19	0.69	-6.4	1.9	5.6 2.1
2	1826		08:46:58.629	+12:37:10.47	11.11	0.37	0.29	0.61	-8.3	2.0	-4.7 2.1
3	1890		08:47:13.795	+12:36:13.50	10.99	1.03	1.05	1.13	1.5	2.0	-26.0 2.1
4	2400	BD + 13 1899	08:47:06.699	+12:34:38.76	11.41	1.11	0.99	1.15	-2.6	2.3	3.2 2.3

4.1. NGC 5385

The vector point diagram for all the candidate member stars having proper motion measurements in the field of NGC 5385 (see Fig. 7 for the identification) from Tycho 2 catalogue is shown in Fig. 8, where panel a) presents the Color-Magnitude Diagram (CMD) derived from our photometry, whereas panel b) shows the proper motions

distribution. Of the 10 brightest stars which presumably define the aggregate (they clearly detach from the bulk of the stars), 7 are plotted with open squares, and these are the stars for which we obtained a radial velocity measurement (see below). However, we found in the Tycho 2 catalogue proper motion for all these 10 stars, and the 3 for which no radial velocity measurement is

Table 5. Photometry and proper motion of the most obvious candidate members in the field of Collinder 21.

ID	TYC 1759-	Name	(J2000:0)	(J2000:0)	V	(B - V)	(U - B)	(V - I)	cos	μ_{α} [mas/yr]	μ_{δ} [mas/yr]
1	450	BD + 26 307	01:50:11.56	+27:01:59.0	8.20	0.45			56.5	1.5	-33.5 0.9
2	1472	BD + 26 304	01:50:01.41	+27:03:40.0	9.80	0.61		0.82	5.1	1.2	-11.0 1.2
3	1025	BD + 26 306	01:50:12.17	+27:07:47.1	9.69	1.75	1.65		-8.7	1.4	-1.1 1.4
4	468		01:50:08.22	+27:07:00.2	11.32	1.39	1.50	1.46	-0.7	2.3	-3.3 2.3
5	994		01:50:13.40	+27:05:57.5	11.17	0.74	0.44	0.95	48.7	1.9	-11.5 1.9
6	462		01:50:27.06	+27:04:09.7	11.32	0.59	0.26	0.83	-6.4	1.9	-8.4 1.9
7	1333		01:50:08.53	+27:01:49.4	11.61	0.89	0.69	1.07	-6.9	1.2	+16.7 0.9
8	266		01:50:05.23	+27:02:00.0	11.21	0.98	0.79	1.14	24.1	1.9	4.3 1.9
9	866	BD + 26 305b	01:50:02.41	+27:06:13.8	10.09	0.85	0.58	1.04	-3.9	1.3	-6.7 1.2
10	1114	BD + 26 305a	01:50:02.08	+27:05:56.6	10.36	1.08	1.08	1.18	-3.6	1.8	-5.0 1.8
11	1749		01:50:18.14	+27:04:28.9	12.34	0.56	0.27	0.78	11.9	2.3	-5.5 2.3
12	272		01:50:23.97	+27:02:23.1	12.81	0.65	0.27	0.86	1.9	1.3	-8.4 2.9

Note: star # 1 was saturated in our photometry, and magnitude and color have been taken from Tycho-2. Stars # 7 and # 12 have been taken from UCAC2 catalog.




Fig. 10. Photometry and proper motions in the region of NGC 2664. a) Color-Magnitude diagram. Open squares are Tycho 2 stars having both proper motion and radial velocity measurements (see panel b) and Table 7), while filled triangles are stars having only proper motion measurements. b) Vector point plot of Tycho 2 proper motions and proper motion errors. Symbols are as in panel a). As for the identification, we are plotting here Tycho 2 numbering. See text for more details.

available are plotted with filled triangles. The remaining stars are mainly plotted with open circles.

It is evident that these stars do not share a common mean tangential motion. In fact individual proper motion vectors range from about 5 to 63 mas yr^{-1} , while the typical uncertainties of the proper motions is about 2 mas yr^{-1} per component. This spread in tangential motion is seen to be in agreement with that shown by field stars in

the surrounding of NGC 5385. We are therefore inclined to suggest that the stars belonging to NGC 5385 simply are part of the local Galactic field.

4.2. NGC 2664

The vector point diagram for all the candidate member stars having proper motion measurements in the field of

Table 6. Spectroscopic results for NGC 5385

ID	TYC 4558-	Date of Observation	S/N	Resolution	Julian Date	Rad. vel [km /s]	Sp. Type	[Fe/H] [dex]	($m - M$) _v [mag]
1	1641	12/07/2002	120	20000	2452467.4011	10:5 1:2	F 6 V	+ 0.2 0.3	7.49
		15/02/2003		20000	2452685.6788	7:5 1:0			
		12/05/2003		20000	2452772.3750	6:9 0:3			
2	1385	12/07/2002	110	20000	2452468.3527	61:8 2:3	F 4 V	+ 0.4 0.3	8.07
		15/02/2003		20000	2452685.5271	62:5 1:0			
		12/05/2003		20000	2452772.4526	62:4 0:3			
3	457	12/07/2002	120	20000	2452468.4014	1:9 1:3	F 8 V	+ 0.9 0.4	7.19
		15/02/2003		20000	2452685.6436	1:1 1:0			
		12/05/2003		20000	2452772.5267	1:5 0:3			
4	399	12/07/2002	120	20000	2452467.4495	2:5 1:8	G 2 V	+ 0.4 0.3	6.51
		15/02/2003		20000	2452685.5377	1:1 1:0			
		12/05/2003		20000	2452772.4272	4:9 0:3			
5	1797	12/07/2002	120	20000	2452468.4493	12:0 0:7	G 6 IV	+ 0.3 0.3	8.17
		15/02/2003		20000	2452685.6092	11:3 1:0			
		12/05/2003		20000	2452772.5043	12:3 0:3			
6	1873	12/07/2002	90	20000	2452469.4313	38:1 0:7	G 1 V	+ 0.5 0.4	7.08
		15/02/2003		20000	2452685.7243	29:4 1:0			
		12/05/2003		20000	2452772.4789	28:7 0:3			
7	217	12/07/2002	90	20000	2452469.3571	3:8 0:7	K 4 V	+ 0.3 0.3	4.87
		15/02/2003		20000	2452685.4985	4:0 1:0			
		12/05/2003		20000	2452772.4015	3:7 0:2			

NGC 2664 (see Fig. 9 for the identification) from the Tycho-2 catalogue is shown in Fig. 10, together with the CMD obtained from our photometry. Also in this case, there is a group of stars sharply detached from the bulk of the stars in the field. These are 4 stars, and are plotted with squares in both the panels.

It is evident that these stars do not share a common mean tangential motion. In fact individual proper motion vectors range from about 4 to 26 m as yr^{-1} , while the typical uncertainties of the proper motions is about 2 m as yr^{-1} per component. Only, we note that stars # 2354 and # 2400 have quite a similar (within the errors) tangential motion, and might constitute a physical pair. The spread in tangential motion is seen also in this case in agreement with that shown by field stars in the surrounding of NGC 2664.

4.3. Collinder 21

The CMD and the vector point diagram for all the candidate member stars having proper motion measurements in the field of Collinder 21 (see Fig. 11 for the identification) from the Tycho-2 catalogue are shown in Fig. 12 panels. 12 bright stars, in the magnitude range $8 < V < 13$, define the aggregate and are plotted in both panels as

open squares. Proper motions are also available for 3 dimmer stars (solid triangles) which bridge (see the CMD in panel a) the brightest group to the bulk of the stars in the field. However these 3 stars have significantly diverse proper motion components (see Table 5). Coming back to the 12 brightest stars, their proper motion vectors range from about 3 to 68 m as yr^{-1} , while the typical uncertainties of the proper motions is about 2 m as yr^{-1} per component. It is evident that these stars do not share a common mean tangential motion. There are two clear pairs. One is the well known visual binary system BD + 26 305ab (stars # 866 and # 1114), and the other is composed by the stars # 1472 and # 272, which however deserve further investigation. Apart from these two pairs, the bulk of the stars exhibit a disordered motion.

5. Spectroscopic analysis

Multi-epoch spectra have been acquired for all the candidate members (23 stars in total for a grand total of 54 spectra) in the aggregates under investigations. The results of the spectroscopic survey are listed in Tables 6 to 8.

Table 7. Spectroscopic results for NGC 2664

ID	TYC 816-	Date of Observation	S/N	Resolution	Julian Date	Rad. vel [km /s]	Sp. Type	[Fe/H] [dex]	(m -M) _v [mag]
1	2354	15/01/2003	160	20000	2452655.5065	26:0 2:0	F5 IV-V	+0.3 0.3	7.70
		12/05/2003		20000	2452772.3168	27:1 0:3			
2	1826	15/01/2003	130	20000	2452656.4174	13:5 1:5			
		06/12/2003		20000	2452979.5724	54:4 3:5			
3	1890	15/01/2003	140	20000	2452655.5844	14:4 1:5	K 0 III	+0.6 0.4	9.79
		12/05/2003		20000	2452772.3446	12:7 0:3			
4	2400	15/01/2003	110	20000	2452656.5643	23:0 1:5	K 1 III	+0.1 0.3	10.41
		06/12/2003		3600	2452979.6014	20:5 4:1			




Fig. 6. Histogram of the surface density of the stars shown in Fig. 5, measured by counts in 0.5 wide annuli centered on $\alpha = 01^{\text{h}}50^{\text{m}}11^{\text{s}}$, $\delta = +27^{\circ}04'26''$. Dots with error bars indicate the mean surface density of stars in a surrounding field centered on $\alpha = 01^{\text{h}}54^{\text{m}}10^{\text{s}}$, $\delta = +27^{\circ}44'00''$, and the expected 1- σ range of statistical fluctuations around this mean for each bin.

5.1. NGC 5385

For this object we provide 3 epochs Echelle spectra for seven bright stars, which turn out to be all dwarfs (see




Fig. 7. Identification map ($8^{\text{h}}:14$ – $8^{\text{h}}:14$), taken from DSS-2) of member candidates of NGC 5385 having proper motion measurements from Tycho 2. North is up, East on the left. The labels give the star numbers assigned in the Tycho-2 catalog (in the sense TYC 4558 < ... >) Data for these stars are given in Tables 3 and 6.

Table 6), but for star # 1797, which seems to be a subgiant. By looking at this table results we find that 2 stars are probably unresolved binary stars (# 1641 and # 1873), whereas all the other stars do not exhibit significant variations in their radial velocity. The studied stars have radial velocity ranging from -62 km/s to $+30$ km/s, and they differ one from the other, thus suggesting that NGC 5385 is not a physical aggregate. There might be some common motion pairs in this sample. Stars # 399 and # 1797 (see

Table 8. Spectroscopic results for Collinder 21

ID	TYC 1759-	Date of Observation	S/N	Resolution	Julian Date	Rad. vel [km /s]	Sp. Type	(m -M) _v [mag]
1	450	04/07/2002	530	20000	2452468.5756	3.2 0.7	F 9 V	3.90
		20/09/2003	300	3600	2452902.5724	7.4 5.5		
		05/12/2003		3600	2452979.1951	42.4 4.8		
2	1472	20/09/2003	360	3600	2452902.5906	5.3 6.2	G 0 V	5.35
		05/12/2003		3600	2452979.2083	11.7 5.7		
3	1025	20/09/2003	330	3600	2452902.6222	55.2 7.8	M 8 III	10.84
		05/12/2003		3600	2452979.4393	68.3 11.4		
4	468	09/10/2003	210	3600	2452922.4483	25.5 4.4	K 2 III	10.52
		05/12/2003		3600	2452979.2983	15.9 10.3		
5	994	09/10/2003	230	3600	2452922.4725	46.9 7.3	G 8 V	5.67
		05/12/2003		3600	2452979.2378	37.5 8.0		
6	462	09/10/2003	210	3600	2452922.5142	42.9 7.4	G 3 V	6.32
		05/12/2003		3600	2452979.3997	24.8 6.4		
7	1333	05/12/2003	190	3600	2452979.5238	48.5 4.6	K 2 V	5.31
		04/01/2004		3600	2453009.3715	57.0 5.2		
8	266	20/09/2003	230	3600	2452902.4968	50.7 4.6	G 8 III	9.86
		04/01/2004		3600	2453009.3979	40.4 5.2		
9	866	20/09/2003	310	3600	2452902.4559	3.5 5.7	G 5 III	9.49
		04/01/2004		3600	2453009.4311	17.2 7.9		
10	1114	20/09/2003	270	3600	2452902.5238	48.5 4.6	K 1 III	9.36
		04/01/2004		3600	2453010.3701	5.2 5.7		
11	1749	20/09/2003	150	3600	2452980.2946	24.8 7.2	F 8 V	8.14
		04/01/2004		3600	2453010.3303	23.0 7.1		
12	272	20/09/2003	120	3600	2452980.3317	33.9 7.3	G 0 V	8.36
		04/01/2004		3600	2453010.2885	41.2 8.3		

Table 3) have marginally consistent proper motion components, and stars # 1641 and # 217 as well. The first couple has also compatible spectral type (see Table 6). Radial velocities however (see Table 6) contradict this hypothesis. Moreover both pair members are located in very different position (see Fig. 7).

We derived stellar distances from the Sun basing on the spectral classification reported in Table 6. The relation between spectral type, luminosity class and absolute magnitude has been taken from the Michigan Spectral Catalogue Project.³ Distance moduli range from 4.87 to 8.17 mag, which turn into distances going from 94 to 430 pc. Schlegel et al (1998) extinction maps provide for NGC 5385 region a reddening of 0.04 mag, which does not significantly alter the derived distances. The significant spread of stars distances rules out the possibility they form a physical group. Metal abundances were derived using the MOOG code

(freely distributed by Chris Sneden, University of Texas, Austin) as described in Carraro et al. (2004). The S/N ratio and the limited resolution prevented us from deriving very precise metallicity estimates, and therefore the value reported in Table 6 must be taken as an indication. We find that within the uncertainties the bulk of the stars possess solar iron abundance, but for star # 457, which exhibits a super solar iron abundance. Within 3 the metallicity distribution looks homogeneous (0.4 - 0.2), although we believe that more precise estimates of the metallicity would probably reveal some spread, which is expected for a random group of stars.

5.2. NGC 2664

For this object we provide 2 epochs Echelle spectra. From the analysis of Table 7 results we find that one star is a spectroscopic binary (# 1826), and due to blending prob-

³ <http://www.astro.lsa.umich.edu/users/hdproj/mosa/absp/absp.html>




Fig. 9. Identification map ($8^{\circ}:14' - 8^{\circ}:14'$, taken from DSS-2) for member candidates of NGC 2664 having proper motion measurements from Tycho 2. North is up, East on the left. The labels give the star numbers assigned in the Tycho-2 catalog (in the sense TYC 816 < ::: >). Data for these stars are given in Tables 4 and 7.

lem s we were not able to derive either stars spectral type or metallicity. The remaining three stars do not exhibit significant variations in their radial velocity. Star # 2354 is a dwarf, while stars # 1890 and # 2400 are giants. The studied stars have radial velocity ranging from +12 km/s to +30 km/s, and they basically differ one from the other, thus suggesting that NGC 2664 is not a physical aggregate. The proper motion pair (# 2354 and # 2400) seem to be actually a pair, since the radial velocities of the two stars do not differ very much (see Table 7).

As for NGC 5385, we computed spectral type based distances, which are listed in the last column of Table 7. The distance moduli range from 7.70 to 10.41 mag., which implies distances ranging from 350 to 1207 pc, which slight corrections due to the interstellar reddening, which in the direction of NGC 2664 amounts to 0.02 mag (Schlegel et al. 1998). The four stars which define the group cannot be considered physically bound. Finally, the possible related stars # 2354 and # 2400 lie too apart to be considered as a physical pair.

In the same table we list estimates of the metal content of three stars, which turns out to be around solar or slightly supersolar for all the stars within the errors.

5.3. Collinder 21

For this object we provide 2 epochs medium resolution spectra for all stars but for # 450, for which we add an Echelle spectrum. 5 stars turn out to be giants, and all the other are dwarfs. Though in this case the errors are much larger, from the analysis of Table 8 results we confirm that this star (# 450) is a binary. Large velocity variations are also shown by the couple # 866-# 1114 (BD + 26 305a,b), which however we are inclined to ascribe to their nature of visual binary. All the other stars show deviating radial velocity, roughly ranging from -20 km/s to +60 km/s, thus suggesting that this circlet is not a physical aggregate. The possible proper motion pair (# 1472-# 272) is not confirmed by radial velocity measurements, and, on top of that, these two stars are located in quite different places (see Fig. 11).

As for the previous objects we have derived stars individual distances based on spectral classification. The distance moduli range from 3.9 to 10.84 mag, which means a distance between 60 and 1470 pc. The reddening in this direction is 0.07 mag (Schlegel et al. 1998), and therefore these estimates do not change significantly due to it. The distance spread is too huge to be consistent with Collinder 21 being a physical aggregate.

Unfortunately, in this case we do not have sufficiently high resolution spectra to derive reasonable estimates of the metal abundance.

6. Discussion and Conclusions

In this paper we have analysed the possibility that NGC 5285, NGC 2664 and Collinder 21 might be POCRs. Our study however yields negative answers for all of them. In fact, star counts and the appearance of the CMDs confirm that we are facing significant stellar overdensities above the general Galactic field.

However, the close scrutiny of the kinematic data (proper motion and radial velocity) provide us with a completely diverse picture. All the agglomerates are not physical clusters since they show very different velocity components, and therefore they force us to consider them simply as projective chance alignment of unrelated field stars.

We found out a number of possible unresolved binary stars or common motion pairs which might deserve further investigation.

The probable binary fraction we found ranges from about 25% in NGC 2664 and NGC 5385 to 18% in Collinder 21, and it is not much different from the typical Galactic disk field binary percentage.

Our findings confirm previous suggestions (Odenkirchen & Soubiran 2002) that an overdensity of stars does not necessarily imply the existence of a physical ensemble, and cast some doubts on the POCRs list provided by Bica et al. (2001), which is purely based on star counts. Nevertheless, work is in progress to define a more efficient OCRs




Fig. 12. Photometry and proper motions in the region of Collinder 21. a) Color-Magnitude diagram. Open squares are Tycho 2 stars having both proper motion and radial velocity measurements (see panel b) and Table 8), while filled triangles are stars having only proper motion measurements. b) Vector point plot of Tycho 2 proper motions and proper motion errors. Symbols are as in panel a) As for the identification, we are plotting here Tycho 2 numbering. See text for more details.

nding criterion (de la Fuente Marcos, in preparation) and to probe the real nature of other POCRs candidates.

Acknowledgements. The entire Asiago technical staff is deeply acknowledged for the kind night assistance over the whole duration of this project. GC thanks Brian A. Skiff for providing continuous and very useful comments. This study made use of Simbad and WEBDA.

References

- Bassino, L.P., Walchhausen, N. & Martnez, R.E. 2000, *A & A*, 355, 138
- Baum e, G., Villanova, S., Carraro, G., 2003, *A & A* 407, 527
- Baumgardt, H. 1998, *A & A*, 340, 402
- Bica, E., Santiago, B.X., Dutra, C.M., et al. 2001, *A & A*, 366, 827
- Carraro, G. 2000, *A & A*, 357, 145
- Carraro, G. 2002, *A & A*, 385, 471
- Carraro, G., Ng, Y.K., Portinari, L. 1998, *MNRAS*, 246, 1045
- Carraro, G., Bresolin, F., Villanova, S., et al. 2004, *AJ*, in press (astro-ph/0406679)
- de la Fuente Marcos, R. 1997, *A & A*, 322, 764
- de la Fuente Marcos, R. 1998, *A & A*, 333, L27
- Desidera, S., Fantinel, D., Giron, E. 2001, *AFOCUS USER MANUAL*
- Dias W.S., Alessi B.S., Moitinho A. & Lepine J.R.D., 2002, *A & A*, 389, 871
- Halbwachs, J.L., Mayor, M., Udry, S., Arenou, F., 2003, *A & A* 397, 159
- Høg, E., Fabricius, C., Makarov, V.V., et al. 2000, *A & A*, 355, L27
- Landolt, A.J. 1992, *AJ*, 104, 340
- Monet D.G., Levine S.E., Casian B., et al. 2003, *AJ*, 125, 984
- Odenkirchen, M., Soubiran, C. 2002, *A & A*, 383, 163
- Patat, F., Carraro, G. 2001, *MNRAS*, 325, 1591
- Pavani, D.B., Bica, E., Dutra, C.M., et al. 2001, *A & A*, 374, 554
- Schlegel, D.J., Finkbeiner, D.P., Davis, M. 1998, *ApJ*, 500, 525
- Villanova, S. 2003a, Master Thesis, Padova University
- Villanova, S., Carraro, G., de la Fuente Marcos, R. 2003b, in "Milky Way Surveys: Structure and Evolution of our Galaxy", proceedings of the 5th Boston University Astrophysics Conference, in press
- Villanova, S., Baum e, G., Carraro, G. & Geminale, A. 2004, *A & A* 419, 149
- Zacharias, N., Urban, S.E., Zacharias, M.I., et al. 2003, *AJ*, in preparation




Fig. 11. Identification map ($8^{\circ}:14 \ 8^{\circ}:14$, taken from DSS-2) for member candidates of Collinder 21 having proper motions measurements from Tycho 2. North is up, East on the left. The labels give the star numbers assigned in the Tycho-2 catalog (in the sense TYC 1759 < ::: >). Data for these stars are given in Tables 5 and 8.