

H I R O C S — a galaxy cluster survey at high redshifts

S. Falter¹, H.-J. Roser¹, H. Hippelein¹, C. Wolf² and E. Bell¹

¹Max Planck Institute for Astronomy, Königstuhl 17, D-69117 Heidelberg, Germany

²Dep. of Physics, Denys Wilkinson Bldg., Univ. of Oxford, Keble Road, Oxford, OX1 3RH, UK

Abstract. Clusters of galaxies are the largest bound gravitational systems in the universe. They outline the large scale structure and thus are able to test predictions of cosmological models. Detailed studies of galaxy populations in clusters at high z offer insights into galaxy evolution as a function of look-back time. Unfortunately, large homogeneous samples of galaxy clusters only exist at redshifts < 0.5 . There are around 1000 candidate clusters with photometric or otherwise estimated redshifts above 0.5, but spectroscopically confirmed clusters are still scarce in this redshift regime. Some clusters with $z > 1$ have recently been identified from X-ray surveys.

1. Survey outline

H I R O C S (= Heidelberg InFRared/Optical Cluster Survey) intends to homogeneously select candidate clusters with redshifts up to 1.5. Our survey is part of a large programme (M A N O S = M P I A for Astronomy Near-infrared Optical Surveys) exploring the evolution of galaxies. To reach out to redshifts of 1.5 we are observing in the infrared (H) supplemented by four optical (B, R, i, z) bands. H I R O C S will cover 11 fields in four elds, three stripes with 3 fields and one with 2 fields (see Table 1). This ensures a sufficient coverage of large scale structure features in the foreseen redshift range. Furthermore, predictions for the number of galaxy clusters above unity from Bartelmann & White (2002) for the SDSS show that our dedicated survey area should yield a statistically significant cluster sample. They predict more than 5 clusters per field with $z > 1$.

With one exception (CADIS-16h field) we have equatorial fields with few bright stars enabling follow-up studies from both hemispheres with the VLT and the LBT (available in the course of 2004). The right ascension of the fields is distributed such that observations are possible during the whole year. Sufficiently low and uniform galactic extinction over the field area is another requirement.

The MUNICS and CADIS fields provide extensive information for fixing the selection function of the reduced H I R O C S filter set. In these fields we have multi-color data sets as well as a large number of spectra for QSOs, galaxies and stars. We chose the COSMOS-10h field because the COSMOS survey will obtain high-resolution images from HST. The H I R O C S-22h field is a new field.

Our survey aims at a limiting magnitude of $L^2 + 1$ for an elliptical galaxy at $z = 1.5$ (5'). Table 2 shows the limiting magnitudes and the required exposure times.

The multi-colour-classification scheme developed for the CADIS and COMBO-17 surveys (Wolf, Eisenheimer & Roser 2001) will classify all objects and assign photometric redshifts. However, the accuracy of the redshifts will be not as accurate though sufficient for cluster detections.

Field name	RA (J2000)	DEC (J2000)	Size	E(B-V)
MUNICS-S2F1	03 ^h 06 ^m 41 ^s .0	00 01 ^o 12 ⁰⁰	3 1	0.080 mag
COSMOS-10h	10 ^h 00 ^m 28 ^s .0	02 12 ^o 21 ⁰⁰	2 1	0.018 mag
CADIS-16h	16 ^h 24 ^m 32 ^s .0	55 44 ^o 32 ⁰⁰	3 1	0.006 mag
HIROCS-22h	22 ^h 00 ^m 00 ^s .0	02 14 ^o 20 ⁰⁰	3 1	0.065 mag

Table 1. HIROCS fields coordinates, sizes and galactic extinction from Schlegel, Finkbeiner & Davis (1998).

	$m_k^?_{corr}$	E-corr	$m^?$	obsLim	t	S/N	t/
B _J	29.0	-5.975	23.1	24.1	1500	19.0	6 ksec
R _J	27.3	-2.878	24.4	25.4	5000	5.3	20 ksec
i	26.7	-2.291	24.4	25.4			20 ksec
z	25.0	-2.480	22.5	23.5	6000	5.2	24 ksec
H	21.2	-1.348	19.9	20.9	3000	5.0	48 ksec

Table 2. Limiting M magnitudes (col. obsLim) and exposure times to cover 1 for the HIROCS fields. For the calculation a standard E galaxy SED is scaled to produce an absolute magnitude of an L²-galaxy in the Gunn r-band of $M_r = -21.75$ mag (Garilli, Macagnani & Andreon 1997, Paolillo, Andreon, Longo, et al. 2001). This spectrum is k-corrected ($z = 1.5$) and corrected for evolutionary effects after Poggianti (1997). The observations are dithered and split into short exposures. The summed integration time per pointing is shown in the column t. The last column incorporates the field of view of the detector and specifies the required time to cover 1. We assume that one hour of exposure time yields a net science exposure time of 3 ksec. Cosmology: $H = 65$, $M = 0.3$, $\Omega = 0.7$.

2. Cluster finding

We will employ a 3-dimensional cluster finding algorithm based on finding overdensities in position on the sky and in redshift space. According to its redshift z_0 derived from the multi-colour-classification the projected Abell radius R_A for each object is calculated and the number of objects N_{obj} within R_A counted. Only objects with velocity differences below a given threshold are added to N_{obj} :

$$\frac{v}{c} = \frac{(1+z_1)^2 - (1+z_0)^2}{(1+z_1)^2 + (1+z_0)^2} \approx 6 \frac{v}{c_{lim}} \quad (2.1)$$

Candidate galaxy clusters are found as excess numbers in N_{obj} which exceed values expected for a random object distribution. From these excess numbers we define an overdensity parameter.

A first test application to the object catalog of the Chandra Deep-Field South (CDFS) observed by COMBO-17 (field size $31^o \times 30^o$) reveals galaxy overdensities at $z = 0.5, 0.7$ and 1.1 (Figure 1).

In the past several other techniques for identifying clusters of galaxies have been developed. We intend to compare the performance of our algorithm with the Voronoi tessellation (Ramella, Boschin, Fadda & Nonino 2001), the matched-filter and the friends-of-friends algorithm. Because of the imaging in *ve* filters we will also be able to apply the red sequence approach established for the Toronto Red-sequence Cluster Survey (Gladstone & Yee 2000) for various colour combinations.




Figure 1. Left: z-histogram shows object enhancements at $z = 0.5, 0.7, 1.1$. Objects are selected by overdensity; Right: Object distribution in the CDFS. The condensations of points indicate candidate galaxy clusters (coordinates in arbitrary units).

3. Project status

Since August 2003 we are collecting the survey data at the Calar Alto 3.5m telescope and since December 2002 with the Wide-Field-Imager (WFI) at the ESO 2.2m telescope. A first impression of the quality of the IR data gives a pipeline-reduced mosaic of 12 H-band images in the HROCS-22h field observed with the NIR wide-field camera OMEGA 2000 on this webpage:

http://www.caha.es/newletter/new_s04a/OMEGA2000_mosaic.pdf

It covers almost $3/4$; the PSF is 0.008 . Each frame is composed of 25 single exposures with 1 minute exposure time, the limit is 19.9 mag (5 σ). The holes around bright objects are due to over-corrected background, as the pipeline, whose result is shown, uses only a local sky determination.

References

- Bartelmann, M. & White, S.D.M. 2002 A & A 388, 732{740.
- Garilli, B., Accagni, D. & Andreon S. 1997 A & A 342, 408{416.
- Gadders, M.D. & Yee, H.K.C. 2000 AJ 120, 2148{2162.
- Paolillo, D.S.F., Andreon, S., Longo, G., et al. 2001 A & A 367, 59{71.
- Poggianti, B.M. 1997 A & A 122, 399{407.
- Ramella, M., Boschin, W., Fadda, D. & Nonino, M. 2001 A & A 368, 776{786.
- Schlegel, D.J., Finkbeiner, D.P. & Davis, M. 1998 ApJ 500, 525S.
- Wolf, C., Eisenheimer, K. & Roser, H.-J. 2001 A & A 365, 660{680.