# Intracluster P lanetary N ebulae as P robes of Intracluster Starlight

John Feldm eier

Case W estern Reserve University, C leveland OH, 44106, USA

A bstract. I review the progress in research on Intracluster P lanetary Nebulae (IPN). Hundreds of IPN candidates have now been found in the Virgo and Formax galaxy clusters, and searches of two nearby galaxy groups have made. From the results thus far, approximately 10{20% of all stars in Virgo and Formax are in an intracluster component, but there are few such stars in galaxy groups. From the spatial distribution of IPN, it appears that the intracluster stars are clustered, in agreement with tidal-stripping scenarios. In Virgo, the IPN have a large line-of-sight depth, which im plies that the bulk of intracluster stars in this cluster derive from late-type galaxies and dwarfs. I also discuss other in portant developments in IPN research such as the detection of intracluster H II regions, a possible detection of IPN in the Com a Cluster, and future observational and theoretical developments.

#### 1 Introduction

W hy do planetary nebulae (PN), which are a late phase of stellar evolution, m ake excellent tracers of intracluster stars, stars between the galaxies inside of a galaxy cluster? An example of the utility of PN under sim ilar circum stances can be seen in Figure 1. On the left is a standard broad-band im age of the fam ous interacting galaxy pair M 51. A lthough there are num erous signs of the interaction at high surface brightnesses, there are m any low er surface-brightness features [5] that are barely visible in this representation.

However, the distribution of PN found in M 51 [20] displayed on the right give additional information on this interacting system. There is a clear tidal tail structure to the west of the secondary galaxy, and there are also signs of an extension of the spiralarm to the south. The lack of PN in the central regions of the galaxies are due to crowding e ects. W hat this illustration rem inds us is that planetary nebulae trace stellar lum inosity, not stellar surface brightness. Because PN are an end-phase of stellar evolution form ost stars between 1{8 solarm asses, the distribution of planetary nebulae closely follows the distribution of stars in galaxies [9]. If there is enough lum inosity in a stellar population for su cient num bers of PN to be present, then we can detect and study that population, regardless of its surface brightness. Intracluster stars, which are believed to be quite lum inous (anywhere between 10% and 70% of the cluster's total stellar lum inosity; [51,41]), but have a very low surface brightness (less than 1% of the night sky in the optical bands; [30,22]) are therefore an ideal target for PN searches. F ig.1. Two in ages of the interacting galaxy pair M 51.0 n the left is a broad-band in age taken with the KPNO 4-m telescope showing the classical high-surface brightness features of this interacting system. On the right shows the location of the 64 planetary nebula candidates found by [20]. Note the clear tidal tail structures that are extrem ely di cult to detect in the broad-band in age, but can be seen easily in the PN distribution.

PN have additional features that make them useful probes of intracluster starlight. Because PN are emission-line objects, any bright PN that can be detected photom etrically can also be observed spectroscopically to high precision ( $12 \text{ km s}^{-1}$ ). In the case of M 51, spectroscopic follow -up of the PN [18] found that the western tidal tail consists of two discrete structures that overlap in projection, one from each galaxy. W ith the high velocity precision of the PN observations, the separation of the two kinematic structures was trivial. A l-though I will not focus on the dynam ical aspects of IPN /PN in this review (see reviews by A maboldi, D ouglas, G erhard, & P eng this conference), it is in portant to state that IPN are the easiest (and perhaps, the only) way to obtain dynam ical inform ation about the intracluster light.

Finally, through the [O III] 5007 P lanetary N ebulae Lum inosity Function (PNLF), extragalactic PN m ake excellent distance indicators (see C iardullo, this conference). The distinctive PNLF can be used in the intracluster environment to gain information on the line-of-sight distribution of the intracluster stars.

W hy study intracluster stars in the st place? Once a curiosity proposed by Zwicky ([62]), intracluster light is potentially of great interest to studies of galaxy and galaxy cluster evolution. The dynam ical evolution of cluster galaxies is complex, and involves the poorly understood processes of cluster accretion and tidal stripping [14]. The intracluster light provides a unique way to study these echanism s.M odern num erical simulations show that the intracluster light [15,A4,55,59,39] has a complex structure, and can be used to gain inform ation on the dynam ical evolution of galaxies and galaxy clusters.

## 2 H istory of Intracluster P lanetary N ebulae R esearch

The history of IPN research begins over a decade ago with the stPN survey of the Virgo cluster [32] (hereafter JCF). In this survey of elliptical galaxies, JCF found 11 PN that were much brighter than the expected [0 III] 5007 PN LF cut-o magnitude. JCF attempted to explain these \overlum inous" PN with a number of hypotheses, but none was entirely satisfactory.

The next step involved spectroscopic follow -up of objects from the JCF survey. During a radial velocity survey of PN in the V irgo elliptical galaxy M 86, [1] found that 16 of the 19 detected PN velocities were consistent with the galaxy's mean velocity ( $v_{radial} = -227$  km s<sup>-1</sup>). The other three planetaries had mean radial velocities of 1600 km s<sup>-1</sup>, more consistent with the V irgo cluster's mean velocity. [1] argued convincingly that these objects were intracluster

planetary nebulae, and it is here that the term rst enters the literature. The \overlum inous" PN candidates were thus naturally explained as a population of intracluster stars in front of the target galaxies. A lm ost simultaneously, the rst search for IPN candidates in the Formax cluster was published [57], and m ore detections of IPN candidates in Virgo quickly followed [36,10,21].

However, a surprise was in the works. Spectroscopic follow-up of the IPN candidates revealed that som e were not IPN, but instead background em ission-line objects with extrem ely high equivalent width [27,34]. This was unexpected, because previous deep em ission-line surveys had found very few such objects [48], though m any have now been detected at fainter m agnitudes [50]. The m ost likely source of the contam ination was found to be Lym an- galaxies at redshifts  $3.12\{3.14, where the Lym an-$  1215 line has been redshifted into the [0 III] 5007 Iters used in IPN searches. However other types of contam inating objects m ay also exist [56,46].

A lthough these contam inants caused som e constemation at st, a num ber of lines of evidence quickly showed that the majority of IPN candidates are in fact, actual IPN.0 bservations of blank control elds with identical search procedures as the PN surveys [11,7] have found that the contam ination fraction is signi cant, but was less than the observed IPN surface density. The surface densities found correspond to a contam ination rate of 20% in the Virgo cluster and 50% in Formax (Formax is more distant than Virgo, so its PNLF is fainter, and therefore further down the contam inating sources lum inosity function). There are still signi cant uncertainties in the background density due to large-scale structure, and to the sm all num bers of contam inating objects found thus far. How ever, deeper and broader control elds are forth coming. Spectroscopic follow -up of IPN candidates [27,11,3], and A maboldi et al., this conference, clearly show large numbers of IPN candidates have the expected [0 III] 5007 and 4959 em ission lines, with a contam ination rate sim ilar to the blank eld surveys. Finally, there is independent evidence of individual intracluster stars in V irgo from observations of individual Intracluster R ed G iant stars (IRG s; [26,17]) from the Hubble Space Telescope.

It is worth reiterating that all deep [0 III] 5007 em ission-line searches will have such contam ination at fainter m agnitudes. In fact, such objects have already been seen in conventionalPN surveys ([37,18] and R om anow sky, private communication). Extragalactic PN researchers should keep this into account. The brightest of such objects have been detected with a m  $_{5007}$  m agnitude of 25.5 [27], though the lum inosity function of the contam inants is still poorly know n.

Currently, with the widespread use of mosaic CCD detectors, and autom ated detection methods derived from DAOPHOT and SExtractor, over a hundred IPN candidates can be found in a single telescope run [47,23,3,25]. IPN candidates are easily identi ed as stellar sources that appear in a deep [D III] 5007 in age, but com pletely disappear in an in age through a liter that does not contain the [D III] line. Currently, over 400 IPN candidates have been detected in the V irgo cluster, and over 100 IPN candidates have been found in the Formax cluster.

Fig.2. Im ages of the V irgo cluster (left), and the Formax cluster (right), with the various published IPN survey elds marked, as well as the two H ST IRG elds. Soon to come are additional elds from Aguerri et al. (2004; V irgo), C iardullo et al. (2004; Formax), Feldm eier et al. (2005; V irgo) and C iardullo et al. (2005; IRG V irgo)

Figure 2 sum marizes the status of the di erent surveys.Despite all of the e ort, to date, only a few percent of the total angular area of V irgo and Formax have been surveyed.Literally thousands of IPN wait to be discovered by 4-m eter class telescopes.

## 3 The Spatial D istribution of the Intracluster Light

If we want to understand the mechanism s that produce intracluster stars, it is useful to compare the intracluster stars' spatial distribution with that of the better-studied components of galaxy clusters: the cluster galaxies, the hot intracluster gas, and the invisible dark matter. Theoretical work predicts very di erent spatial distributions for the intracluster light depending on its production time. For example, ifm ost intracluster stars are rem oved early in a cluster's lifetime [38], the distribution of this di use component will be smooth and follow the cluster potential. How ever, if a signi cant portion of the stars are rem oved at late epochs via galaxy encounters and tidal stripping [51,42], then the intracluster light should be clumpy and have a non-relaxed appearance. In particular, the \harassment" models of [42,43] rst predicted that m any intracluster stars will exist in long ( 2 M pc) tidal tails, which m ay m aintain their structure for G yrs.

W ith regards to angular clustering, there is good evidence that the IPN follow a non-random distribution. [47,25] show evidence for clumps of IPN within their survey elds, and Aguerri et al. (2004) found a strong clustering signal using a standard two-point correlation function analysis (reported in [45]). The scale of the angular clustering ranges from one to ten arcm inutes, (corresponding 5 to 50 kpc assuming a mean Virgo distance of 15 M pc), depending on the eld studied. How ever, no tail-like features have yet been seen in the IPN distribution, although tidal arc features have been seen in broad-band im ages of other galaxy clusters [58,31,6].

W ith the use of the [D III] 5007 PNLF, we can also obtain line-of-sight inform ation on the intracluster stars, a property unique to IPN. A lthough a full analysis of the observed lum inosity functions has not yet been attempted, we can already learn something useful from the brightest IPN candidates in each

eld. G iven the empirical PNLF, there is a maximum absolute magnitude M, that a PN may have in the light of  $[D \ III]$ . If we assume that the brightest observed IPN candidate in each eld has this absolute magnitude, we can derive an upper limit on its distance, and therefore estimate the distance to the front edge of the V irgo intracluster population. This is plotted in F igure 3.As can be clearly seen, there is an o set between the IPN of subclusters A and B : this is

interpreted as due to the di ering distances of these two subclusters. However, the most revealing feature of the distribution of the IPN is in subclum p A. The upper limit distance for most of the elds is well in front of the cluster core. This is partially due to a selection e ect: it is easier to detect an IPN on the near side of V irgo than the far side. Nevertheless, the depth im plied from the m easurements is remarkable. If we take the data at face value, then the IPN distribution has a line-of-sight radius of over 4 M pc. If we com pare this radius to the classical radius of V irgo on the sky (six degrees, or 1.6 M pc), we nd that the V irgo cluster is more than 2.6 tim es as deep as it is wide. V irgo is now here near a spherical cluster: it contains considerable substructure, and is elongated signi cantly along our line of sight.

The great depth derived for the IPN distribution gives an important clue to the parent galaxies of the intracluster stars. Numerical simulations [43,16] show that the majority of intracluster stars are ejected into orbits similar to that of their parent galaxies. In V irgo, the cluster ellipticals are clustered within a radius of 2 M pc from the cluster core [32]. In contrast, the hydrogen de cient spirals found in the V irgo cluster have a radius of 4 M pc or larger [54], and the dwarf ellipticals have a depth of 6 M pc [33]. Therefore, it seems clear that a signi cant portion of V irgo's intracluster stars originate from late-type galaxies whose highly radial orbits take them in and out of the cluster core.

## 4 Converting IPN densities to Lum inosity Densities

In principle, determ ining the am ount of intracluster lum inosity from the observed numbers of IPN is straightforward. Theories of simple stellar populations [49] have shown that the bolom etric lum inosity-specil c stellar evolutionary ux of non-star-forming stellar populations should be 2 10 <sup>11</sup> stars-yr <sup>1</sup>-L <sup>1</sup>, (nearly) independent of population age or initial mass function. If the lifetime of the planetary nebula stage is 25;000 yr, and if the empirical PNLF is valid to 8 m ag below the PNLF cuto , then every stellar system should have 50 10 <sup>8</sup> PN-L <sup>1</sup>. A coording to the empirical PNLF, approximately one out of ten of these PNe will be within 2.5 m ag of M . Thus, under the above assumptions, most stellar populations should have <sub>2:5</sub> 50 10 <sup>9</sup> PN-L <sup>1</sup>. The observed number of IPN, coupled with the PNLF, can therefore be used to deduce the total lum inosity of the underlying stellar population.

In practice, there are a number of system atic e ects that must be accounted for before we can transform the numbers of IPN to a stellar lum inosity [25], which we brie y summarize here.

F irst, although stellar evolution theory originally predicted a constant  $_{2:5}$  value for all non star-forming populations, observations present a more com – plicated picture. [8] found that in a sample of 23 elliptical galaxies, lenticular galaxies, and spiral bulges, the observed value of  $_{2:5}$  never exceeded  $_{2:5} = 50 \ 10^{9} \text{ PN-L}^{-1}$  but was often signi cantly less, with higher lum inosity galaxies having system atically smaller values of  $_{2:5}$ . Since the amount of intracluster



F ig.3. A com parison of the upper lim it distances obtained from the intracluster planetary nebulae to direct distances to V irgo C luster galaxies. At the top are the upper lim it distances (denoted by the open squares) from IPN observations by O kam ura et al. (2002) and A maboldi et al. (2002). Below that are the distances derived from as the overlum inous IPN found in front of M 87 (C iardullo et al. 1998). These upper lim it distances are com pared to the PN LF distances of V irgo ellipticals (denoted by the open circles; Jacoby, C iardullo, & Ford 1990; C iardullo et al. 1998), and C epheid distances to spiral galaxies (denoted by the lled circles; P ierce et al. 1994; Saha et al. 1997; Freedm an et al. 2001). The subcluster of V irgo that each intracluster eld resides in is noted. N ote the great depth of IPN, com pared to that of the elliptical galaxies.

starlight derived is inversely proportional to the  $_{2:5}$  parameter, a large error in the amount of intracluster light can result if this is not accounted for.

By comparing the numbers of  $\mathbb{P}N$  in a eld surrounding a HST W FPC2 eld, with RGB and AGB star counts, [17] found a value of  $_{2:5} = 23^{+10}_{-12}$  10  $^{9}$  PN-

L  $^1$  for V irgo's intracluster population. This observational measurement is bolstered by the new theoretical work of Buzzoni & A maboldi (this conference), which show that the observational value for  $_{2:5}$  is well within the range for models of moderate lum inosity galaxies.

Second, as we have discussed previously, IPN surveys are not pristine: approxim ately 20% of V irgo IPN candidates, and 50% of Formax IPN candidates, are likely to be unrelated background sources. A lthough these objects are subtracted out statistically from all modern IPN surveys, the importance of eld-to-eld variations in the background due to large-scale structure is still unknown.

Finally, the IPN candidates of V irgo have a signi cant line-of-sight depth. Since the conversion between number of PN and lum inosity depends on the shape of the lum inosity function, this depth can change the am ount of intracluster light found from the data. M odels [21] indicate that the di erence between a singledistance m odel (the m ost conservative) and assum ing that the IPN are uniform ly distributed in a sphere of radius 3 M pc (the least conservative), changes the derived intracluster star lum inosity by up to a factor of three. Thus far, all IPN researchers have adopted a single distance m odel in order to be conservative, but this e ect is the least studied at this point.

A fier applying all of these corrections, the intracluster stellar fractions for V irgo vary between 10 and 20% [2,47,3,25], with the errors being dom inated by the system atic e ects. The IRG measurements [26,17] nd som ewhat less intracluster light ( $10\{15\%$ ), but the various results agree within the errors. How ever, it is important to note that each of these studies makes di errent assumptions concerning the calculation of intracluster lum inosity, the am ount of light bound to galaxies, and the contam ination of background sources. For Formax, there are only two measurements of the intracluster star fraction thus far from IPN. [57] report a fraction of up to 40%, but this is before the detection of contam inating sources, and is likely to be an overestim ate. C iardullo (2004, in prep), reports an approximate fraction of 20% , with sim ilar errors as the V irgo results. Regardless of the uncertainties, it is clear that a signi cant fraction of all stars in V irgo and Formax are in an intracluster component.

How does the IPN lum inosity density compare with well-known cluster properties such as radius, or projected galaxy density? [25] has compared these properties in Virgo, and has found little or no correlation. This may be due to Virgo's status as a dynamically young cluster, or due to a selection e ect (IPN researchers must observe where the galaxies are not, to avoid confusion with norm alextragalactic PN). More data will be needed to con im this result.

## 5 Intracluster H II R egions

A recent discovery stemming from  $\mathbb{IPN}$  searches is the detection of H II regions in the far halos of galaxies or in intracluster space. In the course of spectroscopic follow-up of a Virgo  $\mathbb{IPN}$  eld, [29] found an emission-line source that has the properties of a 400 solar mass, 3 M yr compact H II region, over 17 kpc away from the nearest Virgo galaxy (see also [35] for a similar, earlier, underFig.4. A comparison of detected intracluster star fractions from modern measurements, as a function of the velocity dispersion. Note the abrupt change from the cluster environment to the group environment.

appreciated example). There are now a number of other examples of intergalactic star form ation in V irgo and in other galaxy clusters and groups [60,53,52,13,19].

These discoveries have two in portant in plications. First, star form ation can occur at large distances from galaxies, and environments quite dierent than that normally studied. This has in plications for a number of elds of astrophysics, including the origin of metallicity in the intergalactic and intracluster medium, and the possible in situ origin of B stars in the Galactic hab. At this time, it does appear that intracluster star formation is only a small fraction of the total intracluster star production, but the exact am ount is uncertain. Second, these objects are another form of contam ination to pure IPN surveys. D iscouragingly, the only way to separate these objects from IPN is through deep spectroscopy. Lum inous PN have [D III] / H ratios of two or greater [12], where lum inous H II regions do not. From the surveys thus far, these objects appear to be relatively rare component of IPN surveys ( 3%), but more study is needed in this area.

## 6 Intra-group Starlight

A lthough the presence of intracluster starlight in clusters such as V irgo and Formax is now well established, the am ount of 'intra-group' starlight is still uncertain. Theoretical studies predict that ifm ost intracluster stars are rem oved by galaxy collisions [51,42], the fraction of intra-group stars, to storder, should be a sm ooth function of galaxy number density ( $L_{ICL} = N_{Gal}^2$ ). To test this hypothesis, two di erent groups have surveyed the nearby M 81 and Leo groups of galaxies [7,19], using sim ilar detection m ethods as the cluster searches. In both cases, no genuine intra-group PN were found. These non-detections strongly im – plies there is substantially less (4{15 times}) less intergalactic stars in groups than there are in clusters.

W hen com pared w ith otherm easurem ents of intracluster star fractions through m odem deep in aging [4,30,22,24], or through the detection of intracluster supernovae [28], an interesting pattern emerges, plotted in Figure 4. For galaxy clusters, the data is consistent w ith an approxim ate fraction of 20%, albeit w ith large error bars or intrinsic scatter. W hen we m ove to the group environm ent, the fraction abruptly drops, with no sign of any sm ooth decline. This im plies that there is som ething special about the cluster environm ent that prom otes intracluster star production. M ore data w ill be needed to con m and strengthen this result, especially IPN searches of additional galaxy groups.

## 7 IPN in the Com a Cluster?

At this conference, G enhand et al. announced the possible detection of at least one IPN candidate in the C om a C luster. If this candidate is con m ed, it will be the m ost distant individual PN ever detected, at a distance of over 100 M pc. This object was found using a technique of lling the entire focal plane of the 8-m Subaru telescope with a slit m ask through the appropriate [0 III] 5007

Iter, and looking for narrow -line em ission sources.D eep broad-band in aging has in plied that the C om a C lusterm ay have a very high intracluster star fraction, up to 50% [4], so it is plausible that a few IPN could be detected, despite the sm all area surveyed. The most exciting aspect of this observation is that it opens up a way to observe PN at much greater distances than previously thought possible, and makes it possible to place IPN density lim its in more distant galaxy clusters.

#### 8 The Future

Studies of IPN are not even a decade old, and much more work still needs to be done. The most crucial observations needed in the near term are spectroscopic follow-up of a large number of IPN candidates. This would allow us to better determ ine the contam ination rate from background galaxies, avoid any other intracluster H II regions, and most importantly, gain information on the kinematics of intracluster stars. More IPN imaging is needed in V irgo and Fornax at di ering radii and densities, in order to con im the lack of correlations found thus far. IPN surveys of additional galaxy groups are needed to determ ine the lack of intra-group stars in these systems, and to better characterize the steepness of the intracluster production \cli ."

However, the future of IPN research will involve close comparisons of IPN properties to other observations of extragalactic PN and galaxy clusters and close comparison to the modern numerical simulations now being undertaken. Some examples are given below, but it is almost certain that many more will be added in the next few years.

V illaver (this conference), presented the st num erical hydrodynam ical sim - ulations of IPN within a hot intracluster gaseous medium. The results im ply that IPN do survive in the intracluster medium, but the nebula may be spatially distorted.

[40] has just completed the rst phase of a deep CCD in aging survey of the Virgo cluster core using the techniques of ultra-deep surface photom etry. These new observations reach extrem ely faint surface brightnesses ( $_{\rm V}$  29:6) over large angular areas (2 square degrees). Comparing these deep in aging m aps with the IPN observations should be revealing, and allow us to better determ ine the reality of tidal features in both data sets, and to give clear targets for follow-up IPN observations.

HST + ACS broad-band observations are planned of an intracluster eld in Virgo by Summer 2005, with the goal of obtaining the rst color-magnitude diagram of the intracluster stars. By placing tighter constraints on the age and

#### 10 John Feldmeier

lum inosity of the intracluster population, we should be able to better constrain the  $_{2:5}$  parameter, and in prove the accuracy of the total amount of intracluster starlight.

F inally, with the advent of advanced num erical simulations of galaxy clusters, we will be able to compare the IPN data to models of intracluster starproduction. Due to the large angular size of nearby clusters, it will be many years before the majority of the clusters will be surveyed for IPN. Num erical simulations will hopefully allow us to survey more e ciently, and to obtain better quality inform ation on the properties of intracluster starlight.

I would like to thank the conference organizers for allowing me to give this review, and for running an excellent conference. I would also like to thank my collaborators for their years of ort on the science presented here. This work was supported in part by NSF grant AST 03-02030 and NASA grant NAG 5-9377.

#### References

- 1. A maboldi, M ., et al. 1996, Ap. J., 472, 145
- 2. A maboldi, M ., et al. 2002, A . J., 123, 760
- 3. A maboldi, M ., et al. 2003, A . J., 125, 514
- 4. Bernstein, G.M., et al. 1995, A.J., 110, 1507
- 5. Burkhead, M.S. 1978, Ap.J. Suppl., 38, 147
- 6. Calcaneo-Roldan, C., et al. 2000, M. N. R. A. S., 314, 324
- 7. Castro-Rodr guez, N., et al. 2003, Astr. Ap. 405, 803
- 8. Ciardullo, R. 1995, in IAU Highlights of Astronomy, 10, ed. I. Appenzeller (Dordrecht: K luwer), p. 507
- 9. Ciardullo, R., et al. 1989, Ap. J., 339, 53
- 10. Ciardullo, R., et al. 1998, Ap. J., 492, 62
- 11. Ciardullo, R., et al. 2002a, Ap. J., 566, 784
- 12. Ciardullo, R., et al. 2002b, Ap. J., 577, 31
- 13. Cortese, L., et al. 2004, A & A , 416, 119
- 14. D ressler, A. 1984, A R A & A ., 22, 185
- 15. Dubinski, J. 1998, Ap. J., 502, 141
- 16. Dubinski, J., Murali, C., & Ouyed, R. 2000, unpublished preprint
- 17. Durrell, P.R., et al. 2002, Ap.J., 570, 119
- 18. Durrell, P.R., et al. 2003, Ap.J., 582, 170
- 19. Durrell, P.R., et al. 2004, IAU Sym posium, 217, 90
- 20. Feldm eier, J. J., Ciardullo, R., & Jacoby, G. H. 1997, Ap. J., 479, 231
- 21. Feldmeier, J.J., Ciardullo, R., & Jacoby, G.H. 1998, Ap.J., 503, 109
- 22. Feldm eier, J. J., et al. 2002, Ap. J., 575, 779
- 23. Feldm eier, J. J., et al. 2003a, Ap. J. Suppl., 145, 65
- 24. Feldmeier, J.J., et al. 2004a, Ap.J., in press
- 25. Feldmeier, J.J., et al. 2004b, Ap.J., in press
- 26. Ferguson, H.C., Tanvir, N.R., & von Hippel, T. 1998, Nature, 391, 461
- 27. Freem an, K.C. et al. 2000, A SP Conf. Ser. 197: Dynam ics of Galaxies: from the Early Universe to the Present, ed.F.Com bes, G.A.M amon, & V.Charm and aris (San Francisco: A SP), 389.

- 28. Gal-Yam, A., et al. 2003, A J., 125, 1087
- 29. Gerhard, O., et al. 2002, Ap.J.L., 580, L121
- 30. Gonzalez, A.H., et al. 2000, Ap.J., 536, 561
- 31. G regg, M. D., & West, M. J. 1998, Nature, 396, 549
- 32. Jacoby, G.H., Ciardullo, R., & Ford, H.C. 1990, Ap.J., 356, 332
- 33. Jerjen, H., Binggeli, B. & Barazza, F.D. 2004, A.J., 127, 771
- 34. Kudritzki, R.-P., et al. 2000, Ap.J., 536, 19
- 35. Lee, H., Richer, M.G., & McCall, M.L.2000, Ap.JL., 530, L17
- 36. M endez, R.H., et al. 1997, Ap.J.L., 491, 23
- 37. M endez, R.H., 2001, Ap.J., 563, 135
- 38. Merritt, D. 1984, Ap. J., 276, 26
- 39. M ihos, J.C., et al. 2004a, in preparation
- 40. M ihos, J.C., et al. 2004b, in preparation
- 41. Miller, G E. 1983, Ap. J., 268, 495
- 42. M oore, B., et al., 1996, N ature, 379, 613
- 43. Moore, B., Lake, G. & Katz, N. 1998, Ap. J., 495, 139
- 44. M urante, G ., et al. 2004, Ap. J. L., 607, L83
- 45. Napolitano, N.R., et al. 2003, Ap.J., 594, 172
- 46. Norman, C., et al. 2002, Ap. J., 571, 218
- 47. 0 kam ura, S., et al. 2002, P.A.S.J., 54, 883
- 48. Pritchet, C.J. 1994, PASP., 106, 1052
- 49. A.Renzini, A.Buzzoni: In: Spectral Evolution of Galaxies, ed.by C.Chiosi, A.Renzini (Reidel, Dordrecht), p.195 (1986)
- 50. Rhoads, J.E., et al. 2000, Ap.J.L., 545, L85
- 51. Richstone D.O., & Malum uth, E.M. 1983, Ap. J., 268, 30
- 52. Ryan-W eber, E.V., et al. 2004, A.J., 127, 1431
- 53. Sakai, S., Kennicutt, R.C., van der Hulst, J.M., & Moss, C. 2002, Ap. J., 578, 842
- 54. Sanchis, T., et al. 2002, Ap J., 580, 164
- 55. Som m er-Larsen, J., R om eo, A D., Portinari, L.2004, M N R A S., subm itted, available as astro-ph/0403282
- 56. Stem, D., Bunker, A., Spinrad, H., & Dey, A. 2000, Ap. J., 537, 73
- 57. Theuns, T., & Warren, S.J. 1997, M N R A S., 284, L11
- 58. Trentham, N. & Mobasher, B. 1998, M. N. R. A. S., 293, 53
- 59. W illm an, B., et al. 2004, M N R A S., subm itted available as astro-ph/0405094
- 60. Xu, C., Sulentic, J.W., & Tu s, R. 1999, Ap. J., 512, 178
- 61. Yun, M S., Ho, P.T P., & Lo, K.Y. 1994, Nature, 372, 530
- 62. Zwicky, F. 1951, PA SP., 63, 61

This figure "fig1.jpg" is available in "jpg" format from:

http://arxiv.org/ps/astro-ph/0407625v1

This figure "fig2.jpg" is available in "jpg" format from:

http://arxiv.org/ps/astro-ph/0407625v1

This figure "fig4.jpg" is available in "jpg" format from:

http://arxiv.org/ps/astro-ph/0407625v1