# Im proving X M M {N ew ton E P IC pn data at low energies: m ethod and application to the Vela SN R

Konrad Dennerl<sup>1</sup>, Bernd Aschenbach<sup>1</sup>, Ulrich G.Briel<sup>1</sup>, Herm ann Brunner<sup>1</sup>, Vadim Burw itz<sup>1</sup>, Jakob Englhauser<sup>1,2</sup>, Michael J. Freyberg<sup>1,3</sup>, Frank Haberl<sup>1</sup>, Gisela Hartner<sup>1,3</sup>, Anatoli F. Iyudin<sup>1</sup>, Eckhard Kendziorra<sup>4</sup>, Norbert Meidinger<sup>1,2</sup>, Elmar Pfe erm ann<sup>1</sup>,

W olfgang Pietsch $^1$ , Lothar Struder $^{1,2}$ , Vyacheslav E. Zavlin $^{1,5}$ 

<sup>1</sup> Max {Planck {Institut fur extraterrestrische Physik, Giessenbachstrae, 85748 Garching, Germany

<sup>2</sup> M P I{H albleiterlabor, O tto{H ahn{R ing 6, 81 739 M unchen, G erm any

<sup>3</sup> M P I{R ontgentestan lage PANTER, G autinger Strae 45, 82061 Neuried, G erm any

<sup>4</sup> Institut fur A stronom ie und A strophysik, Sand 1, 72076 T ubingen, G erm any

<sup>5</sup> Observatoire A stronom ique, 11 rue de l'Universite, 67 000 Strasbourg, France

# ABSTRACT

High quantum e ciency over a broad spectral range is one of the main properties of the EPIC pn camera on { board XMM -Newton. The quantum e ciency rises from 75% at 0.2 keV to 100% at 1 keV, stays close to 100% until 8 keV, and is still 90% at 10 keV.<sup>1</sup> The EPIC pn camera is attached to an X {ray telescope which has the highest collecting area currently available, in particular at low energies (m ore than 1400 cm<sup>2</sup> between 0.1 and 2.0 keV).<sup>2</sup> Thus, this instrum ent is very sensitive to the low {energy X {ray em ission. How ever, X {ray data at energies below 0.2 keV are considerably a ected by detector e ects, which become m ore and m ore in portant towards the lowest transm itted energies. In addition to that, pixels which have received incorrect o sets during the calculation of the o set m ap at the beginning of each observation, show up as bright patches in low {energy im ages. Here we describe a m ethod which is not only capable of suppressing the contam inations found at low energies, but which also im proves the data quality throughout the whole EPIC pn spectral range. This method is then applied to data from the Vela supernova rem nant.

K eyw ords: XMM {Newton, X {ray astronomy, X {ray detectors, EPIC, pn{CCD, detector noise, calibration, im aging, spectroscopy, Vela SNR

# 1. IN TRODUCTION

The EPIC pn camera on {board XMM {Newton is currently the most sensitive instrument in space for detecting soft X {rays. In order to fully utilize this unique capability, we have developed methods which improve the data quality in particular at low energies. These methods are generally available in SASS v6.0, the most recent version of the XMM {Newton Science Analysis System, in the task epreject.

In Section 2 we describe a method which removes bright patches that become visible in low (energy in ages. This is achieved by correcting the energy of events which are detected in the corresponding pixels. Thus, this method does not only improve the cosmetic quality of low (energy images, but does also improve the spectral quality thoughout the whole bandpass of EP IC pn. A line of straightforward application of this method so that it can also be applied if only the event le is available.

A nother in provem ent, described in Section 3, concerns the suppression of detector noise, which becomes signi cant at energies below 200 eV. This is done by computing the probability that a speci c event m ay be a noise event and by assigning, on a statistical basis, appropriate ags to individual events. Subsequent rem oval of events which were agged as due to detector noise leads to a conspicuous in provem ent of the data quality

Further author inform ation: (Send correspondence to K D .) K D .: E-m ail: kod@mpempgde



F igure 1. Number of events per pixel and M s (=  $10^6$  s) as a function of raw pulse height am plitude [adu] for a 23:1  $1^{0}$  s exposure (obtained during revolution # 462) in full fram e m ode with the liter wheel closed. T hese spectra are shown individually for all EP IC pn quadrants (arranged according to their position on the detector). Note the di erent intensity of detector noise in the four quadrants.

below 300 eV and makes the background subtraction in this energy range more reliable. Additionally, as a by {product, the event les become considerably smaller and easier to handle.

In Section 4, we demonstrate these methods on the example of an XMM EPIC probservation of an area within the Vela supernova remnant. In particular we show how the noise suppression method reveals the presence of soft diuse emission at energies below 200 eV which would otherwise be hidden in the detector noise.

# 2.OFFSET SHIFTS

## 2.1. The Problem

The usual mode of operating the EPIC pn camera consists of taking an oset map immediately before the beginning of an exposure. Ideally, this map contains for each pixel the energy oset (expressed in analog{to{ digital units, adu}. During the subsequent exposure, these osets are subtracted on{board from the measured signals, and only events where the di erence signal exceeds a lower threshold (usually 20 adu) are transmitted to Earth.

The o set map is calculated in the following way: the charge in each pixel is read out one hundred times, the three smallest and the three highest adu values are ignored, and the sum of the remaining values, divided by 94 and truncated to the nearest integer, is computed as the o set of that pixel. This procedure is applied subsequently to regions of 64 4 pixels along each CCD. The reason for excluding the extrem e values is to minimize the in uence of energetic particles on the o set calculation. The highest adu values are ignored because the charge deposited by minimum ionizing particles is usually close to the upper end of the dynamic adu range. However, the charge deposited by a highly ionizing particle may show up as a very small charge, because the large amount of charge generated by such a particle takes a long time to be cleared from the anode. Depending on the number of signal electrons, the clearing goes on during the readout of the next pixels, and the measured adu value is thus slighly reduced. Therefore, also the smallest adu values are ignored.

There are cases where so much charge is deposited by highly ionizing particles that it takes several frames until the original state is restored. It may also happen that several minimum ionizing particles are hitting the



Figure 2.Computing the \residualo set m ap": a) O riginalo set m ap for the observation 0526\_0147511101. The vertical stripes are m ainly caused by di erences in the gain of the individual colum ns. b) Sam e as (a), but after subtracting from each colum n its m edian value. The horizontal stripes are m ainly caused by the common mode e ect (see Sect.2.2.1). c) Sam e as (b), but after subtracting also from each row its m edian value. The remaining patterns in this \residualo set m ap" are a superposition of persistent features and tem porary patches caused by energetic particles.

same pixels during the one hundred readouts. At such occasions the technique of rejecting the three sm allest and the three highest adu values before computing the mean oset is not su cient, and the oset is not computed correctly. As the erroneous adu values are predom inantly toolow, the computed mean oset becomes too sm all. This elect introduces in the alected pixels deviations from the correct osets which range typically from 5 adu to +3 adu, corresponding to energy shifts from 25 eV to +15 eV. As a consequence, the energies of all events in these pixels appear to be shifted by the same amount. The number of alected pixels scales with the intensity of the particle background during the calculation of the oset map and depends also on the readout mode. From our analysis of full frame mode (FF) data, we nd that 54% of all pixels contain correct osets, while for 26% the oset is by 5 eV too high and for 12% it is by 5 eV too sm all. Deviations of this amount (1 adu) are not necessarily all caused by energetic particles, as they may also relect statistical and numerical luctuations in the computation of the oset map. Larger deviations, how ever, can clearly be attributed to energetic particles, as can be seen by their characteristic patterm (e.g. Fig.2c). For 5% the oset is by 10 eV too high and for 2% it is by 10 eV too sm all. Osets of 15 eV and more were found for 1% of the pixels. For 70% of them, the oset was too sm all.

If the o set subtracted on {board was too sm all, then the adu values which are assigned to events in such pixels became too high. Thus, events which have adu values below the lower threshold and which would norm ally have been rejected, may show up in the data set. A sm ost of such events are due to detector noise, which is steeply increasing towards lower energies (Fig.1), any reduction of the lower energy threshold leads to a considerable increase in the num ber of events. Thus, pixels where the o set is too sm all show up as bright patches in in ages accum ulated in the lowest transm itted adu range. Due to the speci c pixel block sam pling applied for the o set m ap calculation (see above), these patches appear as rectangular areas, which are characterized by a brightening in four consecutive pixels along readout direction. D epending on the width of the trail caused by the energetic particles and its orientation with respect to the CCD, the patchesm ay also extend over several consecutive pixels perpendicularly to the readout direction. P ixels with too high o sets are less obvious in low {energy in ages, as they are only som ew hat darker than their environm ent.

W hile the occurrence of bright patches in EPIC pn in ages which are accumulated at low energies (e.g.Fig.4) is immediately obvious, there is also another consequence: a shift in the energy scale over the whole spectral bandpass. This shift degrades the energy resolution for extended objects. For point sources, the X {ray spectrum



F igure 3. Separating the persistent and temporary features in the residual o set m ap: a) Sam e as Fig.2c, but for a di erent observation ( $523_0147510801$ ); note the di erent distribution of m ost patches. b) \R esidual o set reference m ap": this m ap shows for each pixel the m edian of seven residual o set m aps (from revs 522, 523, 525, 526, 527, 528, and 544) and thus contains the persistent features in the residual o set m aps. c) Di erence (a) (b), containing the temporary features for this speci c observation.



F igure 4. Separating the persistent and tem porary features in the 20 adu im age: a) D istribution of all events with 20 adu accum ulated in the 76 ks observation  $523_0147510801$ , norm alized to 1 ks. b) \20 adu reference m ap": this m ap shows for each pixel the m edian of ten residual o set m aps (from revs 344, 349, 522, 523, 525, 526, 527, 528, 544, and 548), each norm alized to 1 ks exposure, and thus contains the persistent features in the 20 adu m aps. c) D i erence (a) (b), containing the tem porary features for this speci c observation. Note the sim ilarity with Fig.3c.

m ay be shifted by som e 10 eV, in m ost cases towards higher energies, if the position of the source happens to coincide with one of these patches. Here we describe a method for restoring the correct energy scale and for rem oving the bright patches in the soft energy images. A lthough this paper concentrates on the low energy properties of EPIC pn data, we note that the correction of the o set shifts improves the energy calibration throughout the full spectral bandpass of the EPIC pn detector.

#### 2.2. Correcting the energy scale

The correction is more straightforward if the oset map is available. If this is not the case, then the osets have to be reconstructed from the spatial and spectral distribution of the transmitted events. These two cases are described in the next sections.

## 2.2.1. If the o set m ap is available

The o set m ap contains for each pixel the adu value which was subtracted from all events in the particular pixel before transm itting this information to ground. As it contains both correct and incorrect o sets, the main goal is to distinguish between both cases. Such a distinction is possible under the assumption that the generic o sets of all pixels (not a ected by electronic e ects during readout) are very sim ilar and that the areas where incorrect o sets were computed occur in isolated patches.

The most obvious property of the o set m ap is a pattern of vertical stripes (Fig.2a). This is caused by the fact that each of the 768 readout columns has its own am pli er, and the am pli cation varies by 4.1% (m s) from column to column. In order to spot the patches where wrong o sets have been applied, the column to column has to be removed. This can be achieved by subtracting the median value from each column. The resulting map (Fig.2b) is dominated by row to row variations, which are mainly caused by the \common m ode" e ect: a tem poral change of the o sets of all the pixels in a CCD row, occurring in a predominantly irregular way. This e ect is present during the calculation of the o set m ap as well as during the subsequent exposure. During the exposure, it is suppressed on {board by subtracting rst the adu o sets contained in the o set m ap, and then the median of all the adu values in a CCD row from these values, individually for each row should also be subtracted from the corresponding row, in order to remove the row to row variation. This leads to a \residual o set m ap" (Fig.2c), which is close to zero except for the pixels where speci c o sets were applied.

However, not all of the o sets in the residual o set m ap are artefacts caused by energetic particles. Comparison of such maps (e.g. Figs.2c and 3a) indicates that they consist of a superposition of both persistent and tem porally variable patterns. By computing for each pixel the median value of several such maps, only the persistent patterns remain (Fig.3b). Subtraction of such a \residual o set reference map" from an individual residual o set map reveals then the tem porally variable content of this \cleaned o set map". Comparison of such maps with in ages accumulated from the low {energy events in the subsequent exposure (see Fig.3a and Sect.2.2.2) shows a clear correlation between the position of the bright patches in the low {energy in age and the pixels which have a non{zero value in the cleaned o set map. This shows that the patches seen in the cleaned o set maps re ect incorrect o sets caused by energetic particles.

As the processed o set m aps contain directly the adu shifts which have to be subtracted from all events in the corresponding pixels in order to reconstruct the correct energy scale, an implementation of this algorithm is straightforward. However, while the correct energy scale can be reconstructed, there is no possibility to recover events which were not transmitted to ground because a too high o set was subtracted so that they fellbelow the lower energy threshold. On the other hand, events where a too low o set was subtracted will show up after the correction with adu values below the lower threshold. In order to restore a hom ogeneous data set, these events should be removed afterwards. Then the gain and CTI correction should be repeated for the whole data set.



F igure 5. Pixel by pixel correlation between the excess counts at 20 adu, norm alized to 1 ks, and the values in the cleaned o set m ap, for all quadrants (arranged according to their position on the detector). While the energy shifts to be applied can only occur as integer multiples of 1 adu ( 5 eV), the excess counts exhibit a much smoother distribution due to their norm alization to 1 ks. Their distribution is illustrated by the thickness of \histogram tubes", which were all expanded to have the same m axim um thickness. The number of pixels which they contain are given at right. This correlation was derived by combining the observations of the Lockm an hole in XMM revolutions 522, 523, 525, 526, 527, 528, and 544 ( 636 ks total exposure time), all taken in fulfimmer mode. For Q 3, four noisy columns were excluded. A vertical red line indicates the median value of the distributions. Shaded areas show the ranges of excess counts for which a certain energy shift should be applied. Note how the brightness of pixels at 20 adu responds di erently in the four quadrants to energy shifts. No such variation was found within the CCD s of the same e quadrant.



F igure 6. Extrapolation of the empirical correlation between excess counts in the 20 adu im age and the energy shift to be applied. The numbers below the horizontal steps specify the boundaries of the interval of excess counts, where the energy scale should be corrected by the adu values given to the left of the horizontal steps. The extrapolation was done by tting a cubic function to the median values of the excess counts found for o set shifts between 2 and + 4 adu (cf.Fig.5); these points are marked by sm all dots. An extrapolation was necessary because corrections by m ore than 4 adu may occur, but they do not occur frequently enough for a direct determ ination of the correlation function.

## 2.2.2. If the o set m ap is not available

Even if the o set m ap is not available (this was the general situation in the past), the correction described above can be achieved, by extracting the necessary information directly from the event le. This is possible because the am ount of noise events is a monotonically and steeply increasing function towards lower energies (Fig.1). As this function is also fairly stable in time, the intensity of a pixel in the lowerm ost energy channel transmitted (20 adu) is expected to be correlated with the shift in the energy scale: if the energy scale in a pixel is shifted by only 1 adu towards higher energies, i.e., if the 20 adu bin contains 19 adu events, then this pixel should contain 1.8 times more events.

However, this correlation is disturbed by the fact that the brightness of a pixel at 20 adu is also in uenced by other factors, in particular by its individual noise properties. In Fig.4a, the lower half of the detector (Q 2 and Q 3) are brighter than the upper one (Q 0 and Q 1). Com parison with Fig.1 shows that this behaviour is caused by the fact that there are more noise events at 20 adu in Q 2 and Q 3 than in Q 0 and Q 1. In addition to the brightness di erences from quadrant to quadrant, brightness variations can also be seen within the quadrants and even within the CCDs (Fig.4a).

Separation of brightness variations caused by shifts in the energy scale from variations caused by other e ects can be achieved with a similar method as applied to the residual o set maps (Sect.2.2.1): from a set of 20 adu in ages, derived from long exposures with no bright sources in the eld of view and normalized to the same exposure time, a reference in age can be computed, containing for each pixel the median value, i.e., its nom inal, temporally constant, 20 adu brightness (Fig.4b). This reference in age can then be subtracted from an individual 20 adu in age, in order to derive the brightness variations characteristic for that particular exposure. The resulting \cleaned 20 adu in age" (Fig.4c) shows a clear spatial correlation with the cleaned o set map of the sam e exposure (Fig.3c). In addition to the spatial correlation, there is also a direct correlation between the brightness of the pixels in the cleaned 20 adu im ages (normalized to the sam e exposure tim e) and the cleaned o set m aps, as expected. This correlation (Fig.5) is di erent between the four quadrants, due to their di erent noise properties. No such di erences were found am ong the CCD s within a quadrant.

The correlation shown in Fig.5 was derived from seven long observations of the Lockm an hole (636 ks in total). This region contains only faint sources, which cannot disturb the oset maps or the 20 adu in ages. Due to the long exposure times, the statistical uncertainties in the number of excess counts at 20 adu are minimized. The statistical quality is good enough to deduce the energy shifts to be applied from the number of excess counts



Figure 7. Oblique view onto the EPIC pn detector, illustrating the spatial variations of detector noise for the fullfram e mode. For each row of each CCD, the noise spectrum in the range 20 { 65 adu is displayed in a color code which extends from 0 (black) to 0.01 events ks  $^{1}$  px  $^{1}$ adu <sup>1</sup> (white); the maximum value is 1.3 events ks <sup>1</sup> px <sup>1</sup> adu <sup>1</sup>. In order to cover the high dynam ic range, the colors change with the square root of the noise. These spectra were derived from 10 exposures with the lter wheel closed (in XMM revolutions 129, 266, 339, 351, 363, 393, 409, 454, 462, 532), yielding a total exposure time of 144 ks. The inform ation displayed here is used for the suppression of the detector noise.

for the range from  $3 \text{ to } + 4 \text{ adu. A s larger shifts do not occur frequently enough for a direct determ ination of the correlation function, they were determ ined by extrapolation (Fig.6).$ 

W ith the correlation derived above, it becomes possible to reconstruct the orset shifts directly from the event le, at least in an approximate way: while orset shifts can occur only at discrete adu steps, the correspondence between the 20 adu brightness and the value of the orset shift is not always unique (Fig.5). In addition, the presence of Poissonian noise in the 20 adu images, in particular for short exposures, limits the sensitivity for locating the pixels where a correction of the energy scale should be applied. The sensitivity of locating such pixels can be increased by rebinning four consecutive rows of the 20 adu image into one row each, in order to utilize the information about how the orset map was calculated on {board. Nevertheless, this method is less accurate than the technique described in the previous section, but it does in prove the overall data quality and can be applied if no or set map is available.

## 3.DETECTOR NOISE

W hile there is practically no detector noise present at energies above 80 adu ( 400 eV), X-ray data below this energy, in particular below 40 adu ( 200 eV), are considerably contam inated by noise events, which become m ore and m ore dom inant towards the lowest transm itted energy channels (Fig.1). Investigations of 40 hours of in{orbit calibration data with the lter wheel closed, taken over a period of m ore than two years, show that the noise properties vary with position and energy, but are fairly stable in time for m ost areas on the detector. This property enables a statistical approach for suppressing the detector noise: with the inform ation about the spatial and spectral dependence of the detector noise, the am ount of events which correspond to the expected noise can be agged and suppressed.

The noise properties were derived from 10 exposures in fullfram emode, with the liter wheel closed, between XMM (New ton revolutions # 129 and # 532, yielding a total exposure time of 144 ks. These measurements were rst corrected for o set shifts (see Sect.2) and then used for accumulating raw spectra below 65 adu, individually for each CCD row (Fig.7). The ne spacing along readout direction was chosen because the noise properties change considerably with distance from the readout node, in particular close to the node. In order to get a su cient number of events, no subdivision was made along the CCD row s. This approach was motivated by the fact that the noise properties do not show a pronounced dependence along this direction (unless there is a bright colum n). As the resulting spectra su ered som ewhat from low count rate statistics, they were sm oothed along

readout direction with a running median liter extending over 5 rows. This smoothing was not applied to the 20 rows closest to the readout nodes, where the spectra contained more counts and where the dependence of the spectra on the distance from the readout nodes is high.

W ith this inform ation, potential noise events in a speci c observation can be suppressed, on a statistical basis, in the following way: for each CCD row and adu bin, the number of events is compared with the corresponding value in the noise data (scaled to the same exposure). A coording to the ratio between the actual number of events and the expected noise contribution, individual events can then be random ly agged. In order to improve the statistics som ewhat, the spectra from the observation to be corrected are internally sm oothed by a running m edian liter along readout direction before the noise contribution is com puted.

This method extends the usable energy range down to the lowest energies transmitted, i.e.  $120 \quad 140 \text{ eV}$ , depending on the position on the CCD. Below this energy, parts of the detector become essentially insensitive due to the low energy threshold applied on {board and the combined e ect of charge transfer loss and gain variations within the 768 ampliers in the EPIC pn camera. In addition to the improvement of the data quality, a rem oval of events which were agged as noise events also makes the les considerably smaller and easier to handle.

Suppressing the detector noise by this m ethod has not only advantages for in ages, but leads also to a m ore reliable subtraction of detector noise in the spectra and thus to an improved spectral quality. A large part of the instrum ental background can be taken into account by the unusal technique of background subtraction, i.e., by determ ining the background spectrum from a di erent region on the detector and subtracting it, appropriately scaled, from the spectrum to be analyzed. However, as the detector noise is not vignetted (in contrast to background com ing through the X {ray telescope} and not distributed hom ogeneously across the detector (F ig.7), the usualbackground subtraction technique m ay lead to a system atic error in the norm alization of the low {energy part of the spectrum . A nother interesting consequence of the noise suppression m ethod described here for spectral studies is that the detector noise can be separated from other background com ponents.



F igure 8. Illustration of the steps for correcting and cleaning the data set for the 33.0 ks observation of the Vela SNR in XMM rev.367. Shown are in ages accumulated from all events in the lowest energy channel transmitted (20 adu). These im ages are dominated by detector noise (cf.Fig.1), despite the presence of the SNR.a) Im age derived from the original data set. b) Im age after correcting the energy scale in speci c pixels. c) Im age after suppressing also the detector noise. Note the di erent color scale, which extends from 0 to 50 events per pixels for (a) and (b), but from 0 to 5 events per pixels for (c).



Figure 9.E ects of the steps illustrated in Fig.8 on a low {energy in age of the Vela SNR. These in age are all accumulated in celestial coordinates (cf.Fig.11) in the (instrum ental) energy range 120 {200 eV.a} O riginal in age, b) after correcting the energy scale in speci c pixels, c) after suppressing also the detector noise. The color scale extends from 0 to 40 events perpixel for (a) and (b), and from 0 to 20 events perpixel for (c). The original in age (a) contains 1:4 10 events, while the cleaned in age (c) contains only 0:5 10 events.



F igure 10.a) Events in the 120 { 200 eV energy range which were rejected as due to detector noise. The color scale extends from 0 to 20 events per pixel. This in age contains 0.9 10 events in total. b) Im age of all events with (instrum ental) energies below 120 eV, after correcting the energy scale in speci c pixels, and displayed in a logarithm ic color scale extending from 0 to 500 events per pixel. This im age shows that not all areas of the detector are sensitive to energies below 120 eV.c) Same as (b), but for energies below 140 eV. The insensitive areas have practically all disappeared.

## 4.APPLICATION TO THE VELA SNR

Here we apply the methods described in Sections 2 and 3 to a 33 ks observation of the Vela supernova rem nant, performed in December 2001 (XMM rev.367) in fullframe mode with the medium liter. The observed region (Fig.11) contains the north {west rim of another supernova rem nant, RXJ0852.0-4622, which was discovered in the ROSAT all(sky survey.<sup>3</sup> Compared to the Vela SNR, which is 2 10 years old, RXJ0852.0-4622 is considerably younger (less than 1.5 10 years, with 680 years as the most probable value<sup>4</sup>) and is emitting more energetic X {rays. In the ROSAT data, it becomes visible only above 1.3 keV. At lower energies, it is outshone by the bright emission of the Vela SNR (Fig.11).

Figure 8a shows the EPIC pn data of this observation accumulated at 20 adu, the lowest energies transmitted. White patches resulting from enroneous energy o sets are clearly seen. A fler applying the local corrections of the energy scale described in Section 2, these patches disappear (Fig.8b). The data, how ever, are still dom inated by detector noise. This noise can be suppressed by m one than one order of m agnitude at 20 adu (Fig.8c) with the m ethod described in Section 3.

A lthough the 20 adu in ages are well suited for performing and verifying the corrections, the spectral bandwidth is too small to see di use X {ray emission from the Vela region. In order to demonstrate the improvement in the low {energy X {ray data, Figs.9a-c show images for the 120 { 200 eV energy band, corresponding to the steps illustrated in Figs.8a-c, The cleaned image (Fig.9c) reveals that there is indeed di use X {ray emission present, which is, however, lost in the detector noise if no correction is applied (Fig.9a). The cleaning in this energy range is substantial: 2/3 of the original events at 120 {200 eV were suppressed as due to detector noise. Fig.10a shows an image of all the events which were been removed, in the same intensity scale as Fig.9c.

W hile these methods make it possible to extend the useful energy range down to 120 eV, there is a constraint which cannot be avoided: the combined e ects of charge transfer losses and variations in the gain of the 768 ampli ers cause an increase of the local low energy threshold (20 adu at readout, corresponding to an instrum ental energy of 100 eV) for speci c areas on the detector. This is illustrated in Fig.10b, which demonstrates that not all areas of the detector are sensitive to energies below 120 eV. Hom ogeneous sensitivity across the whole detector is reached at energies above 140 eV (Fig.10c).

All the images shown so far were obtained at energies below 200 eV. However, the improvement in data quality is not restricted to this energy range. Comparison of the spectra obtained before and after the correction (Fig.12) indicates that the detector noise is suppressed up to 400 eV (corresponding to 80 adu), where it becomes negligible (cf.Fig.1). The high sensitivity of XMM EPIC pn makes it possible to produce several narrow (band images of the Vela SNR region below 1 keV. Figure 13 shows exposure corrected images derived from the corrected and cleaned data set in the energy bands indicated in Fig.12a. These images reveal a stunning variety of complex X (ray em ission, which changes considerably with energy, even in this sm all 0.1 { 1.keV spectral range. Due to the improved spectral resolution of XMM compared to RO SAT, em ission from the north {west rim of the young supernova remnant RX J 0852.0-4622, which is visible in the RO SAT data only above 1.3 keV, is already obvious at energies above 0:7 keV.

### REFERENCES

- 1. R. Hartmann, G. D. Hartner, U. G. Briel, K. Dennerl, F. Haberl, L. Strueder, J. Truemper, E. Bihler, E.Kendziorra, J.E. Hochedez, E. Jourdain, P.D hez, P. Salvetat, J.M. Auerhammer, D. Schmitz, F. Scholze, and G. Ulm, \Quantum e ciency of the XMM pn-CCD camera," in Proc. SPIE Vol. 3765, p. 703-713, EUV, X-Ray, and Gamma-Ray Instrumentation for Astronom y X, Oswald H. Siegmund; Kathryn A. Flanagan; Eds., pp. 703{713, Oct. 1999.
- 2. F. Jansen, D. Lumb, B. Altieri, J. Clavel, M. Ehle, C. Erd, C. Gabriel, M. Guainazzi, P. Gondoin, R. Much, R. Munoz, M. Santos, N. Schartel, D. Texier, and G. Vacanti, \XMM -Newton observatory. I. The spacecraft and operations," A&A 365, pp. L1{L6, Jan. 2001.
- 3. B. A schenbach, \D iscovery of a young nearby supernova rem nant," N ature 396, pp. 141 {142, 1998.
- 4. B. A schenbach, A. F. Iyudin, and V. Schonfelder, \C onstraints of age, distance and progenitor of the supernova rem nant RXJ0852-4622/GRO J0852-4642," A & A 350, pp. 997{1006, 1999.



Figure 11. ROSAT 0.1 {0.4 keV im age of the eastern part of the Vela SNR, obtained during the all{sky survey. The white contour outlines the XMM {Newton EPIC pn eld of view during the observation in rev.367 (cf.Fig.13).



Figure 12. EPIC pn spectra of the full eld of view from the pointing to the Vela SNR, displayed in a linear (a) and logarithm ic (b) scale. The spectra are com posed of all \good" (FLAG = 0) and valid patterns (singles, doubles, triples, and quadruples). The upper curves show the original spectra, while the lower curves are the result of the corrections described in the text. Shaded areas in (a) mark the energy bands which were selected for creating the im ages in Fig.13.



Figure 13. EPIC pn im ages of the region of the Vela SNR containing the north {west rim of RXJ0852.0-4622, observed in XMM rev.367 (cf.Fig.11). The energy ranges indicated below each im age are also shown in Fig.12a. All im ages have been corrected for exposure variations and are composed of all valid patterns, after having applied the processing steps described in the text. Note the pronounced changes of the X {ray appearance in this sm all region of the X {ray spectrum.