D iscovery of an X-ray N ova in M 31

Benjam in F.W illiam s¹, M ichael R.G arcia¹, A lbert K.H.Kong¹, Frank A.Prim in i¹, and Stephen S.M urray¹

ABSTRACT

We have obtained snapshot in ages of an X-ray nova in M 31 from Chandra ACIS-I and the Hubble Space Telescope (HST) Advanced Camera for Surveys (ACS). The Chandra position of the X-ray nova was RA = $00.44.06.68 1.74^{\circ}$, Dec.= $+41.12.20.0 2.31^{\circ}$. A follow-up HST observation 24 days later revealed a source at RA = 00.44.06.81, Dec.= +41.12.24.0 that was B = 25.75 0.05. This optical source faded to B = 27.1 0.1 in 3 m onths. During this time period, the X-ray ux decayed linearly from (3.6 0.2) 10^{-4} to < (6.9 0.09) 10^{-5} ct cm $^{-2}$ s $^{-1}$. The HST identication of an optical source in the same region experiencing an obvious drop in brightness in concert with the X-ray nova suggests that this optical source is the counterpart of the X-ray nova. However, the precision of the X-ray position allows the possibility that the optical source is a nearby variable star. We discuss the in plications of both possibilities.

Subject headings: X-rays: bursts | X-rays: binaries | galaxies: individual (M 31)

1. Introduction

Since the discovery of bright X-ray novae (XRNe), these bright, transient X-ray events have allowed some of the most detailed studies of the properties of the accretion disks of stellar-mass compact objects (e.g. Chen et al. 1997; McClintock & Remillard 2004, and references therein). While XRNe have been observed in both high-mass X-ray binaries (HMXBs; Tanaka & Shibazaki 1996) and low-mass X-ray binaries (LMXBs), most of those occurring in LMXBs are conmed black hole binaries. In 15 Galactic XRNe systems optical follow up has revealed that the accreting object is more massive than 3 M (McClintock & Remillard 2004), and therefore cannot be a stable neutron star (NS). The millisecond

¹Harvard-Sm ithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138; william s@ head.cfa.harvard.edu; garcia@ head.cfa.harvard.edu; akong@ head.cfa.harvard.edu; fap@ head.cfa.harvard.edu; ssm @ head.cfa.harvard.edu

variability and > $10^{38} {\rm erg~s}^{-1}$ X-ray lum inosity further supports the argument that these binary systems contain black holes. Therefore, these objects are among the mest examples of black hole candidates known (Charles 1998), and are of intense interest as sites to study general relativity in the 'strong gravity' regime. The discovery of extragalactic XRNe is therefore a crucial step forward in the study of extragalactic stellar-mass black holes.

W ith X-ray lum inosities ranging from $10^{36}\{10^{40}\ {\rm erg\ s}^{\ 1}\ ({\rm Chen\ et\ al.}\ 1997)$, XRNe can easily be detected by Chandra and resolved from other X-ray sources within host galaxies severalM pc away, making M 31 a prime target for extragalactic studies of XRN with Chandra. In addition, with optical magnitudes ranging from $5^{\ 6}\ {\rm M_{\ V}}^{\ 6}$ 5, it is possible to detect and resolve counterparts for extragalactic XRNe in the Local Group with HST.

Extragalactic XRNe have also been discovered in the M agellanic C louds (W hite & Carpenter 1978), M 32 (Garcia et al. 2000), and M 31 (O shome et al. 2001; Trudolyubov et al. 2001; K ong et al. 2001). Chandra and XMM have allowed the discovery of nearly 50 XRNe in M 31 (Trudolyubov et al. 2001; K ong et al. 2002; D i Stefano et al. 2004; W illiam s et al. 2004); W illiam s et al. in preparation). M ost recently, by combining the power of Chandra with HST, X-ray/optical studies of the XRNe in M 31 have begun, providing new details and comparisons to their Galactic siblings (e.g. W illiam s et al. 2004).

Herein we report the discovery of a new XRN in M 31 with new Chandra ACIS-I data. A long with three X-ray detections of this source, we have obtained contemporaneous HST ACS in ages of the region where the XRN appeared. The X-ray data provide measurements of the XRN luminosity and the decay pattern. The optical data provide constraints on the physical properties of the progenitor binary system.

2. Data

2.1. X -ray D ata

We obtained Chandra ACIS-I im ages of M 31 on 09-N ov-2003, 26-N ov-2003, and 27-D ec-2003. In addition to the 4 ACIS-I chips, the S2 and S3 chips were on during the observations. The observations were performed in \alternating exposure readout", so that every 6th frame had 0.6 seconds of exposure instead of the canonical 3.2 seconds. This mode lowers the elective exposure time by 20%, but it provides a second low exposure image in which bright sources are not piled up. The details of these observations, including target coordinates, roll angle of the telescope, and exposure time, are provided in Table 1.

These 3 observations were all reduced in an identical manner using the X-ray data

analysis package C A0 v3.02. We created exposure maps for the images using the C A0 script merge all, and we found and measured positions for the sources in the image using the C A0 task wavdetect. Each data set detected sources down to (0.3{10 keV}) uxes of 2 10 ct cm so 1. Errors for the X-ray positions were determined using the ABA task imagentroid, which determines the errors of the point spread function (PSF) along each pixel axis independently. The counts from the source are projected onto each axis, and the error in the position is then calculated by dividing the standard deviation of the pixel positions of all of the source counts by the square root of the number of counts. As the pixels in Chandra images are aligned with north up and east to the left, the X position error was taken to be the RA. error, and the Y position error was taken to be the Dec. error.

We aligned the positions of bright (> 100 counts) X-ray sources with known globular cluster counterparts to the positions of the centers of their host globular clusters in the images of the Local G roup Survey (LGS; Massey et al. 2001) using the IRAF task comap. These images have an assigned J2000 (FK5) world coordinate system accurate to 0.25° , and they provided the standard coordinate system to which we aligned allofour data for this project. The alignment errors between the Chandra and LGS are shown for each observation in column AL in Table 2. A lignments allowed for adjustments in pixel scale as well as shifts in X and Y.

We cross-correlated the X-ray source positions of all 3 observations against all previously published X-ray catalogs and the Sim bad 5 database to look for any new, bright X-ray source likely to be an XRN.

The data pertaining to the bright new X-ray source CXOM 31 J004406.7+ 411220 (hereafter s1-86, see W illiams et al. 2004 for naming convention details) at RA = 00.44.06.68, Dec. = 41.12.20.0 (J2000) was then studied in further detail. A lthough the observations were performed in the ACIS-I con guration, s1-86 was detected on the S3 chip in the rst 2 observations and on the S2 chip in the third observation.

We extracted the X-ray spectrum of s1-86 from all 3 observations using the CIAO task

²http://cxc.harvard.edu/ciao/download/scripts/merge_all.tar

³http://cxc.harvard.edu/ciao3.0/down.load/doc/detect_htm lm anual/M anual.htm l

 $^{^4}$ IRAF is distributed by the NationalO pticalA stronomy O bservatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

⁵ http://sim.bad.u-strasbg.fr/

psextract⁶. We then these spectra independently, binning the spectra of the rst two observations so that each energy bin contained 20 counts and binning the spectrum of the third observation so that each energy bin contained 10 counts. These binning factors allowed the use of standard ² statistics when thing the spectra.

We the spectra with an absorbed power-law model and an absorbed disk blackbody model using the CIAO 3.0/Sherpa thing package (Freem an et al. 2001). The best thing model parameters, and the associated thing statistics, are provided in Tables 4 and 5. Results are discussed in x 3.

2.2. Optical D ata

We obtained two sets of HST ACS data, one observed at UT 21:35 on 03-D ec-2003 and one observed at UT 16:38 on 01-M ar-2004. Each of these were pointed at RA = 00:44:07, Dec.= 41:12:19.5. The observations had orientations of 71.75 deg and 32.73 deg respectively. Both observations were taken using the standard ACS box 4-point dither pattern to allow the naldata to be drizzled to recover the highest possible spatial resolution. All exposures were taken through the F435W lter. The total exposure times were 2200 seconds for each data set.

We aligned and drizzled each set of 4 im ages into high-resolution (0.025°) pixel ¹) im ages using the PyRAF⁷ task multidrizzle, which has been optimized to process ACS imaging data. The task removes the cosmic ray events and geometric distortions, and it drizzles the dithered frames together into one nalphotometric image. We processed the relevant sections of the nal images with DAOPHOT-II and ALLSTAR (Stetson et al. 1990) to not and measure the count rates of the sources. Sections (10°) of the images, centered on the X-ray position of the detected XRN, are shown in Figure 1.

We converted the count rates measured on our images to ST magnitudes and VEGA magnitudes using the conversion techniques provided in the ACS Data Handbook⁹. The count rates, ST (F435W) magnitudes, and VEGA (B) magnitudes for the optical source of interest are provided in Table 3 and discussed in \times 4.1.

⁶ http://cxc.harvard.edu/ciao/ahelp/psextract.htm 1

⁷PyRAF is a product of the Space Telescope Science Institute, which is operated by AURA for NASA.

 $^{^{8}}$ m ultidrizzle is a product of the Space Telescope Science Institute, which is operated by AURA for NASA. http://stsdas.stsci.edu/pydrizzle/multidrizzle

 $^{^9}$ http://www.stsci.edu/hst/acs/documents/handbooks/DataHandbookv2/ACS_longdhbcover.html

Finally, we aligned the drizzled ACS im ages with the LGS coordinate system with com ap using stars and clusters common to both images. This alignment was performed independently for each of the $2\ ACS$ data sets to check the alignment consistency. As displayed by the coordinates in Figure 1 and the errors measured by com ap (0.025^{0}), we were able to accurately align the ACS images to the LGS coordinate system. Because the ACIS-I images had been aligned to the same coordinate system, we were able to compare the coordinates of objects the the ACS and ACIS-I images reliably.

3. Results

3.1. X -ray

The rst X-ray observation (0 bsID 4678) provided the rst detection of s1-86 (C X O M 31 J004406.7+ 411220 at R A = 00.44.06.71, Dec. 41.12.20.0 (J2000)). This position is 16^0 east and 4^0 south of the nucleus, about 1^0 east of the eastern spiral arm and in the outskirts of the bulge. Because this source was very bright and did not appear in any previous publication, we assumed that it was an XRN. We were able to not a 10 ks XMM image of this region (0 B S_ID = 0112570601) taken 28 December 2000, which contained this region near the edge of the eld. Assuming a power-law spectrum with a slope of 3.0 and an absorption column of $2 \cdot 10^{21}$ cm⁻², the non-detection of s1-86 in this XMM observation provides a 4 upper-limit to the unabsorbed 0.3 {7 keV luminosity of 2.5 10^{36} erg s⁻¹. Our rst detection of the source, with an unabsorbed 0.3 {7 keV luminosity of 2.4 10^{38} erg s⁻¹, shows that it brightened by a factor of $^> 100$.

The next two observations in our M 31 m on itoring program (0 bs $\!\!$ D s 4679 and 4680) con $\!\!$ m ed the transient nature of the source, as the X-ray $\!\!$ ux decayed by a factor of 7.3 in 48 days. This decay is clearly seen in the X-ray im ages shown in Figure 3, demonstrating that $\!\!$ sl-86 is an XRN.

The three-point X-ray lightcurve from the 3 AC IS-I detections of s1-86 is shown in Figure 2. Although there are only 3 data points, the decay is not consistent with an exponential. The best thing exponential decay is shown on the gure with a dashed line. It has $^2 = 27.28=1$ and an e-folding time of 26.9 days. On the other hand, a linear decay, shown with the solid line, provides an excellent to the data ($^2 = 0.33=1$) with a slope of -9.3 10 6 ct cm 2 s 1 day 1 and a total decay time of 60 days.

To look for short-term variability, we performed a Kolmogorov-Smimov (KS) test between the photon arrival times for s1-86 during its rst detection (ObsID 4678) and a uniform photon arrival distribution (simulating a constant source) containing the same number of

counts. We considered the arrival times for the 758 counts contained in the region of s1-86 as measured by wavdetect. This test returned a P-value of 0.969, which provides no indication of variability during the observation.

Next, the three positional measurements for s1-86 and their errors (given in Table 2) were combined to determ ine the most accurate coordinates for the X-ray source. Table 2 includes the 3 largest sources of positional errors in our analysis. These errors were all included in our determination of the X-ray position of s1-86. The quantity $_{\rm cen}$ corresponds to the positional error measured by immentroid, which is limited by the large PSF at the location of s1-86, $16^{\rm 0}$ o -axis. The quantity $_{\rm AL}$ corresponds to the positional error of the alignment of the AC IS X-ray positions with the LGS coordinate system, measured by comap. The quantity $_{\rm S!~I}$ corresponds to the positional error of sources on the AC IS-S chips when AC IS-I is the aimpoint, as was the case for all 3 of our detections of s1-86; this error was estimated by M arkevitch. He found an error of $1^{\rm 0}$ is typical for such positions. This error is independent of our alignment procedure as none of our alignment GCs were located on the S-chips. We included this error by adding it directly to our totals from the other errors.

The weighted mean of the three positions and combined random errors (excluding $s!\ _{\rm I}$) provided our best measurement of the location of $s!\ _{\rm S}!\ _{\rm I}$) error to both coordinates, the nal (1) R A position error is 1.74° , and the nal (1) Dec. position error is 2.31° . At the distance of M 31 (780 kpc; W illiam s 2003), these errors correspond 6.6 pc and 8.7 pc.

In addition, as a consistency check to our weighted mean position, we co-added the 3 aligned ACIS observations and measured the wavdetect position and immentroid position errors of the combined image of s1-86. This position is RA=00:44:06.75 0.61° , Dec.=41:12:20.2 1.35° , in good agreement with the weighted mean position calculated above. After adding the systematic ($_{\rm S!~I}$) error to both coordinates, the nal (1) RA. position error is 1.61° , and the nal (1) Dec. position error is 2.35° .

Finally, the three X-ray spectra were allwell-t by an absorbed power-law model and reasonably t (though not as well) by an absorbed disk blackbody model. All of the properties and ² values for the ts are provided in Tables 4 and 5.

The power-law ts show no evidence for spectral variability during the decay of the X-ray ux. All of the power-law ts are consistent with a power-law index of 3 and an absorption column of 2 10^{21} cm 2 . The disk blackbody ts suggest that, during the decay of the X-ray ux, there was a decrease in the inner disk tem perature from 0.6 keV to 0.4 keV.

¹⁰ http://asc.harvard.edu/cal/H mm.a/optaxis/platescale/geom_public.html

These ts yield a constant inner disk radius $^{>}$ 50 km and a constant absorption column of 5 10^{20} cm 2 . The tem peratures and radii are similar to those found for AC IS-I observations of the M 31 X-ray transient r2-67 ($T_{\rm in}=0.35~0.05~{\rm keV}$, $R_{\rm in}\cos^{1-2}i^{>}$ 140 km), which had an optical counterpart detected with HST (W illiams et al. 2004).

The best estim ate of the absorption to s1-86 is the weighted m ean of the values m easured from the rst 2 spectral m easurem ents, which is $(2\,2\,0\,3)\,10^{1}\,$ cm 2 (power-law model) or $(5\,2\,0\,9)\,10^{20}\,$ cm 2 (disk blackbody model). The maximum observed absorption-corrected luminosity is $0.9\{2.4\,10^{38}\,$ erg s 1 depending on the model t, so the luminosity alone does not discriminate between an accreting black hole or neutron star.

3.2. Optical

We searched the ACS im ages, aligned to the same coordinate system as the ACIS im ages, for sources within 4.5° (2) of the X-ray position. DAOPHOT found 791 optical sources in the rst ACS observation down to B = 27.8 (the 4 detection limit). DAOPHOT found 799 optical sources in second ACS observation down to B = 27.8 within 4.5° of the X-ray position. With a distance modulus to M 31 of 24.47 and a typical extinction $A_{\rm B}$ 0.4, this detection limit corresponds to M $_{\rm B}$ 2.9.

In addition, we note that the brightest star within $4.5^{\circ\circ}$ of the X-ray position has B = 24.25 0.03, fainter than a counterpart for a high-m ass Be transient. Such transients have optical counterparts with spectral type 0.8.5 (B 2 (N egueruela 1998), $5.0 < M_B < 2.6$, or 19.8.0 < B < 22.2 scaled to the distance and extinction of M 31.

None of the optical sources down to B=26.8 within the 1 error ellipse of the X-ray source position in the rst ACS in age had a signi cantly lower (m 1 mag) optical ux in the second ACS observation. Extending the search to the 2 error ellipse, there is only 1 optical source with B 26.8 in the rst ACS in age that had a signi cantly lower (m 1 mag) optical ux in the second ACS observation. This sole counterpart candidate is located at RA=00:44:06.81 and Dec=+41:12:24.0 (J2000). This position is shown on the ACS in ages of the region of interest in Figure 1. The source ux decreased by 70%, from $m_B=25:75=0:05$ to $m_B=27:1=0:1$, between ACS observations. Further details about this source in both ACS observations are provided in Table 3.

If this fading optical source is indeed the counterpart of s1-86, we can use the simultaneous X-ray and optical luminosities to predict the orbital period for the system using the relation of van Paradijs & McClintock (1994). Since this relation was seen in Galactic XRNe events, there have been many new systems discovered. Using the LMXB catalog of Liu et al.

(2001), we checked the consistency of this relation with more recent X-ray novae, including GRO J1655-40, 4U 1543-47, XTE 1550-564, XTE J1118+480, and V4641 Sgr. With the exception of XTE J1118+480, which was two magnitudes brighter in the optical than the prediction from the relation, these more recent XRNe all the van Paradijs & McClintock (1994) relation. The higher than expected optical luminosity of XTE J1118+480 can be understood in the context of the van Paradijs and McClintock model, which postulates that the optical ux is due solely to X-ray heating. XTE J1118+480 has the lowest X-ray luminosity of all sources in the sample, and is therefore the most likely to reveal optical emission above and beyond that due to X-ray heating.

The van Paradijs & McC lintock (1994) relation requires only simultaneous measurements of the absolute V magnitude and the X-ray luminosity to predict the orbital period of an LM XB system. These measurements can be taken from any part of an outburst when the dominant contributor to the optical ux is X-ray heating; the peak luminosity and outburst amplitude are not required in the application of the relation.

We determ ined the $L_{\rm X}$ / $L_{\rm opt}$ ratio at the time of the rst H ST observation (03-Dec-2003), assuming the X-ray luminosities given by the power-law ts in Table 4. We apply the X-ray luminosity on 03-Dec-2003 as predicted by our linear decay to the lightcurve (1.4 10^8 erg s 1). To determ ine M $_{\rm V}$, we apply the B magnitude of our candidate (25.75 0.05), and correct for the absorption measured in the X-ray spectra ((2.2 0.3) 10^{21} cm 2). This absorption can be converted to $A_{\rm B}=1.6-0.2$ using the conversion of P redehl & Schmitt (1995) and assuming a standard interstellar extinction law. These values give M $_{\rm B}=0.34-0.23$. A ssuming the mean intrinsic B-V colors of LM XBs in the Liu et al. (2001) Galactic LM XB catalog (-0.09 0.14), M $_{\rm V}=0.25-0.27$ for the optical counterpart candidate.

Applying these L_X and M $_V$ values and errors to the relation of van Paradijs & M cC lintock (1994), including the errors on the parameters of the relation, we obtain an orbital period of 1.0 $^{+2.9}_{-0.6}$ day. Applying K epler's law, the inferred disk size is then $0.7^{+0.9}_{-0.4}$ 10^{12} M $_{10}^{1=3}$ cm. These values are reasonable, but pushing the low end, for a large disk that would produce a long, linear X-ray decay like the one observed for s1-86 (K ing & R itter 1998).

If we sim ply apply the X-ray lum inosity and absorption given by the power-law $\,\,$ t from the X-ray detection closest in time to the $\,\,$ rst optical detection (26-N ovem ber-2003; 1.8 $\,$ 10 8 erg s 1 ; see Table 4), rather than interpolating the X-ray lum inosity on D ecember 3 as we have done above, the calculated period is not signicantly a ected (0.5 $^2_{0.7}$ day). However, if we apply the X-ray lum inosity and absorption column values given by the disk blackbody ts in Table 5, the calculated period is 0.3 $^{0.5}_{0.1}$ day. Applying Kepler's law, the inferred disk size is then $\,\,$ 3 $^{+2}_{1}$ $\,\,$ 10 11 M $^{1=3}_{10}$ cm .

If our only plausible optical counterpart candidate is not the true counterpart, then the true counterpart must have been fainter than B = 26.8. Since our (4) detection lim it was B = 27.8, a source fainter than B = 26.8 in the rst im age would only provide a lower-lim it for the magnitude decrease, and the lower-lim it would be less than 1 m ag, even if the source was unseen in the second ACS observation. Applying the same calculations as described above to this conservative B magnitude lim it (if the optical candidate is not the true counterpart) provides an upper-lim it on the orbital period of the system of p < 0.6 day and an upper-lim it on the size of the disk of a < 4.5 10^{11} M $_{10}^{1=3}$ cm. Such values are below those predicted for XRNe with long, linear X-ray decays (King & Ritter 1998; see x 4.1.2).

4. Discussion

4.1. A rgum ents for an optical counterpart

4.1.1. Optical magnitudes

Chen et al. (1997) determ ined that the mean optical decay timescale for Galactic XRNe is a factor of 2.2 longer than the X-ray decay timescale. This mean difference in decay timescales is in agreement with theoretical models of accretion processes, which predict that the optical decay timescales will be longer than X-ray decay times by a factor of 2 $\{4 \text{ (K ing \& R itter 1998)}.$

For s1-86, the total X -ray decay time was 60 days. Since the optical ux of the second detection (01-M ar-2004) of the counterpart candidate was 30% of the 03-D ec-2003 optical ux 88 days earlier, we estimate the optical decay time for the candidate to be 130 days. Our estimate of the optical/X -ray decay time ratio is therefore 2.2.

The decay time ratios for the Galactic XRNe in the Chen et al. (1997) sample cover a wide range, as do more recent Galactic XRNe. In some cases, such as the May 1989 XRNe of 2023+338 (V 404 Cyg), the optical decay time has been observed to be shorter than the X-ray decay time (Kitam oto et al. 1989; Wagner et al. 1991; Chen et al. 1997). However, we note that the optical decay of the source detailed in Table 3 at RA = 00.44.06.81, Dec. = 41.12.24.0 (J2000) lies near the mean of Galactic events and is consistent with theoretical models. This consistency argues in favor of the candidate being the true optical counterpart.

4.1.2. Disk size and orbital period

As discussed in King & Ritter (1998), one may use the decay time of the XRN to predict the size of the accretion disk of the system. This method was applied to the X-ray transient source r2-67 by Williams et al. (2004) and gave an answer consistent with the period prediction from the optical and X-ray luminosities.

We calculated the size of the disk expected for a 10M black hole with a linear decay to approximately zero X-ray ux 60 days after a peak X-ray luminosity of 2.4 10^8 erg s 1 . Similar to the calculation for r2-67 in Williams et al. (2004), the disk size prediction for s1-86 is

$$2 \quad 10^{12} \frac{M_{10}^{1=3}}{2=3T_4^{2=3}} \text{ cm ;}$$

where is the angular momentum transport e ciency, M $_{10}$ is the black hole mass in 10 M units, and T $_4$ is the local disk temperature in units of 10^4 K.W e note that if the true peak X-ray luminosity was higher, the disk size predicted by the model would increase. If we apply the lowest possible peak X-ray luminosity, that given by the disk blackbody to the 09-Nov-2003 data (9 10^{37} erg s 1), the disk size prediction is

$$10^{12} \frac{\text{M}_{10}^{1=3}}{2^{=3}\text{T}_{4}^{2=3}} \text{ cm} :$$

A coording to K epler's law s, the size of the orbit of the secondary is $a=6:3 ext{ } 10^{11} P_1^{2=3} M_{10}^{1=3}$ cm, where P_1 is the orbital period in days. Because the disk cannot be larger than the orbit of the secondary (a), these calculations suggest that the orbital period of this system is > 1 day. This estimate of the orbital period is consistent with the prediction from the van Paradijs & McC lintock (1994) relation when the optical luminosity of the counterpart candidate is applied (see x 32), providing some additional condense that the candidate is the true optical counterpart.

4.2. A rgum ents against an optical counterpart

4.2.1. Positional (dis) agreem ent

On the other hand, the precision of the X-ray position for s1-86 is not optimal. As shown in Table 2 and discussed in x 3, the X-ray position errors include 50 square arcseconds of area within the 2 error ellipse.

The optical candidate is 0.8 away from the R A. of the X-ray position and 1.7 away from the D eclination of X-ray position. Assuming G aussian statistics, the optical source has 9% probability of occupying the same location as the X-ray source; however, we do not rule out the possibility that this is the counterpart because the C handra PSF is known to be complicated far o -axis. In addition, as shown in Figure 3, the location of the optical counterpart candidate appears qualitatively reasonable when placed on the X-ray in ages of s1-86.

4.2.2. A Itemate optical source possibilities

If the optical candidate is not the counterpart of the X-ray transient, the candidate must be a dierent type of optical source that varies by 1.3 m ag in brightness. Such sources are not extremely rare. We searched the overlapping portions of our 2 ACS observations for other sources that became fainter by more than 1 m ag between the rst and second observations. We found 75 such sources in the 9.2752 square arcm inutes of overlapping area. Since our 2 error ellipse covers 0.0140 square arcm inutes, there is an 11% random probability of one of these sources falling in this region, similar to the probability that the candidate occupies the same position as the X-ray source.

There are several types of variable sources that could show the same behavior as the counterpart candidate, including eclipsing binaries, high-amplitude variable stars, and dwarf nova systems. For example, a common type of variable with properties similar to those of the counterpart candidate is RR Lyrae stars. M 31 is known to have $^{>}$ 5 RR Lyrae stars per square arom inute (Brown et al. 2004). At B $_{\rm C}$ 26, assuming a distance modulus of 24.47 (780 kpc) and foreground extinction (A_B = 0.4), the counterpart candidate has M_B $_{\rm C}$ 1, typical of RR_{ab} Lyrae stars, which can vary by 0.5 $^{<}$ m $^{<}$ 1.5 mag (Binney & Merri eld 1998). If the candidate is such a star, its period and phase must have allowed us to observe it near maximum during our rst HST observation and near minimum during our second HST observation. There were 88.8 days between our two HST observations. If this optical source is an RR_{ab} star, its period is likely 0.55 days (Binney & Merri eld 1998). Therefore, the star would have undergone 161.5 cycles between our observations, allowing us to observe it in opposite phases. It is clearly possible that the counterpart candidate is an unassociated variable star.

5. Conclusions

We have reported the detection of a new, bright transient X-ray source in M 31, which we call s1-86 (CXOM 31 J004406.7+ 411220). A fiter aligning the Chandra AC IS-I im ages to the LGS coordinate system, this source is located at R A = 00.44.06.68, D ec.= 41:12:20.0 with nal R A. and D ec. errors of 1.74° and 2.31° , respectively. The X-ray spectral properties of this source as measured from 3 Chandra AC IS-I detections are typical of known G alactic XRNe. Power-law ts indicate a spectral index of 3 and an absorption column of $2.2 ext{ 16}^{\circ}$ cm $^{\circ}$. Disk blackbody ts indicate an inner disk temperature decrease from 0.6 keV to 0.4 keV over 1 m onth and an absorption column of 5.10^{20} cm $^{\circ}$. The 0.37 keV lum inosity was $2.4 ext{ 10}^{38}$ erg s $^{\circ}$ (power-law), or $0.9 ext{ 10}^{38}$ erg s $^{\circ}$ (disk blackbody), depending on the spectral model employed. The X-ray lightcurve exhibits a linear decay, suggesting a large accretion disk.

In two HST ACS observations, we detected only one optical source within $4.5^{\circ\circ}$ of the X-ray position of the XRN that faded in concert with the X-ray source. A fiter aligning the ACS images to the LGS coordinate system, this optical counterpart candidate was located at RA=00.44.06.81 and Dec=+41.12.24.0 (J2000). While the optical brightness and decay timescale suggest that this optical source is the counterpart, the errors in the X-ray source position allow the possibility that the counterpart candidate is a nearby RR_{ab} Lyrae star which we observed near maximum and minimum brightness. If so, no optical counterpart was seen down to B=26.8.

If the candidate is the optical counterpart and the X-ray lum inosity indicated by the power-law spectral t is correct, then the X-ray/optical lum inosity ratio yields predictions for the orbital period of the X-ray binary and the size of the accretion disk. The period prediction is $1.0^{+2.9}_{-0.6}$ day, and the disk size prediction is $0.7^{+0.9}_{-0.4}$ 10^{12} M $_{10}^{1=3}$ cm. These values are on the low side ofm odel predictions for XRNe with long, linear X-ray decays, and the lower X-ray lum inosity given by the disk blackbody spectral tyields even lower values.

If the optical counterpart was not detected, it was at least 1 m ag fainter than the candidate counterpart. This upper-lim it to the optical lum inosity provides lower-lim its on the X-ray/optical lum inosity ratios given by both X-ray spectral models. In both cases, these lower-lim its yield upper-lim its on the orbital period and the disk size that are below model predictions for XRNe with long, linear X-ray decays.

Support for this work was provided by NASA through grant number GO-9087 from the Space Telescope Science Institute and through grant number GO-3103X from the Chandra X-Ray Center. MRG acknowledges support from NASA LTSA grant NAG5-10889.

REFERENCES

- Binney, J., & Merri eld, M. 1998, "Galactic astronomy" (Galactic astronomy / James Binney and Michael Merri eld. Princeton, NJ: Princeton University Press, 1998. (Princeton series in astrophysics))
- Brown, T.M., Ferguson, H.C., Smith, E., Kimble, R.A., Sweigart, A.V., Renzini, A., & Rich, R.M. 2004, AJ, 127, 2738
- Charles, P. 1998, in Theory of Black Hole Accretion Disks, 1
- Chen, W., Shrader, C.R., & Livio, M. 1997, ApJ, 491, 312
- DiStefano, R., et al. 2004, ApJ, 610, 247
- Freem an, P., Doe, S., & Siem iginow ska, A. 2001, in Proc. SP IE Vol. 4477, p. 76-87, A stronom ical Data Analysis, Jean-Luc Starck; Fionn D. Murtagh; Eds., 76
- Garcia, M. R., Murray, S. S., Primini, F. A., McClintock, J. E., & Callanan, P. J. 2000, IAU Circ., 7498, 3
- King, A.R., & Ritter, H. 1998, MNRAS, 293, L42
- Kitam oto, S., Tsunemi, H., Miyam oto, S., Yamashita, K., & Mizobuchi, S. 1989, Nature, 342, 518
- Kong, A., Garcia, M., Murray, S., Primini, F., McClintock, J., & DiStefano, R. 2001, The Astronomer's Telegram, 76, 1
- Kong, A.K.H., Garcia, M.R., Primini, F.A., Murray, S.S., DiStefano, R., & McClintock, J.E. 2002, ApJ, 577, 738
- Liu, Q.Z., van Paradijs, J., & van den Heuvel, E.P.J. 2001, A&A, 368, 1021
- Massey, P., Hodge, P.W., Holmes, S., Jacoby, G., King, N.L., Olsen, K., Saha, A., & Smith, C. 2001, in American Astronomical Society Meeting, Vol. 199, 13005
- McClintock, J.E., & Remillard, R.A. 2004, in Compact Stellar X-ray Sources (astro-ph/0306213)
- Negueruela, I. 1998, A & A, 338, 505
- O shome, J.P., et al. 2001, A & A, 378, 800
- Predehl, P., & Schmitt, J.H.M.M.1995, A&A, 293, 889

- Stetson, P.B., Davis, L.E., & Crabtree, D.R. 1990, in ASP Conf. Ser. 8: CCDs in astronomy, 289
- Tanaka, Y., & Shibazaki, N. 1996, ARA & A, 34, 607
- Trudolyubov, S.P., Borozdin, K.N., & Priedhorsky, W.C. 2001, ApJ, 563, L119
- van Paradijs, J., & McClintock, J.E. 1994, A&A, 290, 133
- Wagner, R.M., Bertram, R., Starr eld, S.G., Howell, S.B., Kreidl, T.J., Bus, S.J., Cassatella, A., & Fried, R. 1991, ApJ, 378, 293
- W hite, N.E., & Carpenter, G.F. 1978, MNRAS, 183, 11P
- W illiam s, B.F. 2003, MNRAS, 340, 143
- W illiam s, B.F., Garcia, M.R., Kong, A.K.H., Prim ini, F.A., King, A.R., DiStefano, R., & Murray, S.S. 2004, ApJ, 609, 735

This preprint was prepared with the AAS $\mathbb{P}T_EX$ macros v5.0.

Table 1. Chandra ACIS-I observations

0 bsID	D ate	R.A. (J2000)	Dec. (J2000)	Roll (deg.)	Exp. (ks)	
4678	09-N ov-2003	00 42 44.4	41 16 08 3	239.53	3.9	
4679	26-N ov-2003	00 42 44.4	41 16 08 3	261.38	3.8	
4680	27-D ec-2003	00 42 44.4	41 16 08.3	285.12	4.2	

Table 2. Position M easurements and Errors of XRN s1-86

${\mathbb D}$	R A . (J2000)	cen	ΑL	S! I	tot	Dec. (J2000)	cen	ΑL	S! I	tot
4678	00 44 06.64	0 . 96 [®]	0.1000	1^{\odot}	1 . 96 [®]	41 12 20.0	1 . 68 [®]	0.1800	1^{\odot}	2 . 69 [®]
4679	00 44 06.73	1.23°	0.09^{00}	1^{\odot}	2.23 ⁰⁰	41 12 20 1	2.40 ⁰⁰	0.19^{00}	1^{∞}	3.41 [®]
4680	00 44 06.85	3 . 79 ⁰⁰	0.12^{00}	1^{\odot}	4.80 ⁰⁰	41 12 19.9	3 . 96 ⁰⁰	0.24^{00}	1^{∞}	4.96 ⁰⁰
M ean	00 44 06.68				00	41.72 20.0				

Table 3. HST ACS F435W photom etry for the candidate optical counterpart of s1-86

D ate	R A .(J2000)	Dec.(J2000)	(⁰⁰) ^a	Rate (ct s ¹)	STm ag ^b	VEGAm ag ^c	
	00:44:06.81 00:44:06.81			1.00 0.05 0.30 0.03			

^aD istance between this source position and the X-ray source position.

^cThe VEGA magnitude of the source is obtained from a system where Vega has a magnitude of 0 at all wavelengths. This magnitude is the best approximation of the Johnson B magnitude of the source.

^bThe ST magnitude of the source is obtained from a system with a at reference spectrum (see http://www.stsci.edu/hst/acs/documents/handbooks/DataHandbookv2 formore details).

Table 4. Hardness Ratios and Power-law Spectral Fits to ACIS-ID etections of XRN s1-86

D ate	0 bsID	C ts ^a	F lux ^b	S lop e ^c	N H	2=	Q ^d	н R -1 ^е	н R −2 ^f	r ^X a
09-N ov-2003	4678	666	5.1 0.2	2.9 0.2	2.25 0.35	39.33/30	0.12	0.11 0.04	-0.58 0.07	2.4
26-N ov-2003	4679	465	3.6 0.2	3.0 0.3	2.13 0.47	21.40/19	0.31	0.02 0.05	-0.69 0.08	1.8
27-D ec-2003	4680	79	0.7 0.1	3.4 0.4	2.0 ^h	3.47/8	0.90	0.10 0.11	-1.14 0.39	0.5

^aThe background-subtracted number of counts in the detection.

 $^{\rm d} T \, he$ probability that this $\,$ t is representative of the true spectrum , determ ined from $\,^2\text{=}\!\text{dof}$.

 $^{\rm e}H$ ardness ratio calculated by taking the ratio of M -S/M + S, where S is the number of counts from 0.3{1 keV and M is the number of counts from 1{2 keV .

 $^{\rm f}$ H archess ratio calculated by taking the ratio of H-S/H+S, where S is the number of counts from 0.3{1 keV and H is the number of counts from 2{7 keV.

 $^{\rm g}{\rm T}\,{\rm he}$ absorption-corrected lum inosity of the source in units of 10^{38} erg s 1 (0.3{7 keV).

 $^{\rm h}$ The absorption was xed to the spectrum from 27-Dec-2003, as N was unconstrained if left as a free parameter.

 $^{^{\}mathrm{b}}$ The exposure corrected 0.3{10 keV ux in units of 10 $^{\mathrm{4}}$ ct cm $^{\mathrm{2}}$ s $^{\mathrm{1}}$.

^cSlope of the best-tting absorbed power law model.

Table 5. Disk Blackbody Spectral Fits to ACIS-ID etections of XRN s1-86

D ate	OBSID	T _{in} a	$R_{in}\cos^{1=2}i^b$	N _H	2=	Q°	L _X d
09-N ov-2003	4678	0.60 0.04	56 9	0.5 0.1	39.37/30	0.12	0.9
26-N ov-2003	4679	0.49 0.05	$74^{+}_{19}^{15}$	0.6 0.2	30.37/19	0.05	0.6
27-D ec-2003	4680	0.37 0.07	65 ^{+ 24} 40	0.5 ^e	4.51/8	0.81	0.2

^aThe tem perature of the inner disk in keV.

 $^{^{\}mathrm{b}}\mathrm{T}\,\mathrm{he}$ radius of the inner disk in km , assum ing the distance to M 31 is 780 kpc and the inclination (i) of the binary is 0 degrees.

 $^{^{\}rm c}\text{T}\,\text{he}$ probability that this $\,$ t is representative of the true spectrum , determined from $^{\rm 2}\text{=}\text{dof}$.

 $^{^{\}rm d}{\rm T}\,{\rm he}$ absorption-corrected lum inosity of the source in units of 10^{38} erg s 1 (0.3{7 keV).

 $^{^{\}mathrm{e}}$ The absorption was xed to the spectrum from 27-Dec-2003, as N was unconstrained if left as a free parameter.




Fig. 1. The relevant sections of the 2 drizzled ACSF435W images are shown. Left panel: the UT 21:35 03-Dec-2003 observation is shown with a black circle marking the position of the source that faded by more than 1 mag. Right panel: the UT 16:33 01-Mar-2004 observation is shown with a black circle marking the position of the source that faded by more than 1 mag. White ellipses mark the 1 (8 pc) and 2 (8 pc) X-ray position errors.




Fig. 2. The lightcurve of s1-86 during our three ACIS-I detections. Dashed curve: the best texponential decay curve, with an e-folding time of 26.9 days (2 = 27.28/1). Solid line: the best tlinear decay curve, with a slope of 9.3 10 6 ct cm 2 s 1 day 1 (2 = 0.33/1).




Fig. 3. Top left panel: The 1 and 2 X-ray position errors are shown with white ellipses on the HST image from 03-Dec-2003. The optical counterpart candidate is indicated with the black circle. Top right panel: The same error ellipses are shown on the ACIS-I image of s1-86 from 09-Nov-2003. Lower left panel: The same error ellipses are shown on the ACIS-I image of s1-86 from 26-Nov-2003. Lower right panel: The same error ellipses are shown on the ACIS-I image of s1-86 from 27-D ec-2003.