

# The discovery of 10 pulsars in an Arecibo drift-scan survey

D. R. Lorimer<sup>1,2</sup>, K. M. S. Alexander<sup>2</sup>, A. S. Fruchter<sup>3</sup>, I. H. Stairs<sup>4</sup>, F. Camilo<sup>5</sup>,  
A. M. Vazquez<sup>6</sup>, J. A. Eder<sup>6</sup>, M. A. McLaughlin<sup>1</sup>, M. S. E. Roberts<sup>7</sup>,  
J. W. T. Hessels<sup>7</sup> and S. M. Ransom<sup>8</sup>

<sup>1</sup> University of Manchester, Jodrell Bank Observatory, Macclesfield, Cheshire, SK11 9DL

<sup>2</sup> University of Virginia, Department of Astronomy, PO Box 3818, Charlottesville, VA 22903, USA

<sup>3</sup> Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

<sup>4</sup> Department of Physics and Astronomy, University of British Columbia, 6224 Agricultural Road, Vancouver, BC V6T 1Z1, Canada

<sup>5</sup> Columbia Astrophysics Laboratory, Columbia University, 550 West 120th Street, New York, NY 10027, USA

<sup>6</sup> Arecibo Observatory, HC 3 Box 53995, Arecibo, PR 00612, USA

<sup>7</sup> Physics Department, McGill University, 3600 University Street, Montreal, QC H3A 2T8, Canada

<sup>8</sup> National Radio Astronomy Observatory, 520 Edgemont Road, Charlottesville, VA 22903, USA

Accepted 2005 March 8; Received 2005 March 7; in original form 2005 February 24

## ABSTRACT

We present the results of a 430-MHz survey for pulsars conducted during the upgrade to the 305-m Arecibo radio telescope. Our survey covered a total of 1147 deg<sup>2</sup> of sky using a drift-scan technique. We detected 33 pulsars, 10 of which were not known prior to the survey observations. The highlight of the new discoveries is PSR J0407+1607, which has a spin period of 25.7 ms, a characteristic age of 1.5 Gyr and is in a 1.8-yr orbit about a low-mass (> 0.2 M<sub>J</sub>) companion. The long orbital period and small eccentricity (e = 0.0009) make the binary system an important new addition to the ensemble of binary pulsars suitable to test for violations of the strong equivalence principle. We also report on our initially unsuccessful attempts to detect optically the companion to J0407+1607 which imply that its absolute visual magnitude is > 12.1. If, as expected on evolutionary grounds, the companion is an He white dwarf, our non-detection implies a cooling age of least 1 Gyr.

Key words: pulsars: general | pulsars: individual (J0407+1607).

## 1 INTRODUCTION

Recycled pulsars | neutron stars that are thought to have been spun up by the accretion of matter from a binary companion | are interesting objects to study. In addition to their application as excellent clocks for high-precision timing experiments (Davis et al. 1985), progress in our understanding of their origin and evolution continues to be made by large-area surveys of the radio sky.

Following Wołszczan's (1991) discovery of two recycled pulsars at high Galactic latitudes, Johnston & Bailes (1991) demonstrated that the local observable population of these objects should be isotropic, and that low-frequency (< 400 MHz) large-area surveys would be an excellent means of probing this population. This prediction was largely borne out by the discovery of 50 recycled pulsars from a number of large-area surveys undertaken in this frequency range by many groups around the world (see, e.g., Camilo 1995 for a

review). More recently, sensitive surveys at higher radio frequencies (> 1.4 GHz) continue to discover millisecond and recycled pulsars in large numbers at intermediate and low Galactic latitudes (Edwards et al. 2001; Faulkner et al. 2004) where interstellar propagation effects limit the Galactic volume surveyed at lower frequencies.

The 305-m Arecibo telescope has played a key role in low-frequency surveys. Between 1993 and 1998, the upgrade to the telescope meant that it was often parked at a fixed azimuth for extended periods. This provided a unique opportunity to survey the sky as it drifts through the stationary telescope beam at the sidereal rate. At 430 MHz, the sky coverage for a declination is  $60 \cos \delta$  deg<sup>2</sup> per day. As a number of groups were interested in drift-scan surveys (Foster et al. 1995; Camilo et al. 1996; Lommen et al. 2000; McLaughlin et al. 2002), the available sky was divided up into ve zones and the zenith angle of the telescope was changed daily to cover each region uniformly.

In this paper, we report on an analysis of data in the regions surveyed by the Space Telescope Science In-

<sup>?</sup> Email: Duncan.Lorimer@manchester.ac.uk

Table 1. Declination strips covered in the ST ScI/NA IC survey. For each strip we list the central zenith angle (ZA), gain (G), system temperature ( $T_{\text{sys}}$ ) of the 430-MHz line feed, approximate limiting flux density ( $S_{\text{min}}$ ), area of sky covered and the number of new and previously known pulsars detected (new + old).

range (deg)	ZA (deg)	G (K/Jy)	$T_{\text{sys}}$ (K)	$S_{\text{min}}$ (mJy)	Area (deg <sup>2</sup> )	Pulsars (new + old)
1{2	17	10.5	92	0.9	83	0+0
6{7	12	13.0	76	0.6	114	2+3
11{12	7	15.2	72	0.5	55	1+5
16{17	2	16.4	68	0.5	287	5+6
21{22	3	16.4	68	0.5	163	2+3
26{27	8	15.0	73	0.5	315	0+5
31{32	13	12.5	80	0.7	116	0+1
36{37	18	10.0	96	1.0	14	0+0

stitute and National Astronomy and Ionospheric Center" (ST ScI/NA IC) collaboration. A preliminary account of this work was given by Xilouris et al. (2000). In Section 2, we describe the survey observations and data analysis procedure. The results, follow-up timing and implications of the ten new pulsars found in the survey, including the new binary pulsar J0407+1607, are detailed in Sections 3 and 4.

## 2 SURVEY OBSERVATIONS AND RESULTS

The survey observations were carried out between April 1994 and March 1995 using the 430-MHz carriage house line feed. While the nominal sensitivity of this receiver for observations close to the zenith is  $16 \text{ K Jy}^{-1}$ , these observations were made at a variety of zenith angles and during periods when the feed was slightly out of focus due to the maintenance work. As a result, the gain of the telescope varied considerably in the range  $10\{16 \text{ K Jy}^{-1}$ . In drift-scan mode, a source passes through the 3 dB points of the primary beam (10 arcmin) in about 40 s. The declination ranges assigned to the ST ScI/NA IC collaboration, area coverage and approximate system parameters are summarised in Table 1. The sky coverage is shown in Fig. 1. An ASCII file containing the exact sky positions observed is available at <http://www.blackwellpublishing.com/products/journals/suppmat/MNR/MNR9005/MNR9005sm.htm>.

The incoming radio signals were sampled by the line feed as two orthogonal circular polarisations which, after amplification and down-conversion, were passed to a 32 250-kHz analogue filterbank spectrometer which sums the two polarisation channels prior to detection with an RC time constant of 333  $\mu$ s. The resulting total-power signals were sampled every 250  $\mu$ s before being digitised with 3-bit precision and written to magnetic tape for off-line analysis.

The data were processed in overlapping 33-s segments using a derivative of the software package developed for earlier Arecibo pulsar searches (Fruchter 1989; Nice 1992; Nice, Fruchter & Taylor 1995). In brief, the data were de-dispersed for 448 different assumed values of dispersion measure (DM) in the range  $0\{504 \text{ cm pc}^{-3}$ . The resulting time series were then Fourier transformed so that the amplitude spectra could be searched for periodic signals buried in the noise. To increase sensitivity to signals with narrow duty cycles (i.e. many harmonics in the Fourier domain) the spec-

Table 2. ST ScI/NA IC survey detection statistics. Listed are the 10 newly discovered pulsars (marked with asterisks), 23 previously known pulsars detected and 6 previously known pulsars which lie within 5 arcmin of a pointing which were not detected. For each pulsar, we give the period (P), DM, 400-MHz flux density ( $S_{400}$ ), offset ( $\theta$ ) from the beam centre, observed S/N and the predicted S/N. Blank S/N entries show the non-detections. Flux densities marked with a y have been estimated from catalogued  $S_{1400}$  values assuming a spectral index of 1.6.

P SR	P (m s)	DM (cm <sup>3</sup> pc)	$S_{400}$ (m Jy)	$\theta$ ( $^\circ$ )	S/N (obs)	S/N (pre)
J0137+1654*	414.7	27	1.4	1.5	16.7	13
J0329+1654*	893.3	42	0.6	1.3	14.9	12
J0407+1607*	25.7	36	10	0.8	48.7	45
J0609+2130*	55.6	39	0.8	1.2	13.6	7
B 0823+26	530.7	19	73	1.2	20.0	1802
B 0919+06	430.7	27	52	2.8	46.4	389
B 0940+16	1087.4	20	7	4.5	16.8	56
B 1530+27	1124.8	15	13	1.9	30.4	157
B 1534+12	37.9	12	36	4.2	46.5	238
J1549+2113*	1262.4	25	0.9	1.2	14.9	23
J1652+2651	915.8	41	11	1.8	23.2	104
J1822+1120*	1787.0	95	0.5	1.4	14.1	7
J1838+1650*	1901.9	34	2.6	1.2	21.0	23
J1848+0604*	2218.6	243	1.0	1.8	11.9	7
B 1900+06	673.5	503	22	7.5	9.8	11
J1900+0634	389.8	323	1.7y	3.8	{	6
B 1904+06	267.2	473	2.8	3.0	{	8
J1905+0616*	989.7	256	0.5	6.6	10.4	7
B 1907+10	283.6	150	50	5.5	27.0	103
J1909+0616	755.9	352	2.4y	6.7	7.6	3
B 1911+11	600.9	100	1	2.1	12.0	6
B 1913+16	59.0	169	4	1.8	12.7	22
B 1913+167	1616.2	63	5	1.3	25.7	44
B 1918+26	785.5	28	6	3.1	33.6	49
J1920+1110	509.8	182	2.8y	3.8	{	17
B 1924+16	579.8	177	8	1.6	17.8	68
B 1929+10	226.5	3	303	8.4	46.4	267
B 1933+16	358.7	158	242	9.4	25.8	150
B 1937+21	1.558	71	240	4.3	30.5	181
J1943+0609	446.2	71	{	2.9	15.6	{
J1946+2611	435.0	165	2	4.6	{	5
J1951+1123	5094.0	31	1	1.8	{	27
J1953+1149	851.8	140	2	4.4	11.8	19
J2002+1637	276.4	94	0.4	2.3	18.1	4
J2040+1657*	865.5	51	0.6	1.4	18.4	7
J2156+2618	498.1	49	2.7	5.6	10.2	21
J2234+2114	1358.7	35	2.6	5.8	12.6	10
B 2303+30	1575.8	50	24	8.7	19.0	56
B 2315+21	1444.6	21	15	2.8	24.8	217

tra were incoherently summed so that up to 2, 3, 4, 8 and 16 harmonics were combined. Signal-to-noise ratios (S/Ns) were computed from the amplitude of each point in the spectrum and the local root-mean-square deviation. The Fourier components of the 100 strongest events were then inverse transformed back into the time domain to compute the S/N




Figure 1. Survey sky coverage equatorial coordinates showing new discoveries (filled triangles) previously known pulsars detected by the survey (circles) and previously known pulsars not detected (squares). Dotted lines show constant Galactic latitudes of  $20^\circ$ .

of the resulting pulse profile. The profiles, periods, S/Ns and best DMs of each candidate were saved for visual inspection.

The sensitivity of the survey was dependent on the telescope's zenith angle and hence varied for the different declination strips. In Table 1 we quote our theoretical minimum detectable flux density ( $S_{\text{min}}$ ) to long-period pulsars with duty cycles of 4% using the radiometer equation (see, e.g., Lorimer & Kramer 2005). This calculation assumes a sky background temperature of 30 K and that the pulsar transits through the beam centre. These values should therefore be taken as lower limits to the true survey sensitivity which degrades for larger background temperatures, pulse widths and shorter pulse periods. Typical values for millisecond pulsars, for example, are expected to be  $> 3$  mJy. A more detailed discussion on the sensitivity of the observing system can be found in Camilo, Nice & Taylor (1996).

The data analysis resulted in the detection of 33 pulsars in  $1147 \text{ deg}^2$  of sky. Ten of these pulsars were previously known prior to the survey observations. In Table 2, we list the S/Ns of the pulsars along with an estimate of the theoretical S/N based on the nominal survey parameters and accounting for position offsets from the centre of the (assumed Gaussian) telescope beam power pattern. In general, for S/Ns below 20, the observed and theoretical values are in reasonable agreement. For some of the bright pulsars (e.g. B0823+06), the detected S/Ns are much less than the theoretical values. We are unsure of the reasons for this, and, unfortunately, the raw data are no longer available. Given our short integration times, it is possible that the observations were affected by adverse scintillation and/or nulling.

Also included in Table 2 are 6 previously known pulsars that lie within 5 arcmin of the searched area but were

not detected. Although one of these pulsars, J1909+0616, made the list of candidates with a S/N = 7.6, this was below the nominal survey threshold S/N of 9 and does therefore not count as a blind detection. Only two of the remaining pulsars have theoretical S/Ns above this threshold. For PSR J1920+1110, we are unsure of its 430-MHz flux density which has been extrapolated from the 1400-MHz value (Morris et al. 2002) assuming a spectral index of 1.6. As this pulsar was not detected in more sensitive 430-MHz observations by Bhat et al. (2004), either the spectrum is flatter than assumed or the pulse is significantly affected by scattering at 430 MHz. The remaining pulsar which is nominally above our detection threshold, J1951+1123, has a period of 5.1 s and is known to null for several periods (Nice 1992). This is the most likely explanation for our non-detection.

In summary, given the uncertainties in pointing and unknown variations in received flux during these observations, this simple analysis of the detection statistics shows that our survey sensitivity is consistent with that reported by other groups (see, e.g., Camilo et al. 1996).

### 3 FOLLOW-UP OBSERVATIONS

Confirmation and follow-up observations of the new discoveries began at end of the Arecibo upgrade in 1998. These made use of the Penn State Pulsar Machine (PSPM; Cadwell 1997), a 128 60-kHz channel analogue Interbank spectrometer which records either continuously-sampled data (‘search mode’) or synchronously-averaged pulse profiles at the predicted pulse period (‘timing mode’).

As in the original survey observations, we used the 430-

Table 3. Parameters for the nine isolated pulsars derived from the TEMPO timing analysis. Figures in parentheses are 1- $\sigma$  uncertainties in the least-significant digits as reported by TEMPO. The arrival times from which these ephemerides are derived are freely available on-line as part of the European Pulsar Network (EPN) database (<http://www.jb.man.ac.uk/pulsar/Resources/epn>).

PSR	R.A. (J2000) (h:m:s)	Dec. (J2000) ( $^{\circ}$ : $'$ : $''$ )	Period, P (s)	Epoch MJD	P- ( $\times 10^{15}$ )	DM ( $\text{cm}^{-3}\text{pc}$ )	MJD range
J0137+1654	01:37:23.88 (1)	+16:54:42.1 (4)	0.4147630265082 (5)	52225	0.01223 (3)	26.6 (4)	51329{53119
J0329+1654	03:29:08.55 (3)	+16:54:02 (2)	0.893319660649 (1)	52225	0.21525 (6)	42.1 (2)	51329{53119
J0609+2130	06:09:58.8861 (2)	+21:30:2.84 (9)	0.05569801392951 (2)	52758	0.000235 (2)	38.73 (2)	52395{53119
J1549+2113	15:49:40.941 (5)	+21:13:26.9 (1)	1.262471311613 (5)	51738	0.8541 (8)	24.8 (2)	51375{52045
J1822+1120	18:22:14.60 (1)	+11:20:56.2 (2)	1.78703680542 (4)	51602	2.346 (4)	95.2 (6)	51330{51873
J1838+1650	18:38:43.062 (4)	+16:50:16.05 (6)	1.90196739931 (1)	51602	2.677 (1)	33.9 (2)	51330{51873
J1848+0604	18:48:54.622 (7)	+06:04:46.8 (3)	2.21860264699 (3)	51602	3.736 (6)	242.7 (7)	51330{51873
J1905+0616	19:05:06.849 (5)	+06:16:16.7 (1)	0.989706023196 (7)	51631	135.232 (1)	256.05 (1)	51330{51931
J2040+1657	20:40:17.865 (3)	+16:57:30.46 (7)	0.865606225032 (4)	51602	0.5949 (6)	50.7 (2)	51330{51873




Figure 2. Integrated 430-MHz pulse profiles for nine of the newly discovered pulsars derived from a phase-coherent addition of the individual PSPM observations. Each profile shows 360 degrees of rotational phase. These profiles are freely available as part of the EPN database. A pulse profile for PSR J0609+2130 has already been published by Lorimer et al. (2004) and is not included here.

MHz line-feed receiver for the bulk of our follow-up work. Following initial confirmation observations which verified the existence of each pulsar in search mode, we generated a preliminary ephemeris which was subsequently used to obtain timing-mode data. The basic observing and analysis procedures used were identical to those described by Lorimer, Camilo & Xilouris (2002).

In order to determine accurate spin and astrometric parameters for each pulsar, we carried out a dedicated campaign of observations using the PSPM in timing mode. For each observation the set of 128 folded pulse profiles (one profile per frequency channel) was appropriately delayed with respect to the centre observing frequency to account for the effects of interstellar dispersion. Standard software tools (Lorimer 2001) implemented this procedure to form a de-dispersed time-tagged integrated pulse profile for each observation. A pulse time of arrival (TOA) for each profile was then obtained by cross-correlating it in the Fourier domain with a high S/N template profile formed from the

sum of many individual observations (Taylor 1992). The final template profiles are shown in Fig. 2.

The TOAs obtained for each pulsar were transformed to equivalent arrival times at the solar system barycentre and fitted to a simple spin-down model using the TEMPO<sup>1</sup> software package. After several iterations, an initial ephemeris was obtained which was then used to form an improved template profile from the sum of each pulse profile appropriately phase-aligned according to the timing model. The resulting high S/N template was then used to obtain TOAs of higher precision which were analysed with TEMPO in a further iteration to produce the final timing solution. Refined values of DM for each pulsar were obtained by forming four 1.92-MHz-wide sub-bands from each PSPM profile. The DM was derived from a fit to the TOAs from each sub-band keeping the spin and astrometric parameters fixed at the final timing solution values. For nine of the new pulsars, the TOAs were adequately fitted using an isolated pulsar spin-down model and the resulting parameters from each fit are presented in Table 3.

In the time since the original survey observations and follow-up, five of the new pulsars have been independently discovered in other surveys. The binary pulsar J0407+1607 discussed below was detected during a survey of gamma-ray error boxes at Parkes (Roberts et al. 2004). PSRs J0609+2130 (Lorimer et al. 2004) and J1549+2113 (Foster et al. 1995; Lewandowski et al. 2004) were discovered in contemporary 430-MHz Arecibo drift-scan surveys. Finally, PSRs J1848+0604 (Faulkner, private communication) J1905+0616 (Morris et al. 2002) were detected by the Parkes multibeam survey. PSRs J1549+2113 and J1848+0604 have no previously published timing solution and are presented for the first time in Table 3. For PSR J0609+2130, we have added an additional year of PSPM TOAs to those already published by Lorimer et al. (2004). The longer time baseline of the resulting ephemeris presented in Table 3 has significantly improved the precision of the derived parameters. For PSR J1905+0616, we have included Jodrell Bank TOAs collected at 1412 MHz by Morris et al. (2002) in our final

<sup>1</sup> <http://pulsar.princeton.edu/tempo>

Table 4. Measured parameters for PSR J0407+1607 from 71 TOAs from MJDs 52304 (January 30, 2002) to 53292 (October 14, 2004). The numbers in parentheses are the 1 $\sigma$  TEMPO uncertainties in the least significant digit quoted.

Parameter	Value
Right ascension (J2000) (h:m:s)	04:07:54.939(3)
Declination (J2000) ( $^{\circ}$ : $^{\prime}$ : $^{\prime\prime}$ )	16:07:16.4(2)
Spin period, $P$ (ms)	25.70173919463(2)
Period derivative, $\dot{P}$ ( $10^{-20}$ )	7.9(3)
Epoch of period (MJD)	52799
Dispersion measure, DM ( $\text{cm}^{-3}$ pc)	35.65(2)
Projected semimajor axis, $x$ (s)	106.45026(2)
Orbital period, $P_b$ (days)	669.0704(1)
Orbital eccentricity, $e$	0.0009368(6)
Longitude of periastron, $\Omega$ (deg)	291.74(2)
Epoch of periastron, $T_0$ (MJD)	52774.69(3)

timing analysis. This has resulted in an improvement on the ephemeris published by Morris et al.

The most interesting of the new discoveries is PSR J0407+1607. The intermediate 25.7-ms spin period of this pulsar suggested that it was either a young or recycled object. Following initially unsuccessful attempts to fit an isolated spin-down model to the data, sets of closely-separated TOAs were used to obtain estimates of the barycentric pulse period for a given epoch. These periods were then fit to a simple Keplerian model which revealed 1.8-yr sinusoidal variations implying that the pulsar was a member of a long-period circular orbit binary system.

While our early PSPM timing-mode observations could be used to obtain a preliminary orbit ephemeris, they were not of sufficient quality or density to obtain a phase-coherent timing solution. In January 2002, we began a regular observing campaign to collect data with the PSPM in search mode. These data were then folded  $\phi$ -line to produce a set of pulse profiles and hence TOAs for a timing analysis. The advantage of this approach is that the data could be re-folded later when the precision of the orbital parameters improved.

Despite this improved approach, small uncertainties in the assumed position resulted in significant covariances between the orbital parameters and positions in TEMPO. Fortunately, as mentioned above, this pulsar was independently discovered by Roberts et al. (2004). Follow-up 1400-MHz Arecibo observations from that survey resulted in a positional determination which was used as a starting point for the final timing analysis. The final phase-coherent timing solution presented in Table 4 is the result of a simultaneous fit for spin, astrometric and binary parameters. The observed minus computed TOA residuals for this solution are featureless with a root-mean-square value of 16  $\mu$ s. The DM was determined by including 1400-MHz observations of the pulsar with the Wideband Arecibo Pulsar Processor (Dowd, Sisk & Hagen 2000) and, as for the other pulsars, keeping all other model parameters constant in the TEMPO fit.

To obtain an estimate of the transverse speed of J0407+1607, we performed a scintillation analysis on an observation at 430 MHz using the Arecibo Observatory Fourier Transform Machine (AOFTM), a fast-dump digital spectrometer which provides 1024 frequency channels across a




Figure 3. Upper panel: full Stokes profile for PSR J0407+1607 at 430 MHz showing total intensity (solid line), linear polarisation (dashed) and circular polarisation (dotted) as a function of pulse phase. Lower panel: position angle versus pulse phase.

10-MHz bandwidth (see, e.g., McLaughlin et al. 2002). Performing a standard autocorrelation analysis on the pulsed intensity as a function of time and radio frequency (for details of this technique see Lorimer & Kramer 2005) we find the characteristic timescale and bandwidth of scintillation to be 843  $\pm$  6 s and 190  $\pm$  2 kHz respectively. Adopting the expressions given by Cordes & Rickett (1998) for a uniform scattering medium, and taking a distance of 1.3  $\pm$  0.2 kpc derived from the DM and the Cordes & Lazio (2002) electron density model, we estimate the transverse speed of PSR J0407+1607 to be 35  $\pm$  5 km s $^{-1}$ . Although this result is model dependent, it implies a transverse speed which lies at the low end of the distribution known for millisecond pulsars (Nice & Taylor 1995). In future it should be possible to measure the proper motion of this pulsar directly by extending the existing baseline of timing observations. The predicted proper motion at this distance is 6  $\pm$  1 mas yr $^{-1}$ . Through simulations with TEMPO, we estimate that a direct proper motion measurement of this magnitude through timing techniques will be possible within 3 yr. The effect of proper motion on the observed semimajor axis should also be measurable on a similar timescale, allowing us to place constraints on the orbital inclination angle in future.

The PSPM and AOFTM data do not provide any information about the polarisation properties of PSR J0407+1607. To investigate this, in March 2002, we observed the pulsar for 78 min at 430 MHz using the Princeton Mark IV data recording system (Stairs et al. 2000) in narrow-band (5 MHz) mode. The Mark IV data were coherently de-dispersed  $\phi$ -line and the data were calibrated using observations with a noise diode of known flux density and polarisation. The total intensity, linear and circularly polarised pulse profiles are shown in Fig. 3. Relative to the total intensity, we find that both the linear polarisation

Table 5. Observed and derived parameters for the ten pulsars discovered in the survey. For each pulsar, we list the Galactic longitude ( $l$ ) and latitude ( $b$ ), the flux density at 430 MHz ( $S_{430}$ ), the pulse width measured at 50 and 10% of the peak intensity (respectively  $W_{50}$  and  $W_{10}$ ), the equivalent width of a top-hat pulse having the same area and peak intensity of the profile ( $W_{eq}$ ), the distance derived from the dispersion measure assuming the Cordes & Lazio (2002) electron density model ( $D$ ), as well as the base-10 logarithms of the 430-MHz luminosity ( $L_{430} = S_{430} D^2$ ), the characteristic age ( $\tau = P/(2\dot{P})$ ), the surface dipole magnetic field ( $B = 3.2 \times 10^9 (P\dot{P})^{1/2}$  G) and the rate of loss of spin-down energy ( $\dot{E} = 3.95 \times 10^{46} P\dot{P}^3$  erg s $^{-1}$ ).

PSR	$l$ deg	$b$ deg	$S_{430}$ m Jy	$W_{50}$ ms	$W_{10}$ ms	$W_{eq}$ ms	$D$ kpc	$\log_{10}$ [ $L_{430}$ (m Jy kpc $^2$ )]	$\log_{10}$ [ $\tau$ (yr)]	$\log_{10}$ [ $B$ (G)]	$\log_{10}$ [ $\dot{E}$ (erg s $^{-1}$ )]
J0137+1654	138.3	{44.6	1.4	14	62	20	1.2	0.3	8.73	10.9	30.8
J0329+1654	168.5	{31.7	0.6	17	37	23	1.8	0.3	7.82	11.6	31.1
J0407+1607	176.5	{25.8	10.2	5.3	6.6	4.8	1.3	1.3	9.71	9.16	32.3
J0609+2130	189.2	{1.0	0.8	1.6	7.4	2.4	1.2	0.06	9.57	9.56	31.7
J1549+2113	34.8	{49.3	0.9	7	22	15	2.1	0.6	7.37	12.0	31.2
J1822+1120	39.8	{11.6	0.5	25	59	27	3.8	0.9	7.08	12.3	31.2
J1838+1650	46.6	{10.5	2.6	72	92	39	2.0	1.0	7.05	12.4	31.2
J1848+0604	37.9	{3.5	1.0	46	73	40	7.0	1.7	6.97	12.5	31.1
J1905+0616	40.1	{0.1	0.5	21	46	24	5.7	1.2	5.06	13.1	33.7
J2040+1657	61.2	{14.8	0.6	16	79	31	3.0	0.7	7.36	11.9	31.6

and modulus of the circularly polarised component have an average value of 13%, typical of other millisecond pulsars (Xilouris et al. 1998; Stairs, Thorsett & Camilo 1999). Also shown in Fig. 3 is the position angle of the linear polarisation as a function of pulse phase. As for many other millisecond pulsars the position angle does not follow a smooth S-shape variation expected from the rotating-vector model of Radhakrishnan & Cooke (1969). A determination of the emission geometry of this pulsar is, therefore, non-trivial.

#### 4 DISCUSSION

Various observed and derived parameters of the new pulsars are summarised in Table 5 where we also include estimates of each pulsar's 430-MHz flux density. These were derived from the PSPM profiles using identical procedures to those described by Lorimer et al. (2002).

##### 4.1 General remarks

In common with results from other Arecibo drift-scan surveys (Foster et al. 1995; Camilo et al. 1996; Lommen et al. 2000; McLaughlin et al. 2002), the new pulsars are relatively faint, nearby objects. Their inclusion in future population syntheses should provide valuable constraints on the low end of the radio pulsar luminosity function. In 1147 deg $^2$  of sky searched, we detected five recycled pulsars: J0407+1607, J0609+2130, B1534+12, B1913+16 and B1937+21. Of the two new discoveries, the binary pulsar J0407+1607 is discussed below, while the isolated pulsar J0609+2130 was already discussed in detail by Lorimer et al. (2004). We simply note here that our detection rate of one recycled pulsar per 230 deg $^2$  is consistent with that of other Arecibo surveys (see Camilo 1995).

The youngest pulsar in our sample, with a characteristic age of 116 kyr is PSR J1905+0616. As noted and discussed by Kramer et al. (2003), this object lies within the bounds of the EGRET source 3EG J1903+0550. An association is unlikely simply due to the large density of pulsars in this

region and hence high probability of a chance alignment. In addition, as noted by Kramer et al., the spin-down energy loss of J1905+0616 is not large enough to be a likely source of the gamma-ray emission seen.

##### 4.2 PSR J0407+1607 and 3EG J0407+1610

PSR J0407+1607 is coincident with the unidentified EGRET source 3EG J0407+1610 (Hartman et al. 1999), a relatively weak gamma-ray source with mean flux  $F = 2.4 \times 10^{-10}$  erg cm $^{-2}$  s $^{-1}$  above 100 MeV (assuming a photon spectral index of 2). While the variability index (see McLaughlin et al. 1996 for a definition) is only 0.19, and consistent with other EGRET sources known to be associated with pulsars, our  $P$ -measurement of J0407+1607 implies a rate of loss of spin-down energy  $\dot{E} = 1.9 \times 10^{32}$  erg s $^{-1}$ . This can be used to test the validity of an association with 3EG J0407+1610 by calculating the required efficiency of conversion of the pulsar's spin-down luminosity into gamma rays. At the DM-derived distance of 1.3 kpc, and assuming a gamma-ray beaming fraction of  $f = 4$  sr, we find an incompatible value of  $\eta = (F D^2 / \dot{E}) = 20$ . A much smaller distance and/or beaming fraction would be required so that  $\eta < 1$ . We therefore conclude that PSR J0407+1607 is most likely not the source of the gamma rays from 3EG J0407+1610.

##### 4.3 PSR J0407+1607 and SEP violations

The orbital parameters of the J0407+1607 binary system are consistent with standard models of binary evolution (see, e.g., Phinney & Kulkarni 1994) in which the radio pulsar has been spun up by the accretion of matter while its companion star was in a red giant phase. Indeed, the observed orbital eccentricity is in excellent agreement with the prediction from Phinney's (1992) relationship between eccentricity  $e$  and orbital period  $P_b$ .

Millisecond pulsar-white dwarf binaries like PSR J0407+1607 make excellent probes of the strong equivalence principle (SEP). The SEP states that free fall

of a body is completely independent of its gravitational self energy. If the SEP is violated, then the ratio of inertial mass to gravitational mass of a falling body differs from unity by an amount  $\epsilon$ . As originally proposed by Damour & Schäfer (1991), and by analogue with Nordvedt's earlier tests of the SEP in the Earth-Moon system (Nordvedt 1968a,b), a violation of the SEP for a pulsar-white dwarf binary would mean that both objects "fall" in the local Galactic gravitational field at a different rate. As a result, a violation would cause a polarisation of the orbital plane toward and parallel to the centre of the Galaxy.

Following Damour & Schäfer's early work, Wex (1997, 2000) derived limits on  $\epsilon$  from an ensemble of binary pulsar-white dwarf systems. As the cure-of-misfit for this calculation is  $P_b = e^2$ , long-period circular orbit binaries like J0407+1607 are particularly important members of the ensemble used in the test. Recently, a robust Bayesian analysis by Stairs et al. (2005) which uses all known binary pulsars relevant for this test including J0407+1607 and a limit on SEP violations of  $\epsilon < 5.5 \times 10^{-3}$ . This new limit is comparable to the weak-field limits found in the Earth-Moon system (Nordvedt 1968a,b).

#### 4.4 The companion to PSR J0407+1607

The observed mass function of PSR J0407+1607 is  $2.89 \times 10^{-3} M_\odot$ . Assuming a mass for the pulsar in the range  $1.3\{1.6 M_\odot$ , the implied companion mass is  $0.2 M_\odot / \sin i$ , where  $i$  is the unknown inclination angle between the plane of the orbit and the plane of the sky. From extensive studies of other millisecond pulsar binaries with similar orbital characteristics to J0407+1607 (for a recent review, see van Kerkwijk et al. 2005) it is most likely that the companion to J0407+1607 is a low-mass white dwarf. An inspection of archival Palomar data in the digitized sky survey archive<sup>2</sup> reveals no optical counterpart down to the plate limit (approximately 19th magnitude), with the closest object in the field being 8 arcsec (50 $\mu$ ) offset from the radio position given in Table 4.

To search for a fainter optical counterpart, the field surrounding PSR J0407+1607 was observed on October 2003 with the 2.3-m Bok telescope at Kitt Peak using the Steward Observatory imaging camera in combination with the Harris Band Filters. Four 27-min B and four 20-min V images were obtained under non-ideal seeing (1.5 arcsec) conditions. All images were bias-subtracted and flat-fielded using the IRAF package<sup>3</sup>. For photometry calibration, we selected all stars in a  $3 \times 3$  arcmin box around the pulsar location from the USNO-B1.0 Catalog<sup>4</sup>.

The detection limit was determined to be magnitude 21 in B. This is in good agreement with the 20.9 magnitude of the star identified as 1061-0042921 in the USNO-B1.0 catalogue detected at the noise level in our images. Four 20-min exposures were aligned and added to produce the final image. We used the published visual magnitude of the nearby stars SA 94 171 and GD 71 (Landoit 1992) to determine the relationship between instrumental and aperture magni-

tudes. From this procedure, we estimate a limiting apparent visual magnitude for detection of 23.

At the nominal distance of  $1.3 \pm 0.2$  kpc, our upper limit corresponds to an absolute visual magnitude of at least 12.1. Assuming the companion to be an H white dwarf, we can use the cooling curves computed by Hansen & Phinney (1998) to place a lower limit on the age of the white dwarf companion. From an inspection of Fig. 13 in their paper, our current non-detection constrains the cooling age to be  $> 1$  Gyr, compatible with the 1.5 Gyr characteristic age of J0407+1607. More stringent constraints should be available soon following deeper observations of this system now underway with the Palomar telescope which will be described elsewhere.

#### ACKNOWLEDGEMENTS

The Arecibo observatory, a facility of the National Astronomy and Ionosphere Center, is operated by Cornell University in a co-operative agreement with the National Science Foundation (NSF). We thank Alex Wolszczan for making the PSPM freely available for use at Arecibo, as well as Paulo Freire, Jim Sheckard, Andrea Lommen, Zaven Arzoumanian, Ramesh Bhat and David Champion for assistance with the PSPM timing observations. DRL is a University Research Fellow funded by the Royal Society. JHS holds an NSERC UFA and is supported by a Discovery Grant. FC is supported by NASA and the NSF. The Digitized Sky Survey was produced at the Space Telescope Science Institute under U.S. Government grant NAG W-2166. The images of these surveys are based on photographic data obtained using the Oschin Schmidt Telescope on Palomar Mountain and the UK Schmidt Telescope. The plates were processed into the present compressed digital form with the permission of these institutions.

#### REFERENCES

- Bhat, N. D. R., Cordes, J. M., Camilo, F., Nice, D. J., Lorimer, D. R., 2004 *ApJ*, 605, 759  
 Cadwell B. J., 1997, PhD thesis, Penn State University  
 Camilo F., 1995, in *Aspar A., Kiziloglu U., van Paradis J., eds, The Lives of the Neutron Stars (NATO ASI Series)*, Kluwer, Dordrecht, 243  
 Camilo F., Nice D. J., Shrauner J. A., Taylor J. H., 1996, *ApJ*, 469, 819  
 Camilo F., Nice D. J., Taylor J. H., 1996, *ApJ*, 461, 812  
 Cordes J. M., Lazio T. J. W., 2002, *astro-ph/0207156*  
 Cordes J. M., Rickett B. J., 1998, *ApJ*, 507, 846  
 Damour T., Schäfer G., 1991, *Phys. Rev. Lett.*, 66, 2549  
 Davis M. M., Taylor J. H., Weisberg J. M., Backer D. C., 1985, *Nature*, 315, 547  
 Dowd, A., Sisk, W., Hagen, J., 2000, in *Kramer M., Wex N., Wielebinski R., eds, 2000, Pulsar Astronomy - 2000 and Beyond*, IAU Colloquium 177 Astronomical Society of the Pacific, San Francisco, 275  
 Edwards R. T., Bailes M., van Straten W., Britton M. C., 2001, *MNRAS*, 326, 358  
 Faulkner A. J., Stairs I. H., Kramer M., Lyne A. G., Hobbs G., Possenti A., Lorimer D. R., Manchester R. N.,

<sup>2</sup> <http://www-gsss.stsci.edu/DSS>

<sup>3</sup> <http://iraf.noao.edu>

<sup>4</sup> <http://www.nofs.navy.mil/data/fhpix>

- McLaughlin M. A., D'Amico N., Camilo F., Burgay M., 2004, *MNRAS*, 355, 147
- Foster R. S., Cadwell B. J., Wolszczan A., Anderson S. B., 1995, *ApJ*, 454, 826
- Fruchter A. S., 1989, PhD thesis, Princeton University
- Hansen B. M. S., Phinney E. S., 1998, *MNRAS*, 294, 557
- Hartman R. C. et al. S. D. B., 1999, *ApJS*, 123, 79
- Johnston S., Bailes M., 1991, *MNRAS*, 252, 277
- Kramer M. et al. 2003, *MNRAS*, 342, 1299
- Landolt A. U., 1992, *AJ*, 104, 372
- Lewandowski W., Wolszczan A., Feiler G., Konacki M., Soltyński T., 2004, *ApJ*, 600, 905
- Lommen A. N., Zepka A., Backer D. C., McLaughlin M., Cordes J. M., Arzoumanian Z., Xilouris K., 2000, *ApJ*, 545, 1007
- Lorimer D. R., 2001, SIGPROC-v1.0: (Pulsar) Signal Processing Programs, Arecibo Technical Memo No. 2001{01 <http://www.jb.man.ac.uk/~drl/sigproc>
- Lorimer D. R., Camilo F., Xilouris K. M., 2002, *AJ*, 123, 1750
- Lorimer D. R., McLaughlin M. A., Arzoumanian Z. A., Xilouris K., Cordes J. M., Lommen A. N., Fruchter A. S., Chandler A. M., Backer D. C., 2004, *MNRAS*, 347, L21
- Lorimer, D. R. & Kramer, M. 2005, *Handbook of Pulsar Astronomy*. Cambridge University Press
- McLaughlin M. A., Mattox J. R., Cordes J. M., Thompson D. J., 1996, *ApJ*, 473, 763
- McLaughlin M. A., Arzoumanian Z., Cordes J. M., Backer D. C., Lommen A. N., Lorimer D. R., Zepka A. F., 2002, *ApJ*, 564, 333
- Morris D. J. et al. 2002, *MNRAS*, 335, 275
- Nice D. J., 1992, PhD thesis, Princeton University
- Nice D. J., Fruchter A. S., Taylor J. H., 1995, *ApJ*, 449, 156
- Nice D. J., Taylor J. H., 1995, *ApJ*, 441, 429
- Nordtvedt K., 1968a, *Phys. Rev.*, 169, 1014
- Nordtvedt K., 1968b, *Phys. Rev.*, 170, 1186
- Phinney E. S., 1992, *Philos. Trans. Roy. Soc. London A*, 341, 39
- Phinney E. S., Kulkarni S. R., 1994, *Ann. Rev. Astr. Ap.*, 32, 591
- Radhakrishnan V., Cooke D. J., 1969, *Astrophys. Lett.*, 3, 225
- Roberts M., Ransom S., Hessels J., Livingstone M., Tam C., Kaspi V., Crawford F., 2004, in Camilo F. & Gaensler B. eds, *Young Neutron Stars and Their Environments*, IAU Symposium 218 Astronomical Society of the Pacific, San Francisco, 415
- Stairs I. H. et al. 2005, *ApJ*, submitted
- Stairs I. H., Splaver E. M., Thorsett S. E., Nice D. J., Taylor J. H., 2000, *MNRAS*, 314, 459
- Stairs I. H., Thorsett S. E., Camilo F., 1999, *ApJS*, 123, 627
- Taylor J. H., 1992, *Philos. Trans. Roy. Soc. London A*, 341, 117
- van Kerkwijk M. H., Bassa, C. G., Jacoby, B. A., Jonker, P. G., 2005, in Rasio F. & Stairs I. H., eds, *Binary Pulsars*, Astronomical Society of the Pacific, San Francisco, 328, 357
- Wex N., 1997, *A & A*, 317, 976
- Wex N., 2000, in Kramer M., Wex N., Więbowski R., eds, 2000, *Pulsar Astronomy - 2000 and Beyond*, IAU Colloquium 177 Astronomical Society of the Pacific, San Francisco, 113
- Wolszczan A., 1991, *Nature*, 350, 688
- Xilouris K. M., Fruchter A., Lorimer D. R., Eder J., Vazquez A., 2000, in Kramer M., Wex N., Więbowski R., eds, 2000, *Pulsar Astronomy - 2000 and Beyond*, IAU Colloquium 177 Astronomical Society of the Pacific, San Francisco, 21
- Xilouris K. M., Kramer M., Jessner A., von Hoensbroech A., Lorimer D., Więbowski R., Wolszczan A., Camilo F., 1998, *ApJ*, 501, 286