# ON THE ECCENTRICITIES AND MERGER RATES OF DOUBLE NEUTRON STAR BINARIES AND THE CREATION OF \DOUBLE SUPERNOVAE".

H.K.Chaurasia Swinburne University of Technology

M .Bailes Sw inburne University of Technology D raft version April 11, 2024

#### ABSTRACT

We dem onstrate that a natural consequence of an asymmetric kick imparted to neutron stars at birth is that the majority of double neutron starbinaries should possess highly eccentric orbits. This leads to greatly accelerated orbital decay, due to the enormous increase in the emission of gravitational radiation at periastron as originally demonstrated by Peters (1964). A uniform distribution of kick velocities constrained to the orbital plane would result in 24% of surviving binaries coalescing at least 10,000 times faster than an unperturbed circular system. Even if the planar kick constraint is lifted,

6% of bound system s still coalesce this rapidly. In a non-negligible fraction of cases it may even be possible that the system could coalesce within 10 years of the nal supernova, resulting in what we might term a \double supernova". For system s resembling the progenitor of PSR J0737{3039A, this number is as high as 9% (in the planar kick model). Whether the kick velocity distribution extends to the range required to achieve this is still unclear. We do know that the observed population of binary pulsars has a decit of highly eccentric system s at small orbital periods. In contrast, the long-period system s favour large eccentricities, as expected. We argue that this is because the short-period highly eccentric system s have already coalesced and are thus selected against by pulsar surveys. This elect needs to be taken into account when using the scale-factor method to estim ate the coalescence rate of double neutron star binaries. We therefore assert that the coalescence rate of such binaries is underestim ated by a factor of several.

Subject headings: pulsars; binary, gravitational radiation, m erger rates

#### 1. INTRODUCTION

Beginning with the detection of the Hulse & Taylor double neutron star (Hulse & Taylor 1975), pulsar astronom ers have unveiled a sm all but growing sam ple of double neutron star binaries over the last thirty years. Owing to the extrem e regularity of their radio pulse em issions, these exotic objects have proven to be invaluable scienti c laboratories, particularly for tests of general relativity, as they are some of the few known objects for which the e ects of gravitational radiation are measurable. In fact, the rst con m ation that gravitational radiation existed was through measurements of the orbital decay of the rst known double neutron star (Taylor & W eisberg 1989).

Recent surveys have started to unveil a better picture of the double neutron star population. The Parkesm ultibeam surveys have discovered three probable double neutron star systems (Lyne et al. 2000; Burgay et al. 2003; Faulkner et al. 2005), and A recibo surveys have detected two m ore (W olszczan 1991; C ham pion et al. 2004). A lthough we are still dealing with sm all num ber statistics, the population is now large enough to see trends em erging.

A phenom enon of crucial in portance to the binary coalescence rate is the kick velocity in parted to a neutron star during its supernova, a dom inant factor in determ ining the eccentricity and coalescence time of the subsequent orbit (H ills 1983; D ewey & C ordes 1987; B ailes 1989). B inary neutron star form ation and evolution also depends greatly upon this little-understood supernova kick, since a double neutron star system m ust survive two such kicks in order to rem ain bound. To some extent, know ledge of the kick velocity distribution can be extracted statistically from m easurem ents of the observed sam ple of double neutron stars (W illem s & K alogera 2004). The m agnitude of the kicks can be inferred by the space velocities of pulsars, which are an order of m agnitude higher than that of their progenitors, the O B stars. Investigations of the pulsar population dem onstrate that the average kick m ust be of order  $400\{500 \text{ km s}^1$  (Lyne & Lorim er 1994; C ordes & Chemo 1998).

Further evidence for pulsar kicks com es from precession e ects observed in the binary pulsar PSR J0045{ 7319 (K aspi et al. 1996) and spin-orbit coupling in the relativistic binaries PSR B1913+16, PSR B1534+12 and PSR J1141{6545 (W eisberg et al. 1989; Stairs et al. 2004; H otan et al. 2004). A ssum ing tidal forces and m ass transfer align the spin and angular m om entum axes prior to the ultim ate supernova explosion, these system s suggest a kick out of the orbital plane has occurred.

B inary pulsars are in some sense fossil records of the progenitor system from which they evolved. We date these systems by the characteristic age of the pulsar, and can therefore extrapolate back to the orbital conditions immediately after the nal supernova explosion. What prohibits us from going further back in time to derive a unique progenitor is the possibility that the new ly-born

E lectronic address: hem ant.chaurasia@ gm ail.com

E lectronic address: m bailes@ sw in edu au

pulsar received an impulsive kick at birth. Since the kick velocity distribution is largely unknown, including its orientation with respect to any reference point, such as the pulsar spin axis, we are left with a rather unsatisfactory fam ily of potential progenitors for any observed neutron star binary.

The relativistic nature of these objects has brought attention to the galactic rate of double neutron star coalescence { the m erger event at the clim ax of a long orbital decay and inspiral due to gravitational radiation. Such events are some of the m ost signi cant anticipated sources of gravitational waves, detectable by interferom eters such as the Laser Interferom eter G ravity-W ave O bservatory (LIGO) (A bram oviciet al. 1992). D ouble neutron star coalescence is also considered a m a pr candidate for explaining the origin of a subset of the mysterious cosm ic gam m a-ray bursts observed by gam m a-ray observatories such as the BAT SE instrum ent aboard the C om pton G am m a-Ray O bservatory (CGRO) (M eegan et al. 1992).

M uch research has been conducted into predicting the double neutron star coalescence rate, using M onte C arlo population synthesis m odels (B elczynski et al. 2002; Tutukov & Yungelson 1993; Lipunov et al. 1997; Portegies Zwart & Spreeuw 1996) or by statistical extrapolation from the observed sam ple (C urran & Lorim er 1995; K alogera et al. 2001; K alogera et al. 2004). M onte C arlo m ethods are currently ham pered by the vastness of the population param eter space and com pounded uncertainties, and statistical m ethods su er from the sm all sam ple size of known double neutron stars.

It is custom any to estim ate the event rate that advanced LIGO will see by estimating what fraction of neutron star binaries we could have detected in pulsar surveys if they were of sim ilar period and lum inosity to those known, and assigning a \scale factor" to each. We then multiply each member of the population by their scale factors and obtain an estimate of the number of sim ilar systems in the Galaxy, thus obtaining a coalescence rate for the M ilky W ay that can be used to infer an event rate for the local Universe by estimating the num ber of Galaxies out to the LIGO detection lim it. We will refer to this method as the scale factor approach. W hilst lim ited by sm all num ber statistics, the growing population of observed neutron star binaries makes us increasingly con dent that coalescence is relatively com m on in the Universe.

The only alternative approach is to attempt to simulate the entire binary pulsar population. Uncertainties in initial conditions and the physics of binary evolution and kicks mean that such estimates vary by more than two orders of magnitude (Belczynski et al. 2002). The population synthesis method is therefore of limited use in estimating the merger rate.

For now, the coalescence rate is better estimated by considering the scale factor method, however when this is done we nd the coalescence rate is dominated by the double pulsar PSR J0737{3039A (Burgay et al. 2003). This serves to demonstrate the heavy in uence of small number statistics on the uncertainties in the scale factor method.

The scale factor method also su ers from the extent to which the relativistic binary pulsar population remains invisible to pulsar surveys. For instance, if there existed a population of neutron stars that were com pletely invisible in the radio we would have no way to estim ate their population from radio observations. Sim ilarly, if only 1% of recycled neutron stars em itted radio waves, we should multiply the scale factors currently in use by 100.

Our best estimates of the inspiral rate require us to think of all the reasons why double neutron stars might be invisible to our surveys, and take them into account. W hat we explore in this paper is a selection e ect that rem oves binary pulsars from our surveys and has received little attention to date. This is the production of highlyeccentric short-period binaries with very short lifetim es, produced via favourably-oriented kicks in the binary progenitor. These systems are not long-lived, and the coalescence rate of the scale-factor approach needs to be corrected for it. W hilst this e ect has been alluded to from time to time (Tutukov & Yungelson 1993), we determ ine its in portance for an arbitrary binary in the presence of a kick velocity that can occupy any of the phase space that will produce a bound orbit, as in this case the problem is scale invariant.

In this paper, we derive the expected distributions of orbital eccentricity for any surviving binary pulsar, if the kick velocity is capable of uniform ly populating the available phase space after the explosion (Section 2). In Sections 3 and 4 we look at the lifetim es of the surviving system s due to gravitational radiation losses and identify a population of binaries that coalesce very rapidly. These results are used to explain the observed distribution of binary pulsar eccentricities (Section 5), which at long orbital periods favour high eccentricities, and at low periods, sm all eccentricities. In Section 6 we argue that the coalescence rate should be revised to take this into consideration, particularly for the double pulsar.

### 2. ECCENTRICITIES

In this section we exam ine the expected orbital eccentricities of relativistic double neutron stars from a purely analytical point of view, avoiding any assumptions about speci cs such as the distributions of stellar masses, kick velocities and orbital separations. Instead, we present a study of the potential eccentricities of an arbitrary double neutron star system in units scaled to the circular orbital speed, such that the conclusions drawn here can be considered common to the evolution of all double neutron star systems. Indeed, all of our results also hold for any two compact objects (black holes, white dwarfs or neutron stars) where the nal compact object (a black hole or a neutron star) is created in a supernova and receives a kick.

We view the e ect of kick velocity through a map of the second neutron star's post-supernova \velocityspace" { our representation of all possible relative velocities which correspond to bound orbits. The boundary of this velocity-space region is circular, with radius equal to the escape speed of the system ( 2 in units of the circular orbital speed). Each point in the map corresponds to one possible orbital velocity vector, where the position (0,1) would correspond to the circular orbital velocity.

To calculate the sem im a jor axis and eccentricity of an arbitrary binary, we derive form ulae from two expressions for the total energy E of the system (Hills 1983), and two expressions for the total angular momentum L of the system (M urray & Dermott 1999) :

$$E = \frac{GMm}{2a}$$
(1)

$$E = \frac{GMm}{R} + \frac{1}{2}v^2$$
 (2)

$$L = vR\cos$$
 (3)

$$L = [GM_{T}a(1 e^{2})]^{\frac{1}{2}}$$
(4)

Here = (M m) = (M + m) is the binary system's reduced mass,  $M_T$  its total mass, a its sem imapr axis, e its eccentricity, v the magnitude of its relative orbital velocity oriented at an angle to the plane de ned by the vector that connects the two stars, and R the initial orbital separation. From these four equations we derive expressions for the sem im a pr axis and the eccentricity of an arbitrary binary, given the initial relative orbital velocity v:

$$a = \frac{R}{2 v^2}$$
(5)

$$e = [1 \quad v^2 \cos^2 (2 \quad v^2)]^{\frac{1}{2}}$$
(6)

These form ulae are applied here to generate a contour m ap of eccentricities in velocity-space. Note that the results are independent of the choice of R. A code was written to uniform ly populate the velocity-space region with grid points, then apply Eq. 6 to calculate the eccentricity corresponding to each grid point (where each grid point represents one possible velocity). The 2D data was nally plotted as a contour map, and is presented here in Fig. 1.

In interpreting this map, note that the relative velocity vector just prior to supernova would be a little larger than 1.0 in the scaled units used here, because of the larger stellar m ass prior to supernova. (For example, a pre-supernova m ass of 2.0 M corresponds to an initial orbital speed of 1.102 units.) A lso, for a sense of scale, note that the relative circular speed of PSR J0737 {3039A is of order 600 km s  $^{1}$ . We operate on the assumption that each system 's nal supernova is accompanied by an asymmetric supernova kick, which has the e ect of displacing the initial relative velocity to a new position in velocity-space. This results in a new eccentricity and sem im a praxis, which can be calculated using the new velocity and Eq. 5 and 6. W e further assume that the outgoing supernova shell's impact with the companion has negligible e ect.

Now, the most striking feature of the map in Fig. 1 is the very small size of the low eccentricity regions. This is surprising, as the currently observed relativistic binaries tend to have eccentricities below 0.3 and therefore must have been kicked into a very speci c region of their postsupernova velocity-space (the third inner-most circle in Fig. 1). This seems an unusually fortuitous coincidence, requiring a very consistently precise kick. Furthermore, with m ean kick magnitudes of order 400{500 km s<sup>-1</sup>, we expect system s to be highly prone to being kicked out of these tiny low eccentricity regions in velocity-space.

The 2D eccentricity data generated for Fig. 1 is presented as a histogram and cum ulative distribution in Fig. 2. Here we have also provided eccentricity data for a 3D



Fig. 1.] Contour map of orbital eccentricity in the postsupernova velocity-space of an arbitrary binary star system, in which one star receives a random ly oriented kick. The velocities are measured in units of that required to place the post-supernova system in a circular orbit { for exam ple, the relative circular speed of PSR J0737{3039A is of order 600 km s<sup>-1</sup>. Prograde orbits have positive tangential velocities. The circular boundary of the region represents the escape velocity of the system, which is <sup>-2</sup> time s the circular speed.



Fig. 2. Eccentricity distribution histogram s are shown for binaries that are produced from random ly-oriented kicks which llthe available phase space for the planar (solid) and general (dashed) cases. In the general case the kicks can be out of the plane. A lso shown is the probability that the naleccentricity will be less than a given value, which is simply the integral of the eccentricity distributions. Note the high probability of eccentric system s resulting from a supernova kick in both cases.

version of our planar velocity-space, uniform by populating a \sphere" of radius  $\overline{2}$ . This allow sus to investigate the e ect of supernova kicks which are not restricted to the orbital plane.

The eccentricity histogram clearly illustrates a heavy bias towards very high eccentricities in the planar velocity-space, with over 50% of all eccentricities greater than 0.8. The distribution is less skewed when we allow for kick velocities out of the orbital plane, but the bias is still towards very high eccentricities. We assert that this illustrates a strong inherent tendency towards highly eccentric systems arising as a consequence of supernova kicks.

This nding is not re ected in the number of highly eccentric relativistic binaries in the observed sam ple of double neutron stars, which tend to have low eccentricities. A nalysing this from the standpoint of kick velocities, the observed systems seem to have been kicked consistently into a very speci c region of their post-supernova velocity-space, the \bullseye" seen in Fig. 1. Our explanation of this tendency is discussed further in the follow ing sections.

#### 3. LIFET MES

In this section we exam ine the probable coalescence times of relativistic double neutron stars, using a velocity-space contour map to illustrate dependence on the supernova kick velocity. As in the previous section, we avoid assumptions about speci cs such as the distributions of stellar masses, kick velocities and orbital separations, and present our results in units of the circular orbital speed and circular coalescence time (the merger time for an arbitrary system with zero eccentricity, denoted as <sub>circ</sub>). These generic units have been chosen for their usefulness in comparing binaries in a scale-invariant manner.

In producing the aforem entioned contour m ap of coalescence times, we rst generated a family of arbitrary double neutron star system s over all possible postsupernova relative orbital velocities. Computationally, this was achieved by uniform ly populating the \velocityspace" with grid-points, where each point represents a relative orbital velocity. These velocities were then used to calculate the corresponding sem im a jor axes and eccentricities, using Eq. 5 and 6. Both of these factors will greatly in uence the rapidity of each system 's orbital decay, with close, highly eccentric binaries being much more volatile than wider, more circular ones. To com pute num erical estim ates of just how rapidly each system will coalesce, each grid point's initial orbital param eters were evolved forward in time by num erically integrating the coupled di erential equations derived by Peters (1964) for describing orbital decay by relativistic gravitational radiation. A Runge-Kutta 4th order algorithm was implemented with an adaptive step size, designed to integrate the equations forward to the precise time when the sem i-major axis of the system drops to zero, thus determining the coalescence time. After iterating through all grid-points, the 2D coalescence time data was used to produce the contourm ap shown in Fig. 3.

The most signi cant feature of Fig. 3 is the large central region within which coalescence times rapidly drop from 0.1  $_{\rm circ}$  (in units of the coalescence time of a circular system) to 10  $^5$   $_{\rm circ}$  and even lower. To put this in perspective, for the orbital parameters of PSR J0737{ 3039A /B, 10  $^5$   $_{\rm circ}$  corresponds to a coalescence time of less than 1,000 years. Fig. 3 thus demonstrates that a double neutron star system which receives a supernova kick of the order of its circular speed has a very high like-lihood of coalescing within an incredibly short time span.

The cumulative distribution of coalescence times is provided in Fig. 4, which shows the uncoalesced per-



Fig. 3. Contour map of coalescence times of binaries (in units of circ) as a function of orbital velocity im mediately after the second supernova. Only prograde orbits are shown, but the diagram is symmetric about the x axis as for Fig 1. The time units are relative to the coalescence time due to gravitational wave emission for a system with a circular orbit. If neutron stars receive a random ly oriented kick after the supernova explosion that produces them, it is possible to populate the entire phase space shown. Binaries with little angular momentum after the explosion are seen here to coalesce in remarkably short times relative to those in near-circular orbits. The consequence of this is that such system s are extrem ely di cult to observe com pared to the system s of sm all eccentricity.



Fig. 4. The fraction of systems that remain uncoalesced as a function of time, when considering a fam ily of arbitrary binaries with all possible post-supernova orbital velocities. The solid curve here illustrates the case where the post-SN velocity is constrained to the pre-SN orbital plane, and the dashed line represents the general case. The unit used for time ( $_{\rm circ}$ ) is the coalescence time of a system which remains in a circular orbit in mediately after the supernova. For a sense of scale, consider that 1  $_{\rm circ}$  is equivalent to 88 M yr when scaled to the param eters of PSR J0737{3039A /B}.

centage of the representative population as a function of time. This Figure dram atically illustrates the high probability of rapidly coalescing double neutron star systems, with up to 24% of all systems coalescing in less than 10  $^4$  circ (equivalent to 8,500 years when scaled to the parameters of PSR J0737{3039}. A firer just 0.1 circ about half of all systems have coalesced. This is an intriguing result, im plying that the coalescence rate of double neutron stars is underestimated. These rapidly coalescing systems would be practically undetectable by m odem pulsar surveys, since they would not live long enough to give su cient opportunity for detection. They would, how ever, be detectable by gravitational wave ob-



Fig. 5.] The evolution of observed eccentricities as a function of time, for the binary population as in Fig. 2. The units are again in term s of the time it would take a circular system to decay due to gravitational radiation (  $_{\rm circ}$ ). A fler just 0.01  $_{\rm circ}$  the eccentricity distribution (dashed) has attened signi cantly as all the shortperiod, highly eccentric system s have already coalesced. A fler one com plete circular coalescence time, the population of surviving systems (dotted) is m uch m ore uniform .

servatories such as LIGO, which will come online in the near future. We might predict that at short orbital periods there will be a decit of highly eccentric systems because these are selected against due to their shorter lifetimes. We now investigate this in the following section.

#### 4. ECCENTRICITY DISTRIBUTION EVOLUTION

Here we analyse the time evolution of the eccentricity distribution of a fam ily of arbitrary double neutron star system s. The computation perform ed here is similar to that perform ed in the prior section { to begin, for every possible post-supernova relative orbital velocity (represented computationally by uniform ly spread grid-points in our \velocity-space"), we calculate an eccentricity and sem i-m a jor axis by applying Eq. 5 & 6. This results in a standard representative fam ily of double neutron star system s, whose orbital parameters we can then num erically integrate forward in time using the coupled di erential equations derived by Peters (1964). In this way, the eccentricity distribution of the whole fam ily of possible double neutron star system s can be computed at birth, and at any time afterwards. Fig. 5 shows our resulting computed eccentricity distributions at three stages in time.

This Figure serves to illustrate the rapid coalescence of the vast majority of highly eccentric systems, which were produced in the greatest numbers when all possible bound orbital velocities were represented in our sample. This produces a rapidly evolving eccentricity distribution which changes on very short timescales (0.01 <sub>circ</sub>), and generally shifts towards lower eccentricities. W hile the initial post-supernova eccentricity distribution may be highly skewed, Fig. 5 demonstrates that it doesn't retain its shape for very long, quickly attening out to a more uniform distribution, from which the observed eccentricity distribution of double neutron star systems could be more plausibly derived.



Fig. 6. Orbital parameters of all currently known eccentric (e > 0:05) binary pulsar systems residing within the disk of our galaxy. We have excluded pulsars in globular clusters, where the form ation histories are quite dienent. Relativistic systems (those which will coalesce within a Hubble time) are indicated by dotted circles, and non-coalescing systems are represented by stars. Note the comparatively low eccentricities of the relativistic binaries. Pulsar inform ation was obtained from the ATNF PulsarCatalogue (M anchester et al. 2005).

#### 5. OBSERVED POPULATION

Here we will exam ine the observed population of double neutron star binaries, particularly in terms of the eccentricity distribution. Fig. 6 shows the sem im a jor axes and eccentricities of all currently known eccentric double neutron star systems residing within the disk of our galaxy.

The majority of relativistic binaries tend to have low eccentricities, which seems fortuitous considering the tiny size of the low eccentricity \bullseye" region in the velocity-space as shown in Fig. 1. To achieve low eccentricities, these relativistic binary systems would have had to receive consistently precise supernova kicks. A ny sm all deviation in kick direction or magnitude would result in higher eccentricities.

Our analysis of uniform ly distributed pulsar kicks nds that surviving binaries should create mostly highlyeccentric system s. W e contend that supernova kick velocities can easily be large enough in m agnitude to kick a binary out of its sm all low -eccentricity \bullseye" and into the dom inant, high-eccentricity region of velocity-space (see Fig. 1). These large supernova kicks can be inferred from the abundance of single pulsars roam ing the galaxy at high velocities (m ean of order 400 {500 km s<sup>1</sup>), m ost likely accelerated by supernova kicks. W ith regards to our simple kick model, it is almost certain that the real kick velocity distribution would not uniform ly populate the available phase space after a supernova explosion, as the kicks in close binaries would have to range from small. velocities to higher than 1000 km s<sup>1</sup> with equal probability. Nevertheless, trying to guess the exact shape and magnitude of the pulsar kick velocity distribution leads to a plethora of models, and we resist this tem ptation. Our ignorance of the kick velocity distribution does not invalidate our results, unless pulsars can never receive a su ciently large kick, of magnitude similar to that of the original pre-supernova orbital velocity. We

are inclined to assume that such large kicks can exist, how ever, as observed pulsar velocities suggest kicks may be imparted with magnitudes up to and exceeding 1000 km s<sup>1</sup> (C ordes et al. 1993).

We propose that the observed population of relativistic double neutron stars exhibits low eccentricities because systems with higher eccentricities have coalesced before they could be detected. An average pulsar is detectable by modern pulsar surveys for approximately 10 M yr { however, we have shown that binary systems have an appreciable likelihood of being kicked into coalescence within 1 M yr or even less. Not only will these systems coalesce rapidly, they will also have orbits so tight that the e ects of \D oppler smearing" are too strong for current search algorithm sto detect them in some surveys. In sum mary, the low eccentricities of the relativistic double neutron star systems suggest that a strong selection effect is in operation, and we claim that accelerated decay due to gravitational radiation is the cause.

Faukneret al. (2005) suggest that there is a correlation reinforced by the discovery of PSR J1756{2251 between orbital eccentricity and spin period of recycled pulsars. Our results suggest that many highly-eccentric systems resulting from close binaries are never seen by pulsar surveys, and that long-period binaries favour high eccentricities. C lose binaries transfer matter when the evolutionary tim escale of the donor is still long. We therefore expect that short-period progenitors will be spun up to shorter periods than the longer-orbital period system s. A natural consequence of this is that the short-period high e system s are selected against, that very few long-period low e system s are born, and therefore a correlation is observed.

## 6. D ISCUSSION: IM PLICATIONS FOR THE MERGER RATE AND \DOUBLE SUPERNOVAE "

W e have seen that the binary pulsar population ism issing highly-eccentric binary pulsars in relativistic orbits, which we attribute to their short observable lifetimes. The most reliable estimates of the merger rate of neutron star binaries come from extrapolation of the observed population, which ignores these m issing binaries that we expect to be formed in large numbers before dying young. Our calculations suggest that factors of several are required to correct the m erger rate for these m issing binaries.

We have also shown that there is a non-negligible potential for coalescence times of the order of months if the pulsar kick is of the same order as the pre-supernova orbital velocity. This may mean that even relatively wide pre-supernova orbits can be driven to coalesce in remarkably short times by receiving an appropriatelyoriented retrograde kick. This phenom enon could reveal itself in the form of a supernova follow ed closely by a second sim ilarly-scaled catastrophic event (the coalescence), where both would be observed at the same position a few months or years apart { what m ight be described as a \double supernova". How soon after the original burst of star form ation in the Universe m ight we expect to see such an event? Population III stars { the ancient stars which were the st to form in the history of the Universe { may have often occurred in binaries where both stars underwent supernovae, subsequently taking part in an ultra-rapid coalescence which could be detected through gravitational radiation em issions (the only known way of observing these objects, otherwise hidden by the Epoch of Reionisation). This idea has been alluded to before (Belczynski et al. 2004), however our ndings em phasize the possibility that such a coalescence could occur very quickly after initial star form ation, allowing future gravitational wave detectors to directly study events that occurred within the st few M yrof star form ation in the Universe.

This paper is the product of a ten-week Summer Vacation Scholarship undertaken by HKC at the Centre for Supercomputing and A strophysics at Sw inburne University, Hawthom, Australia.

#### REFERENCES

- Abram ovici, A., Althouse, W. E., Drever, R.W. P., G& quot,ursel, Y., Kawam ura, S., Raab, F.J., Shoem aker, D., Sievers, L., Spero, R.E., Thorne, K.S., Vogt, R.E., Weiss, R., Whitcomb, S.E., & Zucker, M.E. 1992, Science, 256, 325
- Bailes, M .1989, ApJ, 342, 917
- Belczynski, K., Bulik, T., & Rudak, B. 2004, ApJ, 608, L45
- Belczynski, K., Kalogera, V., & Bulik, T. 2002, ApJ, 572, 407
- Burgay, M., D'Amico, N., Possenti, A., Manchester, R.N., Lyne, A.G., Joshi, B.C., McLaughlin, M.A., Kramer, M., Sarkissian, J.M., Camilo, F., Kalogera, V., Kim, C., & Lorimer, D.R. 2003, Nature, 426, 531
- Cham pion, D. J., Lorim er, D. R., M cLaughlin, M. A., Cordes, J. M., Arzoum anian, Z., W eisberg, J. M., & Taylor, J. H. 2004, M NRAS, in press, astro-ph/0403553
- Cordes, J.M. & Chemo, D.F. 1998, ApJ, 505, 315
- Cordes, J.M., Romani, R.W., & Lundgren, S.C. 1993, Nature, 362, 133
- Curran, S.J. & Lorim er, D.R. 1995, MNRAS, 276, 347
- Dewey, R.J. & Cordes, J.M. 1987, ApJ, 321, 780
- Faulkner, A. J., K ram er, M., Lyne, A. G., M anchester, R. N., M cLaughlin, M. A., Stairs, I. H., Hobbs, G., Possenti, A., Lorim er, D. R., D'Am ico, N., Cam ilo, F., & Burgay, M. 2005, ApJ, 618, L119
- Hills, J.G. 1983, ApJ, 267, 322
- Hotan, A.W., Bailes, M., & Ord, S.M. 2004, ArX iv A strophysics e-prints

- Kalogera, V., Kim, C., Lorimer, D. R., Burgay, M., D'Amico, N., Possenti, A., Manchester, R. N., Lyne, A. G., Joshi, B. C., McLaughlin, M. A., Kramer, M., Sarkissian, J. M., & Camilo, F. 2004, ApJ, 601, L179
- Kalogera, V., Narayan, R., Spergel, D. N., & Taylor, J. H. 2001, ApJ, 556, 340
- Kaspi, V. M., Bailes, M., Manchester, R. N., Stappers, B. W., & Bell, J. F. 1996, Nature, 381, 584
- Lipunov, V.M., Postnov, K.A., & Prokorov, M.E. 1997, 2, 43
- Lyne, A. G., Cam ilo, F., Manchester, R. N., Bell, J. F., Kaspi, V. M., D'Am ico, N., McKay, N. P. F., Craw ford, F., Morris, D. J., Sheppard, D. C., & Stairs, I. H. 2000, MNRAS, 312, 698
- Lyne, A.G. & Lorimer, D.R. 1994, Nature, 369, 127
- M anchester, R.N., Hobbs, G.B., Teoh, A., & Hobbs, M. 2005, AJ, 129, 1993
- M eegan, C.A., Fishm an, G.J., Wilson, R.B., Horack, J.M., Brock, M.N., Paciesas, W.S., Pendleton, G.N., & Kouveliotou, C. 1992, Nature, 355, 143
- M urray, C.D. & Derm ott, S.F. 1999, Solar System Dynamics (Cambridge: Cambridge University Press)
- Peters, P.C. 1964, Phys. Rev., 136, 1224
- Portegies Zwart, S.F. & Spreeuw, H.N. 1996, A & A, 312, 670
- Stairs, I. H., Thorsett, S. E., & Arzoumanian, Z. 2004, Physical Review Letters, 93, 141101
- Taylor, J.H. & W eisberg, J.M. 1989, ApJ, 345, 434
- Tutukov, A.& Yungelson, L.1993, MNRAS, 260, 675

Hulse, R.A. & Taylor, J.H. 1975, ApJ, 195, L51

W eisberg, J.M., Romani, R.W., & Taylor, J.H. 1989, ApJ, 347, Wolszczan, A. 1991, Nature, 350, 688 1030

W illem s, B.& Kalogera, V.2004, ApJ, 603, L101