## Param etrization of K-essence and Its K inetic Term

HuiLi, Zong-Kuan Guo<sup>b</sup>and Yuan-Zhong Zhang<sup>a;b</sup>

<sup>a</sup>CCAST (W orld Lab.), P.O. Box 8730, Beijing 100080, China <sup>b</sup>Institute of Theoretical Physics, Chinese Academy of Sciences, P.O. Box 2735, Beijing 100080, China

## A bstract

We construct the non-canonical kinetic term of a k-essence eld directly from the effective equation of state function  $w_k(z)$ , which describes the properties of the dark energy. A dopting the usual parametrizations of equation of state we numerically reproduce the shape of the non-canonical kinetic term and discuss some features of the constructed form of k-essence.

K eywords: param etrization; equation of state; reconstruction; k-essence; non-canonical kinetic term.

PACS number(s): 98.80 Cq, 98.65 Dx

e-mailaddress: lihui@ itp.ac.cn

Recent observations of type Ia supernovae [1, 2], m easurem ents of the cosm ic m icrow ave background [3] and the galaxy power spectrum [4] indicate the existence of the dark energy an energy component with a strongly negative equation of state (EOS)  $w_{DE}$ to matter. For this purpose, various dynamical scalar eld models have been proposed. Speci cally, a reliable model should explain why the present amount of the dark energy is so small compared with the fundamental scale (ne-tuning problem) and why it is comparable with the critical density today (coincidence problem). Quintessence has been proposed as a candidate for the dark energy component of the universe that would be responsible of the currently observed accelerated expansion [5]-[9]. Generally speaking, quintessence is a spatially homogeneous eld slow-rolling down its potential. More recently, models based on scalar elds with non-canonical kinetic energy, dubbed as k-essence, have emerged 10, 11]. A subclass of models feature a tracker behavior during radiation domination, and a cosm ological-constant-like behavior shortly after the transition to matter domination. As long as this transition seems to occur generically for purely dynamical reasons, these models are claimed to solve the coincidence problem without netuning. What is more, lately it has been found that those two theoretical setups are strongly related to each other and every quintessence model can be viewed as a k-essence model generated by a kinetic linear function [12].

The dark energy is characterized by its equation of state parameter w DE, which is in general a function of redshift z in dark energy models. One could specify a Lagrangian and solve the scalar eld equation for the particular theory [3]. Although it may provide a speci c form of  $w_{DE}$  (z) in terms of scalar eld model parameters which can be directly constrained through data tting [4,15], this physically motivated approach clearly limits us to a model by model test [16]. A part from the feasibility of quintessence potential V () and the EOS  $W_{DE}$  (z) reconstruction from supernova observations [17, 18], a phenomenological param etrization of  $w_{DE}$  (z) is still necessary to compare dierent dark energy models. The key point of this alternative strategy is to assume a more or less arbitrary ansatz for  $W_{DE}$  (z) which is not necessarily physically justi ed but specially designed to give a good t to the observational data. This motivation stimulates several parametrization methods to produce an appropriate parameter space [19]-[24]. Recent study found that it is also workable to drive these param etrizations to construct the quintessence potential form [25] and some general features of these constructions were discussed. In this Letter, we try to carry out this m ethod in constructing the k-essence with pressure p ('; X) directly from the dark energy equation of state function  $w_k(z)$ . When applying this method to four typical param etrizations, the shapes of the kinetic term are num erically obtained and relevant discussions on the resulting construction are also given.

Restricting our attention to a single  $\$ eld  $\$ m odel, the action of the  $\$ k-essence generically  $\$ m ay be expressed as

$$S_r = {}^{Z} d^4x {}^{p} - g - \frac{R}{6} + p(';X);$$
 (1)

where we use units such that 8 G=3 1; and

$$X = \frac{1}{2} (r')^2$$
: (2)

The Lagrangian p depends on the specie c particle theory model. In this paper, we consider only factorizable Lagrangians of the form [26]

$$p = K (')p(X);$$
(3)

where we assume that K(') > 0: To describe the behavior of the scalar eld it is convenient to use a perfect—uid analogy. The role of the pressure is played by the Lagrangian pitself, while the energy density is given by [27]

" = K (') (
$$2X p_{:X} (X) p(X)$$
) (4)

$$K (') \sim (X);$$
 (5)

where  $:::_{X}$  denotes a partial derivative with respect to X. The ratio of pressure to energy density, which we call, for brevity, the k-essence equation-of-state,

$$w_k = \frac{p}{n} = \frac{p}{2X p_x} ; \qquad (6)$$

does not depend on the function K (').

We will consider a spatially at FRW universe which is dominated by the non-relativistic matter and a spatially homogeneous scalar eld. The Friedmann equation can be written as

$$H^2 = _m + ";$$
 (7)

where  $_{\rm m}$  is the matter density. The evolution of k-essence eld is governed by the equation of motion

$$\underline{"} + 3H (" + p) = 0;$$
 (8)

which yields

where z is the redshift which is given by  $1 + z = a_0 = a$  and subscript 0 denotes the value of a quantity at the redshift z = 0 (present). As the constant  $\mathbf{v}_0$  is of no importance for further discussion, we will safely set it to 1 and then regard the energy density  $\mathbf{v}(z)$  as a dimensionless quantity.

So we get "(z) as the function of redshift z, and by m eans of the EOS of k-essence p(z) m ay be well prepared as follows,

$$p(z) = \frac{w_k(z)E(z)}{K(t')};$$
(10)

Once the form of the potential function K (') is specified, '(z) supplied by the Eq. (4) will describe the behavior of (2). With the help of Eq. (4) and Eq. (6), X can also be implicitly expressed as the function of (2):

"(z) = K (') 2X 
$$\frac{dp(z)}{dz} \frac{dz}{dX}$$
  $p(z)$  : (11)

Therefore, the redshift relates two functions p(z) and X(z), and the reconstruction m ay be implemented. Using  $m = m_0 (1+z)^3$  and Eq. (9), the Friedmann Eq. (7) becomes

H (z) = H<sub>0</sub> 
$$_{m 0}$$
 (1 + z)<sup>3</sup> +  $_{k0}$ E (z) ; (12)

where  $_{m\ 0}$   $_{m\ 0}$ =H  $_{0}^{2}$  and  $_{k0}$   $_{k0}$ =H  $_{0}^{2}$ . Due to the hom ogeneity of k-essence X =  $^{\prime}$ \_=2 and therefore,

$$X (z) = \frac{('^{0}(z))^{2} (1+z)^{2} H (z)^{2}}{2};$$
 (13)

where the dot represents derivative with respect to the physical time t and the prime derivative with respect to the redshift z. As a consequence, '\_ '0(z)dz=dt =  $\frac{1}{2}$ X. Because p(x) is only dependent on the variable X, the sign of '\_ is insignicance for the construction process below.

For de niteness we will choose K (') = ' $^2$  [26]. Since dz=dt = (1 + z)H (z) and H (z) is given by Eq. (7), after straightforward calculations the di erential equation for '(z) will be written as the Bernoulli type

$$v^{0}(z) + \frac{1}{1+z} + \frac{H^{0}(z)}{H(z)} + \frac{w_{k}^{0}}{1+w_{k}} + \frac{w_{k}E^{0}(z)}{(1+w_{k})E(z)} + v(z) + \frac{w_{k}}{w_{k}+1}v(z)^{2} = 0;$$
 (14)

where v(z) ' $\stackrel{\circ}{=}$ '.

A fler identifying ' with the dimensionless quantity '  $^{-}$  M  $_{\rm pl}$ , the construction equations are of the same form as Eq. (10) and Eq. (13), which relate the non-canonical kinetic term p(X) to the equation of state function w $_{\rm k}$ (z). Given an elective equation of state function w $_{\rm k}$ (z), Eq. (14) will make these two construction equations (10) and (13) simulate the non-canonical kinetic energy p(X).

Our method relates directly the kinetic terms of k-essence to the equation of state function, and so enables us to construct easily the former without assuming its form. The dark energy properties are well described by the elective equation of state parameter  $w_k$  (z) which in general depends on the redshift z.

Following the previous work [25], we consider the construction process with the following parametrization methods [19]-[24]: a constant equation of state parameter and the other three two-parameter parametrizations,

Case 1: 
$$w_k = w_0$$
 (Ref. [19])




Figure 1: Constructed kinetic term p(X) in the k-essence model.

```
Case 2: w_k = w_0 + w_1 z (Ref. [20])

Case 3: w_k = w_0 + w_1 z = (1 + z) (Refs. [21, 22, 23])

Case 4: w_k = w_0 + w_1 \ln (1 + z) (Ref. [24])
```

we choose  $w_0=0.7$ ,  $w_1=0.2$  and  $r_0=3=7$  for the special construction 28]. This is quite diagram the quintessence case as  $dw_k=dz>0$  is generally required for the form er [10, 29]. This noticeable feature would enable supernova data to distinguish between the two theories, although some ambiguity is sticky to deal with. Just for deniteness we have chosen the initial values of the k-essence eld  $\prime_0=1.0$  and  $v_0=0.1$  at z=0. This degree of freedom of initial value settings will be easily eliminated after appropriate eld rescaling and have no essential in the uence on the general form of p(X). As a matter of fact, the change of the initial value  $\prime_0$  or  $v_0$  just lifts the corresponding curves vertically and keeps their shapes intact. The numerical results are plotted as follows. Fig. 1 shows the constructed non-canonical kinetic term p(X). At low redshift region (z<1, corresponding to x>4), all these models share the same asymptotical behavior, but begin to deviate from this around redshift z>1.

We should note that the param etrizations we adopted here m ight not be of particular signicance but just for concreteness. Since there is little theoretical guidance as to the nature of the dark energy, the phenom enological param etrization of  $w_k(z)$  should be designed as generally as possible unless no observations can electively unravel such excess complication. It has been shown that more than two additional param eters is not viable

for a general twith next generation data but two parameter approximations in current use are reasonable and realistic for the near future observations [29, 30, 31]. On the other hand, due to the lack of data and the large statistical error bars, a wide range of dark energy models seem to have ranges of parameters that the existing data equally well and no particular model stands out as being observationally preferred. As a consequence, any such parametrizations made in potential reconstruction methods may be too restrictive since many dient potential energy functions are conceivable and many of them may give results degenerate with each other. While it would still be dicult to break this observational degeneracy, the challenging goal of making the fundamental physics distinction of the sign of the EOS time variation is achievable in the next generation of experiments [30, 31]. One of the direct rewards relevant to the two parameter approximation then would be that the usual form of k-essence with dw k=dz > 0 could be knocked down. Anyhow, although we have concentrated specially on the k-essence form in Ref. [26], it is obvious that the techniques exploited here may be applied to more general k-essence models as well.

In conclusion, we have used the scheme of constructing the form of dark energy developed in [25] directly from the elective equation of state function  $w_k$  (z). Then we have considered four parametrizations of equation of state parameter and showed that the constructed k-essence model may also be constructed without further dynamical information of the k-essence ed. Current popular parametrizations we have referred to are directly related to the data of type Ia supernovae; therefore, our approach will be a useful tool to reexamine the k-essence models, and the two parameters extracted from the SNe Ia observations will not only help us select the suitable k-essence form or rule out the usual form of k-essence models with great condence in the future, but also may lead to a strong constraint and convenient guidance for the fundamental theory of the perplexing dark energy. The future Supernova/A coeleration P robe with high-redshift observations [32], in combination with the P lanck CMB observation [33], will be able to determine the parameters in the dark energy parametrization to high precision. By precision mapping of the recent expansion history, we hope to learn more about the essence of the dark energy and get a deeper understanding of the dynamics of the universe.

## A cknow ledgem ents

This project was in part supported by National Basic Research Program of China under Grant No. 2003CB716300 and by NNSFC under Grant No. 90403032.

## R eferences

[1] A.G.Riess et al., Astron.J.116 (1998) 1009; S.Perlmutter et al., Astrophys.J.517 (1999) 565.

- [2] Y.W and and M. Tegmark, PhysRev.D 71 (2005) 103513; B.Feng, X.W and and X. Zhang, PhysLett.B 607 (2005) 35; Y.G ond and Y.Z. Zhang, gr-qc/0508053.
- [3] D. N. Spergel et al., Astrophys. J. Suppl. 148 (2003) 175.
- [4] M. Tegm ark et al., Phys. Rev. D 69 (2004) 103501.
- [5] B.Ratra and P.J.E. Peebles, Phys. Rev. D 37 (1988) 3406.
- [6] R.Caldwell, R.D ave and P.J. Steinhardt, Phys. Rev. Lett. 80 (1998) 1582;
- [7] C.W etterich, Nucl. Phys. B 302 (1988) 668.
- [8] I. Zlatev, LM. Wang and PJ. Steinhardt, Phys. Rev. Lett. 82 (1999) 896.
- [9] P.J. Steinhardt, L.W ang and I.Zlatev, Phys. Rev. D 59 (1999) 123504.
- [10] C.Arm endariz-Picon, V.M ukhanov and Paul J. Steinhardt, Phys Rev Lett. 85 (2000) 4438; C.Arm endariz-Picon, V.M ukhanov and Paul J. Steinhardt, Phys Rev. bf D 63 (2001) 103510.
- [11] R.J. Scherrer, PhysRevLett. 93 (2004) 011301; L.P. Chimento and R. Lazkoz, PhysRev.D 71 (2005) 023505; H.Weiand R.G. Cai, PhysRev.D 71 (2005) 043504.
- [12] JM. Aguirregabiria, L.P. Chim ento and R. Lazkoz, astro-ph/0411258.
- [13] P.J.E. Peebles and B. Ratra, The Astrophys. J. 325 (1988) L17.
- [14] G.Chen and B.Ratra, Astrophys. J.Lett. 612 (2004) L1.
- [15] C.R.W atson and R.J.Scherrer, PhysRev.D 68 (2003) 123524.
- [16] S.C. Cindy Ng and D.L.Wiltshire, Phys.Rev. D 64 (2001) 123519.
- [17] D. Huterer and M. S. Tumer, Phys. Rev. D 60 (1999) 081301; T. Chiba and T. Nakamura, Phys. Rev. D 62 (2000) 121301; A. A. Starobinsky, JETP Lett. 68 (1998) 757.
- [18] M. Sahlen, A.R. Liddle and D. Parkinson, astro-ph/0506696; S. Tsujikawa, astro-ph/0508542.
- [19] S. Hannestad and E. Mortsell, Phys. Rev. D 66 (2002) 063508.
- [20] A.R. Cooray and D. Huterer, Astrophys. J. 513 (1999) L95.
- [21] M. Chevallier and D. Polarski, Int. J. Mod. Phys. D 10 (2001) 213.
- [22] E.V. Linder, Phys. Rev. Lett. 90 (2003) 091301.

- [23] T.Padmanabhan and T.R.Choudhury, Mon.Not.Roy.Astron.Soc.344 (2003) 823; H.K. Jassal, J.S.Bagla and T.Padmanabhan, Mon.Not.Roy.Astron.Soc., 356 (2005) L11; H.K. Jassal, J.S.Bagla and T.Padmanabhan, Phys.Rev.D 72 (2005) 103503.
- [24] B.F.Gerke and G.Efstathiou, Mon. Not. Roy. Astron. Soc. 335 (2002) 33.
- [25] ZK.Guo, N.Ohta and YZ.Zhang, PhysRev.D 72 (2005) 023504 335 (2002) 33.
- [26] M. Makquarti, E.J. Copeland, A.R. Liddle and M. Trodden, PhysRev. D 67 (2003) 123503.
- [27] C. Arm endariz-Picon, T. Dam our and V. Mukhanov, Phys.Lett. B 458 (1999) 209.
- [28] V. Barger and Danny Marfatia, PhysLett. B 498 (2001) 67.
- [29] Y.W ang and P.M. Gamavich, Astrophys. J. 552 (2001) 445.
- [30] E.V. Linder and D. Huterer, PhysRev. D 72 (2005) 043509.
- [31] R.R. Caldwell and E.V. Linder, Phys. Rev. Lett. 95 (2005) 141301.
- [32] SNAP hom epage: http://snap.lbl.gov.
- [33] Planck hom epage: http://www.rssd.esa.int/PLANCK.