News from the \D entist's C hair": O bservations of A M 1353-272 with the V IM O S IF U

Peter M .W eilbacher¹ and Pierre-A lain Duc^2

² Service d'astrophysique, CEA Saclay-O me des Merisiers, Bat. 709, 91191 G if sur Y vette cedex, France, paduc@cea.fr

Sum m ary. The galaxy pair AM 1353-272 nicknam ed \The D entist's C hair" shows two 30 kpc long tidal tails. P revious observations using multi-slit m asks showed that they host up to seven tidal dwarf galaxies. The kinem atics of these tidal dwarfs appeared to be decoupled from the surrounding tidal material. New observations of the tip of the southern tidal tail with the V IM O S integral eld unit con m the results for two of these genuine tidal dwarfs but raise doubts whether the velocity gradient attributed to the outerm ost tidal dwarf candidate is real. W e also discuss possible e ects to explain the observational di erence of the strongest velocity gradient seen in the slit data which is undetected in the new integral eld data, but arrive at no m conclusion. A dditionally, low-resolution data covering m ost of the two interacting partners show that the strongest line em itting regions of this system are the central parts.

1 Introduction

Following old ideas about the creation of dwarf galaxies during interaction of giant galaxies and detailed investigations of several nearby examples of these Tidal Dwarf Galaxies (TDGs, Duc et al., 2000, 1997; Duc & Mirabel, 1998; Hibbard et al., 1994), we carried out a st sm all survey of interacting galaxies (W eilbacher et al., 2000) with the aim of better understanding the star-form ation history of TDGs and constraining the number of TDGs that are built per interaction (W eilbacher et al., 2003). During this survey, we studied a system cataloged as AM 1353-272, which we called \The Dentist's Chair" for its peculiar shape. Fig. 1 shows the two components of the system, A', a galaxy with 30 kpc long tidal tails, and B', a disturbed disk galaxy, 160 M pc (H $_0$ = 75 km s 1 M pc 1). W ithin the tails have a distance of D several obvious clum ps with blue optical colors are visible. U sing optical and near-infrared in aging, evolutionary models, and optical spectroscopy, seven of these clumps were classi ed as TDG candidates in form ation (W eilbacher et al., 2002, , marked a' to d' and k' to m'). The largest velocity gradient

¹ A strophysikalisches Institut Potsdam, An der Stemwarte 16, D-14482 Potsdam, Germany, pweilbacher@aip.de

2 Weilbacher & Duc

with an am plitude of > 300 km s¹ appeared in TDG candidate a', at the very end of the southern tidal tail. This raised the question how an object with relatively low lum inosity could exhibit such fast \rotation". How ever, as these observations were done using the multi-slit technique and hence are spatially restricted due to the narrow slit, subsequent observations were planned using an integral eld unit (IFU) to cover more of the tidal tails and view the velocity structure of the TDGs in two dimensions.



F ig. 1. The interacting system AM 1353-272. The two interacting galaxies (capital letters) and the relevant knots in the tidal tails are marked (lower case letters). O verlayed are the original FORS2 slits and the two VIMOS pointings in low and high resolution mode.

2 IFU Observations

O ur new VIMOS data taken at the ESO VLT consists of 1.3 hours of exposure time in high resolution blue mode (eld of view of 13^{00} 13^{00}) and good seeing conditions ($0^{0.7}$), targeted at the tip of the southern tidal tail. This includes the TDG candidates a' to b'. The center of galaxy A', much of the southern tidal tail, and the companion B' were targeted with low-resolution (blue grism, 54^{00} 54^{00}), and observed for 1 hour in mediocre conditions with $2^{0.9}$ 0 seeing. These two pointings are sketched in Fig. 1.



F ig. 2. Results of the low-resolution pointing, the center of AM 1353-272 A is at the coordinates (0,0); at the bottom right, near (30,32) companion B' is visible. A xes labels are in arcseconds.Left:relative H em ission line ux (bright:high ux, dark:low ux).R ight:velocity eld derived from H em ission line (the greyscale bar gives relative velocities in km s¹).

The data were reduced using the ESO pipeline for the VIMOS instrument. We made a small enhancement to the code that allowed us to interpolate the wavelength solution between adjacent spectra on the CCD. This was only used for the low resolution data, where the errors introduced with this method are smaller than the accuracy allowed for by the spectral resolution. In low resolution mode, on the order of 15% of the spectra could not be wavelength calibrated due to overlapping spectral orders. From the nal datacube of extracted and wavelength calibrated spectra, we measured the relative uxes and velocities in each spectral element using Gaussian ts to the brightest usable em ission line. In the low resolution data (0 iii)5007 is the strongest em ission line but at this redshift (z 0:04) it is strongly blended with a sky emission line, so H had to be used instead. In the high-resolution data, [0 iii]5007 is less a ected by the sky-line and is the only line with sufcient S/N for the analysis in the low surface brightness region near the end of the tidal tail.

3 Results

Fig. 2 sum marizes the results that can be derived from the low-resolution pointing using ts to the H emission line. The southern tidal tail in undetected in this exposure and the strongest line emission appears to be in the center of A' and in the two knots at the end of its bar-like central structure (designated b' and b' in W eilbacher et al., 2002). Galaxy B', despite being strongly reddened, also is a strong source of H line emission. As the velocity resolution is on the order of 100 km s⁻¹, in this mode of V IM O S we cannot resolve the velocity structures in individual knots, but the bar-like structure



Fig. 3. Velocity eld as derived from FORS2 observations (black points) and reconstruction of this velocity eld from the VIMOS IFU datacube (grey crosses).

in A'seems to rotate (the eastern end near knot \forall is receding, the western end near knot \forall ' is approaching). The same is true for the companion B'.

To verify our original FORS2 multi-slit observations, we try to \reconstruct" them from the VIMOS datacube. To that end we average over the spaxels approximately in the dispersion direction of the FORS2 observations and derive an average redshift over the slit width. Fig. 3 shows the resulting velocity eld along this articial slit and compares it with the velocity prole of the FORS2 data. From this plot it can be seen that the velocity gradients of knots b' and b' are very well recovered, while the steep slope within knot a' with an amplitude of 350 km s¹ in the FORS2 data appears alm ost at in the reconstructed VIMOS data. We tested several alternative slit positions and curvatures, varied the e ective slit width, and also tried to add slit e ects (velocity o sets due to non-centered emission within the slit) to our reconstruction. None of these changes im proved the m atch for knot 'a'. In fact, slit e ects signi cantly worsened the agreem ent for all three knots. The FORS2 observations were done on these extended objects in 1° . slit, so that it appears unlikely that slit e ects would have a strong contribution to the observed velocity gradient. O ther problem s, like instrum ental exures should have been removed by the data reduction procedure as detailed in W eilbacher et al. (2002). W e are therefore con dent that slit e ects do not play an important role in the FORS2 data. On the other hand, if we assume that the original slit-based data give the correct results it is unclear,

how the V \mathbbm{M} O S data could be awed to hide this one velocity gradient. The

wavelength calibration works well for the high-resolution mode as con med by checks with sky emission lines.

4 Summary & Outlook

We presented a few tentative results for the interacting system called \T he Dentist's Chair" from new observations with the VIMOS integral eld mode: line emission seems to be concentrated within the centers of the interacting partners while the tidal tails them selves are not detected H narrow band slices. Three knots, previously identied as TDG candidates near the end of the southern tidal tail, are detected in [Diii]5007 emission. For two of them the velocity pro les were con rm ed. How ever, the strongest velocity gradient in the outerm ost TDG candidate (knot a') as measured on FORS2 data is not con rm ed by the VIMOS datacube. The reason for this discrepancy is unknown.

To solve this mystery and nd more clues to the origin of the velocity elds seen in this interacting system, further, deeper IFU observations, taken with appropriate dither o sets to facilitate more accurate sky subtraction, are required.W ith other instrum ents like e.g. the GM OS IFU would be possible to cover both the Ca-triplet and H in the same exposures and directly com pare the stellar velocity eld with ionized gas dynam ics. As good S/N is required to detect the absorption lines, this can only be done in the brighter northern tidal tail.

A cknow ledgem ent. PMW received nancial support through the D 3D net project from the G erm an Verbundforschung of BM BF (grant 05AV 5BAA).We are grateful to A na M on real-Ibero and Lise C hristensen for practical hints on IFU data handling. The data we discuss was taken in service m ode at Paranal (ESO Program 074 B-0629).

References

- Duc, P.-A., Brinks, E., Springel, V., et al., 2000, AJ 120, 1238
- Duc, P.-A., Brinks, E., Wink, J.E., & Mirabel, IF., 1997, A&A 326, 537
- Duc, P.-A. & Mirabel, IF., 1998, A&A 333, 813
- Hibbard, J.E., Guhathakurta, P., van Gorkom, J.H., & Schweizer, F., 1994, AJ 107, 67
- Weilbacher, P.M., Duc, P.-A., & Fritze-von Alvensleben, U., 2003, A&A 397, 545
- Weilbacher, P.M., Duc, P.-A., Fritze-von Alvensleben, U., et al., 2000, A&A 358, 819
- Weilbacher, P.M., Fritze-von Alvensleben, U., Duc, P.-A., & Fricke, K.J., 2002, ApJ 579, L79