# The in uence of grain rotation on the structure of dust aggregates

Paszun D<sup>a</sup>, Dom inik C.<sup>a</sup>

<sup>a</sup>A stronom ical Institute "Anton Pannekoek", University of Am sterdam, Kruislaan 403 1098 SJ Am sterdam, The Netherlands

#### A bstract

W e study the e ect of rotation during the collision between dust aggregates, in order to address a m ism atch between previous m odel calculations of B row nian m otion driven aggregation and experim ents.W e show that rotation during the collision does in uence the shape and internal structure of the aggregates form ed. The e ect is lim ited in the ballistic regim e w hen aggregates can be considered to m ove on straight lines during a collision. How ever, if the stopping length of an aggregate becom es sm aller than its physical size, extrem ely elongated aggregates can be produced.W e show that this e ect m ay have played a role in the inner regions of the solar nebula w here densities were high.

Keywords: Origin, Solar System, Interplanetary Dust

#### 1 Introduction

D ust aggregation plays a central role in m ost theories of planet form ation. W hile for the form ation of giant planets, disk instability scenarios continue to be discussed (Boss, 1997), dust aggregation certainly stands at the beginning of the form ation of planetesim als and therefore of the terrestrial planets. Independent of the detailed process that nally form s planetesim als (e.g. W eidenschilling, 1980; Youdin and Shu, 2002; C uzzi et al., 2001), dust grains rst need to grow and settle towards the m idplane of the disk. P lanet form ation therefore starts with the rst steps of dust aggregation, when dust grains inherited by the disk from the interstellar m edium start to collide and grow. This rst step of grow th is governed by B row nian m otion aggregation. In the high density regions of the disk, dust/gas coupling is so tight that the m ain source of relative velocities between grains are the random m otions produced by individual collisions between gas atom s and the grains. Understanding the Brownian motion phase of dust aggregation therefore means understanding the rst step towards planets in disks.

At the very low collision velocities produced by Brownian motion (cm/s), grains always stick and no restructuring is occurring in the collision (D om in ik and T ielens, 1997; Blum and Wurm, 2000). The structure of aggregates formed in this require is therefore indeed a pure probe of the physical processes driving grow th with B row nian m otion of the particles. Theoretically, grow th by B row nian motion was studied for example by K empfet al. (1999). They calculated the motion of micron sized dust grains enclosed in a box of approximately constant dust number density. D i usion, caused by the presence of a gaseous medium, produces relative motion of grains and leads to the growth of dust. The orientation of the (spherically not symmetric) aggregates was not followed during the computations. Instead, in order to random ize the relative orientation during collisions, the orientations of collision partners just before a collision was selected random ly.W ith these initial conditions, the aggregates were left to collide, without considering rotation during the collision. The calculations show a slow growth of the dust aggregates with time. At any time, the box contains a distribution of aggregate shapes which is characterized by a distribution of fractal dimensions. The mean fractal dimension found in the num erical study is around  $D_f = 1.8$ .

On the experim ental side, a series of low-gravity experim ents has been conducted (B lum et al., 2000; K rause and B lum, 2004) in order to study the growth of fractal aggregates under B rownian motion conditions. The results con rm many expected aspects of this growth regime, but also showed unexpectedly elongated aggregates, in contrast with the predictions by K empfet al. (1999).

The authors qualitatively argue that rotation of the aggregates during the collision m ight lead to m ore elongated aggregates, because the probability of achieving contact between two aggregates at larger separations increases if the aggregates rotate.

To compare experimental and theoretical results, it is necessary to quantify the visual impression of elongation. In this study we will be using two diment ways to do so:

(1) O ne can de ne an elongation factor as the ratio of maximum to minimum diameter of aggregate

$$f_{el} = \frac{d_{max}}{d_{min}}$$
(1)

where  $d_{m in}$  is measured in perpendicular direction to  $d_{m ax}$ . Experim entally, this quantity has to be derived from a few, or even a single picture. W hen m easuring this value for m odel aggregates, we therefore choose three di erent projections of the aggregate and determ ine  $d_{m ax}$  and  $d_{m in}$  from these in ages.

(2) We can also derive the fractal dim ension of the aggregates. This quantity can be computed by measuring them ass{radius relation of aggregates of dierent mass m and size R. If that relation follows a power law

$$m / R^{D_f}$$
 (2)

then  $D_f$  is called the fractal dimension of the aggregate. The fractal dimension measured in the experiments is 1.4, and typical values predicted by the numerical calculations are around 1.8, indicating a mism atch.

W hile the idea of particle rotation is appealing as a mechanism to increase the elongation of dust aggregates, this has never been shown quantitatively. In this paper we investigate the in uence of rotation on the structure of aggregates form ed under B rownian motion conditions.

# 2 Themodel

In order to test the in uence of rotation on the growth of aggregates we developed a model for collisions of rotating aggregates. Since we are interested in the low velocity regime, collisions are not energetic enough to cause restructuring. These processes are therefore neglected, and aggregates are treated as rigid entities. Our code calculates the equation of motion of two aggregates set on collision course with each other. When both aggregates get into the rst contact between any two constituent grains, a new aggregate is form ed. The structure of this aggregate is determined by the position and orientation of the two aggregates at the moment of contact { no further changes occur after this moment.

In all our calculations we used the same m onom ers sizes and m asses as in the CODAG experiment (K rause and B lum, 2004). Each m onom er was 1 m in diameter and its mass was 1.0  $10^{15}$  kg. The temperature assumed in our calculations was 300K, again like in the experiment.

In a fully general calculation, one would have to consider an ensemble of particles and follow the growth of this ensemble. However, it has been shown experimentally that the mass distribution function during Brownian motion growth is very narrow, so that a mono-disperse approximation works very well (K rause and B lum, 2004). We therefore proceeded in the following way: Starting with dimers, we rst compute a large number of collisions between dimers, with random initial position, rotational orientation, and in pact parameter. The translational and rotational velocities are taken from a M axwellian distribution for a given temperature T. The resulting quadrum er structures are stored in a database. In the next step, we collide two quadrum ers selected random ly from the database. A database of quadrum er structures is built in this way. In a similar way, we produce aggregates with larger sizes, always containing  $2^n$  grains.



Fig. 1. Schematic sketch of collisions between two aggregates in two cases. a – non (ballistic collision with mean free path of an aggregate shorter than the aggregate size. Velocities change every f.b – ballistic collisions with mean free path longer than size of an aggregate.

In order to solve the problem we have studied the in uence of rotation in two separate cases. The rst one assumes ballistic collisions where the mean free path of an aggregate is longer than its size. The second assumes aggregation in a non {ballistic regime. The stopping length in this case is shorter than the size of the aggregate.

# 2.1 Ballistic collisions

In the ballistic regime, the mean free path of all involved particles is larger than the size of the largest particle. In this case, the center of mass of each particle moves on a straight line during the collision. In physical terms, the ballistic limit is reached in the limit of low (but non {zero, because the gas still needs to cause B rownian motion of the grains) gas densities. This is generally assumed to be the case both in protoplanetary disks, and in microgravity experiments described above.

W hen aggregates are colliding without rotation and on linear trajectories, elongated aggregates are form ed when the rst contact is made between grains on the outside of the aggregate, while the two aggregates are more or less aligned. Depending on the initial orientation, this situation is realized for di erent in pact parameters: If the aggregates are aligned along the direction of relative motion between the colliding aggregates, a small in pact parameter is needed. If the alignment is perpendicular to the collision direction, a large im pact parameter produces the most elongated result, while a small im pact parameter would lead to a more compact structure. If the aggregates are rotating during approach, the chances that the contact will be made early with an elongated geometry increase. For this process to be e cient, the linear speed of grains far away from the center of mass should be comparable or larger than the translational motion. Therefore, the larger the ratio of rotational and translational speeds are, the greater the average elongation of the aggregate becom es.

This can be achieved in two ways. Superthem algrain rotation does occur in interstellar space, where it is responsible for the alignment of non{spherical grains with the galactic magnetic eld and in this way produces the interstellar polarization (Purcell, 1979). With superthem al rotation, a fast circular motion is induced while the linear velocity remains them al. How ever, superthem al rotation relies on sm all forces like non{isotropic absorption and scattering of light (D raine and W eingartner, 1996), or H<sub>2</sub> form ation on speci c sites (Purcell, 1979), to accelerate the grains. The non{them al rotation speeds can only be sustained at extrem ely low gas densities. In fact, the densities prevailing in protop lanetary disks are prohibitive, and superthem al rotation can be ruled out there (O sænkopf, 1993).

On the other hand, the e ective linear translational speed can be slowed down by embedding the grains into gas of high density, leading into the regime of non {ballistic collisions.

## 2.2 Non {ballistic collisions

The mean free path  $l_{o} = v_{th f}$  of a dust grain is the distance an aggregate with mean therm alvelocity  $v_{th}$  can move during one friction time. The friction time of a dust grain is

$$f = \mathbf{w} \frac{\mathbf{m}}{\mathbf{a}} \frac{1}{\mathbf{g} \mathbf{V}_{\mathbf{m}}}$$
(3)

where m,  $_{a}$ ,  $_{g}$  and  $v_{m}$  are mass, aerodynam ical cross section, gas density and m ean therm all velocity of the gas molecule respectively and " is a proportionality coe cient which is around 0:6 (Blum et al., 1996).

If g increases, the mean free path decreases accordingly. When the mean free path becomes similar to the largest aggregate involved in the collision, the assumption of a ballistic collision is no longer valid. Instead, the particle execute a random walk both in linear motion and in rotation. On average, the aggregates will spent a longer time at large distances before moving closer together. This elect increases the chance of creating a contact already at large

distances, i.e. a strongly elongated aggregate. Fig. 1a presents the schematic picture of a non {ballistic collision.

#### 3 Results

In order to investigate the in uence of rotation on the production of elongated aggregates, we calculated growth of rotating aggregates in the range from ballistic to highly non ballistic collisions.

# 3.1 Ballistic collisions with and without therm al rotation



Fig. 2. Distribution of elongation factors, with peak value normalized to unity. From top to bottom : non{rotation case, ballistic therm al rotation, therm al rotation for di erent gas densities  $_{\rm g}$  in g cm  $^3$ . The distributions are shifted vertically for better visibility.

D istributions of  $f_{el}$  for ballistic collisions are plotted in g.2. In order to be able to compare with the results of K empfet al. (1999), we also computed a

case with rotation turned of. The two top diagrams show the non{rotating case and them al rotation case in the ballistic regime. For the non{rotating case, a very narrow distribution results. This distribution peaks at a value 1:6. For therm al rotation, the peak value shifts to  $f_{el}$ fel 2 and the distribution becomes wider. For both cases, a very small fraction of particles with elongation factors 4 and larger are produced, due to fortuitous initial conditions in the computation. Fig. 3 shows the mean elongation factors as a function of aggregates size. Each point is plotted together with its 1 errorbars. The no rotation case clearly has the lowest elongations and, as in the therm al rotation case, the line is alm ost at for bigger sizes, meaning the grow that ects the elongation very weakly. The elongation factors for rotating aggregates are shifted upwards from values of about  $f_{el} = 1.8$  to about  $f_{el} =$ 23. The e ect of rotation is also clearly seen in the fractal dimensions (g.4)which reach level of  $D_{f}$ 1:7 for non{rotating collisions. Therm al rotation causes the fractal dimension to drop to a value of 1.46.

3.2 Rotating aggregates outside of the ballistic lim it

Further calculations were done for higher gas densities. In this case, the friction length  $l_{\rm b}$  becomes comparable to or shorter than the size of the colliding aggregates. While the thermal velocity decreases with increasing size (m ass) of an aggregate, f const for small, non {compact aggregates. Consequently, the mean free path  $l_{\rm b} = fv_{\rm th}$  decreases with increasing particle size. We performed calculations for several dierent gas densities. By tuning the gas density, we can study the transition from the ballistic to the non {ballistic regime.Fig.2 shows the resulting distribution of elongation factors for dierent gas densities.Fig.3 presents the relation between the mean elongation factor and the aggregate size for dierent gas densities.

The mean elongation factor  $f_{el}$  depends strongly on the gas density  $_{g}$  in the transition regime. Increasing the gas density from 10 <sup>6</sup> to a few times 10 <sup>5</sup>g cm <sup>3</sup> strongly broadens the elongation factor distribution and shifts the peak value to  $f_{el}$  4. Extrem e values of up to  $f_{el}$  = 10 are reached in the tail of the distribution function. The elongation factor also depends on size of aggregates: Larger aggregates reach the largest elongations. This is the result of the decrease of the mean free path with increasing aggregate size. The elongation factor in our calculations follows a power law dependence on the gas density. The power index is size dependent. A ggregates consisting of 32 m onom ers follow a power law with index of 0.18, while the index for thosem ade of 16 m onom ers is 0.13. This means that the latter aggregates can reach a mean elongation factor value about 3.4 for gas density  $_{g}$  = 3.88 10  $_{5}$  g cm  $_{3}$ . For this gas density and aggregates bigger than 32 m onom ers the mean elongation



Fig. 3. Mean elongation factor for di erent gas densities.  $_g = 0$  indicates the limiting case for low densities (mean free path much larger than aggregate sizes). The bottom line shows the elongation factor for non{rotation case. Three boxes are measured elongation factors of agglom erates form ed in the CODAG experiment with 1 errorbars (Blum & Krause personal communication).

factor reaches values above 4.5.

Fig. 3 presents also results of the CODAG experiment (B lum & Krause, personal communication). Three points together with their 1 errorbars show the ratio of maximum to minimum diameter of aggregates formed during the experiment. An excellent t with our calculations seems to be  $_{\rm g}$  = 1:94 10  $^6$  g cm  $^3$  or slightly higher. The density used in the experiment was  $_{\rm g}$  = 2:25 10  $^6$  g cm  $^3$  (Krause and B lum, 2004).

W e also calculated the fractal dimension D<sub>f</sub> by tting a power law function to the plot of mean aggregatem assversus mean radius of gyration  $r_g$ . The average was taken over the entire distribution of particles created for the given mass. The fractal dimension as a function of the gas density is shown in Fig.4. For large densities, D<sub>f</sub> appears to follow a power law with index = 0.062. Thus for a gas density  $_g$  = 3.88 10 <sup>5</sup> the fractal dimension reaches the value 1.11. At low gas densities, the fractal dimension asymptotically approaches a value of 1.46, slightly larger than D<sub>f</sub> = 1.41 as observed in the CODAG experiment. Indeed, it turns out that the CODAG experiment is operating close to, but



Fig. 4. Fractal dimension as a function of gas density. The upper x-axis shows the mean free path of a monomer in units of the monomer radius. The horizontal lines show the CODAG measurements (D<sub>f</sub> = 1:4), the limiting result for lim<sub>10</sub>D<sub>f</sub> = 1:46, and the result for non{rotating aggregates D<sub>f</sub> = 1:72. The inset plot shows an example of fractal dimension determination by slope tting in an r<sub>g</sub> versus m plot, at a density of  $_{\rm g} = 324$  10<sup>6</sup>.

not safely within the ballistic limit. A gas density of 2.25  $10^{6}$  g cm<sup>-3</sup> is in fact entirely consistent with D<sub>f</sub> = 1.4. At this density, the mean free path of a particle is only about 1.5 grain diameters. The experiments therefore are not in the ballistic limit, but start to feel the in uence of random walk during the collision. Further increasing the gas density should strongly enhance this e ect, and fractal dimensions very close to unity should show up when  $_{g}$  exceeds a few times 10<sup>-5</sup> g cm<sup>-3</sup>.

### 4 D iscussion

#### 4.1 The ballistic regime

In the ballistic limit, rotation increases the collisional cross-section by producing more opportunities for a contact while two aggregates are passing each other. In the non{rotation case only speci c 'lucky' parameters (appropriate orientation and impact parameter) lead to an elongation, while rotation allows a much larger set of possible collision parameters to be e ective. This e ect leads to more elongated structures, because it produces signi cantly more opportunities for collisions with larger in pact parameters.

#### 4.2 The non-ballistic regime

In the non {ballistic regime, the random walk executed by the collision partners during a collision e ectively reduces the mean (or e ective) relative translational velocity. The typical distance traveled from the original position in a random walk scales with the square root of the number of steps N, we can de ne an e ective translational speed by

$$\langle v \rangle = \frac{p \frac{1}{N}}{N_{f}}$$
 : (4)

If we take  $\stackrel{\text{p}}{N} = L$  where L is size of the aggregate, then the mean e ective translational velocity during random walk is inversely proportional to the gas density  $_{\text{g}}$  (see eq. 5, 3).

$$\langle v \rangle = \frac{l_b^2}{L_f}$$
 : (5)

As the gas density increases, the elective velocity decreases and indeed leads to very long and slow collisions between aggregates. During that time, the aggregates also execute a random walk in rotation and in this way expose di erent parts toward each other. In the limit of large gas densities, this leads to contact being made always at maximum distance, and consequently to fully linear structures.

It is important to remember that we are only computing collisions of equal size aggregates, i.e. pure cluster-cluster aggregation (CCA). This simpli cation will tend to exaggerate the elongation at a given size. If m onomers and small aggregates contribute signi cantly to the growth, slightly m ore compact aggregates are produced. In fact, if a size distribution of impactors is involved in the growth of a target, the growth physics is an intermediate case between pure CCA and particle-cluster aggregation (PCA). As an indication, we can compare the fractal dimension of 1.7 we found for non{rotating aggregates in pure CCA with the fractal dimension of 1.8 found by K empfet al. (1999) for a realistic size distribution.

#### 4.3 In uence on aggregation timescale

E longated aggregates, in uenced by rotation and aggregation in non {ballistic regime, expose a larger surface to the gas and are easier targets for possible collisions with other dust grains. Just like for the particle mass, one can introduce a fractal dimension for the cross-section D and write the relation between aggregate shape and cross-section as

$$/ r_{g}^{D}$$
: (6)

Thus for limiting case of compact agglom erates D = 2 while for linear grains it is 1. To see how the crossection changes with aggregate size, one can com – bine both fractal dimensions D and  $D_f$ :

$$/ (m = m_0)^{\frac{D}{D_f}} :$$
 (7)

The cross-section of aggregates is related to aggregation timescale by

$$t / \frac{1}{nv}$$
 (8)

where t is aggregation time, n is number density of dust grains and v is velocity. We calculated cross-sections of aggregates with dierent fractal dimensions in order to investigate the timescale of aggregation. Fig. 5 shows the timescale divided by the timescale for the most compact aggregates. Each line corresponds to dierent fractal dimension D<sub>f</sub>. The most interesting are D<sub>f</sub> = 1.8 which corresponds to the aggregates formed in the numerical calculations by K empfet al. (1999) and fractal dimension D<sub>f</sub> = 1.4 which was obtained in the CODAG experiment (K rause and B lum, 2004). The timescale is shorter for elongated and open aggregates because of the larger cross-sections. The dierence increases with increasing size of aggregates.

#### 4.4 Application to protoplanetary disks

To assess the relevance of the e ects discussed in the current paper in a protop lanetary disk, we need to compute the stopping length of dust grains as a function of aggregate size and location in the disk. The boundary condition for ballistic and non {ballistic collisions is reached when the mean free path equals the physical size of the dust aggregate approximately given by the radius of gyration:  $l_{p}(g;T) = r_{g}(m)$ . This condition leads to a relation between cluster size and gas density, indicating how large the cluster should be at a



Fig. 5. A ggregation timescale t /  $\frac{1}{nv}$  divided by aggregation timescale for compact agglomerates as a function of mass. Dierent lines correspond to dierent fractal dimensions. From top to bottom D<sub>f</sub> = 3.0; 2.8; 2.5; 1.8; 1.4; 1:1:

given density in order to start feeling the e ect of elongation enhancem ent by non (ballistic collisions.

W e will make the sim plifying assumption that the stopping time of an aggregates is (at a given density) independent of size. This assumption is valid for very small aggregates, and for aggregates with very open structure, i.e. low fractal dimension.

W e use the de nition of the radius of gyration given by eq.2 and transform it into

$$r_{g}(m) = Bm^{\frac{1}{D_{f}}}$$
(9)

where  $B = \frac{1}{A^{1-D}f}$  and A is a proportionality coe cient from eq.2. Then the relation between cluster m ass and gas density relation can be derived as

$$m^{\frac{1}{D_{f}}} = \frac{\mathbf{m}}{B} \frac{3m_{m}}{8} \frac{1}{n_{0}r^{2}} \mathbf{j}$$
(10)

where  $m_m$  is a mean mass of a gas molecule and  $n_0$  is number of monomers in the aggregate. In order to  $nd n_0$  we substitute the cluster mass by  $m = n_0 m_0$ 

where  $m_0$  is the mass of a monom er.

$$n_{0} = \frac{"}{B} \frac{3m_{m}}{8} \frac{m_{0}^{0.5}}{r^{2}} \frac{1}{p_{f}}}{r^{2}} \frac{2D_{f}}{r^{2}}$$
(11)

Eq.11 show show the critical size of a cluster is dependent on the gas density. It also reveals a dependence on m onom er size. W e applied eq. (11) to the H ayashi m odelof the solar nebula (H ayashi et al., 1985). Fig. 6 shows, for the m idplane



Fig. 6. A ggregate m ass for which in the m idplane of the solar nebula the m ean free path  $l_{\rm free}$  equals the radius of gyration  $r_{\rm g}$ . D i erent lines correspond to di erent m onom er radii  $r_0$ . The dotted line represents a m onom er size of  $r_0 = 0.005$  m, the dashed line  $r_0 = 0.05$  m, and the solid line  $r_0 = 0.5$  m. The shaded area below each line show s the ballistic regim e for corresponding m onom er size. The area above each line represents the non {ballistic regim e.At the top of the diagram, we indicate the m idplane density corresponding to the distance from the Sun, according to the H ayashim odel (H ayashi et al., 1985).

density of the model, the critical size of a cluster as a function of distance from the star. Each line represents this relation for a di erent monom er size. Thus aggregates with a fractal dimension of  $D_f = 1.46$ , are formed below the line, while above that line more elongated grains are produced. Each line is a border between the ballistic and non {ballistic regimes for a given monom er size. For

example, at a distance below 1 AU from the star, the gas density strongly in uences the shape of aggregates form ed through Brownian motion, if the aggregates consist of more than a few tens of 1 m grains. Consequently, the CODAG experiment reproduced conditions present in the inner part of the protoplanetary disk while its applicability to the low density regions in the outer disk is limited. At the CODAG density of  $_g = 2.25 \ 10^{-6} \, \mathrm{g\,cm}^{-3}$ , the critical size is about 1.5 m icron sized monom ers.

# 5 Conclusions

W e have studied the e ect of aggregate rotation during collisions. The results show that rotation must de nitely be treated correctly when modeling grow th due to aggregation, or the geometrical structure of the resulting aggregates will be incorrect.

Rotation plays a role because it enhances the probability that two approaching aggregates make contact early on during the collision, between outer constituent grains. For the most simple case of ballistic collisions, during which the colliding aggregates move on a linear path, ignoring aggregate rotation increases the fractal dimension from 1.46 to 1.7 in the limit of pure clustercluster aggregation.

Thise ect becomes strongly enhanced if the density of the surrounding medium becomes so large that the stopping length of an aggregate becomes shorter than the aggregate size, i.e. in the non {ballistic limit.We nd that reducing the stopping length to the monomer radius results in a fractal dimension of 1.25.W hen the stopping length is reduced to one tenth of the monomer radius, the fractal dimension drops to 1.1.

For the solar nebula, we nd that non {ballistic collisions play a role in the innerm ost regions of the disk for even very sm all aggregates m ade of a few m -sized grains. For sm aller m onom ers and/or further away from the Sun, enhanced elongation can be expected for larger aggregates, m ade of a few 100 or 1000 grains. In the outer disk, all collisions m ay be considered ballistic.

A cknow ledgm ents

Thiswork was supported by the Nederlandse Organisatie voor Wetenschapelijk Onderzoek, Grant 614.000.309.We thank J.Blum for an insightful referee report, and for providing data obtained in CODAG microgravity experiment. W e also thank the second (anonym ous) referee for useful comments that lead to a restructuring of the manuscript.

#### References

- Blum, J., Wurm, G., Jan. 2000. Experiments on Sticking, Restructuring, and Fragmentation of Preplanetary Dust Aggregates. Icanus 143, 138{146.
- Blum, J., Wurm, G., Kempf, S., Henning, T., Dec. 1996. The Brownian Motion of Dust Particles in the Solar Nebula: an Experimental Approach to the Problem of Pre-planetary Dust Aggregation. Icarus 124, 441 [451.
- B lum, J., W unm, G., Kempf, S., Poppe, T., K lahr, H., Kozasa, T., Rott, M., Henning, T., Dorschner, J., Schrapler, R., Keller, H. U., Markiewicz, W. J., Mann, I., Gustafson, B. A., Giovane, F., Neuhaus, D., Fechtig, H., Grun, E., Feuerbacher, B., Kochan, H., Ratke, L., ElGoresy, A., Mor ll, G., Weidenschilling, S. J., Schwehm, G., Metzler, K., Ip, W. H., Sep. 2000. Growth and Form of Planetary Seedlings: Results from a Microgravity Aggregation Experiment. Physical Review Letters 85, 2426 (2429.
- Boss, A.P., 1997.G iant planet form ation by gravitational instability. Science 276, 1836{1839.
- Cuzzi, J. N., Hogan, R. C., Paque, J. M., Dobrovolskis, A. R., Jan. 2001. Size-selective Concentration of Chondrules and Other Sm all Particles in Protoplanetary Nebula Turbulence. ApJ 546, 496{508.
- Dom inik, C., Tielens, A.G.G.M., May 1997. The Physics of Dust Coagulation and the Structure of Dust Aggregates in Space. ApJ 480, 647{+.
- D raine, B.T., W eingartner, J.C., Oct. 1996. Radiative Torques on Interstellar G rains. I. Supertherm al Spin-up. ApJ 470, 551{+.
- Hayashi, C., Nakazawa, K., Nakagawa, Y., 1985. Form ation of the solar system. In: Protostars and Planets II. pp. 1100{1153.
- K em pf, S., P falzner, S., H enning, T.K., Oct. 1999. N-Particle-Simulations of Dust G row th. I. G row th D riven by B row nian M otion. Icanus 141, 388{398.
- K rause, M ., B lum, J., Jul. 2004. G row th and Form of P lanetary Seedlings: Results from a Sounding Rocket M icrogravity Aggregation Experiment. Physical Review Letters 93 (2), 021103{+.
- O sænkopf, V., Dec. 1993. Dust coagulation in dense molecular clouds: The formation of u y aggregates. A & A 280, 617{646.
- Purcell, E.M., Jul. 1979. Supratherm al rotation of interstellar grains. ApJ 231, 404 (416.
- W eidenschilling, S.J., Oct. 1980. Dust to planetesim als Settling and coagulation in the solar nebula. Icanus 44, 172 {189.
- Youdin, A.N., Shu, F.H., Nov. 2002. Planetesim alform ation by G ravitational Instability. ApJ 580, 494 [505.