Evolution of the Color-M agnitude Relation in H igh-Redshift C lusters: B lue Early-Type G alaxies and Red Pairs in RDCS J0910+ 5422

S.Met¹, J.P.B lakeslee¹, S.A. Stanford^{2;3}, B.P.Holden⁴, P.Rosati⁶, V. Strazzullo^{20;6}, N.Hom eier¹, M.Postman^{1;5}, M.Franx¹², A.Rettura⁶, H.Ford¹, G.D. Illingworth⁴, S. Ettori¹⁹, R.J.Bouwens⁴, R.Dem arco¹, A.R.Martel¹, M.C lampin⁵, G.F.Hartig⁵, P.
E isenhardt⁷, D.R.Ardila¹, F.Bartko⁸, N.Ben tez¹⁸, L.D.Bradley¹, T.J.Broadhurst⁹, R.A. Brow n⁵, C.J.Burrow s⁵, E.S.Cheng¹⁰, N.J.G.Cross¹⁷, P.D.Feldm an¹, D.A.Golim ow ski¹, T.Goto¹, C.Gronwall¹³, L.Infante¹⁴ R.A.K im ble¹¹, J.E.Krist⁵, M.P.Lesser¹⁵, F.
M enanteau¹, G.R.Meurer¹, G.K.Miley¹², V.Motta¹⁴, M.Siriannf⁵, W.B.Sparks⁵, H.D. Tran¹⁶, Z.J.T svetanov¹, R.L.W hite⁵, & W.Zheng¹

ABSTRACT

The color-m agnitude relation has been determ ined for the RDCS J0910+ 5422 cluster of galaxies at redshift z = 1.106. Cluster m em bers were selected from HST A dvanced C am era for Surveys (ACS) in ages, com bined with ground {based near{IR in aging and optical spectroscopy. P ostm an et al. (2005) m orphological classi cations were used to identify the early-type galaxies.

The observed early (type color(m agnitude relation (CMR) in (i_{775} z₈₅₀) versus z₈₅₀ shows an intrinsic scatter in color of 0.060 0.009 m ag, within 1⁰ from the cluster X (ray em ission center. Both the ellipticals and the S0s show sm all scatter about the CMR of 0.042 0.010 m ag and 0.044 0.020 m ag, respectively. From the scatter about the CMR, a m ean lum inosity (w eighted age $\bar{t} > 3.3$ G yr ($z_f > 3$) is derived for the elliptical galaxies, assuming a simple stellar population m odeling (single burst solar m etallicity). This is consistent with a previous study of the cluster RDCS 1252.9-292 at z= 1.24 (B lakeslee et al.).

Strikingly, the S0 galaxies in RDCS J0910+ 5422 are system atically bluer in $(i_{775} z_{850})$ by 0.07 0.02 m ag, with respect to the ellipticals. The blue S0s are predom inantly elongated in shape; the distribution of their ellipticities is inconsistent with a population of axisym m etric disk galaxies viewed at random orientations, suggesting either that they are intrinsically prolate or there is some orientation bias in the S0 classi cation. The ellipticity distribution as a function of color indicates that the face-on S0s in this particular cluster have likely been classi ed as elliptical. Thus, if anything, the o set in color between the elliptical and S0 populations m ay be even m ore signi cant.

The color o set between S0 and E corresponds to an age di erence of 1 Gyr, for a singleburst solar m etallicity m odel. A lternatively, it could be the result of a di erent star form ation history; a solar m etallicity m odel w ith an exponential decay in star form ation w ill reproduce the o set for an age of 3.5 Gyr, i.e. the S0s have evolved gradually from star form ing progenitors. The color o set could also be reproduced by a factor of 2 di erence in m etallicity, but the two populations would each need to have very sm all scatter in m etallicity to reproduce the sm all scatter in color.

The early (type population in this cluster appears to be still form ing. The blue early-type disk galaxies in RDCS J0910+ 5422 likely represent the direct progenitors of the more evolved S0s that follow the same red sequence as ellipticals in other clusters.

Thirteen red galaxy pairs are observed and the galaxies associated in pairs constitute 40% of the CMR galaxies in this cluster. This nding is consistent with the conclusions of van Dokkum and Tran et al. that most of the early {type galaxies grew from passive red m ergers.

Subject headings: galaxies: clusters: individual (RDCS J0910+5422) { galaxies: elliptical and lenticular | galaxies: evolution

1. Introduction

The Advanced C am era for Surveys (ACS; Ford et al. 2002), by virtue of its high spatial resolution and sensitivity, allows us to study galaxy clusters in great detail up to redshifts of unity and beyond. At these redshifts, galaxy clusters are still assembling and galaxies are evolving towards the populations that we observe today. Recent results from our ACS Interm ediate Redshift C luster Survey (B lakeslee et al. 2003a; Lidm an et al. 2004; D em arco et al. 2005; G oto et al. 2005; H olden et al. 2005; P ostm an et al. 2005) have shown that galaxy clusters at redshift around unity show

³Institute of Geophysics and Planetary Physics, Law rence Liverm ore National Lab, Liverm ore, CA 94551

⁵Space Telescope Science Institute, 3700 San Martin Drive, Baltim ore, M D 21218

⁶European Southern Observatory, Karl-Schwarzschild-Str. 2, D-85748 Garching, Germany

 $^7\,{\rm Jet}$ P ropulsion Laboratory, C alTech, 4800 O ak G rove D rive, P asadena, C A $\,91125$

⁸Bartko Science & Technology, 14520 Akron Street, Brighton,CO 80602.

⁹School of P hysics and A stronom y, TelA viv U niversity, TelA viv 69978, Israel

¹⁰Conceptual Analytics, LLC, 8209 W oburn Abbey Road, G lenn Dale, M D 20769.

 $^{11}{\rm N}\,{\rm A}\,{\rm SA}$ G oddard Space F light C enter, C ode 681, G reenbelt, M D 20771.

 $^{12}\mbox{Leiden}$ O bærvatory, Postbus 9513, 2300 RA Leiden, N etherlands.

¹³D ept. of A stronom y & A strophysics, Penn State University, University Park, PA 16802.

 $^{14}{\rm D}$ ept. de A stronom a y A strof sica, Ponti cia Universidad C at lica, C asilla 306, Santiago 22, C hile.

 $^{15}\mbox{Stew}$ and O bservatory, U niversity of A rizona, Tucson, A Z 85721.

 $^{16}{\rm W}$. M . K eck O bservatory, 65-1120 M am alahoa H w y., K am uela, H I 96743

 $^{17}\mathrm{R}\,\text{oyal}$ O bservatory E dinburgh, B lackford H ill, E dinburgh, E H 9 3H J, U K

¹⁸ Instituto de A strof sica de Andaluc a (CSIC), Camino Bajo de Huetor 50, Granada 18008, Spain

¹⁹ IN AF - O sservatorio A stronom ico, via Ranzani 1, 40127 Bologna, Italy

²⁰D ipartim ento di Scienze F isiche, U niversita Federico II, I-80126 N apoli, Italy m any similarities with local clusters, in terms of galaxy populations and their distribution, but also signi cant di erences in galaxy m orphology, ellipticity, and m ass{lum inosity ratios. The strongest evolution observed in the early (type population is a de cit of a SO population in this sample when compared to lower redshift samples (Postm an et al. 2005). This would give evidence that the form ation of the SO population is still under way in clusters at redshift unity.

One of the most striking similarities is that the tight relation between early-type galaxy colors and lum inosities that applies locally (the color{ m agnitude relation; CMR) is already in place at redshifts as high as z 13 (e.g. Stanford et al. 1997; Mullis et al. 2005). The CMR in local sam ples of galaxy clusters presents universal properties, in terms of scatter and zero point (Bower et al. 1992; van Dokkum et al. 1998, Hogg et al. 2004; Lopez{Cruz et al. 2004; Bell et al. 2004; Bemardiet al. 2005; M cIntosh et al. 2005) that evolve back in time in agreement with passively evolving m odels (E llis et al. 1997; Stanford, E isenhardt, & Dickinson 1998; van Dokkum et al. 2000, 2001; B lakeslee et al. 2003a; Holden et al. 2004; De Lucia et al. 2004; Blakeslee et al. 2005). ACS enables accurate m easurem ent of the scatter around the CMR, with enough precision to seriously constrain galaxy form ation age, which is in possible to obtain from ground { based data (see for example Holden et al. 2004). The measurement of the CMR scatter of the rst cluster in our ACS cluster survey, RX J1252.9-292, perm itted us to constrain the m ean lum inosity {weighted age for the ellipticals to be > 2:6 G yr (z > 2:7) (B lakeslee et al. 2003a), based on simple modeling. In this paper, we extend the results obtained in B lakeslee et al. (2003a) to RXJ 0910+ 5422.

RXJ 0910+ 5422 is part of the ACS cluster survey (guaranteed tim e observation, GTO, program # 9919), that includes eight clusters in the red-shift range at 0.8 < z < 1.3, selected in the X {ray, optical and near{IR (Ford et al. 2004). RXJ 0910+ 5422 was selected from the ROSAT D eep C luster Survey (Rosati et al. 1998) and con rm ed with near-IR and spectroscopic observations by Stanford et al. (2002). Extensive follow up spectroscopy at the Keck Observatory has been carried out in a magnitude limited sam ple reaching K s = 20.0 m ag in the central3 arcm in (Stanford et al.

¹D ept. of P hysics & A stronom y, Johns H opkins U niversity, Baltim ore, M D 21218; sm ei@ pha.jhu.edu

 $^{^{2}\}mathrm{D}$ epartm ent of P hysics, U niversity of C alifornia, D avis, C A 94516

 $^{^{4}\,}L\,ick$ O bærvatory, U niversity of C alifornia, Santa C ruz, C A 95064

al.2005; in preparation). The mean redshift of the cluster was measured to be z = 1.106 (Stanford et al. 2002). In this paper, we combine ACS im aging with ground (based spectroscopy and near{IR im aging to constrain galaxy ages and form ation histories from the study of their color{m agnitude relation. W e discuss the properties of the elliptical (E) and lenticular (SO) populations separately in the light of sim ple galaxy form ation scenarios.

2. Observations

RXJ 0910+5422 was observed in M arch 2004 with the ACSWFC (Wide Field Camera) in the F775W (i_{775}) and F850LP (z_{850}) bandpasses, with total exposure times of 6840 s and 11440 s, respectively. The ACS W FC scale is 0.0500/pixel, and its eld of view is 210^{00} 204^{00} . The APSIS pipeline (Blakeslee et al. 2003b), with a Lanczos3 interpolation kernel, was used for processing the images. The ACS photom etric zero {points (AB system) are 25.654 m ag and 24.862 m ag in i_{775} and z_{850} , respectively (Sirianni et al. 2005). A Galactic reddening of E (B V) = 0.019 towards RXJ 0910+5422 was adopted (Schlegel et al. 1998), with $A_{1775} = 0.039$ and $A_{2850} = 0.029$ (Sirianniet al. 2005). The ACS W FC eld covers an area that at the redshift of this cluster, z = 1:106, corresponds to 1M pc² in the W M A P cosmology (Spergel et al. (2003): m = 0.27, = 0.73, h = 0.71, adopted as our standard cosm ology hereafter). Fig. 1 shows the ACS color im age with X { ray contours from Chandra ACIS (A dvanced CCD Im aging Spectrom eter) data that have been adaptively smoothed (Stanford et al. 2002). Near-IR JK_s and optical i-band im ages were obtained at Palom ar Observatory as de-

Optical spectroscopy of galaxies in RXJ 0910+ 5422 was obtained using the Low Resolution Im aging Spectrom eter (LR IS; Oke et al. 1995) on the Keck 1 and 2 telescopes (Stanford et al. 2005; in preparation). Our typicalerrors in redshift correspond to errors in velocity between 100 and 300 km /s. Objects for spectroscopy were chosen initially from the catalog of objects with K_s < 20:0 m ag (Vega m agnitudes) within the IR imaging area; outside of this area objects were chosen with i > 21 m ag from the iband image to llout masks. Our nal sample

scribed in detail by Stanford et al. (2002).

included 66% of the objects with K_s < 20.0 m ag. Spectra were obtained using the 400 lines mm⁻¹ grating for all runs except for the initial two discovery m asks as reported in Stanford et al. (2002). N ine m ore m asks were observed using LR IS during four runs between January 2001 and February 2003. U sually each m ask was observed in a series of four 1800 s exposures, with sm all spatial osets along the long axis of the slits. On average, the seeing was 0.9⁻⁰⁰. The blue side data were generally not used since the rest fram e wavelengths probed at z = 1:1 fall far to the blue of the spectral features of interest for galaxies in the cluster. In total, 149 redshifts were obtained.

The slit mask data were separated into slitlet spectra and then reduced using standard long-slit techniques. A fringe fram e was constructed for each exposure from neighboring exposures, each o set from the previous by 3^{00} , in an observing sequence for each mask, and then subtracted from each exposure to greatly reduce fringing in the red. The exposures for each slitlet were reduced separately and then co-added. One-dimensional spectra were extracted for each targeted ob ject, as wellas the occasional serendipitous source. W avelength calibration of the 1-D spectra was obtained from arc lamp exposures taken immediately after the object exposures. A relative ux calibration was obtained from long-slit observations of the standard stars HZ44, G191B2B, and Feige 67 (Massey & Gronwall 1990).

3. Object selection and photom etry

SExtractor (Bertin & A mouts 1996) was used to nd objects in the i_{775} and z_{850} in ages and m easure their m agnitudes. Threshold and deblending settings were used as in Ben tez et al. (2004). A 1though we have extensive spectroscopy, the ACS imaging reaches considerably deeper along the cluster lum inosity function. Thus, we have chosen to use colors $(i_{775} - z_{850})$ and $(J K_s)$ to isolate a set of probable cluster m em bers. In Fig. 2, the $(i_{775} - z_{850})$ and $(J K_s)$ colors are shown as a function of galaxy age, using BC03 stellar population m odels, redshifted to z=1:106. Early { type cluster m em bers would have ages of at least 0.5 Gyr, corresponding to $(i_{775} z_{850}) > 0.8 \text{ m ag}$ and $(J K_s) > 1:45 m ag$. At rst, we give a larger color m argin and select as potential cluster

m embers all m orphologically-classi ed early-type galaxies with 0.5 < $(i_{775} z_{850}) < 1.2$ m ag and $(J K_s) > 1.45$ m ag, down to $z_{850} = 24$ m ag (the lim iting m agnitude of P ostm an et al. (2005) m orphological classi cation, that included all clusters in our sample at redshift unity). Our results in this paper are based on this m orphological classi cation and a detailed discussion of the uncertainties in this classi cation can be found in that work. This selected sample includes 38 galaxies within the ACS eld.

Our nalcolors were measured within galaxy e ective radii (\mathbb{R}_{e}) , to avoid biases due to galaxy internal gradients, following the approach in Blakeslee et al. (2003a) and van Dokkum et al. (1998, 2000). R_e values were derived with the program GALFIT (Peng et al. 2002), constraining the Sersic index n 4 (as in Blakeslee et al. 2003a). To remove di erential blurring effects (the PSF is 10% broader in the z_{850} band) each galaxy image in both i_{775} and z_{850} was deconvolved using the CLEAN algorithm (Hogbom et al. 1974). The $(i_{775} z_{850})$ colors were measured on the deconvolved im ages within a circular aperture of radius equal to R_e, or 3 pixels, whichever is larger. 0 urm edian R_e is 5.5 pixels (13kpc at z=1.106). O ur nal results do not change (within the uncertainties) if the e ective radii are calculated via a two component (Sersic bulge + exponential disk) surface brightness decomposition technique using GIM 2D (Marleau & Sim and 1998; Rettura et al., in preparation), that permits us to better t the galaxy light pro le.

The photom etric uncertainties due to at elding, PSF variations, and the pixel-to-pixel correlation for ACS (Sirianniet al. 2005) were estimated by measuring the standard deviation of photom etry in the background for circular apertures in the range of the measured e ective radii. These photom etric errors were added in quadrature to the Poisson uncertainties in the measured uxes for each object. The derived errors in the colors are between 0.01 and 0.03 m ag down to $z_{850} = 24 \text{ m ag}$. SE xtractor MAG_AUTO were used for the z_{850} m agnitude in the color{magnitude relation; these are fairly robust, though may systematically miss a small fraction of the light (Ben tez et al. 2004).

We nally color{selected 34 early{type (E, S0 and S0/a) galaxies with 0.8 < (i_{775} z_{850}) < 1:1 m ag within 2⁰ from the cluster center, taken

as the center of the X {ray emission (Stanford et al. 2002). Im ages of the color{selected galaxies are shown in F ig 3, F ig 4 and F ig 5. M oreover, there are late{type galaxies with lum inosities that are sim ilar to the red{sequence bright early{type galaxies (F ig. 6). O f the 34 color{selected galaxies, 15 are spectroscopically con m ed cluster m em - bers, one (SO/a, with m agnitude $z_{050} = 24.2 \text{ m ag}$) is a con m ed non-m ember, and the others were not targeted for spectroscopy. The selection in (i₇₇₅ z₈₅₀) at z = 1:1 therefore appears to be robust: only one of the 16 selected galaxies with m easured redshifts is a non-m ember. W e expect few of the other 18 to be interloper eld galaxies.

4. Color-M agnitude Relation

The color{magnitude relation for the nal color{selected objects is shown in Fig 7. Red dots are ellipticals, orange squares and stars are S0 and SO/a galaxies, respectively. Sm aller black sym bols represent early {type galaxies that do not lie on the red sequence. Sm all triangles are late-type galaxies. Boxes are plotted around con med cluster m em bers. C on m ed interlopers are circled in the gure. Surprisingly, the two brightest cluster members are not ellipticals, but S0. The brightest of these two galaxies lies 700 kpc (1.2°) from the cluster center, and the other bright SO 300 kpc (0.6°). M oreover, there are late{ at type galaxies with lum inosities that are similar to the red { sequence bright early { type galaxies. Two of them lie on the red { sequence and are con med clusterm em bers, at 80 kpc from the cluster center (see also below in the discussion of the color and morphology distribution as a function of distance from the cluster center).

We tted the following linear color{m agnitude relation to various subsam ples of the galaxies:

$$i_{775}$$
 $z_{850} = c_0 + S \log(z_{850} 23)$ (1)

The solid line in Fig 7 is the t to the color{ magnitude relation for the ellipticals, the black dotted line is the t to the CMR for the SOs, and the dashed{dotted line the t to the full sam – ple, within 2^{0} from the cluster center (see discussion below). The dashed line is the t to the full sam ple of early{type galaxies in RXJ1252.9-292 from B lakeslee et al. (2003a), scaled to this redshift with BC 03 evolved stellar population m odels, with solar metallicity and a form ation age of

2.6 Gyr (since B lakeslee et al. 2003a obtains elliptical mean ages > 2.6 Gyr). The long{dashed vertical line is the magnitude lim it of the morphological classi cation $z_{850} = 24$ mag. The results for di erent morphological samples are given in Table 1.

A robust linear t based on B isquare weights (Tukey's biw eight; Press et al. (1992)) has been used to t the color{magnitude relation. Uncertainties on the parameters were estimated by bootstrapping on 10,000 simulations. The scatter around the twasestim ated from a biweight scale estim ator (Beers, Flynn & Gebhardt 1990), that is insensitive to outliers, in the same set of bootstrap simulations. The internal color scatter ($_{int}$) was m easured in twoways: 1) to the scatter around the t, we have subtracted in quadrature the average uncertainty due to the galaxy color error; and 2) we have calculated the internal scatter for which the 2 of the twould be unity. Both methods give us internal scatters consistent to within a few 0.001 m ag. All galaxies in this sample lie within three sign a from the t.

The X { ray distribution appears to be very sym metric, and largely con ned within 1.5 from the cluster center. W e calculated the CMR zero point and scatter within 1^0 (which corresponds to a scale of 0.5 Mpc at this redshift), and within 1.5(0.7 Mpc), and within 2^0 (1 Mpc), the scale used for the analysis of R X J1252.9-292). A coording to the results in Table 1, the internal color scatter increases when adding populations between 1^0 and 2° , especially for the S0 and S0/a populations, as also observed in local samples (e.g. van Dokkum et al. 1998), with only a small increase in sample size. We will therefore focus on the results obtained for color{selected galaxies within 1⁰ from the cluster center (where 90% of the colorselected galaxies lie). The slope of the elliptical CMR (0:033 0:015) is slightly steeper than the observed slope in R X J1252.9-292 (0:020 0:009), and in Com a when the latter are shifted to the observed colors at z 1:1, using non-evolving BC03 stellar population models, but still consistent within the uncertainties. We do not nd a atter (with respect to Com a) slope as in Stanford et al. (2002). However, the SO sample shows a much shallower slope (0:005 0:023) than the ellipticals, resulting in a much atter slope for ellipticals and S0s together (0:024 0:020). This

can explain why a shallower slope was found in that work, in which elliptical and SOs were not separated. The spectroscopically-selected ellipticalplus SO slope (0:010 0:034) is also attened by the SO population, while the spectroscopicallyselected ellipticals have a slope (0:021 0:046) sim ilar to that of RX J1252.9-292 and Com a. All the di erence in slope are how ever within the uncertainties and are statistically insigni cant.

Using Bruzual & Charlot (2003; BC 03) stellar population m odels, as in B lakeslee et al. (2003a), we derive a constraint on the age of the stellar populations in the galaxies from galaxy colors and the scatter of the CMR (van Dokkum et al. 2001; Blakeslee et al. 2003a). Two simple models have been considered and our conclusions will depend on the chosen models. The rst model is a single burst model, in which galaxies form in single bursts at random times t_f , between the age of the cluster and the recombination epoch. The second is a model with constant star form ation in a range of time between t_1 and t_2 , random ly chosen to be between the age of the cluster and the recom bination epoch. Colors for 10,000 galaxies were simulated with their scatter around the CMR to be dependent on the burst age. In Fig 8, we show the simulated scatters as a function of burst age, with solar, half solar and twice solar metallicity model. We will assume solar metallicity in what follows. From the scatter ($_{int} = 0.042 \quad 0.010$) in the colors of the galaxies classi ed as ellipticals, we obtain ages > 2:1 Gyr (z > 2), with a mean lum inosity {weighted age t = 3:31 Gyr (z_f 3.1), assuming the random single burst model. From the constant star form ation m odel, we obtain ages > 1:6 Gyr (z > 1:7), with a mean lum inosity{ weighted age t = 326 Gyr (z_f 3). This agrees with the conclusion (e.g. Blakeslee et al. 2003a; Holden et al. 2004; Lidm an et al. 2004; De Lucia et al. 2005) that the elliptical population in clusters of galaxies form ed at $z_f > 3$, and has evolved mainly passively until z = 1:1.

In the (U-B) {rest frame (using BC03 stellar population models with solar metallicity and age equal to 4 G yr), a scatter in $(i_{775} z_{850})$ of 0.042 0.010 corresponds to 0.050+ = 0.011. A spointed out in van Dokkum (2000) and B lakeslee et al. (2003a), CMR scatters vary little with redshift. The B lakeslee et al. (2003a) scatter for the elliptical CMR in RDCS 1252{2927 (0.024 0.008), correspond to a scatter of 0:042 0:014 in the (U - B) {rest fram e, indistinguishable within the uncertainties from our result.

The scatter in the CMR for galaxies classi ed as S0 ($_{int} = 0.044$ 0.020) is comparable to the one in the E CMR, but the galaxy colors are bluer and are not compatible with a population that is as old as the ellipticals. All the S0s lie below the elliptical color{magnitude relation. In fact, between the elliptical and the S0 CMR ts there is a zero point di erence of 0.07 0.02 mag, with the S0s being bluer than the ellipticals. O ne of the three S0/a galaxies has a color that is 0.07 m ag redder than the elliptical CMR, and another has a color that is 0.15 bluer than the CMR relation for Es. The inclusion of the S0/a galaxies does not signi cantly change the tted CMR for the S0s.

W hen we consider all galaxies within 2⁰ from the cluster center, the S0 and total early{type slopes are sim ilar, while the color o set in the CMR is still present (0.05 0.02 m ag). The S0 population of this cluster has a very peculiar CMR with respect to the average cluster of galaxies. In fact, for several other studies the CMR of the S0 population has a sim ilar zero point and on average a larger scatter with respect to the elliptical population (van D okkum et al. 1998; B lakeslee et al. 2003a; H olden et al. 2004; D e Lucia et al. 2004), quite di erent from our results. W e will discuss this peculiar behavior in detail in the rest of the paper, including an exam ination of orientation effects on the classi cation

In Fig. 9, the near-IR (Vega magnitudes) and $(i_{775} z_{850})$ (AB m agnitudes) colors are shown compared with single burst stellar population model predictions from BC03. The S0 colors are consistent with young (< 2 G yr) solarm etallicity, orolder (< 3.5 G yr), half solarm etallicity populations. If the di erence in E and SO m ean colors is mainly due to metallicity, then even if the two populations were form ed at the sam e epoch, ellipticals must have been able to retain more metals than the SOs, i.e., they were more massive at a given lum inosity (given the observed m ass{m etallicity relation for early-type galaxies, e.g., Trem ontiet al. 2004, Bernardietal. 2005, and references therein). This would imply higher mass { to { light ratios for the ellipticals with respect to the SOs. However, the lack of strong evolution in the slope and scatter of the CMR from the present out to z 1 suggests that the CMR ismainly the result of a metallicity { mass (i.e. metallicity {magnitude) relation (e.g., Kodama & Arimoto 1997; Kauman & Charlot 1998 Vazdekis et al. 2001; Bernardi et al. 2005). So, at a given magnitude we do not expect large metallicity variations.

If the o set is due to a di erent star form ation history, a model with solar metallicity and with an exponential decay of the star form ation will reproduce the o set at a galaxy mean age of

3.5 G yr. This age is consistent with the sm all scatter observed in the S0 CM R.W e would then be observing galaxies that followed di erent star form ation: single burst and passive evolution for the ellipticals and exponentially decaying star form ation for the S0s. An exponential decay in the star form ation is observed in eld spiral sam ples (R ow an {R obinson 2001}. If this is the case, our S0 population m ight be the evolved product of an old spiral population that was already in place in this cluster when the ellipticals form ed and then gradually lost available gas for star form ation.

If the E vs S0 color di erence is mainly due to a di erence in age, for a solar metallicity and a single burst BC03 template with age 4 Gyr, the color di erence corresponds to an age di erence of 1 Gyr. For clusters of galaxies at z > 1, the cluster members on the red sequence are only a part of all the progenitors of present{day early{ type galaxies. Som e of today's galaxy progenitors would have been bluer than the red sequence at these redshifts (van D okkum & Franx 2001). In the S0 population of this cluster, we may be seeing the transitional progenitor population that in

1 G yr will evolve onto the sam e red sequence as now occupied by the ellipticals.

E ither of the latter two scenarios would be consistent with the Postman et al. (2005) observed de cit of the S0 population of our ACS cluster sample, when compared to lower redshift samples, im plying that part of the S0 population is still form ing in clusters at redshifts around unity.

5. Galaxy shape properties

Since the galaxies classied as S0 in RXJ 0910+ 5422 are found to be system atically bluer (with respect to the red sequence) than the S0 populations observed in previous studies, we wish to examine further the properties of these

galaxies in term s of their shapes and light distributions, and how they compare to the elliptical and spiral samples in this and other clusters. The shape parameters that we will consider are C oncentration and A symmetry (A braham et al. 1996; C onselice et al. 2004), Sersic index n, and galaxy axial ratios.

5.1. A sym m etry and C oncentration

In Fig 10 (left), we compare the Asymmetry A and the Concentration C for ellipticals, SOs, and spirals with $(i_{775} z_{850})$ colors between 0.5 and 1.2 m ag. The Asymmetry parameter is obtained by subtracting a 180{degree rotated im age from each original galaxy in age, sum m ing the residuals and including a correction for the background. The Concentration parameter is de ned as in Abraham et al. (1996) as the sum of the galaxy ux within an aperture $r_{0:3}$ divided the total ux. $r_{0:3}$ is calculated using the SE xtractor t to the galaxies at 1.5 above the background. The obtained sem i{m a pr and sem i{m inor axes from this twere multiplied by 0.3 to derive the $r_{0,3}$ aperture (see also Hom eier et al. 2005b). Early{ type and late{type galaxies lie on di erent regions in this A vs C plane. All our red sequence S0s have A < 0.3 and C > 0.3. All but one have A < 0.2 and C > 0.3. This is the same locus in the A {C plane that is occupied by m ost early { type galaxies in Abraham et al. (1996) and in our Postman et al. (2005) low {redshift sam ple (Fig 10, right). This last sample includes 5 strong lensing clusters observed as part of our ACS GTO program Zw1455+2232 (z=0258), M S1008-1224 (z=0:301), M S1358+6245, CL0016+1654 (z=0.54), and M S J0454-0300 (z=0.55)]. V isual and autom ated classi cation for this cluster in the i_{775} {band for all galaxies with i_{775} < 22:5 was performed by Postman et al. (2005). We conclude that the S0 population presents statistical parameters typical of an early {type population (low asymmetry and high compactness).

5.2. Sersic Indices

Fig 11 plots Sensic index n as a function of galaxy e ective radius R_e , both from our GAL-FIT modeling, for ellipticals, S0s and spirals with $(i_{775} z_{850})$ colors between 0.5 and 1.3 m ag. Red sequence $((i_{775} z_{850})$ color between 0.8 and 1.1 m ag) galaxies are shown by large diamonds.

M ost spirals have n < 2 and m ost early (types have n > 2, as expected. However, the n values do not perm it us to discrim inate between SOs and ellipticals in a unique way, unless they are com bined with goodness {of { t inform ation for the Sersic m odel (e.g. the B um piness param eter introduced by B lakeslee et al. 2005)

6. A xial R atios

6.1. A xial R atio D istribution

In Fig 12 we com pare the apparent axial ratio from SExtractor versus e ective radii for elliptical and S0s with $(i_{775} z_{850})$ colors between 0.5 and 1.3 m ag. The axial ratios have been veriled by using ELLIPROF (the isophotal thing software that is used for Surface Brightness Fluctuations analysis in Tonry et al. 1997 and M eiet al. 2005) on each galaxy image, after the cleaning procedure. As above, red sequence galaxies are shown by large diam onds. The red sequence ellipticals and the bluer SOs have di erent axial ratio distributions, with all red sequence SOs showing axial ratios $\frac{b}{a}$. 0:7, and nearly all red sequence ellipticals with $\frac{b}{a} > 0$:7. A ssum ing axisym metric disks (oblate ellipsoids) viewed with random orientation, and with a Gaussian distributed intrinsic axial ratio (with mean equal to 0:3 0:1 (extrem e thin disk), 0:5 0:1 (early (type galaxy) and 0:75 0:1 (elliptical) (Jorgensen & Franx 1994)), one would expect & 40%, & 60%, and & 90%, respectively, of the S0s to have axial ratios above 0.7. Just 9% (1 out of 11) of the red sequence S0s are observed to have an axial ratio this large (or 22% for the full S0 sample in this cluster eld) (Fig 13; top). The random probability that the SO axial ratios would show such a low fraction with $\frac{b}{2}$ > 0:7 is less than 1%. This is a very simple model, but it points out a lack of round SOs, indicating either that there is som e orientation bias in the classi cation, or that this class of objects is intrinsically prolate in shape. Jorgensen & Franx (1994) found a sim ilar de cit of round SOs in the center of the Com a cluster. They concluded that part of the face (on SOs were classi ed as elliptical galaxies. Fabricant et al. (2000) also found a de cit of round SOs in the cluster CL 1358+ 62, at z=0.33. Their analysis shows that ellipticities and bulge{to{total{light{ratio do not allow us to distinguish elliptical from SO galaxies. The other two ACS GTO clusters at z > 1 (RXJ1252.9-292 and RX J0848+4452) do not show a sim ilar lack of round S0s, as 90% of the red sequence ellipticals (out of 70) and 47% of the S0s (out of 35) have $\frac{b}{a} > 0.7$ (Fig 13; bottom). This bias is also not observed in other clusters of our ACS Interm ediate Redshift C luster Survey (see Fig 5 from Postm an et al. 2005), for which m ore than 40% of the S0 galaxies have axial ratios above 0.7.

The observed peculiarity of the RXJ 0910+ 5422 S0 axial ratio distribution m ight call into question our result above that the SOs have a signi cant color o set with respect to the ellipticals. For instance, if there is a bias in our colorm easurem ent procedure which causes elongated objects to have colors that are too blue, then the coloro set found above may be articial. Such a color bias might occur if the high inclination angles bias our R_e m easurem ents to higher values, and if the SOs becom e progressively bluer at larger radii. W e rst exam ine this possibility, then proceed to discuss resolutions to the peculiarity of the SO axial ratio distribution, with the aim to establish if a misclassi cation of face{on S0s as ellipticals would bias the measurement of the oset between the ellipticaland SOCMR zero points.

6.1.1. Internal color gradients

It is conceivable that our $(i_{775} z_{850})$ colors could be biased by aperture e ects in the nearly edge(on S0s, for which the (possibly) bluer outer disks m ight contribute m one to the galaxy colors than in the rounder ellipticals population. If this e ect were severe enough, it m ight m in it the o set in color of the S0s and ellipticals found above. W e test this possibility here. S0 and elliptical internal color pro les are shown in F ig 14 and F ig 15. The gradients have been calculated with aperture photom etry on the sam e in ages used to calculate our ($i_{775} z_{850}$) colors. C incles are ($i_{775} z_{850}$) colors at di erent radii, the cross the ($i_{775} z_{850}$) color at the e ective radius, used for the CM R.

The S0 colors pro les do not show strong gradients. In particular, the $(i_{775} z_{850})$ colors calculated at the e ective radii are not system atically bluer than colors determ ined at sm aller radii. Most of S0 galaxies have at pro les; one has a blue inward gradient (ACS ID 1621; $z_{850} =$ 23:72 m ag, $(i_{775} z_{850}) = 1 m$ ag). In two galaxies (ACS ID 1393 and 3177) the colors in the central 0.15^{00^2} are redder than the color at the e ective radius. W hen compared with elliptical gradients, on average S0 colors do not appear biased towards higher e ective radii and bluer colors than the ellipticals. We note that three elliptical galaxies show blue inward gradients (ACS ID 1753, 1519, 3323).

6.1.2. O rientation or intrinsic shape: A xial ratios vs (i₇₇₅ z₈₅₀) colors

O rientation biases are known to occur in the classi cation of ellipticals and SOs in local galaxy sam ples. For instance, Rix & W hite (1990, 1992) showed, based on both isophotal and dynam ical modeling, that a large fraction of ellipticals contain a disk component with at least 20% of the light, but which is hidden due to projection e ects. Jorgensen & Franx (1994) found a strong de cit of round SOs in a sample of 171 galaxies in the central square degree of the nearby C om a cluster, and concluded that inclination angle played a large part in the classi cation of Es and SOs. M ichard (1994) proposed that, except for the bright boxy ellipticals without rotational support, early-type galaxies com prise a single class of oblate rotators with orientation being the main criterion for classi cation as either E or S0.0 n the other hand, van den Bergh (1994) explained the predom inance of attened S0s by invoking two distinct subpopulations: bright disky objects interm ediate between ellipticals and spirals, and a fainter population of prolate ob jects.

W e now address the question of whether the galaxies classi ed as S0 in RXJ 0910+5422 are preferentially attened in shape because of an orientation bias in the classi cations or intrinsically prolate shapes. If it is an orientation bias, then this could mean either that (1) face-on SOs have been m isclassi ed as ellipticals because their disks are not apparent, or (2) that edge-on spirals tend to be called S0s because the spiral structure is obscured. Either would result in a predom inance of attened SOs. However, in the form er case, the m isclassi cation of face-on SOs as ellipticals would tend to blur any color separation between the two classes, while in the latter case, a coloro set m ight be introduced between the two classes because of contam ination by bluer spirals. Because we do observe a color o set between galaxies classi ed as E and S0, with the S0s being bluer, it is possible to

bok for the \m issing" population of face-on blue galaxies by exam ining ellipticity versus color. If a population of round blue objects is found, we can then determ ine the nature of the classi cation bias, and whether it biases our color o set m easurem ent.

H istogram s of the axial ratio distributions for ellipticals and SOs are shown in the top panel of Fig. 13. We nd that 60% of the early-type red sequence galaxies have $\frac{b}{a} > 0.7$, but 95% of these low-ellipticity galaxies are classi ed as Es. However, if we split galaxies instead by color, z_{850}) < 0:99 m ag as the separausing (i775 tion point, then we nd that 54% of all galaxies bluer than this separation have $\frac{b}{a} > 0$;7, while 43% of the red sequence galaxies bluer than this $[0:8 < (i_{775} z_{850}) < 0:99 m ag] have <math>\frac{b}{a} > 0:7$ (Fig. 16). Thus, the de cit of round SO galaxies (which are also signi cantly bluer than the mean of the E class) is not found when the early-type galaxies are split based purely on color. This suggests that som e of the bluer round galaxies are the face-on counterparts of those classi ed as SO.

Fig 17 shows axial ratios vs (i775 Z850) color residuals with respect to the total earlytype galaxy CMR relation, for all galaxies in the RXJ 0910+5422 red sequence. Galaxy types are coded with di erent sym bols; using the color residuals in this way takes out the e ect of the magnitude dependence of the colors. There are ve round blue ellipticals, i.e., with $\frac{b}{2} > 0.7$ and on the blue side of the early-type CMR. If these are the face-on counterparts of the blue SOs, then the S0 axial ratio distribution becomes much more in line with expectations for a random ly oriented disk population (35% are rounder than $\frac{b}{a} = 0.7$). Further, we note that four other ellipticals were classi ed E/S0 in Postm an et al. (2005); if these are also taken as face-on SOs, then the axial ratio distribution com es in very close agreem ent with expectations. We conclude that the face-on SOs in RXJ 0910+ 5422 are classi ed as ellipticals, just as in local early-type galaxy sam ples.

From our axial ratio simulations and statistics from our z 1 ACS GTO sample, the number of blue SO galaxies might be between 25 and 40% higher than estimated in section 5.3.1. Moreover, the inclusion of a few blue SOs in the elliptical red{sequence will have the e ect of increasing very slightly the observed CMR scatter for the ellipticals. The uncertainty in the Postman et al. (2005) S0 fraction in RXJ 0910+5422 is significantly larger than 40% and hence a systematic change by this amount for one cluster does not alter any of the results or conclusions in that paper.

Throughout the paper, we continue to use elliptical and SO classi cations from Postm an et al. (2005), but keeping in mind the presence of this possible projection e ect.

 Color trends, velocity dispersion and merging activity

Stanford et al. (2002) have analyzed the X { ray and near-IR properties of this cluster. This system appears to be fairly relaxed based on its regular X { ray pro le; however, they nd indications that the cluster is in an early phase of formation. In fact, the Chandra ACIS data show evidence for tem perature structure, possibly due to an infalling group or mass stream ing along a lam ent. The soft component of the X { ray em ission (0.5{2 keV) dom inates the X {ray center of the cluster, while to the south there is a harder component (2{6 keV) (see Fig. 6 from Stanford et al. 2002). The cluster does not have a centralBCG or cD galaxy, and the X { ray em ission center does not correspond to an optical grouping of galaxies; rather a num ber of lum inous con-

m ed cluster m bers are linearly distributed, at least as projected on the sky (as shown in F ig 1). W e do not observe any strong trend of the galaxy color (i_{775} z_{850}) with distance from the cluster X (ray center (F ig. 18). Surprisingly, though not statistically signi cant, the bluer galaxies are concentrated tow and the cluster center, instead of the outskirts, as in the other ACS interm ediate redshift cluster sam ple (D em arco et al. 2005; G oto et al. 2005; H om eier et al. 2005; P ostm an et al. 2005). This tendency m ight support the hypothesis that we are observing sub-groups of galaxies in an edge(on sheet, e.g. a group of bluer disk galaxies on a redder, older population of ellipticals.

There are 10 con med elliptical and 5 conmed S0 members in the center of the cluster (R < 500 kpc). The average S0 redshift is 1.102 0.002 and the average elliptical redshift is 1.105 0.007 (the given uncertainties are standard deviations around the mean). This indicates small relative velocities (the two redshifts are indistinguishable given the errors) between the classes and is true regardless of the possible classication bias discussed above. Unfortunately, our spectroscopic sample does not perm it us to track the cluster central structure in detail. The average relative velocity between the con med E and SOs of 500 km/s (that also corresponds to the cluster velocity dispersion; see below) is fairly sm all. If merging of two distinct groups of galaxies is happening along the line of sight, we expect much higher velocity dispersions and/or relative velocities between the infalling SOs and the ellipticals.

Stanford et al. (2002) also suggest that active galaxy-galaxy m erging should be observed, based on the X {ray temperature structure. To investigate any on {going dynam ical activity, we calculated the cluster velocity dispersion, and the m erger rate. From the 25 spectroscopically conmed members (all galaxy types included) the line{of{sight rest{fram e velocity dispersion is = 675 190 km /s, using the software RO STAT from Beers, Flynn, & Gebhardt (1990). The available Chandra data give X-ray tem peratures ranging from $kT = 7.2^{+2.2}_{-1.4}$ keV (Stanford et al. 2002) to $kT = 6.6^{+1.7}_{1.3}$ keV (Ettori et al. 2004). W u et al. (1999; see also Rosati et al. 2002) gives the relationship between kT and for relaxed clusters, which predicts that the velocity dispersion corresponding to the measured X -ray tem perature should be 1000 km /s, considerably higher than we have found. Again, in the case of merging groups (along the line of sight) we would also have expected a higher velocity dispersion.

8. Red galaxy pairs

The quality of the ACS data allows us to discern merging activity among the cluster galaxies. If we assume that galaxy pairs with projected separations less than $20h_{70}^{-1}$ kpc are physically associated, we observe 13 associated early (type galaxies, nine galaxies of which lie on a lamentary structure about 100 kpc from the cluster center (Fig. 1). As noted above, RXJ 0910+ 5422 lacks any cD galaxy near the center of the X (ray em ission (see also Fig. 1), but rather has a lamentary group of galaxies around the X (ray center.

The nine early-type interacting galaxies within this lamentary structure (at radius of $100h_{70}^{-1}$ kpc from the X (ray center) include 3

unique pairs (yellow arrows in Fig. 1), plus a galaxy triplet (the three components marked with red arrows in Fig. 1). Each of the 3 pairs consists of a bright elliptical with a smaller companion (all closer than $10h_{70}^{-1}$ kpc), while the triplet is a large E with two smaller SOs (also closer than $10h_{70}^{-1}$ kpc; one of these two SOs is ACS ID 1621, the SO with an inward blue gradient). One of the pairs (the middle pair in the gure) and the two nearest galaxies in the triplet, have essentially zero relative velocity and thus are likely merger candidates. A lso two of the ellipticals with blue inward gradients lie on the central lam ents, and they are both small satellites of a larger galaxy.

The other two pairs, which do not lie on the lam entary structure, are at $250h_{70}^{-1}$ kpc (two E with sim ilar size, both with weak O II em ission (Stanford et al. 2002) and $300h_{70}^{-1}$ kpc (one SO/a and one E of sim ilar size) from the X {ray center, and have relative velocities of 10,000 km/s and 2000 km/s, respectively.

The presence of the low -velocity pairs is consistent with the low velocity dispersion, and provides evidence for the on {going hierarchical growth of the cluster (e.g., van Dokkum et al. 1999). The pairing of the bright ellipticals with sm aller SOs also argues against the view that we are observing a blue SO-dom inated group infalling into a red cluster elliptical population, but rather com plex stellar population evolution within a lam entary structure. The red{sequence SO/a con rm ed m em bers also lie within this lam entary structure.

The observations of a signi cant number of red galaxy pairs in a cluster at z 1 is interesting in the context of the recent notings by van Dokkum (2005) of red galaxy interactions in

70% of 86 early (type galaxies in a selected sam – ple of nearby red galaxies from the MUSYC (Multiw avelength Survey by Yale(Chile; Gaw iser et al. 2005) and the NOAO Deep W ide(Field Survey. This work concluded that most of the ellipticals in local sam ples were assembled by red galaxy(galaxy m ergers, denom inated dry m ergers because they would involve gas(poor early(type galaxies. At higher redshift, Tran et al. (2005b) con m ed red galaxy m ergers rst observed by van D okkum (1999) in MS 1054-03 at z= 0.83. Tran et al. selected m ergers as associated pairs with projected separation less than $10h_{70}^{-1}$ kpc and relative line(of(sight velocities less than 165 km/s. As in RXJ 0910+5422, the red early{type galaxies involved in these m ergers are among the brightest cluster m embers. Their results suggest that m ost early{type galaxies grew from passive red galaxy{ galaxy m ergers. In our sam ple we observe a triplet and three red galaxy pairs with projected distances less than $10h_{70}^{-1}$ kpc. Of those, the triplet and one galaxy pair show zero relative velocity. Of the two other pairs, com posed of a bright and a fainter com panion, redshifts are not available for the faint com panions.

9. Cluster lum inosity function

To obtain a deeper understanding of the RXJ 0910+ 5422 galaxy population, we constructed galaxy lum inosity functions in the follow – ing way. We start with the original Sextractor catalog (described in Section 3). All objects with m agnitudes brighter than $i_{\rm F~775W}$ = 21.1 m ag were considered as foreground objects. N ine of these bright objects are con rm ed non-members. The rem aining seven are objects that do not belong to the red sequence and whose sizes and lum inosities are m uch larger than those of the con rm ed mem – bers, in particular those of the bright red sequence galaxies; therefore they are very unlikely to be at the cluster redshift.

The contribution to the lum inosity function from both foreground and background eld galaxies (hereinafter, the eld) has been estimated from the galaxy counts in a reference eld. The control region is taken from the GOODS-S (Great Observatories O rigins D eep Survey {South; G iavalisco et al. 2004) ACS eld, observed in the same 1ter as the cluster eld. Point{like objects were elim inated in a consistent way in the cluster and in the control eld, by identi cation of the stellar locus in the diagnostic plot of the SE xtractor param eters MAG_AUTO vs FLUX_RAD IUS (the selected objects have FW HM equal to the PSF in the image). Cluster and control eld lum inosity functions were norm alized to the cluster area. Both cluster and eld counts were binned with a bin size of 0.5 m ag. For each bin, the eld counts are subtracted from the cluster counts, taking into account the extensive spectroscopic sample (m ore than 60% of the objects used for the LF determ ination brighter than M have measured redshifts). Known interlopers were excluded from the analysis. The uncertainties in the cluster counts after subtraction of the eld contribution are calculated by adding in quadrature Poissonian uncertainties.

The lum inosity functions are shown in Fig 19. The lled circles with errors are the total background { corrected cluster lum inosity function. We do not include errors from cosm ic variance due to the choice of the background control region. The red sequence elliptical and spheroidal (S0 and S0/a) lum inosity functions are shown respectively in blue and green. The red histogram is the lum inosity function of all early { type galaxies with $color 0.8 < (i_{775} z_{850}) < 1.1 m ag, exclud$ ing con med non {members. The histogram of red sequence galaxies in RXJ1252.9-292 is shown as the dashed red line. The background contribution is very small for the early-type sample. The red arrows show the histogram values after background subtraction. The rest-fram e B m agnitudes are shown on the top of the plot, calculated from colors obtained from the BC03 stellar population model and tem plates (Sbc, Scd) from Coleman, Wu & Weedman (1980). The solid black line is a Schechter function t to the total cluster lum inosity function. It is obtained by calculating the C (Cash 1979) statistic (a maximum likelihood statistic to t data with Poissonian errors) on a grid in the M { plane for each combination of M and : rst the norm alization () is calculated in order to reproduce the observed num ber of galaxies in the observed m agnitude range, then the C statistic is computed as $C = 2 i_{\text{bin}} n_i \ln m_i m_i \ln n_i!$, where n_i is the observed number of galaxies in the ith bin and m i is the num ber of galaxies predicted in that bin by the Schechter function with parameters M , , and

The combination M , which minimizes the C statistic is taken as the best-t. If the C statistic is de ned as above, the 1-2- and 3- con dence levels for M and can be estimated from C = 2:3;6:17;11:8. We obtain M = $22:6^{+0:6}_{0:7}$ m ag and = 0:75 0:4.

M ost of the faint{end population is composed of S0 and S0/a galaxies. The two brightest galaxies in the red sequence are S0. W ith respect to RXJ1252.9-292, a bright population of red sequence ellipticals is m issing in RXJ 0910+5422. How ever, the large Poissonian errors on the bright end of the cluster population prevent us from de nitively excluding the hypothesis that the two clusters could be drawn from the same parent population. Sim ilarly, sm all num ber statistics do not perm it us to study the lum inosity functions of the di erent red and blue faint populations in this cluster.

10. D iscussion and C onclusions

In this paper we have studied the color{ magnitude relations of galaxies in RX J 0910+ 5422 to constrain their ages and form ation histories. Our results show that the color{magnitude relation for the elliptical galaxies is consistent both in slope and scatter with that of RX J 1252.9-292 (B lakeslee et al. 2003a, Lidman et al. 2004) and recent results from H olden et al. (2004) and D e Lucia et al. (2004), con ming that elliptical galaxies in galaxy clusters show a universal color{ magnitude relation consistent with an old passively evolving population even at z 1. From the color{magnitude relation of the ellipticals, we derive a mean lum inosity{weighted age t > 3:3 G yr ($z_f > 3$).

We nd that the SOs in RXJ 0910+ 5422 de ne a color{m agnitude sequence with a scatter sim ilar to that found for the ellipticals, but shifted bluer by 0:07 0:02 m ag. This is peculiar with respect to previous cluster studies, which more typically found that the SOs followed the same CMR as the ellipticals, but with som ew hat larger scatter (Bower et al. 1992; Ellis et al. 1997; Stanford et al. 1997; Lopez{Cruz et al. 2004; Stanford, Eisenhardt, & Dickinson 1998; van Dokkum et al. 2000, 2001; B lakeslee et al. 2003a; Holden et al. 2004). Only one earlier study, van Dokkum et al. (1998) found a signi cantly bluer S0 population. W e exam ine this population of blue SOs in some detail, noting that there is a strong predom inance of attened system s with axial ratios $\frac{b}{a} > 0:7$, and conclude that the face-on members of the population have likely been classi ed as ellipticals. If so, the color o set between the two classes would becom e even m ore signi cant, and the true CMR scatter for the ellipticals would be slightly lower than we have estimated. This peculiarity is not observed in other clusters of our ACS Interm ediate Redshift Cluster Survey, and its amplitude is sm aller than the uncertainties adopted in Postm an et al. (2005).

If the observed color di erence between the ellipticals and SOs is mainly due to metallicity at the sam e age, this would im ply that the redder ellipticals were able to retain more metals than the SOs, i.e., they are more massive. How ever, current data suggest that the CMR is mainly the result of a m etallicity {m ass (i.e. m etallicity {m agnitude) relation (e.g., Kodam a & Arim oto 1997; Kau man & Charlot 1998 Vazdekis et al. 2001; Bernardi et al. 2005). This implies that we do not expect large m etallicity variations at a given m agnitude. If, instead, the o set is mainly due to age, then the im plied age di erence would be 1 G yr for singleburst solarm etallicity BC 03 m odels. It could also result from di erent star form ation histories, with the SOs experiencing a more extended period of star formation. A model with solar metallicity and with an exponential decay of the star form ation reproduces the o set at a galaxy m ean age of 3.5 Gyr.

The blue SOs may comprise a group infalling from the eld onto a more evolved red cluster population, or they m ay be a transitional cluster population not yet evolved all the way onto the elliptical red sequence (van Dokkum & Franx 2001). Assuming passive evolution, they will reach this red sequence after about 1 G yr. High fractions of faint blue late type galaxies were observed in substructures infalling in a main cluster (e.g. Abraham et al. 1996; Tran et al. 2005a), and proposed as the progenitors of faint SOs in clusters. The view of this cluster as a structure still in form ation is supported by X { ray observations of the cluster tem perature structure (Stanford et al. 2002), the lack of a cD galaxy, and its lam entary structure that suggests m erging of substructures. How ever, we derive a small cluster velocity dispersion, unusual for merging substructures. Moreover, the blue SOs in this sample span the same lum inosity range of the bright ellipticals, are distributed towards the center of the cluster, and some of the faintest ones are physically associated with brighter ellipticals belonging to the central lamentary structure. These elements would argue against the bluer SOs being a young group m erging with an existing red cluster population, and support the hypothesis that we are observing a transitional blue S0 population in a cluster core that is still evolving onto the elliptical red sequence. This result is also consistent with the de cit of S0s observed in our ACS cluster sam ple, when com pared to low er redshift sam ples, that in plies that the S0 population is not yet in place but still form ing in clusters at redshifts around unity (Postm an et al. 2005). Interestingly, we also observe in this cluster potential progenitors for bright S0 galaxies: four bright spirals (spectroscopically con m ed cluster m em bers) with z_{850} brighter than 22.5 m ag and (i_{775} z_{850}) between 0.5 and 1.3 m ag.

Red galaxy pairs are also observed. A triplet and three red galaxy pairs have projected distances less than $10h_{70}^{-1}$ kpc, and of those, the triplet and one pair show zero relative velocity. This would be the evidence of red galaxy m ergers at z 1. van Dokkum (2005) and Tran et al. (2005b) have observed m ergers of red galaxies in a nearby elliptical sam ple and in M S 1054-03 at z= 0.83, respectively. They suggested a scenario in which m ost of the early {type galaxies were form ed from passive red galaxy{galaxy m ergers, called dry m ergers, because they involve gas{poor early {type galaxies.

Future papers will analyze the ages and m asses of the cluster m embers using our optical spectroscopy along with new ly obtained Spitzer IRAC im aging. A larger sam ple would be needed to draw m er conclusions about the form ation of SOs.

ACS was developed under NASA contract NAS 5-32865, and this research has been supported by NASA grant NAG 5-7697 and by an equipment grant from Sun Microsystems, Inc. The Space Telescope Science Institute is operated by AURA Inc., under NASA contract NAS5-26555. We are gratefilto K. Anderson, J.M cCann, S.Busching, A. Fram arini, S. Barkhouser, and T. Allen for their invaluable contributions to the ACS project at JHU. We thank W. J.M cCann for the use of the FITSCUT routine for our color in ages. SM thanks Tadayuki K odam a for useful discussions.

REFERENCES

- Abraham, R.G., van den Bergh, S., Glazebrook, K., Ellis, R.S., Santiago, B.X.; Summa, P., Griths, R.E. 1996, ApJS, 107, 1
- Beers, T.C., Flynn, K., & Gebhardt, K. 1990, AJ, 100, 32

Bell, E F. et al. 2004, ApJ, 608, 883

- Ben tez N., Ford, H.Bowens, R.et al. 2004, ApJS, 150, 1
- Bertin, E. & A mouts, S. 1996, A & A S, 117, 393
- Bernardi, M., Sheth, R.K., Nichol, R.C., Schneider, D.P., Brinkmann, J. 2005, 129, 61
- Blakeslee, J.P., et al. 2003a, ApJ, 596, L143
- Blakeslee, J. P., Anderson, K. R., Meurer, G.R., Ben tez, N., & Magee, D. 2003b, ASP Conf. Ser. 295: ADASS X II, 257
- Blakeslee, J.P., et al. 2005, ApJ, in press
- Bower, R.G., Lucey, J.R., & Ellis, R.S. 1992, MNRAS, 254, 589
- BruzualA.,G.& Charbt,S.2003,MNRAS,344, 1000 (BC03)

Cash, W ., 1979, ApJ, 228, 939

- Coleman, G.D., Wu, C.-C., Weedman, D.W., 1980, ApJS, 43,393
- Conselice, C. J. et al. 2004, ApJ, 600, L139
- De Lucia et al. 2004, ApJ, 610, L77
- Dem arco, R., et al. 2005, A & A, 431, 381
- Ellis, R.S., Smail, I., Dressler, A., Couch, W.J., Oemler, A.Jr., Butcher, H., Sharples, R.M. 1997, ApJ, 483, 582
- Ettori,S., et al. 2004, 354, 111
- Fabricant, D., Franx, M., van Dokkum, P. 2000, 539, 577
- Ford, H.C. et al. 2002, Proc. SP E, 4854, 81
- Ford, H.C. et al. 2004, astro-ph/0408165
- Gawiser, E. et al. 2005, ApJS, in press; astroph/0509202
- Giavalisco, M. et al., 2004, ApJ, 600, L93

Goto, T. et al. 2005, ApJ, 621, 188

Hogbom, J.A. 1974, A&AS, 15, 417

- Hogg, D.W., et al. 2004, ApJ, 601, L29
- Holden, B P., et al., 2004, AJ, 127, 248

- Holden, B P, et al, 2005a, ApJ, 620, 83
- Holden, B P., et al, 2005b, ApJ, 626, 809
- Hom eier, N L., et al. 2005, ApJ, 621, 651
- Jorgensen, I., & Franx, M . 1994, ApJ, 433, 553
- Kau man, G. & Charlot, S. 1998, MNRAS, 294, 705
- Kodama, T. & Arimoto, N. 1997, AJ, 320, 41
- Lidman, C., Rosati, P., Demarco, R., Nonino, M., Mainieri, V., Stanford, S.A., Toff, S. 2004, A&A, 416, 829
- Lopez-Cruz, O., Barkhouse, W. A., Yee, H.K.C. 2004, 614, 679
- Marleau, F.R. & Sim ard, L. 1998, ApJ, 507, 585
- Massey, P.& Gronwall, C. 1990, ApJ, 358, 344
- McIntosh, D. H., Zabludo , A J., Rix, H.-W., Caldwell, N. 2005, 619, 193
- Mei, S. et al. 2005, ApJ, 625,121
- M ichard, R. 1994, A & A, 288, 401
- Mullis, C R., Rosati, P., Lamer, G., Bohringer, H., Schwope, A., Schuecker, P., Fassbender, R. 2005, ApJ, 623, L85
- Oke, J.B., et al. 1995, PASP, 107, 375
- Peng, C.Y., Ho, L.C., Impey, C.D., & Rix, H. 2002, AJ, 124, 266
- Postman, M., et al. 2005, ApJ, 623, 721
- Press, W H. et al. 1992, Num erical Recipes, Cam bridge University Press, New York
- Rix, H., & W hite, S.D.M. 1990, ApJ, 362, 52
- Rix, H., & W hite, S.D.M. 1992, MNRAS, 254, 389
- Rosati, P., della Ceca, R., Norman, C., Giacconi, R.1998, ApJ, 492, L21
- Rosatiet al. 2002, ARA & A, 40, 539
- Rowan-Robinson 2001, ApJ, 549, 745
- Sirianni, M., et al. 2005, PASP, in press

- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525
- Spergel, D N., et al. 2003, ApJS, 148, 175
- Stanford, S.A., Eisenhardt, P.R., & Dickinson, M.1998, ApJ, 492, 461
- Stanford, S. A., Elston, R., Eisenhardt, P.R., Spinrad, H., Stern, D., Dey, A. 1997, AJ, 114, 223
- Stanford, S. A., Holden, B., Rosati, P., Eisenhardt, P.R., Stem, D., Squires, G., Spinrad, H. 2002, AJ, 123, 619
- Tonry, JL., Blakesbe, JP., A har, EA. et al. 1997, ApJ, 475, 399
- Tran, K.-V., van Dokkum, P.G., Illingworth, GD., Kelson, D., Gonzalez, A., Franx, M. 2005a, ApJ, 619, 134
- Tran,K.-V.,vanDokkum,P.G.,Franx,M.,Illingworth,GD.,Kelson,D.,Schreiber{Foerster, NM.2005b,ApJ, 627,L25
- Trem ontiet al. 2004, ApJ, 613, 898
- van den Bergh, S. 1994, AJ, 107, 153
- van Dokkum, P.G. 2005, AJ, in press; astroph/0506661
- van D okkum, P.G., Franx, M., Fabricant, D., K elson, D.D., & Illingworth, G.D.1999, ApJ, 520, L95
- van Dokkum, P.G., Franx, M., Kelson, D.D., Illingworth, G.D., Fisher, D., & Fabricant, D. 1998, ApJ, 500, 714
- van Dokkum, P.G., Franx, M., Fabricant, D., Illingworth,G.D., & Kelson,D.D.2000,ApJ, 541,95
- van D okkum, P.G., Stanford, S.A., H olden, B.P., E isenhardt, P.R., D ickinson, M., & Elston, R. 2001, ApJ, 552, L101
- van Dokkum,P.G.& Franx,M.2001,ApJ,553,
 90

Vazdekis et al. 2001, ApJ, 551, L127

Wuetal. 1999, ApJ, 524, 22

T his 2-column preprint was prepared with the AAS ${\rm IAT}_EX$ m acros v5.0.



Fig. 1. The ACS color in age with X (ray contours overlaid. The X (ray observations are from Chandra ACIS-I, over the energy range of 0.5 (2K eV, and were smoothed with a 5 arcsec FW HM G aussian. N orth is up, E ast on the left. In the enlargement, galaxy pairs in the central lam entary structure are shown. Two m ain groups of interacting galaxies lie along the lam entary structure at the center of the cluster. The rst group is shown by the yellow arrow, the second one by the red arrows. Three other red (sequence early (type galaxies are shown by the green arrows.



Fig. 2. $|(i_{775} z_{850})|$ and (J-K) colors as a function of galaxy age. BC 03 solar metallicity, age equal to 4 G yr models are shown with a solid line. The dashed lines are for half solar and the dot{dashed lines for twice solar metallicity. We considered a sample of color{selected galaxies with 0.8 < $(i_{775} z_{850}) < 1.1 \text{ mag}$ and (J K) > 1.45.



Fig. 3. Color in ages of the 20 CMR ellipticals within 2 0 from the cluster center. Two of the CMR faint ellipticals, that are satellites of brighter galaxies, are shown in the same postage in age of their bright companion.



Fig. 4. Color in ages of the 11 CMR S0 galaxies within 2^{0} from the cluster center.







Fig. 6. Color in ages of the four bright spirals with z_{850} brighter than 22.5 m ag and $(i_{775} z_{850})$ between 0.5 and 1.3 m ag, within 2⁰ from the cluster center. These spirals have lum inosities that are similar to those of the red{sequence bright ellipticals.



Fig. 7. The color-m agnitude relations for RXJ 0910+ 5422. Red circles and yellow squares are respectively E and S0 candidates on the red sequence. Yellow stars are S0/a candidates on the sequence. Smaller black circles, squares, and stars are used for the corresponding early (type galaxies that do not lie on the red sequence. Small triangles are late-type galaxies. Boxes are plotted around con med cluster members. Open circles are plotted around con med interlopers. The solid line is the t to the ellipticals, the black dotted line the t to the S0s, and the dashed (dotted line the t to the entire sam ple within 2° . The green dotted is tted to the S0s within 1° . The dashed line is the RXJ1252.9-292 color{m agnitude relation scaled to the redshift and colors for RX 0910+ 5422. The long(dashed vertical line is the magnitude lim it of the m orphological classi cation $z_{850} = 24 \text{ m}$ ag. The S0 CM R is bluer than the elliptical CM R by 0.07 0.02 m ag.



Fig. 8. CMR scatter as a function of galaxy age from simulation obtained using BC03 single burst (on the top) and constant star form ation (on the bottom) stellar population models for solar metallicity (solid line), half solar (dashed line) and twice solar (dash {dotted line). From the measured scatter it is possible to estim ate the mean age of the stellar bursts.



Fig. 9. $|(i_{775} z_{850})|$ (AB) vs (J K) (Vega) color. Filled circles are ellipticals, squares are SOs, and stars SO/as. BC 03 single burst models are shown for half solar (dashed line), solar (solid line), and twice solar (dotted line) metallicities. Ages run from 1.5 to 3.5 G yr, by steps of 0.5 G yr, as diam onds from left to right. The bluer SO colors might be due to younger ages or low erm etallicities.



Fig. 10. A symmetry (A) vsC oncentration (C) parameters for ellipticals (circles), S0 (squares), S0/a (stars) and spirals (triangles) in the $(i_{775} z_{850})$ color range between 0.5 and 1.3 m ag for RX J 0910+ 5422 (left). Large diamonds identify red sequence $[(i_{775} z_{850})$ color between 0.8 and 1.1 m ag] early (type galaxies. On the right, the same parameters are shown for all galaxies in our low (redshift sam ple using the same symbols for di erent galaxy types. E and S0 in RX J 0910+ 5422 show parameters that are characteristic of an early (type population, A < 0.2 and C > 0.3, as most of the low redshift early (type galaxies. This means that S0s were not likely m isclassi ed late(type galaxies.



Fig. 11. Sensic index n as a function of e ective radius R_e from GALFIT is shown for ellipticals (circles), S0s (squares), S0/a (stars) and spirals (triangles) with (i_{775} z_{850}) colors between 0.5 and 1.3 m ag. Red sequence ((i_{775} z_{850}) color between 0.8 and 1.1 m ag) galaxies are shown by large diam onds. E and S0 in RXJ 0910+5422 show a Sensic index 2. This means that S0s were not likely m isclassi ed late{type galaxies.



Fig. 12. A xial ratios vs e ective radii are shown for ellipticals (circles), S0 (squares) in the $((i_{775} z_{850}) color range between 0.5 and 1.3 m ag)$. Large diam onds identify red sequence galaxies with $(i_{775} z_{850}) color between 0.8 and 1.1 m ag$. Stars are S0/as and triangles are spirals.



Fig. 13. On the top, the histogram of axial ratios for the ellipticals (red line) and SOs (blue line) are shown. D ashed lines are for red sequence galaxies ($(i_{775} z_{850})$) between 0.8 and 1.1 m ag). We notice a lack of high axial ratio (low ellipticity) SOs in RXJ0910+5422. On the bottom, we show elliptical (red) and SOs (blue) distributions in RXJ1252.9-292 and RXJ0848+4452. The median of their SO distribution is consistent with that expected for nearly axisymmetric disks viewed at random inclination angles.



Fig. 14. Red sequence S0 color gradients as a function of distance from the galaxy center. The cross is the color calculated at the e ective radius.



Fig. 15. Red sequence elliptical color gradients as a function of distance from the galaxy center. The cross is the color calculated at the e ective radius.



Fig. 16. We show the histogram of the axial ratio of all early-types (black) and early-types with $(i_{775} z_{850}) < 0.99 \text{ mag}$ (blue). Elliptical W hereas there is a lack of high axial ratio (low ellipticity) SOs in RXJ0910+5422, there is no sim ilar lack in total early type or in the blue early-type sam ple.



Fig. 17. The $(i_{775} z_{850})$ color residuals (with respect to the mean of full early-type CMR t) is plotted versus axial ratios for red sequence $((i_{775} z_{850})$ color between 0.8 and 1.1 m ag) ellipticals (circles), S0s (squares), S0/a (stars) and spirals (triangles). While there is a lack of S0s with $\frac{b}{a} > 0.7$, we nd ve ellipticals and one spiral that have blue colors (negative CMR residuals) similar to most S0s, but round shapes $(\frac{b}{a} > 0.7)$, unlike the classi ed S0s. These may be the face-on counterparts of the elongated blue S0 population (see text).



Fig. 18. Red sequence galaxy colors, corrected by the ellipticalCMR, as a function of the distance from the cluster center. Filled circles are ellipticals, squares SOs, stars SO/a, and triangles spirals. Spectroscopically con rm ed m em bers are indicated by boxes. K now n non-m em bers are om itted from the plot. The continuous and the dashed lines show the elliptical and the SO CMR zero point, respectively. M ost of the blue SO lie close to the cluster center, not on the outskirts.



Fig. 19. | The z_{850} lum inosity functions for all galaxies (black circles with errors), as well as red sequence early-type (red), elliptical (blue), and S0+ S0/a (green) galaxies are shown. The faint end of the red sequence lum inosity function is dom inated by S0s and S0/as. The solid line is the t to the total lum inosity function. We obtain M = $22.6_{-0.7}^{+0.6}$ m ag and = 0.75 0.4. Galaxies have been classified m orphologically down to z_{850} = 24 m ag. The histogram of red sequence galaxies in RD S1252.9-292 is shown as the dashed red line, as a comparison. The red arrows show the histogram values after background subtraction.

Table 1:Color{M agnitude Relations _

| Sample | N C ₀ (m ag) | | S lope | | int (m ag) | | |
|------------------------|----------------------------|------|--------|--------|---------------|-------|-------|
| | | | | | | | |
| E+S0+S0/a ¹ | 31 | 0.99 | 0.01 | -0.030 | 0.020 | 0.060 | 800.0 |
| E+S0 ^{1a} | 14 | 1.00 | 0.02 | -0.010 | 0.033 | 0.054 | 0.009 |
| E ¹ | 19 | 1.02 | 0.01 | -0.033 | 0.015 | 0.042 | 0.011 |
| E ^{la} | 10 | 1.02 | 0.04 | -0.020 | 0.044 | 0.047 | 0.022 |
| S0 ¹ | 9 | 0.95 | 0.02 | 0.005 | 0.023 | 0.044 | 0.02 |
| S0+ S0/a ¹ | 12 | 0.95 | 0.02 | -0.007 | 0.027 | 0.057 | 0.015 |
| | | | | | | | |
| E+S0+S0/a ² | 32 | 0.99 | 0.01 | -0.032 | 0.019 | 0.060 | 800.0 |
| $E + S0^{2a}$ | 15 | 0.99 | 0.02 | -0.021 | 0.034 | 0.054 | 0.009 |
| E ² | 19 | 1.02 | 0.01 | -0.033 | 0.015 | 0.042 | 0.011 |
| E ^{2a} | 10 | 1.02 | 0.04 | -0.020 | 0.044 | 0.047 | 0.022 |
| S0 ² | 10 | 0.95 | 0.02 | -0.012 | 0.036 | 0.051 | 0.018 |
| S0+ S0/a ² | 13 | 0.96 | 0.02 | -0.015 | 0.033 | 0.065 | 0.015 |
| | | | | | | | |
| E+S0+S0/a ³ | 34 | 0.99 | 0.01 | -0.036 | 0.018 | 0.059 | 800.0 |
| E+S0 ^{3a} | 15 | 0.99 | 0.02 | -0.022 | 0.035 | 0.054 | 0.009 |
| E ³ | 20 | 1.01 | 0.01 | -0.032 | 0.015 | 0.044 | 0.010 |
| E ^{3a} | 10 | 1.02 | 0.04 | -0.021 | 0.046 | 0.047 | 0.022 |
| S0 ³ | 11 | 0.96 | 0.02 | -0.022 | 0.038 | 0.053 | 0.015 |
| S0+ S0/a ³ | 14 | 0.96 | 0.02 | -0.024 | 0.034 | 0.065 | 0.013 |

_

1:within 1⁰ 2:within 1.5⁰

3: within 2⁰

a: only con medmembers