Full-Polarization Observations of OH M asers in M assive Star-Form ing Regions: II.M aser Properties and the Interpretation of Polarization

Vincent L.Fish¹

Harvard{Sm ithsonian Center for A strophysics

60 Garden Street, Cambridge, MA 02138

National Radio A stronom y O bservatory

P.O.Box O, 1003 Lopezville Road, Socorro, NM 87801

vfish@nrao.edu

and

Mark J.Reid

Harvard{Sm ithsonian Center for A strophysics 60 Garden Street, Cam bridge, MA 02138 reid@cfa.harvard.edu

ABSTRACT

W e analyze full-polarization VLBA data of ground-state, m ain-line OH m asers in 18 m assive star-form ing regions previously presented in a companion paper. From the aggregate properties ofour sources, we con m results previously seen in the few individual sources studied at m illiarcsecond angular resolution. The OH m asers often arise in the shocked neutral gas surrounding ultracompact H II regions. Magnetic elds as deduced from OH maser Zeeman splitting are highly ordered, both on the scale of a source as well as the maser clustering scale of 10^{15} cm . Results from our large sample show that this clustering scale appears to be universal to these m asers. OH m as around ultracom pact H \square regions live 10^4 years and then turn o abruptly, rather than weakening gradually with time. These masers have a wide range of polarization properties. At one extreme (e.g., W 75 N), -components are detected and the polarization position angles of m aser spots show some organization. At the other extrem e (e.g., W 51 e1/e2), alm ost no linear polarization is detected and total polarization fractions can be substantially less than unity. A typical source has properties interm ediate to these two extrem es. In contrast to the wellordered magnetic eld inferred from Zeem an splitting, there is generally no clear pattern in the distribution of polarization position angles. This can be explained if Faraday rotation in a typical OH m aser source is large on a m aser am pli cation length but sm allon a single (e-folding) gain length. Increasing or decreasing Faraday rotation by a factor of 5 am ong di erent sources can explain the observed variation in polarization properties. Pure -components (in theory 100% linearly polarized) have long been sought, but seldom found. We suggest that alm ost all -com ponents acquire a sign cant am ount of circular polarization from low -gain stimulated em ission of a -component from OH appropriately shifted in velocity and lying along the propagation path.

¹Jansky Fellow

Subject headings: m asers | polarization | m agnetic elds | stars: form ation | ISM : m olecules | radio lines: ISM

1. Introduction

Hydroxyl (OH) massers are common in massive star-forming regions (SFRs). Their small size and large Zeem an splitting coe cient allow them to serve as probes of the velocity and magnetic elds on a very small scale. Because OH massers often cluster together in large numbers on subarcsecond scales (e.g., Reid et al. 1980), very long baseline interferom etric (VLBI) techniques are required to identify individual maser features. Due both to this close clustering of maser spots and to the tendency for the two components of a ground-state Zeem an pair to have highly unequal uxes (Cook 1966), the resolution a orded by VLBI is necessary in order to identify most Zeem an pairs (see, for instance, F ish et al. 2003). Likewise, identifying individual maser spots in a cluster is a prerequisite to understanding the linear polarization of OH masers, since blending of the linear polarization from adjacent masers with di erent polarization properties can corrupt the interpretation of the polarization.

O ver a quarter century has passed since the rst O H m aser source was observed with VLBI resolution: W 3(O H) by Reid et al. (1980). Since then, only a few m ore interstellar ground-state O H m aser sources have been observed at VLBI resolution (H aschick et al. 1981; Zheng 1997; Slysh et al. 2001, 2002). In order to understand the range of environments probed by O H m asers in m assive SFRs, we have undertaken a survey of the O H m asers in 18 m assive SFRs with the N ational R adio A stronom y O bservatory's¹ Very Long B aseline A rray (VLBA). The data have already been published as F ish et al. (2005, hereafter P aper I).

In this paper we analyze the results both in terms of relevance to individual sources and as a large collection of 0 H m asers that can shed additional light on the physical processes of 0 H m asers and the interpretation of their polarization. A brief overview of linear polarization theory is provided in x2. In x3, we consider the set of 0 H m asers as a whole to derive statistical properties. In x4, we present plausibility argum ents to attempt to identify the physical processes responsible for the observed properties of 0 H m asers. Finally, we sum marize our results in x5. C om ments on individual sources are included in Appendix A.

2. Linear Polarization Theory

In the presence of a magnetic eld, each of the main-line ${}^{2}_{3=2}$; J = 3=2 O H transitions split into three lines: two elliptically-polarized -components shifted in frequency by the Zeem an elect, and one linearlypolarized -component at the natural frequency of oscillation for the velocity of the emitting material (see, e.g., D avies 1974). Due in part to unequal amplication, this clear pattern of three lines has only been seen once (Hutawarakom, Cohen, & Brebner 2002). To this point, there have been few, if any, unambiguous detections of -components. Offen, a single -component is seen by itself, without an accompanying component polarized in the opposite circular handedness strong enough to be detected. Signi cant linear polarizations, which could be produced by -components, have been seen (Slysh et al. 2002). However, -components associated with magnetic elds with a nonzero perpendicular (line-of-sight) component are in general elliptically polarized, and elliptical polarization is the sum of circular and linear polarizations.

¹The N ational R adio A stronom y O bservatory is a facility of the N ational Science Foundation operated under cooperative agreem ent by A spociated U niversities, Inc.

Thus, signi cant linear polarization fractions can be produced by both - and -components, the latter theoretically being 100% linearly polarized in all instances.

The existence of maser spots with large (but not unity) fractional linear polarization suggests that components may in practice be partially circularly polarized as well. It is dicult to select an observational threshold of the linear polarization fraction that divides -components from -components. Theoretical modelling by G oldreich, K eeley, & K wan (1973b) and G ray & Field (1995) suggests that amplication of -components is stronger than that of -components for & 55, where is the angle of the magnetic eld to the line of sight. This corresponds to a linear polarization fraction 0:50. But since linear and circular polarization fractions add in quadrature, a completely polarized spot is not more linear than circular until the linear polarization fraction exceeds 0.71. Even then it is unclear how -components, which are theoretically to tally linearly polarized, can acquire a circular component of polarization. This issue is discussed in further detail in x4.4.

It is in portant to distinguish between - and -components, because the interpretation of the orientation of the magnetic eld based on the linear polarization position angle is di erent for the two cases. The electric vector for a -component is perpendicular to the two-dimensionalm agnetic eld direction (i.e., on the plane of the sky), while the electric vector for a -component is parallel to the two-dimensionalm agnetic eld direction. Thus, without inform ation as to which maser spots are -components and which are -components, there is a 90 am biguity in the direction of the magnetic eld on the plane of the sky.

3. Results

3.1. A re Zeem an Triplets Ever Seen?

H utaw arakom, C ohen, & B rebner (2002) identi ed a com plete Zeem an triplet in the northernm ost group of OH m asers in W 75 N, the st and only OH Zeem an triplet identi ed. Figure 1 shows a spectrum of a subregion of the northernm ost group of OH m asers. Three m aser lines can be seen at 4.1, 5.7, and 7.3 km s¹. The splitting of the 4.1 and 5.7 km s¹ m aser lines is consistent with a magnetic eld of + 5.5 m G and a velocity of 5.7 km s¹. We con m the existence of this triplet, although the three components are not perfectly aligned spatially. The angular separation between the - and RCP - component is 55 m as, which corresponds to a two-dimensional linear separation of 1.6 10^{15} cm (110 AU). This is roughly the cluster scale of 0 H m asers (see x3.2), though larger than the typical scale of separation between the two - components in a Zeem an pair. It is possible that the - and both - components all are located in the same cloudlet but from amplic cation paths that are not coincident.

The linear polarization fractions of the -components are 16% and 20%, while the corresponding fraction of the -component is 86%. The polarization position angles of the -components at 4.1 and 7.3 km s¹ are 45 and 67, respectively, while the position angle of the -component at 5.7 km s¹ is 54. This roughly ts the theory that -components are linearly polarized with a position angle perpendicular to that of the -components.

The angle of the magnetic eld to the line of sight, , can be derived from the form ula

$$f = \frac{\sin^2}{1 + \cos^2}; \tag{1}$$

where f is the linear polarization fraction of a -component (Goldreich, Keeley, & Kwan 1973a). This angle is 32 and 35 for the two -components of the Zeem an triplet. Equal amplication rates of the - and



Fig. 1. Zeem an triplet in W 75 N. Top: C ontour maps of the three Zeem an components with polarization vectors included. Note that the polarization vectors of the -com ponents are roughly perpendicular to that of the -com ponent. LSR velocities in km s¹ are shown in the upper right of each panel. B ottom : Spectrum of the dotted box region in the upper plots. The feature labelled \artifact" is due to the sidelable of a very strong maser spot outside the region shown. The velocities of the three marked components are consistent with a + 5:5 m G magnetic eld.

-com ponents occur when $\sin^2 = 2=3$, or for 55, with -com ponents favored when the magnetic eld is more highly inclined to the line of sight (Goldreich, Keeley, & Kwan 1973b). Modelling of OH hyper ne populations for conditions typically found at maser sites shows that the amplication of -com ponents falls o rapidly for < 55 (Gray & Field 1995). For smaller, beam ing and competitive gain favor -com ponents, suppressing amplication of -com ponents. Yet the inclination of the magnetic eld to the line of sight is 35, not 55, in the only con med Zeem an triplet.

Figure 2 shows a plot of the fractional linear polarization of m aser spots in the northernm ost cluster of 0 H spots in W 75 N versus the position angle of polarization. The m asers seem to be grouped into two populations. The rst group consists of m asers whose position angle is less than 90. These m asers have a high linear polarization fraction. The second group, with position angles > 90, are mostly circularly polarized. If we interpret these groups as - and -components respectively, the magnetic eld in plied from linear polarization lies at position angle 48 in the region, with a scatter of 24. This would indicate that the direction of the magnetic eld on the plane of the sky is roughly aligned along the NE-SW distribution of m aser spots and the elongation of the continuum source VLA 1.

3.2. Clustering Scale

In W 3(OH), Reid et al. (1980) found that the clustering scale of OH m assers was approximately 10^{15} cm. They calculated a two-point spatial correlation function and found that the probability per unit solid angle of nding another maser spot within angular distance l of a spot decreased sharply for $1 \& 10^{15}$ cm. We performed a similar analysis on each of our sources, and the results for the ten having a large number of maser spots are shown in Figure 3. For all ten sources, the probability P (1) drops sharply between a separation of 0 and 10^{15} cm (67 AU). The other sources in this study show similar behavior, although the plots are \noisier" owing to the smaller number of maser spots in the sources. This evidence argues in favor of a common clustering scale for OH masers in all massive SFRs. The maser spots within these clusters represent paths in the same condensation where the physical conditions are favorable to exponential amplication. If so, the overall extent of these condensations is perhaps a few times as large (150 AU), since paths through the periphery of the condensation are much less likely to produce comparable amplication lengths. In x4.6 we suggest that instabilities in the shocked neutral gas may lead to the formation of such condensations.

3.3. Zeem an Pairs and Component Intensities

In total, we do nd 184 Zeem and pairs in the entire sample set. These Zeem and pairs are listed in Table 21 of Paper I. A histogram of the distribution of implied magnetic eld strengths is given in Figure 4. The distribution rises with increasing magnetic eld strength to about 4 mG, then falls.

Few Zeem an pairs are found that im ply a splitting greater than about 8 m G, and the largest m agnetic eld strength found is 21 m G in W 51 e2. There is theoretical support for the existence of an upper lim it to the strength of the magnetic eld in a Zeem an pair for OH m asers. The collapse of material increases both the density and the magnetic eld; M ouschovias (1976) suggests that the relation could be as steep as n / B^2 , and Zeem an measurem ents of the magnetic eld in molecular clouds are consistent with this relation (e.g., C rutcher 1999). At some density, the rate of collisional deexcitation will be higher than the pump rate (presum ably from radiative excitation), and the population inversion between the lower and upper states will be destroyed. The rate of collisions of H₂ with OH is $n_{\rm H_2}$ h vi, where h vi = 10 ⁹ 10 ¹⁰ cm ³ s ¹, and



Fig. 2. | Linear polarization fractions as a function of polarization position angle (PPA) in the northern cluster of W 75 N. The elongation angle of VLA 1 (43) is marked along with the angle perpendicular to it. The PPAs of highly linearly-polarized spots (left of dotted line) tend to be aligned parallel to the axis of the continuum source, while the PPAs of highly circularly-polarized spots (right of dotted line) are roughly oriented perpendicular. If these groups are interpreted as - and -com ponents respectively, the implied projected magnetic eld direction is at a position angle near 43.



Fig. 3. Two-point spatial correlation function of the massers in ten massive SFRs. P (1) is the probability per unit solid angle that a masser spot can be found at angular distance 1 from any given masser spot. The angular distances have been multiplied by the distance to each source to give the linear separation, plotted as the abscissa. The probabilities P (1) have been norm alized to the greatest value. Note that P (1) drops to e^{1} at a separation of about 10^{15} cm for all ten sources.

the pump rate likely is 0.03 s^{1} , as would be expected for a far-infrared rotational transition (G oldreich 1975). Thus the population inversion will be destroyed by them al collisions when the density is a few times 10^{8} cm³. We note that the range of detected magnetic elds in our Zeem an pairs is a factor of 35 (0.6 to 21 mG), or roughly $\frac{P}{10^{3}}$. If the B² / n scaling law applies throughout this entire range, this im plies that the range of densities sam pled by those m asens in Zeem an pairs is a factor of 10^{3} , or about $n_{H_{2}} = 10^{5}$ m⁸, assuming that the 21 m G magnetic eld in W 51 e2 is near the upper end of magnetic eld strengths allowable before collisional depopulation of the upper state is signicant.

The lack of Zeem an pairs below 0.5 m G m ay be an observational e ect. For such sm all magnetic elds, the splitting is less than a typical line-width, and we make no attempt to identify Zeem an pairs less than 0.5 m G due to the di culty of distinguishing sm all Zeem an shifts from other e ects. For example, consider a right-elliptically polarized -component in a region where the magnetic eld orientation varies along the ampli cation path, as in F igure 5. The linear component of the polarization will be seen as weak emission in LCP. If there is also a velocity gradient along the ampli cation path, the linear polarization component may be shifted in velocity with respect to the circular polarization component. This would manifest itself in our observations as a velocity di erence between the lines seen in the two circularly-polarized feeds: the RCP feed would detect nearly all the emission, while the LCP feed would detect only the weaker, velocity-shifted linear component. For the parameters shown in Figure 5, the shift between LCP and RCP velocity corresponds to an apparent magnetic eld strength of 0.1 m G at 1667 M H z in the Zeem an interpretation, although broader lines and m ore extrem e variations of the linear polarization fraction across a linew idth can produce larger apparent shift between the components.

Figure 6 shows a histogram of the separation between the -com ponents in each Zeem an pair for which the -com ponent separation is less than 10^{15} cm. Zeem an pairs with larger separations are identified when unam biguous, but they have been excluded from the present consideration in order not to introduce bias. The rapid fallo of the number of pairs identified with increasing component separation suggests that the dearth of identifiable Zeem an pairs at larger separations is real. Note that Figure 6 has not been normalized by area; were it to be so normalized, it would fall even faster. If the distribution of Zeem an component separations were uniform ly random, a plot of the unnormalized number of identifiable Zeem an components versus component separation would be an increasing function of separation (at least up to a distance beyond which pairing is no longer unam biguous). This is certainly not observed, providing further evidence that the 10^{15} cm clustering scale is a physically signi cant scale over which physical param eters are su ciently coherent to provide an environm ent conducive to maser activity.

Thus it appears that the spatial separation of the -components in a Zeem an pair is generally a factor of several smaller than the size of the cluster containing the pair. Nevertheless, there are some pairs of m aser spots polarized predom inantly in opposite circular senses whose separation is comparable to or exceeds 10^{15} cm. It is possible that these are not true Zeem an pairs of -components from the same m asing subregion but rather two oppositely-polarized -components from di erent Zeem an pairs, of which only one component is seen in each. Such a situation could arise if ampli cation at each maser site favors only one sense of circular polarization, as described below. The magnetic elds implied by these Zeem an \cousins" would be less accurate due to two e ects. First, the system is velocity at each maser site may be di erent due to turbulence. Reid et al. (1980) calculate that a typical intracluster turbulent velocity is 0.6 km s¹. This corresponds to an elective Zeem an splitting of 1.0 m G in the 1665 M H z transition and 1.7 m G in the 1667 M H z transition. Second, the magnetic eld strength m ay be di erent at the maser sites. This di erence is generally less than 1 m G (see x3.5). The magnetic eld strengths in plied by Zeem an cousins should therefore be accurate to better than 2 m G.



Fig. 4. Histogram of magnetic eld strengths in plied by Zeem an splitting. The solid and dotted lines plot the same data but with the bins shifted by 0.5 mG. The dropo near zero is due to observational limits. The in plied total density is shown at the top, assum ing $n_{H_2} = 2 - 10^6 \text{ cm}^{-3}$ at 4 mG as suggested for W 3(O H) (Reid, M yers, & Bieging 1987) and an n / B² scaling law. The 19.8 and 21.0 m G elds in W 51, not included in this plot, would correspond to a density of 5 -10^7 cm^{-3} under these assum ptions.



Fig. 5. Velocity shift due for a right elliptically-polarized line with FW HM 0.5 km s¹ whose linear polarization fraction varies across the linewidth as shown in the top panel. The linear fraction could vary in this manner (for instance) due to a bend in the magnetic eld along the ampli cation path. The bottom panel shows the line as would be detected by RCP and LCP feeds. The LCP line is shifted 0.03 km s¹ with respect to the RCP line.

Figure 7 shows the ratio of the LCP and RCP uxes for the -components in each Zeem an pair. We observe that this ratio occasionally reaches values near 100. This is probably not a hard upper lim it but a result of observational constraints. M as spots weaker than 50 to 100 m Jy are too weak to be detected in our survey; m as spots stronger than 10 Jy are rare.

A simple test to determ ine whether our data have unknown system atics is to compare the number of Zeem an pairs with greater ux in the LCP and RCP components. There are 119 Zeem an pairs in the 1665 M H z transition and 65 in the 1667 M H z transition. The LCP component is stronger than the RCP in 69 of the 1665 M H z pairs and 44 of the 1667 M H z pairs. If Zeem an pairs with stronger LCP and RCP components are equally common, we would expect the number of Zeem an pairs in each polarization to be 595 5.5 at 1665 M H z and 32.5 4.0 at 1667 M H z². The deviations from these values are not statistically signi cant. The ratio of intensities of the -components in a Zeem an pair does not appear to be substantially di erent for the 1665 M H z transitions.

Figure 8 shows a histogram of the ux ratios of Zeem an components. There is not an appreciable di erence between the 1665 and 1667 M H z transitions for ux ratios less than 10. The larger number of high ux ratios (> 10) for the 1665 M H z transition appears real, although it m ay partially be due to a selection e ect. A sF igure 9 shows, the brighter component in a typical 1665 M H z Zeem an pair is brighter than that of a 1667 M H z pair. The detection lim it in our survey varies som ew hat by source but is approximately 0.1 Jy. Thus, the identication of a Zeem an pair with ux ratio x requires that the stronger component have a ux density higher than 0.1 x Jy. The relative scarcity of 1667 M H z Zeem an pairs with a ux ratio greater than 10 can be explaned by the paucity of Zeem an components with a ux density greater than 1 Jy. O nly in one-third of cases does a 1667 M H z Zeem an pair include a component stronger than 1 Jy, while over half of 1665 M H z pairs include a component above this threshold. At higher ux density thresholds, the di erence become s m ore extrem e.

The relative absence of 1667 M Hz Zeem an pairs with large ux ratios, noted previously in W 3(0 H) (W right, G ray, & D iam ond 2004), is consistent with the picture that m aser transitions with sm aller Zeem an splitting coe cients tend to have Zeem an pairs in which the -com ponents are m ore equal in intensity (e.g., M oran et al. 1978; C aswell & Vaile 1995). C ook (1966) theorized that correlated velocity and m agnetic eld gradients could be the cause of unequal spot intensities in Zeem an pairs. D eguchi & W atson (1986) argued that even absent a m agnetic eld gradient, a velocity gradient alone is su cient to produce unequal intensities. Velocity gradients (either alone or in com bination with m agnetic eld gradients) are less likely to disrupt am plication of only one of the -com ponents of a Zeem an pair for the 1667 M Hz transition than for the 1665 M Hz transition. M easured line widths for m aser spots at 1667 and 1665 M Hz are sim ilar, but the Zeem an splitting coe cient is sm aller for the 1667 M Hz transition than for the 1665 M Hz transition. The m agnetic eld strength required to produce an eld ective velocity shift greater than the linew idth of a com ponent is therefore greater for 1667 M Hz m asers than for 1665 M Hz m asers.

3.4. M aser Spot Statistics

We nd a total of 342 spots stronger in RCP and 351 in LCP at 1665 MHz, as well as 178 in RCP and 185 in LCP at 1667 MHz. As expected, there does not appear to be a preference for masers to appear

²For a binom ial distribution, ² = N pq, where N is the sample size and p and q are the probabilities of each outcome ($S_{LCP} > S_{RCP}$ and $S_{LCP} < S_{RCP}$).

preferentially in one circular polarization than the other, though occasionally an individual source m ay have a preponderance of m aser spots in one polarization, as is the case in W 75 S, in which we nd 35 RCP and 19 LCP spots in the 1665 M H z transition. For a source with 54 spots, we would expect 27 4 spots in each polarization, so the deviation seen in W 75 S is not signi cant.

There are nearly twice as many spots detected at 1665 M Hz than at 1667 M Hz, although one source (G 40.622 0.137) has one more spot in 1667 M Hz than at 1665 M Hz. This ts with theoretical modelling, which shows that while 1665 M Hz and 1667 M Hz masing offen occur under the same physical conditions, the area of physical parameter space conducive to am pli cation is larger for 1665 M Hz m asers than for 1667 M Hz masers (C ragg, Sobolev, & G odfrey 2002).

3.5. M agnetic Field Structure

M agnetic elds as determ ined from Zeem an splitting of OH m asers are predom inantly ordered in m assive SFRs (e.g., B aart & C ohen 1985; G arc a-B arreto et al. 1988). In all sources with the possible exception of G 43.796 0.127, the line-of-sight direction of the m agnetic eld (i.e., either tow ard or away from the Sun) is either constant for all Zeem an pairs or shows only one organized reversal in which there exists a line that can be drawn that separates the side of the SFR where the m agnetic eld is positive from the side where it is negative.

The relative consistency of magnetic eld strengths in clusters of 0 H masers argues in favor of an organized eld structure. When multiple Zeeman pairs are seen in the same cluster, the range of eld strengths is rarely greater than 2 mG (i.e., 1 mG) and often signi cantly less. In no case does the sign of the magnetic eld change between two Zeeman pairs in the same cluster of spots within $3 \, 10^{15}$ cm. Figure 10 shows the fractional variation of magnetic eld strength measurements in clusters compared with the source as a whole. The intracluster variation in the magnetic eld strength is generally smaller than the intrasource variation. If variations in magnetic eld strength are due to variations in density, this suggests that density uctuations within a cluster may also be smaller than uctuations on the scale of the masing region of a massive SFR.

G iven the uniform ity of magnetic eld direction in the line-of-sight dimension, it is som ew hat surprising that linear polarization vectors in the same cluster, when converted to magnetic eld directions, are often quite disordered. Table 1 shows the relative variation of polarization position angle (PPA) as a function of maser spot separation. This statistic has a range of [0,90], since a PPA of angle x is equivalent to one of angle x + 180 and the di erence cannot exceed 90. For pairwise separations shown in the rst column of Table 1, the rms of the di erence in PPA between the two maser spots was calculated. The statistic was applied only to spots with a linear polarization fraction less than 0.707 (equal parts linear and circular for a totally polarized m aser). This is designed to choose only -components. Any statistic comparing the magnetic eld direction at both - and -components would have to account for the natural 90 di erence arising from the PPAs of - and -components in the same magnetic eld. A sample of maser spots with a uniform random distribution of PPAs would have an rm s of 52. The rm s value of the PPA di erence between maser spots is roughly constant to within the errors for all maser spot separations. Even at the sm allest scale (< 10¹⁴ cm) the rms in PPA di erences is consistent with a random distribution. Given the regularity of magnetic eld direction both on source and cluster scales, the PPA di erences cannot be due to m agnetic eld variations within a cluster. Probably Faraday rotation is large enough in most sources, even on AU scales, to scram ble the linear polarization directions. See x42 for further discussion of the possible e ects of Faraday rotation.

3.6. Relation of OH M asers to the H II Region

The association between OH m assers and UCH II regions has been noted by many authors, including D ieter, W eaver, & W illiam s (1966) and M ezger & Hoglund (1967). Our larger sam ple of OH m assers mapped with m illiarcsecond resolution allows us to con m this nding. For each continuum source with nearby m assers, we determ ined an ellipse whose m a pr and m inor axes best m atched the overall extent of continuum em ission (full width at zero power for a 4 detection). The center of the H II region was taken to be the center of the ellipse, and the radius of the H II region was taken to be the geom etric m ean of the sem im a pr and sem im inor axes. The distance of each OH m asser from the center of the H II region, in units of H II region radii, was computed. Two possible sources of error are the uncertainty in m ap registration between the OH m assers and continuum im ages and the uncertainty in the assignment of the center of continuum em ission in each source. The form er is estim ated to be 0° . (1) by Argon, Reid, & M enten (2000). The latter m ay vary depending on source structure. For a circular H II region, we estim ate a 10% error, which would correspond to a 1 error of 0° . If or a typical 1° UCH II region, resulting in a total error of 0° . So the combination of the two e ects. For large (e.g., G 5.886 0.393) or inregularly-shaped (e.g., G 35.577 0.029) H II regions, the error in the estim ate of the center of continuum em ission m ay be slightly higher.

The distribution of the distance of 0 H m asers, norm alized by area, from the center of the H II region is shown in Figure 11. M asers in G 9.622+0.195, G 34.257+0.154, and C ep A were treated as containing three, two, and four continuum sources respectively, and m asers distances were calculated from the nearest source. Several sources were excluded from this analysis because no nearby continuum source was detected: the northern grouping of G 9.622+0.195, G 40.622 0.137, S269, M on R 2, and G 351.775 0.538. A dditionally, W 75 N and the two southeastern continuum sources of C ep A were excluded since it is not always clear which of several continuum sources to associate a m aser spot with.

Including ON 2 N and G 40.622 0.137, 50% of maser spots are located within 1.5 radii of a UCH II region; this num ber grows to 58% when these two sources are excluded. The distribution of maser spots in these two sources is clearly o set from the H II region, suggesting that the masers may be associated with a second, undetected continuum source nearby. The large peak near 0.5 H II region radii in Figure 11 is due mainly to the western cluster of maser spots in G 5.886 0.393. Overall, it appears that the distribution of O H masers peaks near the center of the H II region, consistent with G aray, Reid, & M oran (1985). The tail of the distribution of maser spots at several radii from the UCH II region may represent spots associated with another star, not with the nearest detectable continuum source, as in ON 2 N and W 51 (e1 and e2).

These results are not consistent with a random distribution of OH m aser spots within a shell around the H II region. A uniform random distribution would peak at a projected radius $r_{proj} > 1$ (in H II region radii). For density distributions falling o as r¹ or r², the peak of the distribution will be at $r_{proj} = 1$. In all cases, the distribution of OH m asers would be expected to double across $r_{proj} = 1$ because m asers located behind the H II region would be obscured by the UCH II region, which is optically thick at = 18 cm. We see no evidence for this discontinuity at 1 H II region radius in our data.

Figure 12 shows the distribution of projected linear distances of OH masers from the center of the associated UCH II region. 82% of maser spots are located within 13000 AU of the center of the H II region. A loo shown is the assumed dynam ical age of the masers for an expansion speed of 3 km s¹. This is consistent with the speed measured in W 3(OH) from both proper motion of OH masers (B loem hof, Reid, & M oran



Fig. 6. Histogram of identi ed Zeem an component separations. Note the rapid fallo of the distribution of -component separations in a Zeem an pair. This suggests that the components of a Zeem an pair are generally separated by a distance less than the 10^{15} cm (67 AU) clustering scale (see x3 2). Ten Zeem an pairs with component separations exceeding 80 AU are not shown. The plot is not norm alized by area; the fallo would be much faster if it were.

Separation (cm)		Numberof Pairs ^a	PPA mrs	Standard E rror ^b
0 10 ¹⁴		25	45 : 7	9 : 1
1014	10 ¹⁵	189	46:6	3:4
1015	10 ¹⁶	566	49:6	2:1
1016	10 ¹⁷	1707	51 : 8	1:3

Table 1. Variation of PPA with Pairwise Maser Separation

Note. | Table of m s variations in the PPA between pairs of m as spots for all sources. Only pairs of spots whose linear polarization fraction is less than 0.707 are considered.

 $^{\rm a}{\rm N}$ um ber of pairs of maser spots with linear polarization fractions less than that shown in the column headings such that the separation between the maser spots is less than that shown in the rst column.

^bStandard error of the mean: $m = \sum_{n=1}^{p} \overline{N}$.



Fig. 7. | Flux ratios of -com ponents in Zeem an pairs. D ata for 1665 M H z pairs are represented by open triangles, while data for 1667 M H z pairs are represented by lled squares. D otted lines are drawn at ratios of 3 and 1=3. The 19.8 and 21.0 m G elds in W 51 are not included.



Fig. 8. Histogram of ux ratios in Zeem an pairs. The ux ratio, determ ined by taking the ratio of the stronger ux to the weaker ux in a Zeem an pair, is binned by factors of 1.5. The dotted line shows data for the 1665 M Hz transition, and the solid line shows data for the 1667 M Hz transition. The data for the two transitions are scaled by the total number of Zeem an pairs identied in the transition. The histogram s are similar for ux ratios less than 10, but 25 of the 28 pairs with ux ratios greater than 10 are in the 1665 M Hz transition.



Fig. 9. Histogram of peak ux densities in Zeem an pairs (peak ux density in either LCP or RCP). Flux densities are binned by increments of 1 Jy. For clarity, points stronger than 40 Jy are suppressed. These consist of three points at 1665 M Hz (72, 163, 239 Jy) and two at 1667 M Hz (41, 85 Jy). The stronger component of a 1665 M Hz pair tends to be stronger than that of a 1667 M Hz pair.



Fig. 10. Plot of magnetic eld strength variation in clusters (sm aller than $3 \ 10^{15}$ cm) and sources. The fractional variation (m s/m ean) of the magnetic eld strength β jfor each cluster of two orm one Zeem and pairs is plotted against the fractional error for the source as a whole. G 9.622+ 0.195 and G 34.257+ 0.154 are each treated as two separate sources. Open squares indicate data points for the middle source in G 9.622+ 0.195, G 35.577 0.029, G 40.622 0.137, and S269, in which all Zeem and pairs are found in a single cluster. The dotted line indicates equal fractional errors in cluster and whole-source magnetic eld strength. 39 of the 43 lled squares fall below this line, indicating that the magnetic eld variation in a cluster is less than the variation in the source as a whole.

1992) and direct expansion of the UCH II itself (K awam ura & M asson 1998). The dynam ical age of m ost m asers is less than 2 10^4 yr.

W hile the number of OH m aser spots cuts o at a dynamical age of several 10^4 yr, individual OH m asers do not appear to fade appreciably during this time. Figure 13 shows the relation of maser power per bandwidth (i.e., ux density normalized to a constant distance) to the separation between the UCH II region center and the masers. Figure 14 shows the mean power per bandwidth and standard error of the mean for the same data. The maser power per bandwidth appears to be constant with distance from the center of the H II region. If the distance of the masers from the H II region is indeed correlated with their age, OH masers do not become system ically brighter (or fainter) with age, at least not in the 4 10^4 yr timescale our data span. Figure 15 shows the relation of maser power per bandwidth to the size of the associated UCH II region. Since H II regions undergo expansion, their size is a measure of the age of the system . A gain, there does not appear to be a correlation between the maser power per bandwidth and the age of the system . How ever, we do not see any maser spots located more than 30000 AU (0.15 pc) from the center of the associated UCH II region (including G 351.775 0.538, for which the nearest UCH II region is several arcseconds away).

Habing & Israel (1979) have observed that 0 H m assers are not seen around H II regions once they leave the ultracom pact phase (d < 0.15 pc). Indeed, not only are 0 H m assers not seen around \com pact H II (CH II) regions" (0.1 < d < 1 pc), they are not seen at com parable radii around ultracom pact H II regions. The lack of 0 H m assers at large distances from the associated ultracom pact H II region was rst noted by Habing et al. (1974), who suggested that 0 H m aser phenom ena disappear at a radius of 15000 AU (0.07 pc). Our larger sam ple size at much higher angular resolution indicates that there is a sharp cuto at about twice this radius. It is possible that the physical conditions (such as tem perature and density) responsible for m aser activity do not exist at large radii. A lternatively, the ionization front catches up to the shock front as the H II region expands into an environm ent whose density decreases with radius, thereby destroying the 0 H m asers, which are believed to exist in the region between the ionization and shock fronts (see x4.6).

3.7. Relation of OH M asers to the Surrounding M aterial

In order to interpret the bulk motions of OH massers, it is necessary to determ ine the velocity of the massive star or stars being formed. Frequently this is determined through hydrogen recombination line observations, which give information on the velocity of the ionized H II region surrounding the central star. But recombination lines are not well suited to the task. Recombination lines are subject to sometimes severe D oppler and impact broadening. Even at high frequency, di erent recombination lines can be biased by a few km s¹ (Berulis & Ershov 1983; Sam s, M oran, & Reid 1996) due to di ering optical depths in an expanding H II region.

We have observed the (J;K) = (1;1) line of ammonia in order to provide a context for the OH maser observations. Reid, M yers, & Bieging (1987) argue, based on the similarity of distribution of NH₃ absorption and OH maser emission in W 3(OH), that the two species are found in the same clumps of material. The physical conditions they deduce from NH₃ (1;1) and (2;2) observations are consistent with the physical conditions necessary for OH maser activity. Also, the velocity of peak absorption in NH₃ closely matches the average velocity of the OH masers. Unfortunately, in many sources NH₃ absorption is not clearly detected, and NH₃ emission velocities must be used instead. NH₃ emission is a less reliable indicator of the velocity of the central star, since emission usually traces motions on a larger scale (offen > 1⁰). In the region of W 3(OH), ammonia emission velocities di er from the average OH maser velocity by approximately 4 km s¹ (W ilson,



Fig. 11. | Histogram of distances between OH m asers and UCH II regions shown in units of equal area. M asers that cannot be identified as unambiguously associated with a particular H II region have been excluded; see x3.6 for details. We nd that 58% of OH m asers appear within 1.5 radii of the H II regions, suggesting that OH m asers in m assive SFRs with H II regions are indeed spatially associated with them. The bump at 2.7 H II radii is due primarily to the masers in ON 2 N; it is probable that these masers are not actually associated with the H II region shown in Figure 23 of Paper I.A small tail of the distribution out to 6 H II radii has been suppressed for clarity.



Fig. 12. Histogram of absolute distances between OH m asers and UCH II regions. The data are as in Figure 11 but plotted in distance units and not norm alized by area. The expansion age, de ned as radius divided by expansion speed, shown at the top would be appropriate for expansion at 3 km s¹, as measured for W 3(OH).



Fig. 13. Distribution of maser power per bandwidth with radius. All ux densities are multiplied by the square of the distance to the source in kiloparsecs. The horizontal axis shows the distance of each maser spot from the center of the associated UCH II region. Multiple H II regions in the same source (e.g., G 9.622+0.195) are considered independently. Data are not plotted when no nearby UCH II region is seen (e.g., S269 and G 351.775 0.538) or when it is unclear which H II region to match maser spots with (W 75 N). There does not appear to be a correlation between the power per bandwidth and the distance of maser spots from the UCH II region. The slight dip near zero radius is due to the inclusion of C ep A, whose proximity allowed detection of spots of weaker norm alized ux density than for other sources.



Fig. 14. Distribution of mean maser power per bandwidth with radius. The plot shows the mean of the maser ux normalized to a distance of 1 kpc as well as the standard error of the mean $(m \approx \frac{P}{N})$. Errorbars are not symmetric because a linear average is plotted on a logarithm ic scale. The data are binned by units of 2000 AU. The mean power per bandwidth and the distance of maser spots from the UCH II region is constant with distance, although large deviations are possible due to source-to-source di erences. See Figure 13 for more details.



Fig. 15. Distribution of maser power per bandwidth with H II region size. The horizontal axis shows the geometric mean of the sem im a pr and sem im inor axes of the UCH II region, converted to linear distance. See Figure 13 for more details. There does not appear to be a correlation between the size of the H II region and the power per bandwidth of the maser spots surrounding it. Since the size of an H II region is a measure of its age, this suggests that OH m asers do not become e system ically fainter over the range of ages of the H II regions in our sam ple. As in Figure 13, the dip near zero radius is due to Cep A.

G aum e, & Johnston 1993). It is possible that system ic biases of a few km s¹ are introduced using emission velocities, but there is no way to obtain the radial velocity of the star to greater accuracy.

The spectra in F igures 40 to 42 of P aper I are provided for regions of emission or absorption located roughly coincident with the extent of O H m asers on the sky. It is important to remember that while O H m asers exist primarily near a UCH II region, the neutral NH_3 m ay exist at a wide range of radii. Thus the inferred NH_3 velocities could in principle be a ected by motion of material quite distant from the H II region. The NH_3 velocity m ay be shifted from the rest velocity of the star by an am ount comparable to the velocity dispersion of the molecular cloud. From the virial theorem, the velocity dispersion is

$$v = \frac{r}{\frac{GM}{2R}};$$
 (2)

where G is the gravitational constant, M is the total mass, and R is the radius of the sphere. For a cloud with M = 100M and a radius of 0.1 pc, v 1.5 km s¹.

W hen the velocity of the nearby NH₃ is measured, it generally falls toward the middle of the range of OH maser velocities, as shown in Figures 40 to 42 of Paper I. There are some minor exceptions to this rule. In G 35.577 0.029 and M on R2, the NH₃ velocity is near an extrem um of OH maser velocities. (In ON 1 the OH masers fall into two disjoint groups at 2:5 6 km s¹ and 13 17 km s¹, and NH₃ emission detected in a broad region located 5° to 30° north of ON 1 (not shown in the panel) falls near 11 km s¹.)

Figure 16 shows a histogram of the di erences between OH m aser velocities and the adopted NH_3 velocity for all sources having detected NH₃. Of the 926 OH m aser spots, 51.7% are blueshifted with respect to the NH₃ velocity, and 48.3% are redshifted. The median velocity dierence is 0.30 km s¹ with an rm s of 1.89 km s¹, while half the di erences fall within the range 3.52 to + 3.01 km s¹. Since a zero di erence falls com fortably within this range, we cannot con dently state that 0 H m asers are consistently blueshifted or redshifted with respect to the surrounding m aterial, as m ight be expected if a single type of m otion, such as expansion or contraction, dom inates OH m aser kinem atics. W hen only m aser spots located within one projected H II region radius are considered (N = 261), the median velocity di erence is + 0.22 km s¹ with an m s of 1:90 km s¹ (Figure 17), with half the di erences falling in the range 2:58 to +3:98 km s¹. In either case, there does not appear to be a detectable di erence between the 0 H and NH_3 velocities to within our errors. If expansion dom inates the dynam ics of the masing regions, the masers projected atop the UCH II region (and therefore in front of it, since UCH II regions are in general optically thick at = 18on) should be blueshifted with respect to the large-scale am bient m aterial. However, the opposite appeared to be true of W 3(OH) (Reid et al. 1980), although later proper motion measurements of the OH masers de nitively established that they are expanding (B loem hof, Reid, & M oran 1992). It is worth pointing out how ever that in G 43.796 0.127, where nearly all of the massers are projected against the UCH II region, the OH m asers are preferentially blueshifted with respect to the NH_3 emission, mildly suggestive of expansion.

Because of the aforem entioned possibility of system ic errors of a few km s¹ in determ ining the radial velocity of the central star from NH_3 velocity m easurements, we cannot identify whether a single type of m otion, such as gravitational infall or slow expansion, dom inates the kinematics of OH m asers in m assive star-form ing regions. We can in general rule out kinematic m odes in which the OH m asers would be m oving at tens of km s¹ or m ore, such as a freely expanding H II region at & 10 km s¹.



Fig. 16. | Histogram of di erences between OH and NH₃ velocities for those sources in Figures 40 to 42 of Paper I with an NH₃ velocity indicated. The median velocity di erence is 0:30 km s¹ with a standard deviation of 1.89 km s¹. The mean is 0:29 km s¹ with a standard error of the mean (rm s/ $\frac{1}{N}$) of 0.06 km s¹, assuming all masers have independent velocities. Taking clumping into account, the standard error of the mean could be higher by a factor of 3.



Fig. 17. Histogram of di erences between OH and NH₃ velocities for those maser spots located within one radius of the associated UCH II region. The median velocity di erence is +0.22 km s¹ with a standard deviation of 1.90 km s¹. The mean is 0:79 km s¹ with a standard error of the mean of 0.12 km s¹.

4. Discussion

4.1. Saturation

Interstellar O H m asers are most likely saturated (R eid et al. 1980). The saturation tem perature, T_s , for O H m asers is given by

$$T_{s} = \frac{h}{2kA} \frac{4}{A}$$
(3)

(Reid & Moran 1988), where A is the Einstein coe cient, is the decay rate, and is the solid angle of beam ing. For = 0.03 s⁻¹ as typical for a far-infrared rotational transition likely pumping the maser, the saturation temperature is $(2 \ 10^8)^{-1}$ K. The most compact maser component from the space-VLBI observation of G 34.257+0.154 by Slysh, Voronkov, & Val'tts (2002) has a beam ing angle 0.01, which corresponds to a maximum saturation temperature $T_s \ 2 \ 10^{10}$ K. For the most part, the brightness temperatures (T_B) listed in Tables 2 through 20 of Paper I are below this value. But these are apparent brightness temperatures calculated from the undeconvolved spot size, which are likely scatter broadened for most sources.

M any of our spots appear to be partially resolved, as shown in Table 2. However, several caveats apply to the deconvolved spot parameters. First, the measured spot size may be larger than the physical spot size due to interstellar scattering. Second, large deconvolved spot sizes may be the result of misidentifying spatially blended maser spots as a single spot. Third, determining the deconvolved spot size of a small maser spot is less accurate than for a large maser spot, because the deconvolved spot size is obtained from di erencing two larger numbers (the squares of the undeconvolved spot size and the beam size). The net of all three e ects is that maser spot sizes are probably smaller than that calculated from deconvolved spot size would underestim at the actual T_B by the square of the ratio of the apparent spot size to the unbroadened spot size.

For a typical FW HM spot size of 3 m as in W 75 N (which has very little scatter broadening), $T_s = 5 \ 10^9$ K.This corresponds to a ux density of S = $2kT_s$ spot², where spot is the solid angle subtended by the maxer spot. Taking spot¹⁶ sr, the saturation value S 7 10^{25} erg cm² s¹ Hz¹ sr¹, or 0.07 Jy. This is near our detection limit, so nearly all spots that we detect are partially saturated if the spot sizes for W 75 N are typical.

4.2. Faraday Rotation

Faraday rotation can complicate the interpretation of linear polarization in two ways. First, external Faraday rotation between a maser and the observer will cause the polarization position angle (PPA) of linear polarization to rotate, making interpretation of the magnetic eld direction on the plane of the sky more di cult. Second, internal Faraday rotation along the ampli cation path may decrease the linear polarization fraction of the radiation, completely circularizing it if the Faraday rotation is strong enough (Goldreich, K eeley, & K wan 1973a). Since this also reduces the elective gain length for linear polarization, Faraday rotation may also prevent otherwise highly linearly-polarized maser components from being amplied to the limits of detectability. Thus, spots with a large linear polarization fraction (e.g., -components and

-com ponents where the magnetic eld is near the plane of the sky) may be suppressed relative to spots with a small linear polarization fraction (e.g., -com ponents where the magnetic eld is directed along the line of sight).

	D istance	N um ber of	M edian D econvolved	90% Range of Spot Sizes	
Source	(kpc)	Spots	Spot Size (m as)	Min (mas)	M ax (m as)
G5.886 0.393	3.8	98	19.66	12.44	30.33
G9.622+0.195	5.7	38	16.44	10.86	21.05
G10.624 0.385	4.8	14	23.55	19.48	28.96
G 34 257+ 0.154	3.8	88	5.46	0.00	13.50
G 35.577 0.029	10.5	15	16.22	12.39	19.55
G40.622 0.137	22	39	19.68	15.57	24,22
G43.796 0.127	9.0	60	7.96	0.00	13.11
W 51 el	7.0	97	10.51	6.62	20.07
W 51 e2	7.0	94	9.53	5.19	15.23
ON 1	3.0	49	2.68	0.00	7.30
КЗ 50	8.7	17	31.46	23.45	34.82
ON 2N	5.6	73	25.29	13.69	37.11
W 75 S	2.0	65	4.17	1.57	7.75
W 75 N	2.0	120	3.14	0.00	8.90
Cep A	0.7	62	6.77	1.95	12.96
NGC 7538	2.8	30	6.83	3.19	11.20
S269	3.8	19	6.08	1.23	9.47
M on R2	0.9	27	7.86	2.20	13.49
G 351.775 0.538	22	50	67.40	45.35	76.07

Table 2. Deconvolved Spot Sizes

Note. | Spot sizes m ay be overestim ates. See x4.1 for details.

External Faraday rotation in the interstellar medium between a maser and the observer would cause a rotation of the PPA of the linear polarization of each spot, given by

$$RM = 8:1 \quad 10^5 \quad n_e B_k \, dl;$$
 (4)

where RM is the rotation measure in rad m², n_e in cm³, B_k is the component of the magnetic eld parallel to the direction of propagation in G, and dl in is the di erential path length along the line of sight in pc (Thom pson, M oran, & Swenson 2001). In some regions, such as the northern cluster in W 75 N, the linear polarization vectors are predom inantly aligned along the line of maser spots (see x3.1). An RM of about 10 rad m² would produce a rotation of the PPAs in a source of 20. A rotation of the polarization vectors by an am ount greater than this would cause the vectors to no longer appear to be aligned with larger structures, unless the rotation was near a multiple of 180. A coording to the ATNF PulsarC atalogue (M anchester et al. 2005)³, the only pulsar with known rotation measure located within 10 of W 75 N at comparable heliocentric distance is B 2021+ 51, for which the RM is 6:5 rad m² (M anchester 1972).

Internal Faraday rotation over the region of ampli cation m ay destroy linear polarization in both – and –components, possibly suppressing –components altogether. The Faraday rotation over a region with average electron density n_e and parallelm agnetic eld strength B_k is

$$= 0:05 \frac{n_e}{1 \text{ cm}^3} \frac{B_k}{1 \text{ mG}} \frac{L}{10^{14} \text{ cm}} \frac{1}{18 \text{ cm}}^2;$$
(5)

where is the wavelength of the transition. For a typical ground-state (= 18 cm) OH m as B5mG.The e ective am pli cation length L is likely to be less than the clustering scale due to velocity coherence. A crude estimate is that L D v= V, where D = 10 15 cm is the diameter of the masing cloud, v = 0.2 km s 1 is a typical maser line width, and V 2 km s¹ is a reasonable velocity shift across the cloud based on observations of W 3 (O H) (Reid et al. 1980) and theoretical modelling (Pavlakis & Kyla s 1996). Thus, for an elective amplication length $L = 10^{14}$ cm, an electron density of about 300 cm³ would be su cient to produce a rotation of 90 along the path of ampli cation. For H $_2$ densities of 10^5 to 10^8 as is typical in OH m asing regions (Cragg, Sobolev, & Godfrey 2002), this would require a fractional ionization $(n_e = n_{H_2})$ of 3 10^{6} to 3 10^{3} . This is higher than the ionization rate that would be expected from cosm ic-ray ionization alone (Shu 1992), but consistent with the 10 4 that occurs in the C II regions around H II regions where OH m asers m ay exist (Stemberg & Dalgamo 1995; Garc a-Barreto et al. 1988). Ionized carbon and to a lesser extent sulfur may play an important role in producing free electrons, due to their abundance and ease of photoionization. W hile the hydrogen in the H II region absorbs all the ultraviolet photons with energies greater than 13.6 eV, m any softer photons pass through undisturbed. Stemberg & Dalgamo (1995) calculate that the ionization fraction m ay be slightly greater than 10 4 in the C II region, located around the H $\,$ II region, and about 10 5 in the S $\,$ II region, in turn located around the C $\,$ II region. Based on their m odels of photon-dom inated regions as well as the locations of 0 H m aser spots just outside H II regions, it is likely that they exist near or embedded in the C II regions.

We can form a consistent picture of linear polarization in OH m assers if the am ount of Faraday rotation in a typicalm aser source is near a critical point, i.e., such that the product $n_e B_k L$ several 10^{17} cm² mG. Based on maser line widths and brightness tem peratures, the am plication length is typically at least 20 unsaturated gain lengths and probably greater for highly saturated m assers (Reid & M oran 1988). A typical

 $^{^3{\}rm T}$ he catalogue is available on line at http://www.atnf.csiro.au/research/pulsar/psrcat .

m aser spot has a signi cant am ount (> 1 rad) of Faraday rotation over the am pli cation length but a sm all am ount (< 1 rad) over a single gain length. In this case, som e linear polarization will survive am pli cation, but Faraday rotation scram bles the PPA of the linear polarization, so it will not be simply interpretable as a m agnetic eld direction. If the Faraday rotation is a factor of 5 sm aller, the Faraday rotation over the am pli cation length will be sm all, so high linear polarization fractions m ay be observed, and the PPA m ay still be correlated with the m agnetic eld direction. On the other hand, if the Faraday rotation is a factor of 5 larger, the Faraday rotation over a gain length can be large, and linear polarization fractions will approach zero (G oldreich, K eeley, & K wan 1973a). In cases where the Faraday rotation per gain length is signi cant, interpretation of the PPA will be di cult because m aser am pli cation will stim ulate em ission in the orthogonal linear m ode as well (M elrose & Judge 2004). Signi cant generalized Faraday rotation m ay also circularize -com ponents.

An example of a source with small internal Faraday rotation over the entire amplication length is W 75 N, in which -components are detected in abundance, especially in the northernmost group of maser spots (see x3.1). As Figure 2 shows, - and -components are easily identiable in this group based on the PPA of the linear polarization. It is interesting to note that modelling of the OH masers in W 75 N by G ray, Hutawarakom, & C ohen (2003) indicates that the maser amplication length is several orders of magnitude smaller than that typically assumed in other sources, although the density is also higher. For the range of ionization fractions given above, the resulting Faraday rotation would be less than 1 radian over the amplication length.

At the other extrem e is W 51 el and e2, in which practically no linear polarization whatsoever is detected. This is consistent with signi cant Faraday rotation along a gain length, which would suppress the ampli cation of -components and circularize the otherwise elliptically-polarized -components. Such Faraday depolarization m ay also explain certain m aser features that are seen with similar ux densities in RCP and LCP but without any detected linear polarization, such as spots 18, 24, and 36 in Table 9 of Paper I. It is highly unlikely that the lack of linear polarization in W 51 can be due to chance alignments of the m agnetic eld in an extrem ely narrow cone oriented toward or away from us at each m aser site, producing

-components that are purely circularly polarized, because the magnetic eld is seen to reverse line-of-sight direction across the source. Thus, the inclination of the magnetic eld to the line of sight must take on values intermediate to the 0 and 180 required for pure-circular masers in the absence of Faraday rotation.

A medium range of Faraday rotation would be enough to partially (but not totally) circularize components. The observed circular polarization fraction will in general be a complicated function of the electron density, maser gain length, and angle of propagation with respect to the magnetic eld direction, but the presence of Faraday rotation will strictly increase the circular polarization fraction compared to the case in which no Faraday rotation is present (Field & Gray 1994). Straightforward application of equation (50) of Goldreich, Keeley, & Kwan (1973a) without accounting for Faraday rotation will cause the inclination of the magnetic eld to the line of sight to be underestim ated.

Faraday rotation m ay also explain why linear polarization vectors appear to be disorganized in some sources. If a large electron density is required for Faraday rotation along the ampli cation path in the interior of a masing cloud, it is likely that the electron density is high exterior to the masing cloud as well. This would rotate the apparent PPA of the maser emission. Fluctuations in the electron density, possibly caused by density inhom ogeneities or anisotropy of the ionizing radiation eld, could cause emission from adjacent maser spots to be Faraday rotated by di erent amounts. If this is indeed the case, reconstructing the magnetic eld orientation in the plane of the sky is a di cult task, and reconstructing the full three-dimensional orientation of the magnetic eld may be nearly in possible.

Scatter broadening of some sources in plies strong density uctuations and a high column density of electrons along the radiation propagation path. Since the scatter broadening is proportional to the distance between the source and the scattering screen (e.g., Boyd & W emer 1972), a screen of electrons near the source is unlikely to increase the angular size as much as a cloud of electrons several kiloparsecs away in the G alactic plane. The lack of correlation between scattered size and Faraday depolarization provides further evidence that the scattering is external to the masing regions. W 51 is not a particularly scatter-broadened source, yet essentially no linear polarization is detected. G 351.778 0.538 is heavily scatter-broadened, but several spots with a high degree of linear polarization are seen. Figure 18 shows a plot of the linear polarization of the linear polarization is detected. There does not appear to be a correlation of the linear fraction with the size of the observed maser spots. This suggests that the electron screen responsible for scattering is G alactic in origin.

4.3. Total Polarization

Of the maser spots in our sample, 97% are at least 75% polarized. Although Tables 2 to 20 of Paper I do not list the total polarization fraction explicitly, it can be approximated by noting that the total polarization fraction is $Q^2 + U^2 + V^2 = I$, where Q;U;V; and I are the Stokes parameters. Stokes I and V can be obtained from the sum and di erence, respectively, of the listed RCP and LCP ux densities of spots, and the linear ux density gives $Q^2 + U^2$. In 86% of the maser spots, LCP and RCP uxes are not both detected, in plying that the spot is circularly polarized to the limits of detectability. For other spots, frequently $Q^2 + U^2 + V^2 = I^2$, indicating that many spots are nearly 100% polarized, as shown in Figure 19. A portion of the discrepancy from equality in the above equation can be explained by a variety of factors. As can be seen in Tables 2 to 20 of Paper I, when the sam e maser spot is seen in both RCP and LCP emission, the position and velocity of the peak emission may be slightly di erent in both. Blending of strong adjacent maser spots can also make determination of t parameters di cult. Furtherm ore, the linear polarization fraction of a maser spot may vary across a spot size.

Nevertheless, there are certainly spots that appear to be only partially polarized (Figure 19). Unpolarized emission (Q = U = V = 0) would appear as equal ux densities in the LCP and RCP feeds with no detectable linear polarization. In W 51 el, for example, there are a number of maser spots where the LCP and RCP ux densities are nearly equal and centered at roughly the same position and velocity. Since there is nearly no detected linear polarization in the OH masers in this source, it is possible that these maser spots are only partially polarized. These maser spots could also be produced by masing in sites where the magnetic eld splitting is small compared to the line width (G oldreich, K eeley, & K wan 1973a). Our data do not provide su cient resolution to tell whether the spots with roughly equal RCP and LCP uxes are due to two 100% polarized spots that are at slightly di erent positions and velocities or whether they are due to a single spot that is not 100% polarized.

4.4. Overlap of M asing C lum ps

Theoretically, -components are favored form agnetic elds inclined 55 to the line of sight (Goldreich, Keeley, & Kwan 1973b; Gray & Field 1995) and -components are favored for smaller angles. Consider an ensemble of maser sites, each threaded by an independent, random ly-oriented magnetic eld. The fraction



Fig. 18. Plot of linear polarization fraction as a function of spot size. The spot size is taken to be the geometric mean of the undeconvolved spot tmajor and m inor axes. The relative lack of spots with a high linear polarization fraction at large spot sizes may be due to blending of adjacent spots within the beam. The linear polarization fraction does not appear to be otherwise correlated with the observed spot size.



Fig. 19. Totalpolarization of maser spots. Only spots with nonzero observed uxes in all three of RCP, LCP, and Linear ($\overline{Q^2 + U^2}$) are shown, which excludes 942 spots. Positive circular polarization fraction corresponds to positive Stokes V (i.e., RCP ux greater than LCP ux). Curves showing 25%, 50%, 75%, and 100% totalpolarization are drawn. M ost maser spots are > 75% polarized.

of maser sites for which amplication of -components is favored is

$$2 \frac{1}{4} = 0 = 0 = 0$$
 sin d d = 0.426; (6)

where the factor of two accounts for the possibilities of the magnetic eld being oriented toward or away from the observer. Accounting for the fact that two -com ponents are produced for each -com ponent by Zeem an splitting, -com ponents would comprise 2 0.426 = (1 + 0.426) = 60% of total maser spots.

As discussed in x3.5, the magnetic eld in any individual source is highly ordered, and the distribution of magnetic eld orientations at maser sites is not oriented in a uniform ly random direction. But for a large sample of sources distributed throughout the Galactic plane, it is plausible that the range of magnetic eld directions sampled will approach a uniform random sample. However, in our sample a large majority of m aser spots have a sm all or zero linear polarization fraction. Two-thirds of m aser spots have no detectable linear polarization at all. As discussed in x42, amplication of -components may be reduced by Faraday rotation along the ampli cation path. Two additional e ects pointed out by Elitzur (1996) may explain the larger fraction of -components we detect. First, although an inclination of 55 divides angular phase space into two distinct regions in which - and - com ponents dom inate, the relative amplication of - and -components is larger for < 55, where -components dominate, than for > 55, where -components dominate. In the unsaturated regime, the ratio of the absorption coe cients for - and -components, = , reaches a minimum of 0.5 at = 90. For comparison, = reaches a value of 2 at 39 and grows in an unbounded manner as ! 0 (see x3 of E litzur). Unsaturated -components should on average be weaker than unsaturated -components, so the percentage of -components above a reasonable detection threshold would be higher than the 60% expected based on an analysis of the sizes of angular phase space alone. Second, as m asers saturate, com petitive gain will favor the stronger com ponent. Saturated -com ponents will reduce the absorption coe cient by a factor of 3, but saturated -components only reduce by a factor of 2 (Elitzur 1996). Thus, -com ponents should be even more num erous than -com ponents even if the masers are saturated.

However, it is clear that some $-components are seen in the^2_{3=2}; J = 3=2 m aser transitions. Some m aser spots exhibit a high degree of linear polarization as would be expected for <math>-components,$ and the distribution of PPA in some sources strongly suggests that -components are seen (see xA 13). Nevertheless, m aser spots that we believe are -components are not 100% linearly polarized. Since m f = 0 radiation is inherently linearly polarized, circular polarization must be generated externally. We especulate that these -components with nonzero circular polarization arise from the superposition of two m asing clumps along the line of sight. If the emission from a -component spot intersects a region of 0 H appropriately shifted in velocity, it may stimulate emission in a mode. Since the incident radiation from the rst cloud (i.e., the -component) is highly amplied and therefore bright, it can strongly stimulate the second cloud, since the incident linear polarization will be seen by the second cloud as a superposition of the two opposite-handed circular modes. Even if the amplication in the second cloud is very weak, a signi cant amount of circular polarization can be added, and the radiation will no longer be completely linearly polarized, as shown in Appendix B. The distinction between a -and a -component may be blurred if there is & 1 gain length of m aterial in the second cloud.

This circularization of a bright -component due to an extrem ely weak -component requires that the weak cloudlet be in front of the bright maxer from the observer's perspective. If the -component is behind the -component, the propagation path of radiation passes rst through the weak () cloudlet and then through the strong -component. The radiation eld that the cloudlet amplies, whether background continuum or

its own spontaneous emission, is much weaker than in the case where the radiation from the -component stimulates emission from the -component, so the superposition of spots would be indistinguishable from an isolated -component with no surrounding material.

In principle the reverse scenario could occur as well: a -component stimulates emission in the -m ode from a smaller cloud of 0 H gas at the appropriate velocity for amplication. This would have the e ect of adding extra linear polarization to a -component. Since -components are in general elliptically polarized (i.e., have a nonzero linear polarization fraction), it may not be possible to distinguish observationally between a -component that has stimulated weak emission in the -m ode from a second maser clump and one that has not. The linear polarization fraction of a -component is a function of the inclination of the magnetic eld to the line of sight (G oldreich, K eeley, & K wan 1973a), so this e ect could lead to overestim ation of the magnetic eld inclination at 0 H maser sites.

It is probable that the overlap ofm aser components along the line of sight would add circular polarization to -componentsmore system ically than it would add linear polarization to -components. Form ost common bulk material motions (e.g., infall, out ow, rotation), the radial component of the velocity eld will change monotonically along a ray from a maser spot to the observer. If the change in radial velocity along the line of sight exceeds the Zeem an splitting between a - and -component (1.2 km s^1 at 1665 MHz and 0.7 km s¹ at 1667 MHz for a 4 mG magnetic eld), the radiation from the -component may stimulate weak amplication from OH in a mode. But a -component could only stimulate emission from the mode of a cloud of OH along the line of sight if the change in radial velocity were in the same sense as the Zeem an splitting of the -component.

Given that only a small column density of OH along the line of sight between a maser and the observer is required to add signi cant circular polarization to the observed maser, it is likely that a large fraction of -components will be misidenti ed as -components. Unless there is an abrupt outer edge to the radial distribution of OH in a massive star-forming region, the radiation from many -components will stimulate weak emission in a -m ode of the surrounding OH.

4.5. E longated A rrangem ents of M aser Spots

In several sources, 0 H m assers are found in elongated law entary arrangements. For instance, there is a line of masser spots near the origin in the W 75 S m ap shown in Figure 25 of Paper I, and the massers in W 75 N (Figure 27 of Paper I) appear to be oriented primarily along two perpendicular axes. In the lam entary arrangement of masser spots seen in the northern grouping of W 75 N, the sky projection of the magnetic eld as deduced from the PPA of the linear portion of the polarization implies that the magnetic eld may be aligned predominantly along the line of elongation.

O fien there is a velocity gradient along the elongation, such as in W 75 S or the m asers in a NE/SW line in ON 2 N. These lines of m asers with velocity gradients are common in CH_3OH . Norris et al. (1993) observed 10 sources for which the m aser spots were distributed mostly in a line with the m ajor axis of the distribution several times greater than the m inor axis. Plots of the velocity of the maser spots versus the m ajor axis o set are generally distributed into two quadrants rather than tightly along a straight line (N orris et al. 1998). The authors speculated that the m asers are tracing circum stellar disks and that the deviation from a straight line in the velocity-m ajor axis plots is due to maser amplication at di erent radii in the same e circum stellar disk.

M ore recent observations do not seem to favor the interpretation of these m aser arrangements as circum stellar disks, however. First, D e Buizer (2003) boked for $H_2 = 1 - 0 S(1)$ emission in m assive SFRs for which circum stellar disks were suspected on the basis of collinear⁴ distributions of m ethanol m asers. Since m olecular hydrogen is a diagnostic of shocked out ows, it was expected that the H_2 emission would be oriented prim arily perpendicular to the putative disks. Instead, the H_2 emission in alm ost all of the sources for which it was detected was preferentially oriented parallel to the line of m ethanol m asers. D e Buizer suggested that the m asers were instead tracing an out ow . Second, the proper motions of m aser spots in the two linear structures in G 9.62+0.20 are directed prim arily perpendicular to the structures M inier et al. (2001). This suggests that in at least som e cases m ethanol m asers are shocks rather than circum stellar disks. W hether this applies to sim ilar arrangements of 0 H m asers is not yet well established. It is believed that the lifetim es of m ethanol and hydroxylm asers overlap but are not identical (R eid 2002). It is therefore possible that structures delineated by 0 H m asers trace a di erent evolutionary phase of form ing high-m ass stars than do C H₃O H m asers.

An alternate possibility to the disk hypothesis is that these elongated arrangements of maser spots may simply be a result of the motion of material threaded by magnetic elds. This could be a result of 1) collapse in the early stages of star formation or 2) shock-driven out ows in later stages. In case 1), as clouds of 0 H and other material fall inward, they will draw the magnetic eld inward with them. This will tend to align eld lines with the material elongations, provided there is enough angular momentum to avoid spherical collapse. In case 2), a shock propagating outward from the boundary of the UCH II region compresses material ahead of it, leading to elongations along the shock front.

The drift speed for am bipolar di usion of a magnetic eld out of a maser cloud is

$$v_{\rm d} = 0.6 \quad \frac{B}{5 \quad 10^{3} \text{ G}}^{2} \quad \frac{n}{10^{6} \text{ cm}^{3}}^{2} \quad \frac{r}{10^{15} \text{ cm}}^{1} \quad \frac{1}{2 \quad 10^{9} \text{ cm}^{3} \text{ s}^{1}}^{1} \quad \frac{x_{\rm e}}{10^{5}} \quad \text{km s}^{1}; \quad (7)$$

where r is the radius of a maser spot, is the ion-neutral collision rate coe cient, and x_e is the ionization fraction (B lack & Hartquist 1979). The drift speed is less than a typical shock speed (> 5 km s¹), especially if carbon is ionized in any substantial fraction. Thus, the magnetic eld will be dragged along and compressed by the shock, resulting in a eld oriented parallel to the shock front and therefore along the material elongation. Unless the shock is totally planar, there may be velocity gradients along the elongations. The slight curvature to some of these elongations may also be explained by the expansion of a spherical shock front or a planar shock into an inhom ogeneous medium. This is in contrast to models explaining elongated arrangements of maser spots as disks, where little curvature would be expected if preferentially seen edge-on. Nevertheless, some curvature along the elongations may be produced if a disk is inclined.

4.6. Is There a Connection Between M aser C lusters and Shocks?

A sm entioned in x3.2, there is a characteristic maser clumping scale of 10^{15} cm. Additionally, maser clusters tend to be concentrated on or near the periphery of H II regions. This is not always the case in complicated sources such as W 75 N, but more often than not these maser clusters appear near the boundary of an H II region, especially given that we observe three-dimensional distributions of masers in projection.

 $^{{}^{4}}W$ e will refer to structures of maser spots aligned along the same line as collinear to avoid any possible confusion with linear polarization.

O ccasionally even individual clusters are elongated in a lamentary manner, as in W 3(OH) (Reid et al. 1980).

It is possible that these clusters of OH m asers form in shocked neutral gas outside the ionization boundary. The initial growth phase of a UCH II region involves an R-type ionization front (see, e.g., K ahn 1954). When the speed of the ionization front slows to twice the sound speed of the ionized m aterial, a transition occurs and the ionization front changes to a weak D-type (Shu 1992). This is characterized by the existence of two separate fronts: an ionization (I) front and a shock (S) front that precedes it. K aw am ura & M asson (1998) directly measured the expansion speed of the UCH II region, i.e., the I front, in W 3(OH) to be 3 5 km s¹. Since the speed of the I front is less than the sound speed in the ionized m aterial (10 km s¹), the ionization front must be of D-type in W 3(OH).

O there have theorized that m asers near a UCH II region appear in the shocked neutralm aterial between the I and S fronts (e.g., Baldwin, Harris, & Ryle 1973; Elitzur & de Jong 1978). Theoretical calculations suggest that I-S fronts are inherently prone to instabilities (Vandervoort 1962). Less clear is the exact mechanism of instability growth, although there is no shortage of candidates (Dyson et al. 2002). Giuliani (1979) found that the slab of m aterial between the I and S fronts was unstable to oscillatory transverse perturbations. The wavelength of fastest perturbation growth was found to increase with time. V ishniac (1983) obtained similar results and additionally suggested that unstable sm all-scale fragmentation would eventually allow neutral gas to be swept up behind the fragm ents and lead to Rayleigh-Taylor instabilities, although it is unclear whether this would occur at the time and size scales of ultracom pact H II regions. Pottasch (1958) argued based on the evolutionary sequence of perturbation shapes of the bright rims in di use nebulae that the Rayleigh-Taylor instability alone cannot be the cause of ngering, although his analysis was based on older, lower-density H II regions. Vishniac also speculated that magnetic elds might produce elongated, lamentary structures. Two-dimensional simulations by Garc a-Segura & Franco (1996) demonstrate the appearance of this hydrodynam ical instability regardless of the density structure of the neutral gas and details of the radiative cooling law included. The wavelength of the fastest-growing perturbations increases with time, with interfragment spacings on the order of several times 100 AU near the base of the ngers and approximately 100 AU near the tips for UCH II ages near 10^4 years, as shown graphically in their Figure 6. Note that this is consistent with both the clustering scale and the dynam ical age of 0 H m asers, as discussed in xx32 and 3.6.

A ssum ing that some clusters near UCH II regions form in the shocked neutral medium between the I and S fronts, two projection-related factors would imply that clusters should be more frequently found at the periphery of H II regions rather than atop them, an e ect not seen (see x3.6). First, velocity gradients are likely to be higher along the ngers of shocked neutral material than across them . Amplication lengths should therefore be longer on average for ngers pointing in the plane of the sky, where the velocity gradient is tangential in projection, than for ngers pointing toward us (i.e., projected near the center of the H II region). Still, potential path lengths are longer along mgers rather than across them, and it is unclear whether velocity gradients are large enough to favor amplication across rather than along the ngers. Second, the magnetic eld threading the neutral material will be dragged along with it. Field lines will be folded such that a eld line entering the nger from the neutral material will exit the nger back into the neutral material after bending through 180. This would imply an elective reversal of the line-of-sight direction for lines of sight along ngers. How ever, intra-cluster magnetic eld line reversals are never seen. Indeed, the magnetic elds deduced from OH maser in clusters superposed atop H II regions in ON 1 and W 3(OH) suggest that there is a consistent line-of-sight eld direction in those sources even atop the H II region. It is possible that OH masing clumps occur near the \palm s" of the ngers or that the ngers of

neutralm aterial containing the OH m assers are not very long. In either case, the bend in the m agnetic eld lines could be much less than 180, consistent with our lack of detection of line-of-sight eld reversals in these clusters.

Large-scale (> 1000 AU) collinear m aser structures, such as the NE/SW line in ON 2 N, probably cannot be explained by shock instabilities around the UCH II region because they are larger in scale than the H II region itself. It is still possible that these structures occur in neutral gas that has been shocked by another source in the star-form ing com plex. There is often a velocity gradient along these structures, which could be explained by a curved or decelerating shock.

4.7. Relation of OH M asers to G alactic M agnetic Fields

Noting that line-of-sight directions obtained from Zeem an splitting in eight G alactic OH m aser sources were consistent with a clockwise G alactic eld, D avies (1974) postulated that OH m asers in m assive starform ing regions traced the G alactic m agnetic eld. Follow-up studies by Reid & Silverstein (1990) and Reid & M enten (1993) supported this claim but were suggestive of a more complicated G alactic eld structure. Subsequent analyses employing this technique on ever-larger sample sizes have indicated that correlations with the G alactic magnetic eld may exist (B audry et al. 1997; F ish et al. 2003), but detailed probing of the G alactic eld with thism ethod remains elusive. O bærvational limitations of thism ethod consist of unknown n distances to many of the sources as well as possibly incorrect magnetic eld data resulting from inadequate spatial resolution to unam biguously identify Zeem an pairs of maser features, since few sources have been observed at VLBI resolution. A dditionally, magnetic eld inform ation has thus far been limited to the sign of the line-of-sight eld orientation (i.e., whether the magnetic eld points in the hem isphere tow ard or away from the Sun). This may be insu cient to accurately probe a predom inantly toroidal G alactic magnetic eld at low er G alactic longitudes, in the direction of the magnetic of massive star-form ing regions (as well as most of the spiral structure of the G alaxy).

If the magnetic eld orientation in massive star-forming regions is correlated with the Galactic eld, the processes of high-m ass star form ation m ust not tightly wrap the magnetic eld con guration despite the rotation and collapse necessary to produce the central condensation, a proposition for which there is theoretical support (e.g., Li & Shu 1996; Allen, Li, & Shu 2003). As discussed in x3.5, an ordered m agnetic eld can be inferred from the regularity of the line-of-sight direction of the magnetic eld over large portions of the source and the eld strengths inferred from Zeem an pairs within the same clusters of 10¹⁵ cm. Furtherm ore, the magnetic elds deduced from OH Zeem an splitting in massive SFRs separated by distances on the order of a kiloparsec show a preference to be co-aligned (Fish et al. 2002). A num erical investigation of the collapse of rotating, magnetized, isotherm all cloud cores suggests that collapse can occur without introducing a signi cant twist to the magnetic eld (Allen, Li, & Shu 2003). These authors nd that the maximum pitch angle of the magnetic eld is approximately 20 along a ridge of accreting material (see their Figure 4). Inward from this ridge the magnetic eld resists wrapping, while outward from the ridge the wrap from di erential rotation is sm all. Simulations by M atsum oto & Tom isaka (2004) con m that the magnetic eld of a collapsing core maintains alignment with the magnetic eld of the parent cloud. They nd that a young star's magnetic eld is inclined no more than 30 from that of the parent cloud for a weak initial eld strength (20 G at a density of 2:6 10^4 cm³), with much better alignment when the initial eld strength is greater.

If these models are correct, the magnetic eld orientation before collapse might be partially preserved in

the material surrounding the core. Since the elds around new ly-form ed massive stars are ordered (see x3.5), this suggests that magnetic eld orientations deduced from OH maser Zeem an splitting may be indicative of the Galactic magnetic eld. A VLBI survey of OH masers in massive star-forming regions would eliminate Zeem an pairing ambiguity and possibly allow for three-dimensional modelling of the ambient magnetic eld in the few sources in which the Faraday rotation is small enough that the full magnetic eld orientation can be inferred from the observed linear polarization fraction and PPA of maser spots. If accurate distances can be obtained as well, as through trigonom etric parallaxes of higher frequency maser transitions, OH masers may prove to be a useful tool for probing the Galactic magnetic eld.

5. Sum m ary of Interstellar O H M aser P roperties

G round-state OH m assers typically cluster on a scale of 10^5 cm, providing evidence that their distribution is linked to a process with an inherent scale, as opposed to turbulence (which is generally scale-free). The magnetic eld strengths implied by Zeem an splitting suggest that OH m assers occur in regions of density 10^5 to several 10^7 cm 3 . OH m assers are found preferentially near the UCH II region in m assive SFRs. Their distribution around UCH II regions suggest an expansion age of 10^4 years for typical expansion velocities. OH m assers do not appear to be system ically shifted from the velocity of the associated star by m ore than a few km s¹, although possible exceptions exist, as in G 5.886 0.393 and W 75 N VLA 2. Taken together, these pieces of evidence support the theory that m ost OH m asers occur in the shocked neutral gas between the ionization and shock fronts of UCH II regions. The distribution of maser uxes with distance from the central UCH II region suggests that OH m asers turn o abruptly rather than weakening gradually after 10^4 years.

Som e O H m asers are seen far from or without any associated H II region. It is unclear whether these m asers are pumped by a starwith an associated weak, undetected hypercompact H II region or whether they are shock-excited without an ionization front. In som e sources (e.g., W 75 S), O H m asers appear to trace a collinear structure with a velocity gradient. These form ations probably trace shock fronts rather than protostellar disks.

M agnetic elds are ordered in massive SFRs, lending observational support to theories that indicate that the ambient magnetic eld direction may be preserved during massive star formation. Nearly all sources show either a consistent line-of-sight magnetic eld direction or a single reversal of the line-of-sight direction across the source. W ithin a maser cluster of size 10^{15} cm, line-of-sight magnetic eld direction reversals are never seen, and the eld strengths deduced from Zeem an splitting are alm ost always consistent within 1 mG.

We do see both $-and -components, including a \Zeem an triplet" in W 75 N (see x3.1). But OH maser$ spots that are 100% linearly polarized, as theoretically expected of <math>-components, are extrem ely rare. There is a range of sources with qualitatively di erent linear polarization properties. At one extrem e (as in W 75 N) high linear polarization fractions are seen, and the PPAs show some correlation with observed structures and probable magnetic eld directions. In most sources the linear polarization fractions are much less than 1 and PPAs cannot be easily interpreted as magnetic eld directions. At the other extrem e are sources such as W 51 el and e2, in which little or no linear polarization is detected and the total polarization fraction of som e maser spots is much less than unity.

The wide range of polarization properties observed in OH masers may be explained by a combination of Faraday rotation and overlap of maser components. If OH masers are indeed near or embedded in C II

regions, the electron density m ay be high enough that the m asers are near a critical point of Faraday rotation. A typical maser spot likely has large (> 1 rad) Faraday rotation over the entire ampli cation length, but not over a single gain length of the maser. If Faraday rotation is a factor of 5 lower, the total Faraday rotation along the ampli cation path m ay be small enough such that the PPAs are still roughly aligned with the magnetic eld lines. On the other hand, if Faraday rotation is a factor of 5 larger, the Faraday rotation per gain length could exceed 1 rad, destroying linear polarization and depolarizing the maser. Even if the Faraday rotation is small enough to allow ampli cation of a 100% linearly polarized -component, its polarization may be partially circularized by one of the -m odes of a weakly-inverted clump of OH between the maser site and the observer. This is likely a very important e ect, as only a modest inversion and a small column density of OH are required to add signi cant circular polarization to a -component.

Theoretically the linear polarization fractions and directions of maser components can be used to determ ine the full, three-dimensional orientation of the magnetic eld at masing sites. But the interpretation of PPAs may be very dicult in sources for which the amount of Faraday rotation along the propagation path between the source and the observer is unknown. Inferring a magnetic eld orientation in the plane of the sky also requires unambiguous identication of - and -components, due to the 90 di erence in PPA response to a magnetic eld. Zeem an pairs provide the surest method of identifying -components, but their polarization properties may be too contam inated by internal Faraday rotation and maser overlap to perm it interpretation of the inclination of the magnetic eld to the line of sight.

We thank M.D.Gray, M.Elitzur, and J.P.M acquart for helpful comments in preparation. Facility: VLBA

A. Notes on Individual Sources

A.1. G 5.886 0.393

There is a reversal of the line-of-sight direction of the magnetic eld across the source. All Zeem an pairs in the south of the source indicate a negative magnetic eld (i.e., oriented in the hem isphere pointing toward the Sun), while all pairs in the north of the source indicate a positive magnetic eld.

N early all of the m aser spots identi ed in the northeastern cluster constitute a component of a Zeem an pair. In total, there are eight Zeem an pairs in the cluster { four each in the 1665 and 1667 M H z transitions. The magnetic eld strengths are consistent, ranging from 1.2 to 2.0 m G in the cluster. The center (m aterial) velocities of the Zeem an pairs range from 8.6 to 10.0 km s¹. This is in excellent agreem ent with C aswell (2001), who nd a 6035 M H z Zeem an pair in this region centered at 9.96 km s¹ with a splitting of 1.49 m G.

We can de ne the Zeem an pairing e ciency as twice the number of Zeem an pairs divided by the total number of maser spots (in both polarizations and transitions) in a region. In the limiting case where every maser spot is a -component in a detectable Zeem an pair, the Zeem an pairing e ciency would be 100%. For the northeastern cluster (Figure 3 of Paper I), the Zeem an pairing e ciency is 84%. The pairing e ciency in the western half of the source is only 16%. We note that the velocities of maser spots range from 2.6 to 15.8 km s¹ in the western half of the source This is a much larger range than for the northeastern cluster,

eld also produces Zeem an pairs more e ciently.

Z ijlstra et al. (1990) observe O H em ission from 45 km s¹ to 17 km s¹ and interpret this em ission as tracing a bipolar out ow . O ur observations span only the upper end of this velocity range. W e see redshifted em ission extending to the southwest of the western group of masers, in general agreem ent with Z ijlstra et al. They do not see em ission at 8.6 to 10.0 km s¹ in the eastern half of the source, probably due to the large channel width (2.2 km s¹ velocity equivalent) and beam size of their observations. They also do not see em ission at the appropriate velocities corresponding to the isolated 2.4 and +1.2 m G Zeem an pairs we detect. At 6035 M H z, C aswell (2001) nds em ission in the velocity range corresponding to our observations and absorption from 25 to +2 km s¹, supporting the m odel of Z ijlstra et al.

Fekt et al. (2003) detect a candidate O -type ionizing star at $= 2:^{0.6}$ $0^{0.2}$; $= 0:^{0.0}$ $0:^{0.2}$ in Figure 1 of Paper I. The star's location is coincident with the isolated 1.2 m G Zeem an pair to within positional errors. The brighter RCP component of this Zeem an pair has a ux density of 2.37 Jy. Since the H II region is optically thick at = 18 cm (A erbach et al. 1996), it is unlikely that direct amplication of stellar radiation is in portant. Is the spatial coincidence of this maser spot with the projected location of the star due to chance, or is amplication favored due to the stellar radiation? N ine other main-line maser features in G 5.886 0.393 are brighter than this feature, indicating that the observed ux density is not strongly a ected by whether the maser is projected atop the star. How ever, the observed ux density of strong (> 1 Jy) m aser spots m ay be relatively insensitive to initial conditions if they are at least partially saturated. Due to interstellar scattering, it is possible to get only a low er lim it for maser brightness tem peratures, but it is probable that the brightest maser spots in this source are saturated (see x4.1).

A.2. G 9.622+0.195

The G 9.62+0.19 com plex contains several UCH II regions, as well as a hot m olecular core (C esaroni et al. 1994). We detect OH m asers around sources D, E, and G in the nom enclature system of G aray et al. (1993) and Testi et al. (2000). At source E, we nd two Zeem an pairs which both indicate a positive m agnetic eld. At source G, we nd two Zeem an pairs which both indicate a negative m agnetic eld. Few OH m aser spots and no Zeem an pairs are identi ed at source D, while no m aser activity at all is seen near the hot core (source F), located between sources D and G.

W e do not see the isolated 1665 M H z R C P m aser detected between sources E and G in Argon, Reid, & M enten (2000). This maser spot has likely weakened below our detectability threshold in the 10 years since their observations. W ater maser emission is seen at this site and associated with sources D, E, and G (H ofner & Churchwell 1996).

A.3. G10.624 0.385

O verall evidence indicates that G 10.624 0.385 is undergoing collapse. O bservations of N H₃ show that the molecular material is rotating and falling inward, with rapid spiral motions inward of about 0.05 pc (Ho & Haschick 1986; Keto, Ho, & Haschick 1987). The plane of rotation cuts through the center of the H II region and is oriented approximately 20 west of north (Keto, Ho, & Haschick 1988). Coherent inward

motions are also seen within the ionized gas in the H II region (K eto 2002).

We detect relatively few maser spots in G10.624 0.385. Em ission falls into three regions: a clump to the east, a clump 2^{00} to the west of the previous clump, and an isolated maser spot to the northwest. We nd only one Zeem an pair, indicating a magnetic eld of 6 m G in the easternm ost clump. The west clump appears arclike, with a length of about 150 mas. This is the only clump to show any linear polarization, most of which is oriented roughly perpendicular to the arc. None of the maser spots we detect lies along the plane of rotation.

The ground-state m asers we detect span a velocity range of 2:4 to + 3:3 km s¹. This is in agreement with the velocity span of excited states of OH: 2:0 to 1:5 km s¹ in $_{1=2}^{2}$; J = 1=2 em ission (G ardner & M art n-P intado 1983), 2:5 to +1:0 km s¹ in $_{3=2}^{2}$; J = 7=2 absorption (F ish, R eid, & M enten 2005), and 1:9 to 0:4 km s¹ in $_{3=2}^{2}$; J = 9=2 absorption (W alm sley et al. 1986).

A.4. G 34.257+0.154

The G 34.3+0.2 com plex contains several UCH II regions. Most prominent is the \com etary" H II region, labelled C in the nom enclature of G aum e, Fey, & C laussen (1994). The morphology of region C can be explained as a bow shock due to the supersonic relative motion of the exciting source with respect to the surrounding medium (Reid & Ho 1985; M ac Low et al. 1991). Region C may be composed of more than one continuum component (Sew ib et al. 2004). Two fainter com ponents, labelled A and B by Reid & Ho (1985), are located to the southeast and northeast of region C, respectively. The com plex has a kinem atic distance of 3.8 kpc (G alactic center distance $r_0 = 10$ kpc) (Reifenstein et al. 1970).

A blueshifted out ow extending to the northwest is seen by Hatchell, Fuller, & M illar (2001). Com ponents A and C were detected in the mid-infrared, but component B was not, suggesting that it is deeply embedded and therefore very young (Campbell et al. 2000). Based on this and the spectral index of com – ponent B (0:9 0:4, from G aum e, Fey, & C laussen 1994), Hatchell et al. conclude that component B is the source of the out ow.

Unlike other sources, there is not a single line that can be drawn across the entirety of G 34.257+0.154 which separates regions of positive and negative magnetic eld. However, such lines can be drawn for the masers associated with regions B and C separately. Region B contains many Zeem an pairs with positive magnetic eld near the continuum source and a single Zeem an pair with negative magnetic eld to the northeast of the source. All of the Zeem an pairs ahead of the bow shock in region C imply a negative magnetic eld except for a small region near the H II region to the south. There is good qualitative agreem ent between the magnetic eld measured here and similar VLBA observations of the source in 1995 by Zheng, Reid, & M oran (2000). In the northerm half of the source, the eld directions are identical between the two observations, and eld magnitudes agree to better than 0.3 m G where they overlap. In the center and south, we nd larger eld magnitudes than reported by Zheng et al. Line-of-sight eld directions are again in agreem ent, although Zheng et al. nd Zeem an pairs in plying magnetic elds of + 0.5 and 0.5 m G near the origin of F igure 9 of P aper I. At this same location, G asiprong, C ohen, & Hutawarakom (2002) nd only one Zeem an pair im plying a magnetic eld of 5.0 m G. In this region, we infer magnetic elds of 0.6,

5:1, 5:7, and 6:0 m G from four Zeem an pairs. At 6035 M H z, C aswell & Vaile (1995) and C aswell (2001) nd three Zeem an pairs indicating a magnetic eld of 4 m G. Their observations do not have the necessary angular resolution to determ ine which H II region the 6035 M H z are associated with, but their results are largely consistent with magnetic elds obtained from the com etary region C, especially in the north.

G asiprong, C ohen, & Hutawarakom (2002) also observed linear polarization with the M ERLIN array. It is di cult to compare our results directly with theirs, since the $0.0^{10}16$ resolution a orded by M ERLIN is insu cient to separate distinct m aser components with very di erent linear polarization position angles. Nevertheless, for the brighter spots it is possible to identify a m aser spot in our data that corresponds to a sim ilar spot in the G asiprong study. Polarization position angles generally agree to 10 or 20.

A.5. G35.577 0.029

This source contains two UCH II regions. All of the maser activity appears to be associated with the western H II region. We detect maser spots only on the western limb of the western H II region with the exception of one isolated spot to the east. Due to registration uncertainties, it is not clear whether this spot is located directly atop the H II region or on the eastern limb. We detect three Zeem an pairs in the western clump of emission, with all three implying a magnetic eld in the range of 4 to 6.3 mG.

A.6. G40.622 0.137

There is a large ($1^{(0)}$) cluster of OH m aser em ission centered approximately 1.5 away from the only detected H II region. We detect two Zeem an pairs in the same region, consistent with a eld strength of approximately 6 mG. Caswell & Vaile (1995) nd one Zeem an pair in 6035 MHz OH maser em ission implying a magnetic eld of + 1:7 mG, suggesting that there may be a reversal of the line-of-sight direction of the magnetic eld in this source. Methanol and water masers are also seen within less than $1^{(0)}$ of the reference position (Forster & Caswell 1989; Beuther et al. 2002).

A.7. G43.796 0.127

The X-band continuum maps show two sources { a bright source to the northwest, and a weak source to the southeast. All maser emission lies atop the northwest source. A total of seven Zeem an pairs were identied. Five of these indicate a positive magnetic eld, and two indicate a negative magnetic eld. Unlike in other sources in which a reversal is seen, it is not possible to draw a single straight line such that the magnetic eld on each side of the line has a uniform line-of-sight direction. A Zeem an measurement of 6035 MHz (excited-state) OH emission implies a magnetic eld of + 3:6 mG (Caswell & Vaile 1995). Three measurements of the Zeem an eld of H_2 masers, which trace a higher range of densities, imply a magnetic eld of 13:3 to 46:1 mG (Sarma et al. 2002).

We have adopted an ammonia velocity of 452 km s¹ for G 43.796 0.127. The NH₃ spectrum itself (see Figure 41 of Paper I) is complicated, and it is di cult to identify which line is the main line and which are hyper ne lines. We identify the line at 452 km s¹ as the main line because its velocity most closely matches that of the CS J = 7! 6 velocity of 44.3 km s¹ (Plume, Ja e, & Evans 1992). Since the critical density of CS J = 7! 6 is 2 10^7 cm³, we feel con dent that it is tracing the same high-density material as the NH₃ emission.

A.8. W 51

We nd a total of 46 Zeem an pairs near sources el and e2, making W 51 the most prolic massive SFR in terms of the number of Zeem an pairs in our survey. As previously reported (Argon, Reid, & Menten 2002), source e2 contains two Zeem an pairs implying the strongest magnetic elds ever seen in interstellar OH massers: 19.8 mG and 21 mG.

Source el show sa reversal of the line-of-sight eld direction, which points tow and the Sun in the northern half of the source and away from the Sun in the southern half. Source e2 is the clearest example yet that shows the extent to which magnetic elds are ordered in massive SFRs. All 22 Zeem an pairs indicate a positive magnetic eld. Although there is a huge variation in the strength of the magnetic eld across the source, multiple Zeem an pairs in each cluster have consistent eld strengths to within about 1 m G.

W 51 is remarkable among our source sample as having almost no detectable linear polarization. Three maser spots near the origin in source el have linear polarization fractions of 1% to 2%. No linear polarization was detected for any other spot in source el or any spot at all in source e2. The plausibility of circularization due to high Faraday rotation along maser amplication paths is discussed in x42.

A.9. ON 1

The maser spots appeared to be grouped into three regions. The masers in the northern group have velocities near 4 km s¹. The masers in the central group are located at about 13 14 km s¹. The southern maser spots fall primarily along an extended collinear feature. The velocities in this line range from 13 to 15 km s¹. OH masers are not seen at intermediate velocities. Methanol masers show a similar velocity structure (Szym czak, Hrynek, & Kus 2000).

Zeem an splitting in ON 1 is everywhere consistent with a magnetic eld pointing in the hem isphere toward the Sun. We do not nd any unambiguous Zeem an pairs in the northern group. At 6031 and 6035 MHz, Desmurs & Baudry (1998) nd four Zeem an pairs im plying magnetic elds from 3.6 to 6.3 mG. These maser spots appear to fall slightly north and west of the northern group of maser spots detected in the ground-state transitions here and in Argon, Reid, & Menten (2000), but their center velocities fall between 13.7 and 15.3 km s¹ as compared with a velocity range of 3.2 to 6.2 km s¹ in the northern ground-state group. It is unclear whether this 10 km s¹ di erence re ects a large velocity gradient in the northern part of the source or whether the 6031 and 6035 MHz em ission com es from a di erent area, re ecting registration uncertainties between the various sets of observations. Two 13441 MHz Zeem an pairs at 14.1 and 0.3 km s¹ indicate magnetic elds of 3.8 and 8.3 mG, respectively (Fish, Reid, & Menten 2005).

Since registration uncertainties of a few tenths of an arcsecond m ay exist between the continuum in age and our maser spot m aps, one possible interpretation is that the northern and southern maser groups are located on the limb of the UCH II region, while the center group is projected onto the H II region. W hile the precise locations of these groups relative to the H II region along the line of sight is unknown, the center group must be located in front of the H II region, since the H II region is optically thick at 18 cm (Zheng et al. 1985). The authors also noted an arcm inute-scale gradient of 11 2 km s¹ pc¹ in NH₃ em ission with a velocity of 12 km s¹ near the H II region (com parable to that of the southern group of 0 H masers) and an H 76 recombination velocity of 5:1 2:5 km s¹, blueshifted with respect to the NH₃ em ission. They concluded that the motions in ON 1 were consistent with infall and rotation.

The velocities of the three groups of maser spots we observe may be consistent with infall and rotation

on a smaller angular scale as well. If the north and south maser groups are at limbs of the rotation, the implied rotation speed would be 5 km s¹ centered at 9 km s¹ and roughly aligned with the direction of rotation noted by Zheng et al. The center maser group might then be infalling at 5 km s¹ as well. At this radius, 5 km s¹ corresponds to the freefall velocity for a 20 M star, so if net rotation is also sustained, rapidly spiralling infall must be occurring. Nevertheless, this could explain why the RCP maser emission in the northerm group is seen farther from the center of the H II region than the LCP emission. Because the magnetic eld splits the RCP emission to a lower LSR velocity than the LCP emission, the coherent path length is larger farther away from the center of the H II region.

A.10. K3 50

This H II region has a diam eter of over 0.1 pc, which is large for an ultracom pact H II region. M asers are found only to the north and east of the H II region. The line-of-sight magnetic eld direction points toward the Sun at all maser groups. However, due to the lack of maser emission to the south and west of the H II region, we cannot conclusively rule out a magnetic eld reversal across K 3 50. U sing the E elsberg 100 m telescope, Baudry et al. (1997) nd two 6035 M H z Zeem an pairs indicating eld strengths of 53 and 9.1 m G centered at 18:68 and 19:44 km s¹, respectively. These are redshifted compared to the Zeem an pairs we identify, whose center velocities range from 22:30 to 19:79 km s¹. Two components at 20:1 to 20:2 km s¹ and 25:0 to 25:5 km s¹ are seen in absorption in the 2 $_{3=2}$; J = 7=2 lines (F ish, Reid, & M enten 2005).

A.11. ON 2N

Eleven Zeem an pairs have been identied in ON 2 N, all indicating a positive magnetic eld. All of the OH maser emission is located to the south and west of the UCH II region, in the same area as the H_2O maser emission (Hofner & Churchwell 1996).

There are three groups of m aser spots arranged roughly in a line with a position angle 35 east of north beginning at the H II region. This line of spots exhibits a velocity gradient with the most redshifted emission toward the southwest. A cluster of m aser spots with a large velocity dispersion is o set from this line and elongated perpendicular to it. A lm ost all spots with any detectable linear polarization, as well as all spots with a high linear polarization fraction, are o set from the line of m asers.

A.12. W 75 S

We have identi ed 13 Zeem an pairs around the UCH II region in W 75 S. There is a reversal of the line-of-sight magnetic eld direction across this SFR. Seven Zeem an pairs to the east indicate a magnetic eld pointing toward the Sun, while six to the west indicate a magnetic eld in the opposite direction. A dditionally, eld strengths within maser clusters are remarkably consistent. The six Zeem an pairs in the western cluster imply eld strengths of 5:6 to 6:6 m G, and the six Zeem an pairs to the southeast of the H II region im ply eld strengths of 3:8 to 5:3 m G.

A collinear arrangement of maser spots exists near the origin in Figure 25 of Paper I. An enlargement of this region is shown in Figure 20, along with a best-t line. There is a velocity gradient along this line

with the velocity increasing to the north. If these massers are interpreted as tracing a circum stellar disk in K eplerian rotation, the radius of the disk is at least 400 AU, and the central mass is at least 6 M \therefore See x4.5 for further discussion of the possible interpretations of this structure.

A.13. W 75 N

The region of 0 H m asing in W 75 N coincides with three continuum sources, identi ed from north to south as VLA 1, VLA 2, and VLA 3 by Torrelles et al. (1997). Based on an elongation of VLA 1 (at position angle 43) and the spectral index of the source, the authors conclude that there is an ionized, biconical, partially optically thick jet em anating from the source. In addition to this ow, there is a larger, 3-pc ow oriented at position angle 62:5 that does not appear to be driven by the out ow in VLA 1 (Shepherd, Testi, & Stark 2003).

W 75 N contains 0 H m aser spots distributed prim arily along two axes. A long the north-south axis the m agnetic eld is oriented away from the Sun, while along the east-west axis the eld is oriented toward the Sun. At the intersection of these two axes there are two Zeem an pairs, each indicating a di erent sign of the m agnetic eld. It appears that m ost m asers are associated with VLA 1. How ever, the dynam ics of the cluster at the limb of VLA 2 (F igure 28 of Paper I) are unlike the rest of the m asers, suggesting that these m asers are indeed associated with VLA 2.

O ther maser species exist in W 75 N as well. H_2O masers are seen close to the H II regions, with spots distinctly on the limbs of VLA 2 and VLA 3 as well as along the position angle of the jet in VLA 1 (Torrelles et al. 1997). CH_3OH masers are seen primarily as an extension to the north of the north-south axis of OH masers, and they are distributed in a line with position angle 42 (M inier, Conway, & Booth 2001). Thus, the CH_3OH masers appear to be associated with VLA 1.

The region around VLA 2 contains a large number of OH maser spots at a wide range of velocities. The di erence in velocity between the most blueshifted and most redshifted spot in this region is 34 km s¹ (E llder 1973; Hutawarakom, Cohen, & Brebner 2002). Our bandwidth only covered about 21 km s¹ of this range. We detect maser emission in both the highest and lowest usable velocity channel at which we observed. A large velocity dispersion in H₂O masers is also seen in this region (Torrelles et al. 2003). Based on these large velocity dispersions, the location of VLA 2, and the steeply rising spectrum of continuum emission, Hutawarakom, Cohen, & Brebner (2002) conclude that VLA 2 is the source of the large-scale molecular out ow. Slysh, Val'tts, & Migenes (2001) interpret the maser spots in the north-south axis as being a disk centered at VLA 1.

Since the observations detailed in Paper I, a 1665 M H z m aser in W 75 N has become the strongest ever detected, reaching a ux of approximately 1 kJy (A lakoz et al. 2005). They nd a 750 Jy RCP feature at 1.8 km s¹, as well as two other new features near 0 and 1 km s¹. All three features are predom inantly linearly polarized and are therefore likely -components or -components where the magnetic eld is oriented close to the plane of the sky. These are o set by about 0.5 km s¹ compared to the bright features in Paper I. It is possible that these features are new or that the masers near VLA 2 are accelerating. This latter possibility cannot be ruled out because of the nature of the masers near VLA 2. They are observed to span a velocity range 34 km s¹ wide and appear to be associated with an out ow (E llder 1973; H utaw arakom, C ohen, & B rebner 2002). If the out ow is decelerating, it is possible that masers entrained in the ow may appear at slightly di erent velocities between epochs.

A.14. Cep A

Cep A is a complex molecular cloud condensation. Hughes & Wouterloot (1984) detected no fewer than 14 H II regions in the complex, and subsequent observations have uncovered even more radio continuum sources (Hughes 1988; Curiel et al. 2002). HW 2, the brightest continuum source in Figure 30 of Paper I, appears to contain at least four compact sources (Hughes, Cohen, & Garrington 1995) and is believed to be the source of two therm al jets (Rodr guez et al. 1994; Hughes 2001). The 6 cm radio jet is oriented at a position angle of 44 and is observed to have a projected velocity of 950 150 km s¹ (Rodr guez et al. 2001). Water masers are seen associated with this jet (Torrelles et al. 1996), and their proper motions suggest the presence of at least three distinct sites of star form ation within a projected 200 AU radius (Torrelles et al. 2001). To the south in Figure 30 of Paper I are three continuum sources: HW 3c, HW 3div, and HW 3dii in the nom enclature of Hughes, Cohen, & Garrington (1995) and Torrelles et al. (1998). Source HW 3c show s evidence of multiple components (Hughes, Cohen, & Garrington 1995). Water masers are seen around HW 3dii and the nearby source HW 3di, which is not detectable in our X-band in age (Torrelles et al. 1998). We detect O H masers around HW 2, HW 3c, and HW 3div, as well as a cluster of masers between HW 3div and HW 3dii and a lone maser not near any continuum source.

Bartkiew icz et al. (2005) identify seven Zeem an pairs at 1665 M H z and two at 1667 M H z w ith M ERLIN. Four of these (Z_2 , both Z_7 's, and Z_8) agree w ith our notings in terms of central velocity and magnetic eld strength to w ithin the errors expected from velocity resolution. Pairs Z_5 and Z_6 also agree w ith the magnetic eld strength we nd in the respective maser clusters, although several maser spots are blended together at these locations in the MERLIN beam. We do not nd counterparts for pairs Z_1 and Z_3 , and we do not have the velocity coverage necessary to observe Z_4 .

A.15. NGC 7538

NGC 7538 is a complex star-forming region. The continuum source in Figure 32 of Paper I, known as IRS 1, contains a core of two compact components and a larger, spherical region to the south (Turner & Matthews 1984; Campbell 1984). Scoville et al. (1986) argued for the existence of an ionized stellar wind out ow based on the spectral index of millim eter continuum emission and for a possible disk oriented east-west based on 13 CO emission. G aum e et al. (1995) nd that the distribution of continuum emission is clum py and suggest that photoionization from the central star is responsible for this emission. There is a bipolar, high-velocity CO ow around IRS 1 (Fischer et al. 1985), possibly collim ated by a denser ring of material seen in CS (K awabe et al. 1992). This out ow, as well as other out ows and stellar winds in the IRS 1-3 region, may be driving the expansion of a molecular half-shell (X u, Zheng, & Jiang 2003).

NGC 7538 is also rem arkable in the variety of maser species detected. Among these are rare maser species such as H_2CO (Forster et al. 1980), $^{14}NH_3$ (M adden et al. 1986), and $^{15}NH_3$ (M auersberger, W ilson, & Henkel 1986). H_2O masers (Genzel & Downes 1976; K ameya et al. 1990) and CH_3OH masers from a variety of transitions (e.g., W ilson et al. 1984, 1985; Batrla et al. 1987) are seen in NGC 7538 as well. OH maser emission has been seen in the 1665, 1667, and 1720 M Hz² $_{3=2}$; J = 3=2 lines (e.g., Downes 1970; D ickelet al. 1982; Hutawarakom & Cohen 2003), the 6035 M Hz² $_{3=2}$; J = 5=2 line (G uilloteau et al. 1984), and the 4765 M Hz² $_{1=2}$; J = 1=2 line (Palmer, G archner, & W hiteoak 1984). Hutawarakom & Cohen nd one Zeem an pair in each of the 1667 and 1720 M Hz transitions im plying magnetic elds of 1:7 and 2:0 m G, respectively. W e nd no Zeem an pairs at 1667 M Hz, but we do detect a Zeem an pair of + 0:7 m G

at 1665 M H z, suggesting that there is a reversal of the line-of-sight direction of the m agnetic eld across the

source.

A.16. S269

We have found three Zeem an pairs in S269, consistent with a magnetic eld of 4:0 to 4:5 mG. O therwise, S269 is one of the simplest sources in our study. It exhibits few maser spots. There is not much linear polarization of maser emission in this source. The VLA survey was unable to detect any continuum emission, nor was ammonia emission found. And the range of velocities of maser emission is a mere 3:9 km s¹.

Them agnetic eld appears to be oriented toward the Sun everywhere across the region of maser emission. AllRCP emission, excluding weak features associated with the linearly-polarized component of two strong left-elliptically polarized masers, occurs at lower velocity than the LCP emission. Figure 21 shows the spectrum of the 1665 MHz emission. When the spectra are corrected for Zeem an splitting (\dem agnetized") for a 4.0 mG magnetic eld, the total velocity range spanned by the RCP and LCP features decreases from 3.9 km s^1 to 1.5 km s^1 .

S269 also exhibits a high degree of variability. C legg (1993) notes a 1665 M H z m aser of 16 Jy LCP ux at 17.9 km s¹ in 1991.5, dim inishing to 1 Jy by 1992.1. In 1991.6, A rgon, Reid, & Menten (2000) nd that this maser has a ux of 7.5 Jy LCP. The closest feature we nd to matching this is a 1665 M H z m aser of 0.11 Jy LCP ux at 17.8 km s¹. We detect this feature in the 17.76 and 17.93 km s¹ velocity channels, but the lack of detection in a third channel prevents us from being able to accurately determ ine the velocity and linew idth of this feature.

A.17. M on R 2

The M on R2 m olecular cloud is one of the nearest high-m ass star-form ing regions, but it contains alm ost no stars of spectral type earlier than B1 (Hughes & Baines 1985). An unusual property of M on R2 is that m aser em ission from the 4765 M Hz ($_{1=2}^{2}$; J = 1=2; F = 1 ! 0) transition of O H is stronger than from the ground-state ($_{3=2}^{2}$; J = 3=2) set of transitions, suggesting that the physical conditions are denser and hotter than norm ally seen for ground-state O H m asers, as noted by Sm its, C ohen, & H utaw arakom (1998). They detect two m asers at 10.65 km s¹ with linear polarization position angles of 13 and 14. This is in excellent agreem ent with the brightest m aser feature we detect, coincident with the brightest 4765 M H z m aser to within registration uncertainties, which has linear polarization with a position angle of 13. The velocity of this 1665 M Hz m aser feature is 10.29 km s¹ when corrected for the Zeem an splitting of the 2:6 m G m agnetic eld at the site. The 4765 M Hz m asers in M on R2 are highly variable, doubling in strength in less than 19 days and reaching a peak of nearly 80 Jy before \disappearing" (Sm its 2003). Sm its nds that 1665 and 1667 M Hz em ission is much less variable, varying relatively sm oothly with changes in ux density not exceeding a factor of two over a tim escale of m ore than four years.

A.18. G351.775 0.538

G 351.775 0.538 contains what was the strongest known interstellar OH maser spot (400 Jy in LCP) until the recent are in W 75 N (A lakoz et al. 2005). Caswell & Haynes (1980) rst noted that the brightness

of this maser spot is highly variable, and it has been monitored frequently since then (see MacLeod & Gaylard 1996). Ground-state masers have previously been seen at velocities as low as 27.8 km s^1 (Argon, Reid, & Menten 2000) and as high as 7 km s^1 (MacLeod & Gaylard 1996). Our VLBA observations covered only the top half of this range.

There is a reversal of the line-of-sight magnetic eld direction across the source, as has been previously noted at 1665 and 1667 M H z by Argon, R eid, & M enten (2002) and F ish et al. (2002). This reversal is seen at 1720 M H z as well, where C aswell (2004) nds magnetic elds of + 3 and 6 m G.C aswell & Vaile (1995) nd a 3:3 m G eld at 6035 M H z.

Because G 351.775 0.538 is a low Declination source, (u;v)-coverage is poor, especially along northsouth baselines. The synthesized beam and spot sizes as listed in Table 20 of Paper I are thus very large. This may explain the separations of Zeem an components for G 351.775 0.538 (see Table 21 of Paper I), which are larger than for other sources.

B. Maser Overlap Polarization Calculation

In this appendix we consider the polarization properties expected of a -component that stimulates weak amplication from a -mode of a clump of OH between the rst maser and the observer, as described in x4.4. It is helpful to analyze the radiation in terms of the Stokes parameters, which are de ned in terms of the electric elds in the radiation as follows:

$$I = \langle x_{x} \rangle + \langle y_{y} \rangle$$

$$Q = \langle x_{x} \rangle + \langle y_{y} \rangle$$

$$U = \langle x_{y} \rangle + \langle x_{y} \rangle$$

$$V = i(\langle x_{y} \rangle + \langle x_{y} \rangle)$$
(B1)

They can also be written in terms of the electric elds in the two senses of circular polarization:

$$I = \frac{1}{2} (I_{rr} + I_{ll})$$

$$Q = \frac{1}{2} (I_{rl} + I_{lr})$$

$$U = \frac{1}{2} (I_{rl} \ I_{lr})$$

$$V = \frac{1}{2} (I_{rr} \ I_{ll});$$
(B2)

where $I_{rr} = \langle r_r \rangle$, etc.

Consider a simple case of weak maser amplication in the absence of Faraday rotation. We will start with radiation that is 100% linearly polarized in the x-direction, as could be produced by a -component. The Stokes parameters of the radiation are

$$I = I_0; Q = I_0; U = 0; V = 0;$$
 (B3)

so the polarization fractions (m $_{\rm L}$ linear, m $_{\rm C}$ circular, and m $_{\rm T}$ total) are

$$m_{\rm L} = 1; m_{\rm C} = 0; m_{\rm T} = 1;$$
 (B4)

Now suppose that this radiation is fed into a second, weak maser spot shifted in velocity such that emission is stimulated in the RCP -mode. The amplication factor is such that the ux density in the RCP mode is multiplied by a factor of 2. Then

$$I_{rr} \quad ! \quad 2I_{rr} = 2I_{0}$$

$$I_{11} \quad ! \quad I_{11} = I_{0}$$

$$I_{r1} \quad ! \quad \frac{p}{2}I_{0}$$

$$I_{1r} \quad ! \quad \frac{p}{2}I_{0}; \quad (B5)$$

where the factors of p = 1 are due to the fact that if the ux density increases by a factor of 2, the electric eld am plitude increases by a factor of p = 1. Substituting these values into equation (B2) results in the following:

$$I = \frac{1}{2}(2I_{0} + I_{0}) = \frac{3}{2}I_{0}$$

$$Q = \frac{1}{2}(\stackrel{p}{2}I_{0} + \stackrel{p}{2}I_{0}) = \stackrel{p}{2}I_{0}$$

$$U = \frac{1}{2}(\stackrel{p}{2}I_{0} \stackrel{p}{2}I_{0}) = 0$$

$$V = \frac{1}{2}(2I_{0} \quad I_{0}) = \frac{1}{2}I_{0}:$$
(B6)

Converting equation (B6) to polarization fractions, we obtain

$$m_{\rm L} = \frac{p \frac{Q^2 + U^2}{I}}{I} = \frac{2^{p} \frac{Z}{2}}{3} \quad 0.943$$

$$m_{\rm C} = \frac{V}{I} = \frac{1}{3} \quad 0.333$$

$$m_{\rm T} = \frac{p \frac{Q^2 + U^2 + V^2}{Q^2 + U^2 + V^2}}{I} = \frac{q}{m_{\rm L}^2 + m_{\rm C}^2} = 1.000: \quad (B7)$$

The net e ect is that the maser is still 100% polarized, but the linear polarization fraction has dropped to less than unity and the circular polarization fraction is reasonably large. For amplication of a single circular mode by a factor of n, the linear polarization fraction is

$$m_{\rm L} = \frac{2^{\rm p} \bar{n}}{1+n} :$$
 (B8)

The circular and linear polarization fractions become equal at n 6, or approximately 1.8 e-fold amplication lengths for an unsaturated maser.

REFERENCES

- A erbach, A., Churchwell, E., Acord, J.M., Hofner, P., Kurtz, S., & DePree, C.G. 1996, ApJS, 106, 423
- A lakoz, A.V., Slysh, V.I., Popov, M.V., & Val'tts, I.E. 2005, Ap. Lett., 31, 375
- Allen, A., Li, Z.-Y., & Shu, F.H. 2003, ApJ, 599, 363
- Argon, A.L., Reid, M.J., & Menten, K.M. 2000, ApJS, 129, 159
- Argon, A.L., Reid, M.J., & Menten, K.M. 2002, in IAU Symposium 206, Cosm ic MASERs: From Protostars to Blackholes, ed.V.Migenes and M.J.Reid (San Francisco: ASP), 367

- Baart, E.E., & Cohen, R.J. 1985, MNRAS, 213, 641
- Baldiyn, J.E., Harris, L.S., & Ryle, M. 1973, Nature, 241, 38
- Bartkiew icz, A., Szym czak, M., Cohen, R.J., & Richards, A.M. S. 2005, MNRAS, 361, 623
- Batrla, W., Matthews, H.E., Menten, K.M., & Walmsley, C.M. 1987, Nature, 326, 49
- Baudry, A., Desmurs, J.F., Wilson, T.L., & Cohen, R.J. 1997, A&A, 322, 255
- Berulis, I.I., & Ershov, A.A. 1983, Soviet Ast. Lett., 9, 341
- Beuther, H., Walsh, A., Schilke, P., Sridharan, T.K., Menten, K.M., & Wyrowski, F. 2002, A&A, 390, 289
- Black, J.H., & Hartquist, T.W. 1979, ApJ, 232, L179
- Bloem hof, E.E., Reid, M.J., & Moran, J.M. 1992, ApJ, 397, 500
- Boyd, R.W., & Wemer, M.W. 1972, ApJ, 174, L137
- Cam pbell, B. 1984, ApJ, 282, L27
- Campbell, M.F., Garland, C.A., Deutsch, L.K., Hora, J.L, Fazio, G.G., Dayal, A., & Homan, W.F. 2000, ApJ, 536, 816
- Caswell, J.L. 2001, MNRAS, 326, 805
- Caswell, J.L. 2004, MNRAS, 349, 99
- Caswell, J.L., & Haynes, R.F. 1980, IAU Circ. 3509
- Caswell, J.L., & Vaile, R.A. 1995, MNRAS, 273, 328
- Cesaroni, R., Churchwell, E., Hofner, P., Walm sley, C.M., & Kurtz, S. 1994, A&A, 288, 903
- Clegg, A.W. 1993, Lecture Notes in Physics, Berlin Springer Verlag, 412, 279
- Cook, A.H. 1966, Nature, 211, 503
- Cragg, D.M., Sobolev, A.M., & Godfrey, P.D. 2002, MNRAS, 331, 521
- Crutcher, R.M. 1999, ApJ, 520, 706
- Curiel, S., et al. 2002, ApJ, 564, L35
- Davies, R.D. 1974, in IAU Symposium 60, Galactic Radio Astronomy, ed.F.J.Kerr and S.C.Simonson III (DordrechtReidel), p.275
- DeBuizer, J.M. 2003, MNRAS, 341, 277
- Deguchi, S., & Watson, W.D. 1986, ApJ, 300, L15
- Desmurs, J.F., & Baudry, A. 1998, A&A, 340, 521
- Dickel, H.R., Rots, A.H., Goss, W.M., & Forster, J.R. 1982, MNRAS, 198, 265
- Dieter, N.H., W eaver, H., & W illiam s, D.R.W . 1966, S&T, 31, 132
- Downes, D. 1970, Astrophys. Lett., 5, 53
- D raine, B.T. 1980, ApJ, 241, 1021
- Dyson, J.E., W illiam s, R.J.R., Hartquist, T.W. & Pavlakis, K.G. 2002, Rev.M exicana Astron. Astro s.(SC), 12, 8
- Elitzur, M. 1996, ApJ, 457, 415
- Elitzur, M., & de Jong, T. 1978, A & A, 67, 323
- Ellder, J. 1973, Research Report No. 116, Research Lab. of Electronics and Onsala Space Obs.
- Feldt, M ., et al. 2003, ApJ, 599, L91
- Field, D., & Gray, M.D. 1994, A&A, 292, 271
- Fischer, J., Sanders, D.B., Simon, M., & Solomon, P.M. 1985, ApJ, 293, 508

- Fish, V.L., Reid, M.J., Argon, A.L., & Menten, K.M. 2002, in IAU Symposium 206, Cosmic MASERs: From Protostars to Blackholes, ed.V.Migenes and M.J.Reid (San Francisco: ASP), 371
- Fish, V.L., Reid, M.J., Argon, A.L., & Menten, K.M. 2003, ApJ, 596, 328
- Fish, V.L., Reid, M.J., & Menten, K.M. 2005, ApJ, 623, 269
- Fish, V.L., Reid, M.J., Argon, A.L., & Zheng, X.W. 2005, ApJS, 160, 220 (Paper I)
- Forster, J.R., & Caswell, J.L. 1989, A&A, 213, 339
- Forster, J.R., Goss, W.M., Wilson, T.L., Downes, D., & Dickel, H.R. 1980, A&A, 119, 84, L1
- Garay, G., Reid, M. J., & Moran, J.M. 1985, ApJ, 289, 681
- Garay, G., Rodr guez, L.F., Moran, J.M., & Churchwell, E. 1993, ApJ, 418, 368
- Garc a-Barreto, J.A., Burke, B.F., Reid, M.J., Moran, J.M., Haschick, A.D., & Schilizzi, R.T. 1988, ApJ, 326, 954
- Garc a-Segura, G., & Franco, J. 1996, ApJ, 469, 171
- Gardner, F.F., & Mart n-Pintado, J. 1983, A & A, 121, 265
- Gasiprong, N., Cohen, R.J., & Hutawarakom, B.2002, MNRAS, 336, 47
- Gaume, R.A., Fey, A.L., & Claussen, M.J. 1994, ApJ, 432, 648
- Gaume, R.A., Goss, W.M., Dickel, H.R., Wilson, T.L., & Johnston, K.J. 1995, ApJ, 438, 776
- Genzel, R., & Downes, D. 1976, Nature, 262, 564
- Giuliani, J.L. 1979, ApJ, 233, 280
- Goldreich, P. 1975, in Les Houches Session XXVI: Physique Atom ique et Moleculaire et Matiere Interstellaire, ed. R. Balian, P. Encrenaz, and J. Lequeux (North-Holland Publishing: Amsterdam), 409
- Goldreich, P., Keeley, D.A., & Kwan, J.Y. 1973a, ApJ, 179, 111
- Goldreich, P., Keeley, D.A., & Kwan, J.Y. 1973b, ApJ, 182, 55
- Gray, M.D., & Field, D. 1995, A&A, 298, 243
- Gray, M.D., Hutawarakom, B., & Cohen, R.J. 2003, MNRAS, 343, 1067
- Guilloteau, S., Baudry, A., Walm sley, C.M., Wilson, T.L., & Winnberg, A. 1984, A&A, 131, 45
- Habing, H.J., Goss, W.M., Matthews, H.E., & Winnberg, A. 1974, A&A, 35, 1
- Habing, H.J., & Israel, F.P. 1979, ARA & A, 17, 345
- Haschick, A.D., Reid, M.J., Burke, B.F., Moran, J.M., & Miller, G. 1981, ApJ, 244, 76
- Hatchell, J., Fuller, G.A., & Millar, T.J. 2001, A&A, 372, 281
- Ho, P.T.P., & Haschick, A.D. 1986, ApJ, 304, 501
- Hofner, P., & Churchwell, E. 1996, A & A S, 120, 283
- Hughes, V.A. 1988, ApJ, 333, 788
- Hughes, V.A. 2001, ApJ, 563, 919
- Hughes, V.A., & Baines, J.G.N. 1985, ApJ, 289, 238
- Hughes, V.A., Cohen, R.J., & Garrington, S.1995, MNRAS, 272, 469
- Hughes, V.A., & Wouterbot, J.G.A. 1984, ApJ, 276, 204
- Hutawarakom, B., & Cohen, R.J. 2003, MNRAS, 345, 175
- Hutawarakom, B., Cohen, R.J., & Brebner, G.C. 2002, MNRAS, 330, 349
- Kahn, F.D. 1954, Bull. A stron. Inst. Netherlands, 12, 187
- Kameya, O., Morita, K.-I., Kawabe, R., & Ishiguro, M. 1990, ApJ, 355, 562

- Kawabe, R., Suzuki, M., Hirano, N., Akabane, K., Barsony, M., Najita, J.R., Kameya, O., & Ishiguro, M. 1992, PASJ, 44, 435
- Kawamura, J.H., & Masson, C.R. 1998, ApJ, 509, 270
- Keto, E. 2002, ApJ, 568, 754
- Keto, E.R., Ho, P.T.P., & Haschick, A.D. 1987, ApJ, 318, 712
- Keto, E.R., Ho, P.T.P., & Haschick, A.D. 1988, ApJ, 324, 920
- Li, Z.-Y., & Shu, F.H. 1996, ApJ, 472, 211
- MacLeod, G.C., & Gaylard, M.J. 1996, MNRAS, 280, 868
- MacLow, M., van Buren, D., Wood, D.O.S., & Churchwell, E. 1991, ApJ, 369, 395
- Madden, S.C., Irvine, W.M., Matthews, H.E., Brown, R.D., & Godfrey, P.D. 1986, ApJ, 300, L79
- M anchester, R.N. 1972, ApJ, 172, 43
- M anchester, R.N., Hobbs, G.B., Teoh, A., & Hobbs, M. 2005, AJ, 129, 1993
- M atsum oto, T., & Tom isaka, K. 2004, ApJ, 616, 266
- M auersberger, R., W ilson, T.L., & Henkel, C. 1986, A&A, 160, L13
- Melrose, D.B., & Judge, A.C. 2004, Phys. Rev. E, 70, 056408
- Mezger, P.G., & Hoglund, B. 1967, ApJ, 147, 490
- M inier, V., Booth, R.S., Ellingsen, S.P., Conway, J.E., & Pestalozzi, M.R.2001, in Proceedings of the 5th European VLBI Network Symposium, eds. J.E. Conway, A.G. Polatidis, R.S. Booth, & Y.Pihlstrom, Chalmers University of Technology, Gothenburg, Sweden, 178
- M inier, V., Conway, J.E., & Booth, R.S. 2001, A&A, 369, 278
- Moran, J.M., Reid, M.J., Lada, C.J., Yen, J.L., Johnston, K.J., & Spencer, J.H. 1978, ApJ, 224, L67
- Mouschovias, T.C. 1976, ApJ, 207, 141
- Norris, R.P., et al. 1998, ApJ, 508, 275
- Norris, R.P., Whiteoak, J.B., Caswell, J.L., Wieringa, M.H., & Gough, R.G. 1993, ApJ, 412, 222
- Palmer, P., Gardner, F.F., & Whiteoak, J.B. 1984, MNRAS, 211, 41P
- Pavlakis, K.G., & Kyla s, N.D. 1996, ApJ, 467, 309
- Plume, R., Jae, D.T., & Evans, N.J., II 1992, ApJS, 78, 505
- Pottasch, S.R. 1958 Rev. M od. Phys., 30, 1053
- Rand, R.J., & Kulkami, S.R. 1989, ApJ, 343, 760
- Reid, M.J. 2002, in IAU Symposium 206, Cosm ic MASERs: From Protostars to Blackholes, ed.V.M igenes and M.J.Reid (San Francisco: ASP), 506
- Reid, M. J., Haschick, A. D., Burke, B. F., Moran, J. M., Johnston, K. J., & Swenson, G. W., Jr. 1980, ApJ, 239, 89
- Reid, M.J., & Ho, P.T.P. 1985, ApJ, 288, L17
- Reid, M.J., & Moran, J.M. 1988, in Galactic and Extragalactic Radio Astronomy, ed.G.Verschuur & K.I.Kellermann (Berlin Springer), 255
- Reid, M.J., & Menten, K.M. 1993, LNP Vol. 412: A strophysical Masers, 412, 137
- Reid, M. J., M yers, P. C., & Bieging, J. H. 1987, ApJ, 312, 830
- Reid, M. J., & Silverstein, E.M. 1990, ApJ, 361, 483
- Reifenstein, E.C., Wilson, T.L., Burke, B.F., Mezger, P.G., & Altenho, W.J. 1970, A&A, 4, 357

- Rodr guez, L.F., Garay, G., Curiel, S., Ram rez, S., Torrelles, J.M., Gomez, Y., & Velazquez, A. 1994, ApJ, 430, L65
- Rodr guez, L.F., Torrelles, J.M., Anglada, G., & Mart, J. 2001, Rev. Mexicana Astron. Astro s., 37, 95
- Samma, A.P., Troland, T.H., Cnutcher, R.M., & Roberts, D.A. 2002, ApJ, 580, 937
- Sam s, B.J. III, M oran, J.M., & Reid, M.J. 1996, ApJ, 459, 632
- Scoville, N.Z., Sargent, A.I., Sanders, D.B., Claussen, M.J., Masson, C.R., Lo, K.Y., & Phillips, T.G. 1986, ApJ, 303, 416
- Sew ilo, M., Churchwell, E., Kurtz, S., Goss, W. M., & Hofner, P. 2004, ApJ, 605, 285
- Shepherd, D.S., Testi, L., & Stark, D.P. 2003, ApJ, 584, 882
- Shu, F.H. 1992, The Physics of A strophysics v.2: G as D ynam ics (University Science: Sausalito, CA)
- Slysh, V. I., et al. 2001, MNRAS, 320, 217
- Slysh, V. I., Migenes, V., Val'tts, I. E., Lyubchenko, S. Yu., Horiuchi, S., Altunin, V. I., Fom alont, E. B., & Inoue, M. 2002, ApJ, 564, 317
- Slysh, V. I., Val'tts, I.E., & Migenes, V. 2001, Astronom y Reports, 45, 942
- Slysh, V. I., Voronkov, M. A., & Valtts, I. E. 2002, in IAU Symposium 206, Cosm ic MASERs: From Protostars to Blackholes, ed. V. Migenes and M. J. Reid (San Francisco: ASP), 105
- Sm its, D.P.2003, MNRAS, 339, 1
- Sm its, D.P., Cohen, R.J., & Hutawarakom, B.1998, MNRAS, 296, L11
- Stemberg, A., & Dalgamo, A. 1995, ApJS, 99, 565
- Szym czak, M ., H rynek, G ., & Kus, A . J. 2000, A & A S, 143, 269
- Testi, L., Hofner, P., Kurtz, S., & Rupen, M. 2000, A&A, 359, L5
- Thom pson, A.R., Moran, J.M., & Swenson, G.W., Jr. 2001, Interferom etry and Synthesis in Radio Astronomy (2d ed.; New York: W iley-Interscience)
- Torrelles, J.M. et al. 2001, ApJ, 560, 853
- Torrelles, J.M. et al. 2003, ApJ, 598, L115
- Torrelles, J.M., Gomez, J.F., Garay, G., Rodr guez, L.F., Curiel, S., Cohen, R.J., & Ho, P.T.P. 1998, ApJ, 509, 262
- Torrelles, J.M., Gomez, J.F., Rodr guez, L.F., Curiel, S., Ho, P.T.P., & Garay, G. 1996, ApJ, 457, L107
- Torrelles, J.M., Gomez, J.F., Rodr guez, L.F., Ho, P.T.P., Curiel, S., & Vazquez, R. 1997, ApJ, 489, 744
- Tumer, B.E., & Matthews, H.E. 1984, ApJ, 277, 164
- Vandervoort, P.O. 1962, ApJ, 135, 212
- Vishniac, E.T. 1983, ApJ, 274, 152
- W alm sley, C.M., Baudry, A., Guilloteau, S., & W innberg, A. 1986, A&A, 167, 151
- W ilson, T.L., Gaume, R.A., & Johnston, K.J. 1993, ApJ, 402, 230
- W ilson, T.L., W alm sley, C.M., M enten, K.M., & Herm sen, W. 1985, A&A, 147, L19
- W ilson, T.L., W alm sley, C.M., Snyder, L.E., & Jewell, P.R. 1984, A&A, 134, L7
- W right, M . M ., G ray, M . D ., & D iam ond, P. J. 2004, M N R A S, 350, 1272
- Xu, Y., Zheng, X.W., & Jiang, D.R. 2003, Chinese J.A stron. A strophys., 3, 133
- Zheng, X.-W. . 1997, Chinese Astron. Astrophys., 21, 182
- Zheng, X., Reid, M.J., & Moran, J.M. 2000, A&A, 357, L37

Zheng, X.W., Ho, P.T.P., Reid, M.J., & Schneps, M.H. 1985, ApJ, 293, 522

Zijlstra, A.A., Pottasch, S.R., Engels, D., Roelfsem a, P.R., Te Lintel Hekkert, P., & Um ana, G. 1990, MNRAS, 246, 217

This preprint was prepared with the AAS ${\tt IAT}_EX$ m acros v5.2.



Fig. 20. Left: Enlargement of collinear maser arrangement in W 75 S. The maser spots are tightly grouped along a line, as shown. Right: P bt of radial velocity v_{LSR} versus distance along major axis. The line of best t, representing a velocity gradient, is shown.



Fig. 21. Top: Observed spectra of 1665 M H z em ission from S269. RCP em ission is shown in bold, and LCP em ission is shown in norm alweight. The two weak RCP bumps at the same velocities as the strong LCP features at 16.0 and 16.8 km s¹ are due to the elliptical polarization of the LCP features. Bottom : The same spectra when corrected for a 4.0 mG m agnetic eld. Note that the velocity range is more than halved com pared to the observed spectra.