

A SUCCESSFUL TARGETED SEARCH FOR HYPERVELOCITY STARS

Warren R. Brown¹, Margaret J. Geller, Scott J. Kenyon, Michael J. Kurtz
 Smithsonian Astrophysical Observatory, 60 Garden St, Cambridge, MA 02138
 Submitted to ApJ Letters

ABSTRACT

Hypervelocity stars (HVSs) travel with velocities so extreme that dynamical ejection from a massive black hole is their only suggested origin. Following the discovery of the first HVS by Brown and collaborators, we have undertaken a dedicated survey for more HVSs in the Galactic halo and present here the resulting discovery of two new HVSs: SDSS J091301.0+305120 and SDSS J091759.5+672238, traveling with Galactic rest-frame velocities at least $+558 \pm 12$ and $+638 \pm 12$ km s⁻¹, respectively. Assuming the HVSs are B8 main sequence stars, they are at distances 75 and 55 kpc, respectively, and have travel times from the Galactic Center consistent with their lifetimes. The existence of two B8 HVSs in our 1900 deg² survey, combined with the Yu & Tremaine HVS rate estimates, is consistent with HVSs drawn from a standard initial mass function but inconsistent with HVS drawn from a truncated mass function like the one in the top-heavy Arches cluster. The travel times of the five currently known HVSs provide no evidence for a burst of HVSs from a major in-fall event at the Galactic Center in the last 160 Myr.

Subject headings: Galaxy: kinematics and dynamics | Galaxy: center | Galaxy: stellar content | Galaxy: halo | stars: early-type

1. INTRODUCTION

All galaxies with bulges probably host massive black holes (MBHs) in their centers. Hills (1988) first showed that a three-body interaction involving a MBH and a stellar binary can eject one member of the binary with $> 1,000$ km s⁻¹ velocity. Hills called stars ejected with $> 1,000$ km s⁻¹ velocities "hypervelocity stars." Hypervelocity stars (HVSs) are thus a natural consequence of the presence of a massive black hole in a dense stellar environment.

Brown et al. (2005) reported the first discovery of a HVS: a $g^0 = 19.8$ late-B type star, 110 kpc distant in the Galactic halo, traveling with a Galactic rest-frame velocity of at least $+709 \pm 12$ km s⁻¹ (heliocentric radial velocity $+853$ km s⁻¹). Photometric follow-up revealed that the object is a slowly pulsating B main sequence star (Fuentes et al. 2006). Only interaction with a MBH can plausibly accelerate a 3 M_⊙ main sequence B star to such an extreme velocity.

Our HVS discovery inspired a wealth of work from both observers and theorists. Edelmann et al. (2005) report a 8 M_⊙ main sequence B star, 60 kpc distant, traveling with a Galactic rest-frame velocity of at least $+548$ km s⁻¹ that may be a HVS ejected from the LMC. Hirsch et al. (2005) report a helium-rich sublumino O star, 20 kpc distant, traveling with a Galactic rest-frame velocity of at least $+717$ km s⁻¹ that is probably a HVS ejected from the Galactic Center. Holley-Bockelmann et al. (2006) suggest that outliers in the velocity distribution of intracluster planetary nebulae around M87 may be HVSs. In this paper, we report the discovery of two more HVSs from our ongoing HVS survey.

With HVSs now an observed class of objects, it is important to define true HVSs. Run-away B stars located many kpc above the Galactic plane have long been

known, but their velocities are typically < 200 km s⁻¹ and they are very probably bound to the Galaxy. HVSs, on the other hand, are unbound. More importantly, the classical supernova ejection (Blaauw 1961) and dynamical ejection (Poveda et al. 1967) mechanisms that explain run-away B stars cannot produce ejection velocities which exceed 300 km s⁻¹ (Leonard 1993; Gualandris et al. 2005). Thus we define a HVS as an unbound star with an extreme velocity that can be explained so far only by dynamical ejection associated with a MBH.

HVSs are important tools for understanding the nature and environs of MBHs: Holley-Bockelmann et al. (2006) predict that a thin torus of ejected HVSs is the signature of two MBHs forming a tight binary. Ginsburg & Loeb (2006) suggest that the stars on highly eccentric orbits around SgrA* may be the former companion stars to HVSs ejected by the MBH. Levin (2005) shows that an intermediate mass black hole (IMBH) on a circular in-spiral into the Galactic Center produces an isotropic burst of HVSs; an IMBH on an eccentric in-spiral produces broad jets of HVSs. Gualandris et al. (2005) find that HVSs produced from stellar binary encounters with single MBHs have higher ejection velocities than HVSs from binary MBHs. Gnedin et al. (2005) show that the distance and full space motion of HVSs can provide significant constraints on the shape and orientation of the Galactic dark matter halo. Yu & Tremaine (2003) expand Hill's original analysis and show that single star encounters with binary MBHs produce 10 times more HVSs than stellar binary encounters with single MBHs.

Our paper is organized as follows. In §2 we describe our survey target selection and observations. In §3 we present the new HVSs. We conclude in §4 by discussing what the observed set of HVSs implies about their origin and the nature of the Galactic Center.

2. DATA

2.1. Target Selection

Electronic address: wbrown@cfa.harvard.edu

¹ Clay Fellow, Harvard-Smithsonian Center for Astrophysics

HVSs ought to be rare: Yu & Tremaine (2003) predict there should be 10^3 HVSs in the entire Galaxy. Thus, in any search for HVSs, survey volume is important. Solar neighborhood surveys have not discovered HVSs because, even if they were perfectly complete to a depth of $d = 1$ kpc, there is a 0.1% chance of finding a HVS in such a small volume. Finding a new HVS among the Galaxy's 10^{11} stars also requires selection of targets with a high probability of being HVSs. Our observational strategy is two-fold. Because the density of stars in the Galactic halo drops off as approximately r^{-3} , and the density of HVSs drops off as r^{-2} (if they are produced at a constant rate), we target distant stars where the contrast between the density of HVSs and indigenous stars is as large as possible. Secondly, the stellar halo contains mostly old, late-type stars. Thus we target faint B-type stars, stars with lifetimes consistent with travel times from the Galactic center but which are not a normally expected stellar halo population. This strategy makes sense because 90% of the $K < 16$ stars in the central 10^{50} of the Galactic Center are in fact normal main sequence B stars (Eisenhauer et al. 2005).

We use SDSS photometry to select candidate B stars by color. To illustrate the color selection, Fig. 1 shows a color-color diagram of every star with $17.5 < g_0^0 < 18.5$ and B- and A-type colors in the SDSS DR4 (Adelman-McCarthy et al. 2006). We use de-reddened colors computed from $E(B-V)$ values obtained from Schlegel et al. (1998). The dashed box indicates the selection region used by Yanny et al. (2000) to identify BHB candidates. Interestingly, there is a faint group of stars with late B-type colors extending up the stellar sequence towards the ensemble of white dwarfs with $(u^0 - g^0)_0 < 0.5$. We select candidate B stars inside the solid parallelogram defined by: $0.42 < (g^0 - r^0)_0 < 0.27$ and $2.67(g^0 - r^0)_0 + 1.33 < (u^0 - g^0)_0 < 2.67(g^0 - r^0)_0 + 2.0$.

We observed a complete sample of 79 candidate B stars in the 1900 deg^2 region of the SDSS DR4 bounded by $7^h 40^m < \text{RA} < 10^h 50^m$ and $\text{Dec} > 15^\circ$. Figure 2 displays the locations of the objects on the sky. The Hirsch et al. (2005) HVS is located in this region on the sky, but it is not a part of our survey because its photometric colors lie far outside of our selection box. Our sample of candidate B stars is 100% complete in the magnitude range $17.0 < g_0^0 < 19.5$.

2.2. Spectroscopic Observations

Observations were obtained 2005 December 3-5 with the Blue Channel spectrograph on the 6.5m MMT telescope. The spectrograph was operated with the 832 line/mm grating in second order, providing 1.2 Å spectral resolution and wavelength coverage 3660 Å to 4500 Å. Exposure times ranged from 5 to 30 minutes and were chosen to yield $S/N = 15$ in the continuum at 4000 Å. Comparison lamp exposures were after obtained after every exposure.

Radial velocities were measured using the cross-correlation package RVSAO (Kurtz & Mink 1998). The average uncertainty is 12 km s^{-1} . We correct the heliocentric velocities to the local standard of rest (Hogg et al. 2005) and remove the 220 km s^{-1} solar reflex motion. Thus all velocities reported here are in the Galactic rest frame, indicated v_{RF} . The velocities of the candi-




Fig. 1. Color-color diagram showing our target selection, illustrated with every star in the SDSS DR4 $17.5 < g_0^0 < 18.5$. For reference, BHB/A stars are located in the dashed box (Yanny et al. 2000). Candidate late B-type stars are seen extending up the stellar sequence towards the ensemble of white dwarfs, and are selected by the solid parallelogram. The original HVS1 (Brown et al. 2005) is marked by plus sign, and the new HVSs are marked by open squares.

date B stars are indicated by color in Fig. 2.

2.3. Selection Efficiency and Unusual Objects

Our sample of 79 targets is composed of 61 late B-type stars, 16 faint DA white dwarfs, and 2 low- z galaxies. We derive spectral types of the late B-type stars based on line indices described in Brown et al. (2003); the types range from B6 to A1. Thus our target selection is 77% efficient for selecting stars of late B spectral type. Changing the color selection edge from $(g^0 - r^0)_0 = 0.42$ to 0.38 would eliminate more than half of the white dwarfs and one of the galaxies, and would increase the target selection efficiency for late B-type stars to 90%. We plan to publish the faint white dwarfs and other unusual objects in a future paper describing the HVS survey as a whole.

3. HYPERVELOCITY STARS

Our targeted search for HVSs uncovered two new HVSs, SDSS J091301.0+305120 (hereafter HVS4) and SDSS J091759.5+672238 (hereafter HVS5). Figure 3 plots a histogram of Galactic rest-frame radial velocity for the 61 late B-type stars in our sample. Ignoring the HVSs, our sample has a velocity dispersion of 114 km s^{-1} consistent with a Galactic halo population. HVS4 and HVS5 are traveling with $v_{RF} = +558 \pm 12$ and $+638 \pm 12 \text{ km s}^{-1}$ (heliocentric radial velocities $+603$ and $+543 \text{ km s}^{-1}$), respectively, and are 5- outliers from this distribution. The escape velocity of the Galaxy is approximately 300 km s^{-1} at 50 kpc (Wilkinson & Evans 1999); thus HVS4 and HVS5 are clearly unbound to the Galaxy.

The new HVSs are not physically associated with any other Local Group galaxy. HVS4 and HVS5 are located




Fig. 2. A sky map, in Galactic coordinates, showing the complete sample of 61 late B-type stars. Dotted lines indicate the 1900 deg² region of the SDSS DR4 we observed. Radial velocities, in the Galactic rest frame, are indicated by the color of the solid squares: purple is -300, green is 0, and red is +300 km s⁻¹. The new HVSs are completely off this color scale and are marked by open squares; HVS1 (Brown et al. 2005) and HVS2 (Hirsch et al. 2005) are marked by plus signs.

at $(l; b) = (194.8; 42.6)$ and $(146.3; 38.7)$, respectively (see Fig. 2). HVS4 is separated from Leo A by 10 on the sky, but the galaxy is in the distant background 800 kpc away (Dolphin et al. 2002). Even if HVS4 were a $M_V = -6$ supergiant at the distance of Leo A, the star's radial velocity differs from Leo A by 575 km s⁻¹. The closest galaxy to HVS5 is the Ursa Minor dwarf 31 away, yet the velocity difference is even more extreme at 730 km s⁻¹.

Both HVS4 and HVS5 have spectral types consistent with B8, however our low-resolution spectra do not allow us to determine exact stellar parameters. Stars of this spectral type are likely blue horizontal branch (BHB) stars or main sequence B stars/blue stragglers. Two of the three previously reported HVSs are main sequence B stars (Fuentes et al. 2006; Edelmann et al. 2005). We note that the Balmer line widths of HVS4 and HVS5 are too broad to be consistent with those of B1 supergiants. If we assume the HVSs are BHB stars rather than B stars, their blue colors means they are hot, extreme BHB stars and thus they are intrinsically very faint. The $M_V(BHB)$ relation of Clowley et al. (2002) yields $M_V(BHB) = +2.0$ and $+1.5$ and heliocentric distance estimates $d_{BHB} = 20$ and 18 kpc for HVS4 and HVS5, respectively. In the BHB interpretation, the volume we effectively survey is much smaller than in the B star interpretation. Thus the BHB interpretation requires more than an order of magnitude larger production rate for HVSs. Because two of the previous HVS are B stars and because the B star interpretation implies a lower production rate probably consistent with Yu & Tremaine (2003), we assume that HVS4 and HVS5 are B stars for the purpose of discussion. The ultimate discriminant will come from higher resolution, higher signal-to-noise spectroscopy.

We estimate the luminosity of a B8 star from the Schaller et al. (1992) stellar evolution tracks for a 4 M star with $Z = 0.02$. Such a star spends 160 Myr on the main sequence and produces 400 $L_\odot$ at $T_{\text{eff}} = 13,000$.




Fig. 3. Velocity histogram of the late B-type stars. Two new HVSs are identified moving away with Galactic rest-frame velocities at least +558 and +638 km s⁻¹.

This luminosity corresponds to $M_V(B8) = -0.9$ (using bolometric correction -0.80 (Kenyon & Hartmann 1995)) and heliocentric distances $d_{B8} = 75$ and 55 kpc for HVS4 and HVS5, respectively. Assuming the Sun is 8 kpc distant from the Galactic center, the Galactocentric distances of HVS4 and HVS5 are thus $r = 80$ and 60 kpc, respectively.

If HVS4 and HVS5 originate from the Galactic Center, the travel times to their present locations are approximately 140 and 90 Myr, respectively, consistent with the 160 Myr main sequence lifetime of a 4 M star. If the HVSs are BHBs or blue stragglers, their lifetimes are considerably longer and the travel time constraint is relaxed.

Our radial velocities provide only a lower limit to the HVS's true space velocities. HVS4 and HVS5 have bright apparent magnitudes ($g^0 = 18.40$ and 17.93, respectively) and are present in the USNOB1 (Monet et al. 2003) and GSC 2.3 (B. McLean, 2006 private communication) catalogs. However, the HVSs are listed with no measurable proper motions, consistent with the estimated distances.

Table 1 summarizes the properties of all we know HVSs. The columns include HVS number, Galactic coordinates $(l; b)$, apparent magnitude g^0 , minimum Galactic rest-frame velocity v_{RF} (not a full space velocity), heliocentric distance estimate d , travel time estimate from the Galactic Center t_{GC} , and catalog identification. We note that the B9 M_V estimate was incorrect in Brown et al. (2005) and should be $M_V(B9) = -0.3$. Thus the correct distance and travel time estimates to HVS1 are $d = 110$ kpc and $t_{GC} = 160$ Myr, as indicated.

4. DISCUSSION

The very existence of the HVSs places interesting limits on the stellar mass function of HVSs, the origin of massive stars in the Galactic Center, and the history of stellar interactions with the MBH. Interestingly, all we know HVSs are moving with positive radial velocity, consistent with the picture of a Galactic Center origin.

In principle, we can constrain the stellar population from which the HVSs originate by combining predictions of HVS rates with the results of our survey. Yu & Tremaine (2003) predict a HVS rate of 10^5 yr⁻¹ ejected by a single MBH at the Galactic Center. A HVS moving 600 km s⁻¹ travels 120 kpc in 200 Myr, thus the Yu & Tremaine (2003) rate implies 2000 HVSs of all types to a depth of 120 kpc. Our $g^0 < 19.5$ magnitude-limited survey reaches the same depth $d = 120$ kpc for

B8 stars and we find two B8 HVSS over our 1900 deg² region, implying 43 ± 31 B8 HVSS over the entire sky. We ask whether this number of B stars is consistent with the number expected from a standard initial mass function (IMF). A Salpeter IMF (Salpeter 1955), integrated over the mass range 0.2–100 M_⊙ and normalized to 2000 stars, contains 26 stars between 3 and 5 M_⊙. A Scalo IMF (Scalo 1986), similarly calculated, contains 13 stars between 3 and 5 M_⊙. The IMF predictions are systematically lower than the observed frequency implied by the two B stars in our survey, but because of small number statistics there is no significant inconsistency.

In the Galactic center, there is some indication that the IMF is top heavy. For example, Stolte et al. (2005) study the mass-segregated Arches cluster and argue for truncation at masses less than 6 M_⊙. Integrating the Stolte et al. (2005) mass function = 0.86 over the mass range 3–100 M_⊙ and normalizing it to 2000 stars results in 750 3–5 M_⊙ stars, an order of magnitude more late-B HVSS than implied by our observations. Thus our observations already indicate that the HVS parent population is not composed entirely of clusters like Arches or the Yu & Tremaine (2003) HVS rate is an underestimate.

HVS travel times from the Galactic Center constrain the history of stellar interactions with the MBH. Assuming the new HVSS are B stars, travel time estimates for the known HVSS are spread rather uniformly between 30 and 160 Myr. Thus there is as yet no evidence for a burst of HVSS from a major in-fall event at the Galactic Center in the last 160 Myr. The current constraints are less clear if HVS4 or HVS5 are not B stars. If new discoveries of HVSS continue to show no evidence for coherent bursts of HVSS, theories of in-falling massive star clusters

(Gerhard 2001; Kim & Morris 2003) possibly containing an IMBH (Hansen & Milosavljević 2003) to transport B stars near the MBH may not be applicable to the Galaxy.

It is interesting that the northern HVSS all have similar $b \approx 40^\circ$ (see Fig. 2). An in-spiraling binary black hole produces a unique distribution of HVSS on the sky (Levin 2005; Holley-Bockelmann et al. 2006). Detailed theoretical predictions of the distribution of HVSS on the sky ejected by a single MBH at the Galactic Center would be an important testbed for future HVS observations.

HVSS are becoming important tools for understanding MBHs. Discovering additional HVSS in a well-defined volume will provide better constraints on the origin of HVSS and the nature of the Galactic Center. Proper motion measurements of HVSS, measured with the Hubble Space Telescope, the Global Astrometric Interferometer for Astrophysics, or the Space Interferometry Mission, may provide significant constraints on the shape of the Galaxy's dark matter potential (Gnedin et al. 2005). The distribution of HVSS in velocity and space may constrain the history of stellar interactions with the MBH. Identifying HVSS around other galaxies (Holley-Bockelmann et al. 2006) is also an exciting prospect. We are continuing our radial velocity survey of every late B-type star in the SDSS.

We thank J. McAffee for his assistance with observations obtained at the MMT Observatory, a joint facility of the Smithsonian Institution and the University of Arizona. This work was supported by W. Brown's Clay Fellowship and the Smithsonian Institution.

REFERENCES

- Adelman-McCarthy, J. K. et al. 2006, preprint (astro-ph/0507711)
 Blaauw, A. 1961, *Bull. Astron. Inst. Netherlands*, 15, 265
 Brown, W. R., Allende Prieto, C., Beers, T. C., Wilhelm, R., Geller, M. J., Kenyon, S. J., & Kurtz, M. J. 2003, *AJ*, 126, 1362
 Brown, W. R., Geller, M. J., Kenyon, S. J., & Kurtz, M. J. 2005, *ApJ*, 113, L33
 Clewley, L., Warren, S. J., Hewett, P. C., Norris, J. E., Peterson, R. C., & Evans, N. W. 2002, *MNRAS*, 337, 87
 Dolphin, A. E., Saha, A., Claver, J., Skillman, E. D., Cole, A. A., Gallagher, J. S., Tolstoy, E., Dohm-Palmer, R. C., & Mateo, M. 2002, *AJ*, 123, 3154
 Edelmann, H., Napiwotzki, R., Heber, U., Christlieb, N., & Reimers, D. 2005, *ApJ*, 634, L181
 Eisenhauer, F. et al. 2005, *ApJ*, 628, 246
 Fuentes, C. I., Stanek, K. Z., Gaudi, B. S., McLeod, B. A., Bogdanov, S., Hartman, J. D., Hickox, R. C., & Holman, M. J. 2006, *ApJ*, 636, L37
 Gerhard, O. 2001, *ApJ*, 546, L39
 Ginsburg, I. & Loeb, A. 2006, preprint (astro-ph/0510574)
 Gnedin, O. Y., Gould, A., Miralda-Escudé, J., & Zentner, A. R. 2005, *ApJ*, 634, 344
 Gualandris, A., Zwart, S. P., & Sapior, M. S. 2005, *MNRAS*, 363, 223
 Hansen, B. M. S. & Milosavljević, M. 2003, *ApJ*, 593, L77
 Hills, J. G. 1988, *Nature*, 331, 687
 Hirsch, H. A., Heber, U., O'Toole, S. J., & Bresolin, F. 2005, *A & A*, 444, L61
 Hogg, D. W., Blanton, M. R., Roweis, S. T., & Johnston, K. V. 2005, *ApJ*, 629, 268
 Holley-Bockelmann, K., Sigurdsson, S., Mihos, C. J., Feldmeier, J. J., Ciardullo, R., & McBride, C. 2006, preprint (astro-ph/0512344)
 Kenyon, S. J. & Hartmann, L. 1995, *ApJS*, 101, 117
 Kim, S. S. & Morris, M. 2003, *ApJ*, 597, 312
 Kurtz, M. J. & Mink, D. J. 1998, *PASP*, 110, 934
 Leonard, P. J. T. 1993, in *ASP Conf. Ser.* 45, *Luminous High-Latitude Stars*, ed. D. D. Sasselov, 360
 Levin, Y. 2005, preprint (astro-ph/0508193)
 Monet, D. G. et al. 2003, *AJ*, 125, 984
 Poveda, A., Ruiz, J., & Allen, C. 1967, *BoL. Obs. Tonantzintla Tacubaya*, 4, 860
 Salpeter, E. E. 1955, *ApJ*, 121, 161
 Scalo, J. M. 1986, *Fundamentals of Cosmic Physics*, 11, 1
 Schaller, G., Schaerer, D., Meynet, G., & Maeder, A. 1992, *A & A*, 96, 269
 Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
 Stolte, A., Brandner, W., Grebel, E. K., Lenzen, R., & Lagrange, A.-M. 2005, *ApJ*, 628, L113
 Wilkinson, M. I. & Evans, N. W. 1999, *MNRAS*, 310, 645
 Yanny, B. et al. 2000, *ApJ*, 540, 825
 Yu, Q. & Tremaine, S. 2003, *ApJ*, 599, 1129

TABLE 1
HYPERVELOCITY STARS

ID	l deg	b deg	g^0 mag	v_{RF} km s ⁻¹	d kpc	t_{gc} Myr	Catalog
HVS1	227.3	31.3	19.8	+ 709	110	160	SD SS J090745.0+ 024507 ¹
HVS2	176.0	47.1	18.8	+ 717	19	32	U S 708 ²
HVS3	263.0	-40.9	16.2	+ 548	61	100	HE 0437-5439 ³
HVS4	194.8	42.6	18.4	+ 558	75	140	SD SS J091301.0+ 305120
HVS5	146.3	38.7	17.9	+ 638	55	90	SD SS J091759.5+ 672238

References. | (1) Brown et al. (2005); (2) Hirsch et al. (2005); (3) Edelmann et al. (2005)