Tailem ission from a ring-like jet: its application to shallow decays of early afterglows and GRB 050709

Yuan-Chuan Zou and Zi-Gao Dai

Department of Astronomy, Nanjing University, Nanjing 210093, China.

Received 2006 January 30; accepted

A bstract Sim ilar to the pulsar, the magnetic axis and the spin axis of the gam ma-ray burst source may not lie on the same line. This may cause a ring-like jet due to collimation of the precessing magnetic axis. We analyze the tailem ission from such a jet, and nd that it has a shallow decay phase with temporal index equal to 1=2 if the Lorentz factor of the ejecta is not very high. This phase is consistent with the shallow decay phase of some early X-ray afferglow detected by swift. The ring-like jet has a tail cusp with sharp rising and very sharp decay. This elect can provide an explanation for the re-brightening and sharp decay of the X-ray afferglow of GRB 050709.

Keywords: gamma rays:bursts { X-rays:general

# 1 IN TRODUCTION

It is wellknown that pulsars originate from the core collapse of massive stars. The average angle between the spinning axis and the magnetic axis of pulsars is about 27 (Leahy, 1991). Sim ilarly, the spin axis and magnetic axis of the central engine of the gam ma-ray burst may not lie on one line. A sthe ejecta may be collimated by the magnetic axis, while the magnetic axis is processing, so the ejecta may be in a spiral shape at rst (Fargion & G rossi, 2005, and reference therein). A s the diversity of the velocities of the ejecta [as assumed in the standard reballm odel of gam ma-ray bursts (P iran, 2005)], the spiral ejecta ejected at di erent times will collide and merge into one whole shell at last. These collisions just produce internal shocks of gam ma-ray burst. At last, these collisions make the ejecta merge into a ring-shaped jet. Even if the the ejecta is conical, the baryon-loaded region still be ring-like (Eichler & Levinson, 2003). G ranot (2005) and Eichler

<sup>?</sup> E-m ail: zouyc@nju.edu.cn, dzg@nju.edu.cn

& G ranot (2005) have analyzed afterglows from ring-like jets. It has also been used to interpret the h  $_{peak}$  E<sub>iso</sub> relation (E ichler & Levinson, 2004).

Tailem ission plays an important role at the times when shocks disappear. The tem – poral index is (2 + ) for a cone-shaped jet, where is the spectral index of the em ission (K um ar & Panaitescu, 2000; Y am azakiet al., 2005). Considering the zero point e ect of time, the light curves can be steeper during a short period (N ousek et al., 2005; Zhang et al., 2005; W u et al., 2006).

Nousek et al. (2005) and Zhang et al. (2005) have also shown that a shallow decay with index of about 1=2 follows the steep decay for most X -ray afterglows. For the X - ray afterglow of short burst G R B 050709, there is an unexpected high - ux point followed by a very steep decay (Fox et al., 2005). These two observations can both be explained naturally by considering the tail em ission of ring-like jets. In x2 we give the expressions of tail em ission from a ring-like jet. In xx3 and 4, the shallow decay and X -ray afterglow of G R B 050709 are analyzed respectively. At last, we sum marize our results in x5.

### 2 MODEL

C onsidering several ring-like sub-jets em itted from the central engine, they merge into one whole ring-like jet accompanied with internal shocks. This nalring with uniform energy density and sharp edges expands with Lorentz factor , as sketched in Fig. 1. A ssum ing the radiation from the ring-like jet begins and ceases at radius  $R_c$  (and correspondingly at time  $t_c$ ) suddenly, we calculate the tail em ission from high latitudes of the ring. The relation is  $R_c' 2^{-2}$ ct, where is the mean Lorentz factor of the internal shocks.

The relation between the latitude angle and the observed time t is

$$R_{c}(1 \cos t) = c(t dt) = (1 + z);$$
 (1)

where z is the cosm obgical redshift. N eglecting the depth of the ejecta and the em ission from time equal arrival surface, and de ning the em issivity  $I^0_{0}$  per unit area in the com oving frame, which is uniform in the whole ring, the ux density in the observer's frame is

f 
$$(t > t_c) = \frac{I_0^0}{4 D_L^2} D^2 \frac{dS}{dt = (1 + z)};$$
 (2)

where  $D_L$  is the lum inosity distance,  $D = 1 = [(1 \quad p = 1 = 2 \cos)]$  is the Doppler factor, and dS is the em itted area during a period dt.

At early times when < w, the tail emission is the same as the case of an on-axis conical jet, which has been investigated by many authors (Kumar & Panaitescu, 2000; Fan & W ei, 2005). There are two limiting cases: for 1 = 1,

$$f(t > t_c) / t';$$
 (3)

and for 1 = 1

$$f(t > t_c) / t^{(2+)};$$
 (4)



Fig.1 Sketch of a ring-like jet at a distance  $R_c$  from the central engine, where the main emission is just ceased, while an observer begins to receive the tail emission from high latitudes. <sub>p</sub> is the half opening angle of the ring. <sub>w</sub> is the half width angle. is the latitude of tail emitted region corresponding to the observed time t.

where t t d. Here we consider the emission as a single power-law pro le  $I_c^0 / c^0$ , which is valid for the high frequency emission  $c^0 > m \operatorname{ax}(c_c^0; m^0)$ , where  $c_c^0$  is the cooling frequency and  $m^0$  is the typical frequency of synchrotron emission.

In the case > w, the width of the ring can be neglected, and the ux density

$$f(t > t_{c}) / D^{(2+)} \frac{\sin_{p} \cos(-2)}{\frac{r}{\sin_{p}} \cos(-2)} :$$
(5)  
$$\sin_{p} 1 \frac{\sin(-2)}{\sin_{p}} :$$

There are two limiting cases in which equation (5) can be simplied. For 1= ,

$$f(t > t_c) / t^{1=2};$$
 (6)

and for 1 = 1

$$f(t > t_c) / t^{(5=2+)}$$
: (7)

### 3 SHALLOW DECAY OF EARLY X-RAY AFTERGLOW

Statistics of the early X -ray afferglow shas shown that there is a shallow decay phase with tem poral index about 1=2 (N ousek et al., 2005; Zhang et al., 2005). This corresponds to the case: 1 = > > w, and can be described by equation (6), where the tem poral index is just 1=2. As the general shallow decay lasts from  $10^{\circ}$   $10^{\circ}$  s to  $10^{3}$   $10^{\circ}$  s (F ig. 1 in Zhang et al., 2005), this gives limits: the Lorentz factor of the em itting shell < 7:3 (1 + z)<sup>1=2</sup>R $_{c;16}^{1=2}$  t $_{3:5}^{1=2}$ , and the width of the ring-like jet w < 1:4  $10^{\circ}$  (1 + z)  $^{1=2}$ R $_{c;16}^{1=2}$  t $_{2:5}^{1=2}$ .

(The conventional donation  $Q = Q_k$  10<sup>t</sup> is used throughout this paper.) This in plies that the shallow decay component originates from the shocked shells with low Lorentz factors, while these shocks may be formed due to ejected sub-shells with dimensional Lorentz factors.

This model can answer the following questions:

Firstly, why is there a steep decay before the shallow decay appears in general case? In Zhang et al. (2005), it is general that the tem poral index of this steep decay is less that

3. The answer is that the two power law decays originate from two di erent emitting shells with di erent Lorentz factors. The steep decay corresponds to the greater Lorentz factor shell, which satis es  $1 = \langle ,$  and the tem poral indices are (2 + ) or (5 = 2 + ) corresponding to equations (4) and (7) respectively. The steep decay m ay become steeper because of the zero time selection e ect (W u et al., 2006).

Secondly, why is there no spectral evolution before and after the break time from the shallow decay phase to the steep decay phase. This is also mentioned with spectral index value 1 in Zhang et al. (2005). It is believed that, after the break, the afterglow becomes a \norm al" afterglow. It is possible that the tailem ission phase and the \norm al" afterglow emission phase are both in the case  $_{\rm X}$  > f  $_{\rm m}$ ;  $_{\rm c}$ g (corresponding to the spectral index p=2) and thus the spectra are the same.

Thirdly, since the shallow phase and the steep phase originate from di erent sources, how to understand the conjunction at the break time (also can be seen in Fig.1 in Zhang et al., 2005)? A stime goes on, the case converts from 1 = > > w to > 1 = > w, and then the light curve of the tail em ission decay has a tem poral index (5=2+). This is steeper than the \norm al" afterglow with tem poral index 12. Some time later, the \norm al" afterglow will exceed the tail em ission de nitely, as in the case GRB 050525a (Fig.1 in Nousek et al., 2005) (at about 3000s, there is a steep decay). However, GRB 050315 (Vaughan et al., 2005) can be classi ed into the case that \norm al" afterglow exceed the shallow tail em ission before the tail em ission breaks to steep phase.

## 4 X-RAY AFTERGLOW OF GRB 050709

GRB 050709 is a short burst with duration 0.3 s, and ve points of X-ray emission after the burst were obtained by Sw iff and Chandra (Fox et al., 2005). Figure 2 shows the t by assuming that the latter four points are the tailemissions from the rst point, with parameters  $R_c = 7:7$   $10^{16}$  cm, = 15:5, = 1:1,  $_p = 0:5$  and  $_w = 0:005$ . As the rst X-ray point occurs at time about 100s, the radius r'  $2^{-2}$  ct= (1 + z) ' 5:2  $10^{6}$   $_2^{-2}$  t<sub>2</sub> cm, is consistent with the value of the parameter  $R_c$ . As a short burst has less total energy than a long burst does, the ejected shell can be decelerated quickly. The Lorentz factor at  $R_c$  is '  $E_{iso} = (R_c^3 nm_p c^2)$ '  $26E_{iso;50}^{-12} n_1^{-12} R_{c;16:5}^{-3=2}$ , where the external medium density



Fig.2 Tailemission t to the X-ray emission from GRB 050709. The ve observed data are taken from Fox et al. (2005).

n is chosen equal to 1 cm  $^3$  because the host is a star-form ing galaxy. Therefore, the parameters chosen to t the X-ray data are reasonable for this short burst.

We can see four stages for this tail light curve: rst, a horizontal phase corresponds to the case  $<_w < 1 = ;$  second, a shallow decay with tem poral index 1=2 corresponds to the case  $_w < < 1 = ;$  third, a sharper decay with tem poral index (2:5+ ) corresponds to the case  $_w < 1 = < ;$  and nally, a tail cusp with sharp rising and very sharp decay, which comes from the end of the ring.

We should note that the solid line doesn't t the data very well, especially that the tail cusp of the model can't reach to the observed data. This shortage may be overcome by considering a non-uniform ring-like jet or some other mechanism. However, its unique feature, which the emission after the tail cusp decays very sharply, is consistent with the last two observed points. On the other hand, it is possible that the second, third and fth points in this gure belong to the \norm al" afterglow from an external shock.

# 5 CONCLUSIONS

Enlightened from pulsars, we suggest that the magnetic axis and the spin axis of a gam ma-ray burst source point to di erent orientations. The ejecta along the magnetic

axis will form a ring nally.G amm a-ray emission will be observed if the observer locates in the solid angle of the ring.W e have investigated the tail emission from a ring-like jet.W e nd that the early shallow decay phase and the late re-brightening of the X-ray emission of GRB 050709 can be explained.

Note that the shallow decay phase is only possible in the low Lorentz factor cases. For the case  $1 = \langle w \rangle$ , only the steep one appears. As the tail emission from the shells with high Lorentz factors decays very quickly, the main emissions will be dominated by the slower shells at later times.

YCZ thanks helpful discussions with Jia W ang and Xuefeng W u. This work was supported by the National Natural Science Foundation of China (grants 10233010 and 10221001).

#### References

Eichler D., Granot J., 2006, ApJ, 641, L5
Eichler D., Levinson A., 2003, ApJ, 596, L147
Eichler D., Levinson A., 2004, ApJ, 614, L13
Fan Y. Z., W eiD. M., 2005, MNRAS, 364, L42
Fargion D., Grossi M., 2005, MNRAS submitted (astro-ph/0504638)
Fox D. B., FrailD. A., Price P. A., et al., 2005, N ature, 437, 845
Granot J., 2005, ApJ, 631, 1022
K um ar P., Panaitescu A, 2000, ApJ, 541, L51
Nousek J. A., K ouveliotou C., G rupe D., et al., 2005, ApJ in press (astro-ph/0508332)
Leahy D. A., 1991, MNRAS, 251, 203
P iran T., 2005, Rev. M od. Phys., 76, 1143
Vaughan S., Goad M. R., Beardm ore A. P., et al., 2006, ApJ, 638, 920
W u X. F., Dai Z. G., W ang X. Y., et al., 2006, ApJ submitted (astro-ph/0512555)
Yam azaki R., Tom a K., Ioka K., Nakam ura T., 2005, MNRAS in press (astro-ph/0509159)

Zhang B., Fan Y.Z., Dyks J., et al., 2005, ApJ in press (astro-ph/0508321)

6